Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
United States												
All races, both sexes	51	2	-	-	-	2	-	1	-	2	-	3
Male	22	1	-	-	-	-	-	1	-	1	-	2
Female	29	1	-	-	-	2	-	-	-	1	-	1
White, both sexes	39	1	-	-	-	1	-	-	-	1	-	1
Male	17	1	-	-	-	-	-	-	-	1	-	1
Female	22	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	11	1	-	-	-	1	-	1	-	1	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	6	1	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Florida												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Illinois												
All races, both sexes	6	2	-	-	-	-	-	-	-	1	-	-
Male	5	1	-	-	-	-	-	-	-	1	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	1	-	-	-	-	-	-	-	1	-	-
Male	5	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Kentucky												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Massachusetts												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Minnesota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
North Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Texas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Washington												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Wisconsin												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
United States												
All races, both sexes	2	3	4	2	1	5	4	4	9	5	2	-
Male	1	1	2	2	1	3	2	2	2	1	-	-
Female	1	2	2	-	-	2	2	2	7	4	2	-
White, both sexes	1	2	2	1	1	5	4	4	9	4	2	-
Male	1	-	1	1	1	3	2	2	2	1	-	-
Female	-	2	1	-	-	2	2	2	7	3	2	-
Black, both sexes	1	1	2	1	-	-	-	-	-	1	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Colorado												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	2	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	2	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Florida												
All races, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Illinois												
All races, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Kentucky												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	-	1	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	1	-
White, both sexes	-	-	-	-	-	-	1	-	1	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Missouri												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
North Carolina												
All races, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Texas												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Washington												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
United States												
All races, both sexes	62	3	2	1	-	1	1	-	1	2	5	5
Male	40	3	1	-	-	1	-	-	1	1	4	3
Female	22	-	1	1	-	-	1	-	-	1	1	2
White, both sexes	57	2	2	-	-	1	1	-	1	1	4	5
Male	37	2	1	-	-	1	-	-	1	-	3	3
Female	20	-	1	-	-	-	1	-	-	1	1	2
Black, both sexes	3	1	-	-	-	-	-	-	-	1	1	-
Male	3	1	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	10	1	-	-	-	-	1	-	-	-	1	1
Male	8	1	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	1	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Florida												
All races, both sexes	10	-	1	-	-	1	-	-	-	1	-	-
Male	6	-	-	-	-	1	-	-	-	1	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	1	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	1	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	1	-
Male	3	1	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	1	-
Male	3	1	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Hawaii												
All races, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Nebraska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Ohio												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Texas												
All races, both sexes	3	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
United States												
All races, both sexes	4	2	3	4	5	5	6	1	8	2	1	-
Male	4	1	2	3	4	2	3	-	5	1	1	-
Female	-	1	1	1	1	3	3	1	3	1	-	-
White, both sexes	4	1	3	4	5	5	6	1	8	2	1	-
Male	4	1	2	3	4	2	3	-	5	1	1	-
Female	-	-	1	1	1	3	3	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	2	-	1	1	-	1	-	-	1	-	-
Male	-	1	-	1	1	-	1	-	-	1	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	1	-	1	-	-	1	-	-
Male	-	1	-	1	1	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Colorado												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Florida												
All races, both sexes	1	-	2	1	-	1	-	-	2	-	-	-
Male	1	-	1	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
White, both sexes	1	-	2	1	-	1	-	-	2	-	-	-
Male	1	-	1	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Missouri												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
North Carolina												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Ohio												
All races, both sexes	-	-	1	-	1	-	-	-	1	-	-	-
Male	-	-	1	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	-	-	1	-	-	-
Male	-	-	1	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Texas												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
United States												
All races, both sexes	827,741	543	221	142	226	478	944	1,525	2,688	5,074	10,624	19,230
Male	396,612	301	116	81	129	296	605	1,025	1,794	3,414	7,105	13,153
Female	431,129	242	105	61	97	182	339	500	894	1,660	3,519	6,077
White, both sexes	710,632	343	154	97	155	330	653	987	1,727	3,382	7,561	13,887
Male	339,084	186	77	59	99	210	435	676	1,197	2,346	5,289	9,850
Female	371,548	157	77	38	56	120	218	311	530	1,036	2,272	4,037
Black, both sexes	98,102	174	57	34	58	137	254	477	851	1,505	2,696	4,751
Male	47,600	100	35	15	28	80	147	305	517	932	1,557	2,880
Female	50,502	74	22	19	30	57	107	172	334	573	1,139	1,871
American Indian 1/, both sexes	3,518	10	2	3	2	5	14	11	30	48	133	185
Male	1,863	5	2	2	-	1	6	7	20	36	95	130
Female	1,655	5	-	1	2	4	8	4	10	12	38	55
Asian or Pacific Islander 2/, both sexes	15,489	16	8	8	11	6	23	50	80	139	234	407
Male	8,065	10	2	5	2	5	17	37	60	100	164	293
Female	7,424	6	6	3	9	1	6	13	20	39	70	114
Alabama												
All races, both sexes	16,378	12	4	6	5	9	25	56	101	158	306	524
Male	7,768	7	3	2	2	7	17	40	66	102	181	331
Female	8,610	5	1	4	3	2	8	16	35	56	125	193
White, both sexes	12,330	8	2	2	4	7	17	32	55	86	186	315
Male	5,821	6	2	-	1	5	13	24	32	59	121	210
Female	6,509	2	-	2	3	2	4	8	23	27	65	105
Black, both sexes	3,992	3	2	4	1	2	8	22	45	71	120	206
Male	1,919	1	1	2	1	2	4	14	33	42	60	119
Female	2,073	2	1	2	-	-	4	8	12	29	60	87
American Indian 1/, both sexes	23	1	-	-	-	-	-	-	-	1	-	2
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	15	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	2	1	-	-	1
Male	20	-	-	-	-	-	-	2	1	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Alaska												
All races, both sexes	877	1	-	-	1	2	2	2	2	12	16	27
Male	490	-	-	-	-	1	2	-	-	7	13	18
Female	387	1	-	-	1	1	-	2	2	5	3	9
White, both sexes	658	-	-	-	-	2	-	1	-	5	9	18
Male	378	-	-	-	-	1	-	-	-	3	8	13
Female	280	-	-	-	-	1	-	1	-	2	1	5
Black, both sexes	24	-	-	-	-	-	-	-	-	2	5	-
Male	14	-	-	-	-	-	-	-	-	1	3	-
Female	10	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	155	1	-	-	1	-	2	1	1	4	2	8
Male	81	-	-	-	-	-	2	-	-	2	2	4
Female	74	1	-	-	1	-	-	1	1	2	-	4
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	1	1	-	1
Male	17	-	-	-	-	-	-	-	-	1	-	1
Female	23	-	-	-	-	-	-	-	1	-	-	-
Arizona												
All races, both sexes	13,902	12	6	5	5	12	21	31	43	90	154	282
Male	7,115	8	4	1	2	7	15	21	32	70	110	196
Female	6,787	4	2	4	3	5	6	10	11	20	44	86
White, both sexes	13,090	10	5	4	4	7	18	22	34	73	127	248
Male	6,636	6	3	1	2	3	12	17	24	54	92	173
Female	6,454	4	2	3	2	4	6	5	10	19	35	75
Black, both sexes	405	1	-	-	-	5	-	6	5	8	6	18
Male	257	1	-	-	-	4	-	2	4	7	5	13
Female	148	-	-	-	-	1	-	4	1	1	1	5
American Indian 1/, both sexes	302	1	1	1	-	-	3	2	3	8	19	13
Male	174	1	1	-	-	-	3	2	3	8	12	7
Female	128	-	-	1	-	-	-	-	-	-	7	6
Asian or Pacific Islander 2/, both sexes	105	-	-	-	1	-	-	1	1	1	2	3
Male	48	-	-	-	-	-	-	-	1	1	1	3
Female	57	-	-	-	1	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Arkansas												
All races, both sexes	9,925	11	5	2	5	8	15	26	51	85	159	297
Male	4,767	5	3	2	3	3	11	16	35	52	104	198
Female	5,158	6	2	-	2	5	4	10	16	33	55	99
White, both sexes	8,430	6	3	2	3	6	8	15	39	51	110	198
Male	4,057	3	3	2	3	3	5	11	29	37	72	136
Female	4,373	3	-	-	-	3	3	4	10	14	38	62
Black, both sexes	1,443	5	2	-	2	1	7	11	11	32	49	95
Male	684	2	-	-	-	-	6	5	5	14	32	59
Female	759	3	2	-	2	1	1	6	6	18	17	36
American Indian 1/, both sexes	27	-	-	-	-	1	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	1	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	1	2	-	1
Male	15	-	-	-	-	-	-	-	1	1	-	1
Female	10	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	86,484	49	18	16	29	38	84	134	246	510	1,066	1,916
Male	42,171	27	12	12	14	28	55	101	168	361	741	1,346
Female	44,313	22	6	4	15	10	29	33	78	149	325	570
White, both sexes	72,142	35	15	11	21	31	64	101	175	358	804	1,439
Male	35,064	19	10	9	12	22	43	76	127	249	569	1,036
Female	37,078	16	5	2	9	9	21	25	48	109	235	403
Black, both sexes	7,033	7	1	2	4	6	14	18	45	92	169	314
Male	3,403	5	1	1	1	5	8	13	23	67	104	186
Female	3,630	2	-	1	3	1	6	5	22	25	65	128
American Indian 1/, both sexes	385	-	-	1	-	-	-	1	-	7	12	19
Male	216	-	-	1	-	-	-	1	-	6	9	16
Female	169	-	-	-	-	-	-	-	-	1	3	3
Asian or Pacific Islander 2/, both sexes	6,924	7	2	2	4	1	6	14	26	53	81	144
Male	3,488	3	1	1	1	1	4	11	18	39	59	108
Female	3,436	4	1	1	3	-	2	3	8	14	22	36

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Colorado												
All races, both sexes	8,578	7	5	4	4	6	19	17	26	47	115	193
Male	4,060	3	3	4	3	2	10	9	18	32	78	146
Female	4,518	4	2	-	1	4	9	8	8	15	37	47
White, both sexes	8,178	7	5	3	4	4	16	15	23	44	107	174
Male	3,853	3	3	3	3	1	10	7	16	29	75	133
Female	4,325	4	2	-	1	3	6	8	7	15	32	41
Black, both sexes	268	-	-	-	-	2	2	2	3	3	4	15
Male	147	-	-	-	-	1	-	2	2	3	1	12
Female	121	-	-	-	-	1	2	-	1	-	3	3
American Indian 1/, both sexes	38	-	-	-	-	-	1	-	-	-	3	2
Male	20	-	-	-	-	-	-	-	-	-	2	1
Female	18	-	-	-	-	-	1	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	94	-	-	1	-	-	-	-	-	-	1	2
Male	40	-	-	1	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	1	2
Connecticut												
All races, both sexes	9,797	2	1	2	-	5	9	22	19	51	80	175
Male	4,448	2	-	2	-	2	8	18	15	39	56	122
Female	5,349	-	1	-	-	3	1	4	4	12	24	53
White, both sexes	9,140	2	1	2	-	5	8	16	14	42	63	132
Male	4,113	2	-	2	-	2	7	12	13	31	47	90
Female	5,027	-	1	-	-	3	1	4	1	11	16	42
Black, both sexes	586	-	-	-	-	-	1	6	5	9	17	41
Male	297	-	-	-	-	-	1	6	2	8	9	30
Female	289	-	-	-	-	-	-	-	3	1	8	11
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Delaware												
All races, both sexes	2,361	3	1	-	-	2	1	3	11	16	30	68
Male	1,129	1	1	-	-	-	1	1	5	13	21	45
Female	1,232	2	-	-	-	2	-	2	6	3	9	23
White, both sexes	1,980	1	-	-	-	1	1	3	4	9	21	45
Male	933	-	-	-	-	-	1	1	2	7	14	32
Female	1,047	1	-	-	-	1	-	2	2	2	7	13
Black, both sexes	370	1	1	-	-	1	-	-	7	7	9	23
Male	192	-	1	-	-	-	-	-	3	6	7	13
Female	178	1	-	-	-	1	-	-	4	1	2	10
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1,921	1	2	-	2	3	5	4	7	15	26	59
Male	912	-	2	-	1	2	3	3	6	6	17	35
Female	1,009	1	-	-	1	1	2	1	1	9	9	24
White, both sexes	441	-	-	-	-	-	2	-	-	3	5	7
Male	228	-	-	-	-	-	1	-	-	1	4	6
Female	213	-	-	-	-	-	1	-	-	2	1	1
Black, both sexes	1,464	1	2	-	2	3	3	4	7	12	21	52
Male	674	-	2	-	1	2	2	3	6	5	13	29
Female	790	1	-	-	1	1	1	1	1	7	8	23
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Florida												
All races, both sexes	57,533	34	15	13	16	23	49	83	135	274	609	1,137
Male	29,014	19	8	8	9	14	29	54	105	179	408	778
Female	28,519	15	7	5	7	9	20	29	30	95	201	359
White, both sexes	51,128	18	10	10	12	14	27	47	85	189	420	812
Male	25,916	9	5	7	9	10	20	31	65	127	301	591
Female	25,212	9	5	3	3	4	7	16	20	62	119	221
Black, both sexes	6,066	16	3	2	4	9	22	33	46	84	179	308
Male	2,917	10	2	1	-	4	9	21	36	52	98	173
Female	3,149	6	1	1	4	5	13	12	10	32	81	135
American Indian 1/, both sexes	48	-	-	-	-	-	-	-	2	-	-	2
Male	23	-	-	-	-	-	-	-	2	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	291	-	2	1	-	-	-	3	2	1	10	15
Male	158	-	1	-	-	-	-	2	2	-	9	12
Female	133	-	1	1	-	-	-	1	-	1	1	3
Georgia												
All races, both sexes	22,175	25	7	5	10	24	39	77	106	219	445	742
Male	10,837	15	4	3	5	16	19	50	64	145	285	497
Female	11,338	10	3	2	5	8	20	27	42	74	160	245
White, both sexes	15,742	12	6	3	5	11	20	34	44	95	242	415
Male	7,718	7	4	2	2	8	11	23	28	66	172	298
Female	8,024	5	2	1	3	3	9	11	16	29	70	117
Black, both sexes	6,215	13	1	2	5	12	17	43	60	121	196	320
Male	3,016	8	-	1	3	8	7	27	34	77	109	195
Female	3,199	5	1	1	2	4	10	16	26	44	87	125
American Indian 1/, both sexes	30	-	-	-	-	1	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	188	-	-	-	-	-	2	-	2	3	6	6
Male	85	-	-	-	-	-	1	-	2	2	4	3
Female	103	-	-	-	-	-	1	-	-	1	2	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Hawaii												
All races, both sexes	3,150	-	1	1	-	1	5	8	13	21	42	80
Male	1,757	-	-	1	-	1	3	5	8	11	21	52
Female	1,393	-	1	-	-	-	2	3	5	10	21	28
White, both sexes	786	-	-	-	-	-	1	1	1	5	8	16
Male	445	-	-	-	-	-	1	-	1	2	5	10
Female	341	-	-	-	-	-	-	1	-	3	3	6
Black, both sexes	21	-	-	-	-	-	1	-	1	1	-	2
Male	12	-	-	-	-	-	-	-	1	1	-	1
Female	9	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2,339	-	1	1	-	1	3	7	11	14	34	62
Male	1,299	-	-	1	-	1	2	5	6	8	16	41
Female	1,040	-	1	-	-	-	1	2	5	6	18	21
Idaho												
All races, both sexes	3,386	2	2	2	1	1	6	6	9	15	37	75
Male	1,722	1	1	1	1	1	4	5	7	9	23	50
Female	1,664	1	1	1	-	-	2	1	2	6	14	25
White, both sexes	3,328	2	2	1	1	1	6	6	9	15	35	72
Male	1,690	1	1	-	1	1	4	5	7	9	22	48
Female	1,638	1	1	1	-	-	2	1	2	6	13	24
Black, both sexes	8	-	-	1	-	-	-	-	-	-	-	1
Male	7	-	-	1	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	-	-	-	-	-	-	-	-	2	2
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Illinois												
All races, both sexes	35,372	23	8	2	9	26	42	87	112	201	426	806
Male	16,517	16	3	-	6	21	26	58	77	136	278	561
Female	18,855	7	5	2	3	5	16	29	35	65	148	245
White, both sexes	29,733	11	5	2	6	16	27	54	65	111	288	543
Male	13,663	8	2	-	5	12	16	34	49	78	203	400
Female	16,070	3	3	2	1	4	11	20	16	33	85	143
Black, both sexes	5,217	11	3	-	2	9	15	31	46	88	131	252
Male	2,642	7	1	-	1	8	10	23	28	57	71	153
Female	2,575	4	2	-	1	1	5	8	18	31	60	99
American Indian 1/, both sexes	20	1	-	-	-	-	-	-	-	-	2	-
Male	10	1	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	402	-	-	-	1	1	-	2	1	2	5	11
Male	202	-	-	-	-	1	-	1	-	1	3	8
Female	200	-	-	-	1	-	-	1	1	1	2	3
Indiana												
All races, both sexes	19,067	22	3	2	3	11	23	40	85	120	301	478
Male	9,114	11	2	1	-	7	14	25	51	76	210	336
Female	9,953	11	1	1	3	4	9	15	34	44	91	142
White, both sexes	17,509	12	3	2	3	7	18	36	59	96	255	409
Male	8,301	6	2	1	-	6	13	24	36	63	175	290
Female	9,208	6	1	1	3	1	5	12	23	33	80	119
Black, both sexes	1,512	10	-	-	-	4	5	4	26	23	44	67
Male	785	5	-	-	-	1	1	1	15	12	33	44
Female	727	5	-	-	-	3	4	3	11	11	11	23
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	-	1	2	1
Male	22	-	-	-	-	-	-	-	-	1	2	1
Female	17	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Iowa												
All races, both sexes	9,639	2	2	-	1	2	4	13	13	38	86	152
Male	4,438	1	1	-	-	1	2	13	8	25	55	103
Female	5,201	1	1	-	1	1	2	-	5	13	31	49
White, both sexes	9,479	1	1	-	1	2	4	12	13	29	83	143
Male	4,354	-	-	-	-	1	2	12	8	20	52	99
Female	5,125	1	1	-	1	1	2	-	5	9	31	44
Black, both sexes	134	1	1	-	-	-	-	-	-	7	3	7
Male	67	1	1	-	-	-	-	-	-	3	3	3
Female	67	-	-	-	-	-	-	-	-	4	-	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	1	-	2	-	2
Male	17	-	-	-	-	-	-	1	-	2	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	8,135	7	2	1	1	7	7	17	20	32	86	154
Male	3,713	5	1	-	-	3	4	9	13	17	55	104
Female	4,422	2	1	1	1	4	3	8	7	15	31	50
White, both sexes	7,638	4	1	1	-	5	7	12	17	27	75	125
Male	3,461	3	-	-	-	1	4	6	11	14	47	86
Female	4,177	1	1	1	-	4	3	6	6	13	28	39
Black, both sexes	399	3	1	-	1	1	-	4	2	5	9	19
Male	192	2	1	-	-	1	-	3	1	3	6	11
Female	207	1	-	-	1	-	-	1	1	2	3	8
American Indian 1/, both sexes	61	-	-	-	-	-	-	-	-	-	2	8
Male	41	-	-	-	-	-	-	-	-	-	2	6
Female	20	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	1	-	1	1	-	-	2
Male	19	-	-	-	-	1	-	-	1	-	-	1
Female	18	-	-	-	-	-	-	1	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Kentucky												
All races, both sexes	13,471	1	5	3	3	10	18	39	58	115	232	399
Male	6,493	-	2	1	2	4	11	27	40	74	171	293
Female	6,978	1	3	2	1	6	7	12	18	41	61	106
White, both sexes	12,553	1	5	3	3	7	15	31	46	96	191	356
Male	6,036	-	2	1	2	3	9	20	32	65	142	267
Female	6,517	1	3	2	1	4	6	11	14	31	49	89
Black, both sexes	877	-	-	-	-	3	3	8	11	18	37	41
Male	438	-	-	-	-	1	2	7	7	9	26	25
Female	439	-	-	-	-	2	1	1	4	9	11	16
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	1	1	3	2
Male	17	-	-	-	-	-	-	-	1	-	3	1
Female	19	-	-	-	-	-	-	-	-	1	-	1
Louisiana												
All races, both sexes	13,120	13	8	4	3	7	24	32	82	111	249	440
Male	6,453	6	6	-	1	4	13	18	38	80	158	289
Female	6,667	7	2	4	2	3	11	14	44	31	91	151
White, both sexes	9,157	4	5	-	1	4	16	14	38	54	127	227
Male	4,508	2	4	-	-	2	9	10	19	41	90	160
Female	4,649	2	1	-	1	2	7	4	19	13	37	67
Black, both sexes	3,896	9	3	4	2	3	8	18	44	57	119	211
Male	1,904	4	2	-	1	2	4	8	19	39	66	127
Female	1,992	5	1	4	1	1	4	10	25	18	53	84
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	45	-	-	-	-	-	-	-	-	-	2	1
Male	26	-	-	-	-	-	-	-	-	-	2	1
Female	19	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Maine												
All races, both sexes	3,754	-	2	1	1	3	7	8	13	19	37	87
Male	1,853	-	2	1	1	2	4	5	9	12	27	68
Female	1,901	-	-	-	-	1	3	3	4	7	10	19
White, both sexes	3,728	-	2	1	1	3	7	8	12	17	34	84
Male	1,834	-	2	1	1	2	4	5	8	10	25	66
Female	1,894	-	-	-	-	1	3	3	4	7	9	18
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	3
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	1	2	2	-
Male	7	-	-	-	-	-	-	-	1	2	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	14,568	10	5	2	7	9	17	36	58	122	228	402
Male	6,841	4	2	-	4	6	12	29	40	78	142	266
Female	7,727	6	3	2	3	3	5	7	18	44	86	136
White, both sexes	10,451	6	2	1	3	5	9	17	24	58	120	217
Male	4,862	3	1	-	3	3	6	16	16	39	83	153
Female	5,589	3	1	1	-	2	3	1	8	19	37	64
Black, both sexes	3,863	4	2	1	4	4	8	17	31	63	105	181
Male	1,842	1	1	-	1	3	6	12	22	38	56	111
Female	2,021	3	1	1	3	1	2	5	9	25	49	70
American Indian 1/, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	240	-	1	-	-	-	-	1	3	1	3	4
Male	131	-	-	-	-	-	-	1	2	1	3	2
Female	109	-	1	-	-	-	-	-	1	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Massachusetts												
All races, both sexes	17,087	4	2	2	3	8	16	24	32	64	167	265
Male	7,663	2	1	2	2	7	14	17	26	46	119	203
Female	9,424	2	1	-	1	1	2	7	6	18	48	62
White, both sexes	16,293	2	2	2	2	7	15	19	28	50	147	234
Male	7,266	1	1	2	1	6	13	14	24	37	105	181
Female	9,027	1	1	-	1	1	2	5	4	13	42	53
Black, both sexes	596	1	-	-	1	1	1	5	4	12	17	25
Male	304	-	-	-	1	1	1	3	2	7	12	16
Female	292	1	-	-	-	-	-	2	2	5	5	9
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	181	1	-	-	-	-	-	-	-	2	3	5
Male	87	1	-	-	-	-	-	-	-	2	2	5
Female	94	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	31,241	14	6	7	5	13	30	52	104	196	390	728
Male	14,807	8	2	3	2	10	21	36	70	133	274	475
Female	16,434	6	4	4	3	3	9	16	34	63	116	253
White, both sexes	26,374	8	4	7	4	8	15	25	59	123	271	510
Male	12,410	5	1	3	2	7	12	18	45	86	202	349
Female	13,964	3	3	4	2	1	3	7	14	37	69	161
Black, both sexes	4,573	4	2	-	1	5	13	26	42	71	115	206
Male	2,264	2	1	-	-	3	8	17	25	46	70	120
Female	2,309	2	1	-	1	2	5	9	17	25	45	86
American Indian 1/, both sexes	149	1	-	-	-	-	-	-	2	1	3	9
Male	66	-	-	-	-	-	-	-	-	1	2	4
Female	83	1	-	-	-	-	-	-	2	-	1	5
Asian or Pacific Islander 2/, both sexes	145	1	-	-	-	-	2	1	1	1	1	3
Male	67	1	-	-	-	-	1	1	-	-	-	2
Female	78	-	-	-	-	-	1	-	1	1	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Minnesota												
All races, both sexes	10,723	6	3	3	1	6	19	15	27	61	121	199
Male	5,170	2	1	-	1	3	10	6	15	39	84	152
Female	5,553	4	2	3	-	3	9	9	12	22	37	47
White, both sexes	10,337	4	2	3	-	3	14	14	21	52	104	169
Male	4,957	1	1	-	-	2	7	6	12	33	73	131
Female	5,380	3	1	3	-	1	7	8	9	19	31	38
Black, both sexes	188	1	-	-	-	3	4	1	3	4	10	15
Male	98	1	-	-	-	1	3	-	2	3	6	10
Female	90	-	-	-	-	2	1	1	1	1	4	5
American Indian 1/, both sexes	103	-	-	-	-	-	1	-	2	1	5	8
Male	60	-	-	-	-	-	-	-	1	-	4	8
Female	43	-	-	-	-	-	1	-	1	1	1	-
Asian or Pacific Islander 2/, both sexes	95	1	1	-	1	-	-	-	1	4	2	7
Male	55	-	-	-	1	-	-	-	-	3	1	3
Female	40	1	1	-	-	-	-	-	1	1	1	4
Mississippi												
All races, both sexes	10,348	6	-	4	4	6	22	33	60	85	194	350
Male	4,848	4	-	1	4	3	14	17	39	56	124	219
Female	5,500	2	-	3	-	3	8	16	21	29	70	131
White, both sexes	6,900	5	-	1	1	2	7	11	28	44	88	162
Male	3,202	3	-	1	1	2	5	7	21	31	58	108
Female	3,698	2	-	-	-	-	2	4	7	13	30	54
Black, both sexes	3,417	1	-	3	3	4	15	22	32	40	106	184
Male	1,629	1	-	-	3	1	9	10	18	24	66	110
Female	1,788	-	-	3	-	3	6	12	14	16	40	74
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	1	-	3
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Missouri												
All races, both sexes	19,281	10	4	3	7	11	24	21	66	117	254	482
Male	8,923	5	1	1	3	7	16	17	42	75	169	334
Female	10,358	5	3	2	4	4	8	4	24	42	85	148
White, both sexes	17,230	8	3	3	7	9	15	14	47	81	198	379
Male	7,970	4	-	1	3	6	11	11	30	53	136	271
Female	9,260	4	3	2	4	3	4	3	17	28	62	108
Black, both sexes	1,953	2	1	-	-	2	8	5	19	35	54	98
Male	907	1	1	-	-	1	5	4	12	22	32	58
Female	1,046	1	-	-	-	1	3	1	7	13	22	40
American Indian 1/, both sexes	35	-	-	-	-	-	-	1	-	-	-	2
Male	18	-	-	-	-	-	-	1	-	-	-	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	63	-	-	-	-	-	1	1	-	1	2	3
Male	28	-	-	-	-	-	-	1	-	-	1	3
Female	35	-	-	-	-	-	1	-	-	1	1	-
Montana												
All races, both sexes	2,492	-	-	-	1	-	1	3	7	15	30	56
Male	1,270	-	-	-	1	-	1	2	7	10	23	39
Female	1,222	-	-	-	-	-	-	1	-	5	7	17
White, both sexes	2,396	-	-	-	1	-	1	2	5	13	26	49
Male	1,218	-	-	-	1	-	1	1	5	9	20	35
Female	1,178	-	-	-	-	-	-	1	-	4	6	14
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	90	-	-	-	-	-	-	-	2	1	4	7
Male	49	-	-	-	-	-	-	-	2	1	3	4
Female	41	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	1	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Nebraska												
All races, both sexes	4,783	3	2	-	2	1	7	7	11	23	57	94
Male	2,155	1	2	-	1	1	4	4	8	20	39	63
Female	2,628	2	-	-	1	-	3	3	3	3	18	31
White, both sexes	4,603	2	1	-	1	1	5	5	11	18	52	78
Male	2,064	1	1	-	1	1	3	3	8	15	35	53
Female	2,539	1	-	-	-	-	2	2	3	3	17	25
Black, both sexes	142	1	1	-	1	-	1	2	-	5	1	13
Male	73	-	1	-	-	-	-	1	-	5	1	8
Female	69	1	-	-	1	-	1	1	-	-	-	5
American Indian 1/, both sexes	23	-	-	-	-	-	1	-	-	-	4	2
Male	13	-	-	-	-	-	1	-	-	-	3	2
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Nevada												
All races, both sexes	6,301	4	1	1	2	4	9	20	41	48	119	194
Male	3,494	1	1	1	-	4	7	12	29	31	76	131
Female	2,807	3	-	-	2	-	2	8	12	17	43	63
White, both sexes	5,545	4	-	1	-	4	6	13	31	33	95	160
Male	3,092	1	-	1	-	4	5	8	21	21	64	113
Female	2,453	3	-	-	-	-	1	5	10	12	31	47
Black, both sexes	478	-	1	-	-	-	1	5	5	11	20	21
Male	255	-	1	-	-	-	1	3	3	8	9	12
Female	223	-	-	-	-	-	-	2	2	3	11	9
American Indian 1/, both sexes	64	-	-	-	-	-	1	1	2	1	1	4
Male	30	-	-	-	-	-	-	1	2	1	1	2
Female	34	-	-	-	-	-	1	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	214	-	-	-	2	-	1	1	3	3	3	9
Male	117	-	-	-	-	-	1	-	3	1	2	4
Female	97	-	-	-	2	-	-	1	-	2	1	5

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
New Hampshire												
All races, both sexes	3,228	3	1	-	2	5	3	-	7	13	34	68
Male	1,536	2	1	-	1	1	2	-	4	9	23	53
Female	1,692	1	-	-	1	4	1	-	3	4	11	15
White, both sexes	3,206	3	1	-	2	5	3	-	7	13	33	66
Male	1,524	2	1	-	1	1	2	-	4	9	22	52
Female	1,682	1	-	-	1	4	1	-	3	4	11	14
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
New Jersey												
All races, both sexes	24,608	12	3	4	7	6	23	41	61	130	253	442
Male	11,393	8	1	2	5	5	12	33	41	92	180	304
Female	13,215	4	2	2	2	1	11	8	20	38	73	138
White, both sexes	21,347	6	2	3	7	5	15	21	42	80	165	302
Male	9,803	3	-	2	5	4	9	18	29	59	135	217
Female	11,544	3	2	1	2	1	6	3	13	21	30	85
Black, both sexes	2,794	6	1	1	-	1	8	19	18	43	78	129
Male	1,348	5	1	-	-	1	3	14	11	26	36	78
Female	1,446	1	-	1	-	-	5	5	7	17	42	51
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	442	-	-	-	-	-	-	1	1	7	9	10
Male	232	-	-	-	-	-	-	1	1	7	8	8
Female	210	-	-	-	-	-	-	-	-	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
New Mexico												
All races, both sexes	4,437	4	1	1	-	3	5	8	19	27	52	96
Male	2,185	4	-	1	-	2	3	6	11	20	39	55
Female	2,252	-	1	-	-	1	2	2	8	7	13	41
White, both sexes	4,158	4	1	-	-	3	5	8	17	23	40	83
Male	2,042	4	-	-	-	2	3	6	10	16	29	45
Female	2,116	-	1	-	-	1	2	2	7	7	11	38
Black, both sexes	95	-	-	-	-	-	-	-	1	1	3	3
Male	42	-	-	-	-	-	-	-	1	1	2	3
Female	53	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	162	-	-	1	-	-	-	-	1	3	8	9
Male	89	-	-	1	-	-	-	-	-	3	7	6
Female	73	-	-	-	-	-	-	-	1	-	1	3
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	60,421	27	14	9	17	26	41	84	130	269	577	1,090
Male	27,315	18	7	7	10	19	33	53	85	190	381	768
Female	33,106	9	7	2	7	7	8	31	45	79	196	322
White, both sexes	50,864	17	9	5	13	19	23	52	79	163	382	736
Male	22,960	12	4	5	8	14	19	33	49	124	269	540
Female	27,904	5	5	-	5	5	4	19	30	39	113	196
Black, both sexes	8,188	9	5	3	4	7	15	29	42	95	178	314
Male	3,623	6	3	2	2	5	11	18	27	57	98	197
Female	4,565	3	2	1	2	2	4	11	15	38	80	117
American Indian 1/, both sexes	60	-	-	-	-	-	-	-	1	1	-	2
Male	20	-	-	-	-	-	-	-	1	-	-	1
Female	40	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	1,309	1	-	1	-	-	3	3	8	10	17	38
Male	712	-	-	-	-	-	3	2	8	9	14	30
Female	597	1	-	1	-	-	-	1	-	1	3	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
North Carolina												
All races, both sexes	23,625	25	12	5	8	19	35	45	67	194	399	618
Male	11,580	17	2	4	6	10	20	30	45	131	265	426
Female	12,045	8	10	1	2	9	15	15	22	63	134	192
White, both sexes	18,169	10	8	3	2	10	20	23	33	103	234	359
Male	8,902	6	2	2	2	7	11	17	24	75	167	258
Female	9,267	4	6	1	-	3	9	6	9	28	67	101
Black, both sexes	5,143	13	3	2	6	9	14	21	34	90	151	244
Male	2,496	10	-	2	4	3	9	12	21	56	87	157
Female	2,647	3	3	-	2	6	5	9	13	34	64	87
American Indian 1/, both sexes	223	1	-	-	-	-	1	-	-	1	10	14
Male	129	-	-	-	-	-	-	-	-	-	7	10
Female	94	1	-	-	-	-	1	-	-	1	3	4
Asian or Pacific Islander 2/, both sexes	90	1	1	-	-	-	-	1	-	-	4	1
Male	53	1	-	-	-	-	-	1	-	-	4	1
Female	37	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2,096	-	1	1	1	-	2	4	4	9	16	28
Male	1,009	-	1	-	1	-	1	2	1	5	10	19
Female	1,087	-	-	1	-	-	1	2	3	4	6	9
White, both sexes	2,027	-	-	1	1	-	2	3	1	8	14	25
Male	971	-	-	-	1	-	1	2	-	4	8	17
Female	1,056	-	-	1	-	-	1	1	1	4	6	8
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	66	-	1	-	-	-	-	1	3	1	2	3
Male	36	-	1	-	-	-	-	-	1	1	2	2
Female	30	-	-	-	-	-	-	1	2	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Ohio												
All races, both sexes	36,528	25	9	4	4	19	40	60	101	205	402	849
Male	17,194	13	4	4	4	15	23	40	71	140	269	578
Female	19,334	12	5	-	-	4	17	20	30	65	133	271
White, both sexes	32,486	18	7	3	4	13	36	41	79	147	318	660
Male	15,183	10	3	3	4	10	20	25	57	104	224	475
Female	17,303	8	4	-	-	3	16	16	22	43	94	185
Black, both sexes	3,916	7	2	1	-	6	4	19	22	57	81	186
Male	1,945	3	1	1	-	5	3	15	14	35	43	101
Female	1,971	4	1	-	-	1	1	4	8	22	38	85
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	2	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	103	-	-	-	-	-	-	-	-	1	1	3
Male	54	-	-	-	-	-	-	-	-	1	-	2
Female	49	-	-	-	-	-	-	-	-	-	1	1
Oklahoma												
All races, both sexes	12,579	9	4	-	3	8	11	27	45	72	159	272
Male	6,106	6	1	-	1	4	7	20	36	49	94	193
Female	6,473	3	3	-	2	4	4	7	9	23	65	79
White, both sexes	11,078	6	3	-	2	6	10	17	31	48	120	225
Male	5,335	4	-	-	-	3	6	14	25	38	68	160
Female	5,743	2	3	-	2	3	4	3	6	10	52	65
Black, both sexes	795	1	1	-	1	1	1	7	9	18	18	23
Male	395	-	1	-	1	-	1	5	7	7	8	17
Female	400	1	-	-	-	1	-	2	2	11	10	6
American Indian 1/, both sexes	654	2	-	-	-	1	-	2	5	6	21	24
Male	355	2	-	-	-	1	-	1	4	4	18	16
Female	299	-	-	-	-	-	-	1	1	2	3	8
Asian or Pacific Islander 2/, both sexes	52	-	-	-	-	-	-	1	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Oregon												
All races, both sexes	9,437	3	-	2	3	4	10	14	19	43	72	158
Male	4,708	1	-	1	2	3	2	8	12	30	50	113
Female	4,729	2	-	1	1	1	8	6	7	13	22	45
White, both sexes	9,135	2	-	2	3	4	9	12	14	41	66	140
Male	4,557	1	-	1	2	3	1	6	8	28	48	99
Female	4,578	1	-	1	1	1	8	6	6	13	18	41
Black, both sexes	105	-	-	-	-	-	-	1	5	-	-	6
Male	61	-	-	-	-	-	-	1	4	-	-	6
Female	44	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	65	1	-	-	-	-	-	-	-	1	3	4
Male	31	-	-	-	-	-	-	-	-	1	1	2
Female	34	1	-	-	-	-	-	-	-	-	2	2
Asian or Pacific Islander 2/, both sexes	132	-	-	-	-	-	1	1	-	1	3	8
Male	59	-	-	-	-	-	1	1	-	1	1	6
Female	73	-	-	-	-	-	-	-	-	-	2	2
Pennsylvania												
All races, both sexes	43,708	20	10	5	10	14	32	57	92	186	430	788
Male	20,128	7	6	3	7	3	23	33	56	133	294	562
Female	23,580	13	4	2	3	11	9	24	36	53	136	226
White, both sexes	39,794	18	7	3	6	10	20	40	61	139	324	605
Male	18,204	6	4	1	4	3	13	22	39	104	224	439
Female	21,590	12	3	2	2	7	7	18	22	35	100	166
Black, both sexes	3,726	2	3	2	4	4	12	17	29	44	97	174
Male	1,821	1	2	2	3	-	10	11	15	26	63	115
Female	1,905	1	1	-	1	4	2	6	14	18	34	59
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	176	-	-	-	-	-	-	-	2	3	9	9
Male	93	-	-	-	-	-	-	-	2	3	7	8
Female	83	-	-	-	-	-	-	-	-	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Rhode Island												
All races, both sexes	3,350	1	1	-	1	2	1	5	4	9	26	50
Male	1,493	-	1	-	-	2	1	4	3	6	19	38
Female	1,857	1	-	-	1	-	-	1	1	3	7	12
White, both sexes	3,250	1	1	-	-	2	1	4	4	7	24	48
Male	1,447	-	1	-	-	2	1	3	3	4	18	36
Female	1,803	1	-	-	-	-	-	1	1	3	6	12
Black, both sexes	69	-	-	-	1	-	-	1	-	1	2	1
Male	29	-	-	-	-	-	-	1	-	1	1	1
Female	40	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12,152	15	4	1	1	8	20	29	60	103	218	410
Male	6,188	7	2	-	1	5	13	24	36	61	153	267
Female	5,964	8	2	1	-	3	7	5	24	42	65	143
White, both sexes	8,557	5	2	1	1	3	12	16	24	55	117	244
Male	4,429	2	1	-	1	3	8	13	14	37	91	173
Female	4,128	3	1	1	-	-	4	3	10	18	26	71
Black, both sexes	3,535	10	2	-	-	4	8	13	35	47	101	163
Male	1,732	5	1	-	-	2	5	11	21	24	62	93
Female	1,803	5	1	-	-	2	3	2	14	23	39	70
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	1	-	-	1	-	-	2
Male	15	-	-	-	-	-	-	-	1	-	-	1
Female	17	-	-	-	-	1	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
South Dakota												
All races, both sexes	2,390	2	-	-	1	-	4	3	4	9	24	49
Male	1,154	2	-	-	1	-	1	3	3	6	16	38
Female	1,236	-	-	-	-	-	3	-	1	3	8	11
White, both sexes	2,276	-	-	-	1	-	1	2	3	8	16	42
Male	1,090	-	-	-	1	-	1	2	2	6	11	32
Female	1,186	-	-	-	-	-	-	-	1	2	5	10
Black, both sexes	12	1	-	-	-	-	-	1	-	-	1	1
Male	10	1	-	-	-	-	-	1	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	100	1	-	-	-	-	3	-	1	1	7	6
Male	52	1	-	-	-	-	-	-	1	-	4	5
Female	48	-	-	-	-	-	3	-	-	1	3	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19,363	19	5	2	4	13	25	40	98	168	361	601
Male	9,474	10	1	1	2	6	19	31	70	114	227	409
Female	9,889	9	4	1	2	7	6	9	28	54	134	192
White, both sexes	16,313	11	4	2	3	9	14	27	65	118	259	437
Male	7,985	5	-	1	1	3	13	23	47	84	162	311
Female	8,328	6	4	1	2	6	1	4	18	34	97	126
Black, both sexes	2,985	8	1	-	1	3	11	12	32	50	100	159
Male	1,450	5	1	-	1	2	6	7	22	30	63	93
Female	1,535	3	-	-	-	1	5	5	10	20	37	66
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	3
Male	10	-	-	-	-	-	-	-	-	-	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	1	-	1	1	-	2	2
Male	29	-	-	-	-	1	-	1	1	-	2	2
Female	22	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Texas												
All races, both sexes	51,464	54	23	11	18	43	75	105	225	378	829	1,518
Male	25,051	30	13	8	10	25	48	63	149	254	537	954
Female	26,413	24	10	3	8	18	27	42	76	124	292	564
White, both sexes	44,359	44	17	6	12	32	60	72	149	270	608	1,126
Male	21,525	24	9	5	8	19	40	42	103	181	411	721
Female	22,834	20	8	1	4	13	20	30	46	89	197	405
Black, both sexes	6,532	8	6	4	5	11	15	33	73	102	206	368
Male	3,223	4	4	2	2	6	8	21	44	69	118	214
Female	3,309	4	2	2	3	5	7	12	29	33	88	154
American Indian 1/, both sexes	84	-	-	-	1	-	-	-	2	1	3	7
Male	42	-	-	-	-	-	-	-	1	1	2	6
Female	42	-	-	-	1	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	489	2	-	1	-	-	-	-	1	5	12	17
Male	261	2	-	1	-	-	-	-	1	3	6	13
Female	228	-	-	-	-	-	-	-	-	2	6	4
Utah												
All races, both sexes	3,916	2	1	-	2	4	7	8	11	22	38	74
Male	1,967	-	-	-	2	3	5	2	10	13	28	52
Female	1,949	2	1	-	-	1	2	6	1	9	10	22
White, both sexes	3,816	2	1	-	2	4	6	7	10	20	33	71
Male	1,915	-	-	-	2	3	4	2	9	11	23	51
Female	1,901	2	1	-	-	1	2	5	1	9	10	20
Black, both sexes	23	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	1	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	-	-	1	1	1	1	2	1
Male	30	-	-	-	-	-	1	-	1	1	2	-
Female	29	-	-	-	-	-	-	1	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Vermont												
All races, both sexes	1,617	2	-	1	1	2	-	1	3	6	17	28
Male	774	1	-	-	1	-	-	1	3	4	14	23
Female	843	1	-	1	-	2	-	-	-	2	3	5
White, both sexes	1,613	2	-	1	1	2	-	1	3	6	17	28
Male	772	1	-	-	1	-	-	1	3	4	14	23
Female	841	1	-	1	-	2	-	-	-	2	3	5
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	18,822	22	9	3	7	25	24	32	72	134	279	514
Male	9,100	11	6	2	3	17	20	23	46	86	196	361
Female	9,722	11	3	1	4	8	4	9	26	48	83	153
White, both sexes	14,679	11	4	1	3	16	16	22	36	77	173	317
Male	7,049	5	2	1	1	12	15	17	24	54	130	228
Female	7,630	6	2	-	2	4	1	5	12	23	43	89
Black, both sexes	3,887	11	5	1	3	9	7	7	33	55	103	185
Male	1,903	6	4	-	2	5	4	3	19	30	65	124
Female	1,984	5	1	1	1	4	3	4	14	25	38	61
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	240	-	-	1	1	-	1	3	3	2	2	12
Male	142	-	-	1	-	-	1	3	3	2	1	9
Female	98	-	-	-	1	-	-	-	-	-	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Washington												
All races, both sexes	14,540	5	1	-	1	6	13	16	47	102	140	268
Male	7,067	4	-	-	1	2	7	13	30	68	99	196
Female	7,473	1	1	-	-	4	6	3	17	34	41	72
White, both sexes	13,532	4	-	-	1	5	10	12	38	77	113	231
Male	6,528	3	-	-	1	2	6	10	24	51	81	166
Female	7,004	1	-	-	-	3	4	2	14	26	32	65
Black, both sexes	362	1	1	-	-	-	1	1	3	13	11	19
Male	198	1	-	-	-	-	-	-	2	9	5	16
Female	164	-	1	-	-	-	1	1	1	4	6	3
American Indian 1/, both sexes	154	-	-	-	-	1	1	1	1	1	6	8
Male	73	-	-	-	-	-	-	1	-	1	4	7
Female	81	-	-	-	-	1	1	-	1	-	2	1
Asian or Pacific Islander 2/, both sexes	492	-	-	-	-	-	1	2	5	11	10	10
Male	268	-	-	-	-	-	1	2	4	7	9	7
Female	224	-	-	-	-	-	-	-	1	4	1	3
West Virginia												
All races, both sexes	6,933	3	1	-	1	2	7	10	19	40	75	152
Male	3,301	1	1	-	1	1	4	7	13	23	49	102
Female	3,632	2	-	-	-	1	3	3	6	17	26	50
White, both sexes	6,709	2	1	-	1	2	7	10	17	40	74	146
Male	3,193	1	1	-	1	1	4	7	12	23	48	97
Female	3,516	1	-	-	-	1	3	3	5	17	26	49
Black, both sexes	217	1	-	-	-	-	-	-	2	-	1	5
Male	103	-	-	-	-	-	-	-	1	-	1	4
Female	114	1	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Wisconsin												
All races, both sexes	15,371	6	1	1	3	9	14	17	39	67	146	263
Male	7,297	5	1	-	2	6	10	10	25	41	92	172
Female	8,074	1	-	1	1	3	4	7	14	26	54	91
White, both sexes	14,682	4	1	-	2	7	12	14	25	55	127	227
Male	6,983	3	1	-	2	4	9	10	20	37	86	152
Female	7,699	1	-	-	-	3	3	4	5	18	41	75
Black, both sexes	546	1	-	1	-	2	1	3	13	8	16	31
Male	247	1	-	-	-	2	-	-	5	2	3	18
Female	299	-	-	1	-	-	1	3	8	6	13	13
American Indian 1/, both sexes	79	-	-	-	-	-	-	-	-	2	3	3
Male	36	-	-	-	-	-	-	-	-	1	3	1
Female	43	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	64	1	-	-	1	-	1	-	1	2	-	2
Male	31	1	-	-	-	-	1	-	-	1	-	1
Female	33	-	-	-	1	-	-	-	1	1	-	1
Wyoming												
All races, both sexes	1,307	-	-	1	-	2	2	3	3	8	15	33
Male	650	-	-	1	-	-	1	1	3	5	8	20
Female	657	-	-	-	-	2	1	2	-	3	7	13
White, both sexes	1,283	-	-	1	-	1	2	3	2	7	13	33
Male	637	-	-	1	-	-	1	1	2	4	8	20
Female	646	-	-	-	-	1	1	2	-	3	5	13
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	1	-	-	1	1	1	-
Male	10	-	-	-	-	-	-	-	1	1	-	-
Female	5	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
United States												
All races, both sexes	28,344	37,777	43,477	51,134	68,521	101,411	140,359	149,366	110,313	45,367	9,936	41
Male	19,803	25,916	28,453	31,951	39,507	53,631	65,169	58,661	33,918	10,039	1,515	30
Female	8,541	11,861	15,024	19,183	29,014	47,780	75,190	90,705	76,395	35,328	8,421	11
White, both sexes	20,717	28,671	34,294	40,861	56,627	87,595	124,734	136,245	101,369	41,519	8,691	33
Male	14,952	20,138	22,854	25,990	33,154	47,042	58,675	54,009	31,343	9,166	1,313	24
Female	5,765	8,533	11,440	14,871	23,473	40,553	66,059	82,236	70,026	32,353	7,378	9
Black, both sexes	6,821	8,000	7,951	8,660	10,048	11,337	12,648	10,335	7,060	3,176	1,064	8
Male	4,298	5,008	4,825	4,959	5,353	5,337	5,129	3,449	1,856	637	147	6
Female	2,523	2,992	3,126	3,701	4,695	6,000	7,519	6,886	5,204	2,539	917	2
American Indian 1/, both sexes	225	276	310	368	360	425	451	336	234	70	20	-
Male	162	189	190	220	200	213	182	124	62	12	5	-
Female	63	87	120	148	160	212	269	212	172	58	15	-
Asian or Pacific Islander 2/, both sexes	581	830	922	1,245	1,486	2,054	2,526	2,450	1,650	602	161	-
Male	391	581	584	782	800	1,039	1,183	1,079	657	224	50	-
Female	190	249	338	463	686	1,015	1,343	1,371	993	378	111	-
Alabama												
All races, both sexes	732	918	1,007	1,176	1,573	2,144	2,589	2,553	1,670	655	155	-
Male	487	603	663	719	903	1,059	1,088	882	461	128	17	-
Female	245	315	344	457	670	1,085	1,501	1,671	1,209	527	138	-
White, both sexes	463	585	711	864	1,172	1,654	2,056	2,117	1,361	539	94	-
Male	319	383	474	547	694	829	872	742	373	105	10	-
Female	144	202	237	317	478	825	1,184	1,375	988	434	84	-
Black, both sexes	263	329	291	310	393	484	528	431	303	115	61	-
Male	163	217	185	172	206	229	213	139	86	23	7	-
Female	100	112	106	138	187	255	315	292	217	92	54	-
American Indian 1/, both sexes	2	1	2	2	2	4	2	2	2	-	-	-
Male	2	1	1	-	1	-	1	-	-	-	-	-
Female	-	-	1	2	1	4	1	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	3	-	6	2	3	3	4	1	-	-
Male	3	2	3	-	2	1	2	1	2	-	-	-
Female	1	1	-	-	4	1	1	2	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Alaska												
All races, both sexes	66	74	54	71	101	112	135	107	59	25	8	-
Male	48	46	42	47	66	76	51	45	23	4	1	-
Female	18	28	12	24	35	36	84	62	36	21	7	-
White, both sexes	50	54	38	52	76	85	106	84	51	21	6	-
Male	36	35	29	35	52	60	44	39	19	3	1	-
Female	14	19	9	17	24	25	62	45	32	18	5	-
Black, both sexes	2	4	1	-	3	2	1	1	-	2	1	-
Male	1	3	1	-	2	1	1	-	-	1	-	-
Female	1	1	-	-	1	1	-	1	-	1	1	-
American Indian 1/, both sexes	13	11	13	12	18	22	20	15	8	2	1	-
Male	10	5	10	7	10	15	5	5	4	-	-	-
Female	3	6	3	5	8	7	15	10	4	2	1	-
Asian or Pacific Islander 2/, both sexes	1	5	2	7	4	3	8	7	-	-	-	-
Male	1	3	2	5	2	-	1	1	-	-	-	-
Female	-	2	-	2	2	3	7	6	-	-	-	-
Arizona												
All races, both sexes	489	639	811	941	1,250	1,767	2,381	2,521	1,677	634	123	8
Male	366	430	559	606	749	941	1,147	1,078	582	156	29	6
Female	123	209	252	335	501	826	1,234	1,443	1,095	478	94	2
White, both sexes	434	582	737	865	1,172	1,671	2,277	2,439	1,632	609	114	6
Male	323	390	505	567	700	895	1,089	1,040	565	146	24	5
Female	111	192	232	298	472	776	1,188	1,399	1,067	463	90	1
Black, both sexes	36	33	36	35	48	40	47	42	18	15	4	2
Male	29	25	27	21	34	19	31	20	7	6	1	1
Female	7	8	9	14	14	21	16	22	11	9	3	1
American Indian 1/, both sexes	15	20	30	34	18	39	38	29	19	6	3	-
Male	12	13	24	15	10	23	17	12	7	2	2	-
Female	3	7	6	19	8	16	21	17	12	4	1	-
Asian or Pacific Islander 2/, both sexes	4	4	8	7	12	17	19	11	8	4	2	-
Male	2	2	3	3	5	4	10	6	3	2	2	-
Female	2	2	5	4	7	13	9	5	5	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Arkansas												
All races, both sexes	380	503	616	691	910	1,220	1,557	1,640	1,165	474	104	1
Male	260	358	404	419	499	608	723	592	352	106	14	-
Female	120	145	212	272	411	612	834	1,048	813	368	90	1
White, both sexes	278	387	500	587	753	1,076	1,387	1,497	1,029	410	84	1
Male	198	276	326	370	422	553	652	541	315	90	10	-
Female	80	111	174	217	331	523	735	956	714	320	74	1
Black, both sexes	98	110	110	100	153	140	163	140	131	63	20	-
Male	60	77	73	49	75	54	69	49	36	15	4	-
Female	38	33	37	51	78	86	94	91	95	48	16	-
American Indian 1/, both sexes	3	2	3	2	2	1	5	1	4	-	-	-
Male	2	1	3	-	1	-	1	1	-	-	-	-
Female	1	1	-	2	1	1	4	-	4	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	3	2	2	3	2	2	1	1	-	-
Male	-	4	2	-	1	1	1	1	1	1	-	-
Female	1	-	1	2	1	2	1	1	-	-	-	-
California												
All races, both sexes	2,802	3,866	4,174	5,010	6,738	10,064	15,013	16,345	12,075	5,113	1,171	7
Male	1,976	2,644	2,742	3,103	3,908	5,422	7,150	6,819	4,115	1,235	188	4
Female	826	1,222	1,432	1,907	2,830	4,642	7,863	9,526	7,960	3,878	983	3
White, both sexes	2,090	2,918	3,250	3,856	5,399	8,232	12,852	14,279	10,657	4,544	1,004	7
Male	1,514	2,035	2,191	2,432	3,186	4,489	6,172	6,023	3,630	1,065	151	4
Female	576	883	1,059	1,424	2,213	3,743	6,680	8,256	7,027	3,479	853	3
Black, both sexes	474	579	543	628	699	816	904	817	565	250	86	-
Male	302	357	323	339	374	424	383	268	156	54	9	-
Female	172	222	220	289	325	392	521	549	409	196	77	-
American Indian 1/, both sexes	21	33	28	47	35	58	51	50	13	8	1	-
Male	16	25	12	33	19	29	26	18	4	1	-	-
Female	5	8	16	14	16	29	25	32	9	7	1	-
Asian or Pacific Islander 2/, both sexes	217	336	353	479	605	958	1,206	1,199	840	311	80	-
Male	144	227	216	299	329	480	569	510	325	115	28	-
Female	73	109	137	180	276	478	637	689	515	196	52	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Colorado												
All races, both sexes	314	377	450	528	695	1,020	1,385	1,589	1,160	514	103	-
Male	228	281	281	324	381	536	615	634	354	103	15	-
Female	86	96	169	204	314	484	770	955	806	411	88	-
White, both sexes	293	347	411	495	643	971	1,332	1,548	1,138	499	99	-
Male	213	259	259	304	355	510	593	616	347	100	14	-
Female	80	88	152	191	288	461	739	932	791	399	85	-
Black, both sexes	15	26	28	27	34	24	29	29	12	11	2	-
Male	11	18	16	17	18	16	13	11	4	2	-	-
Female	4	8	12	10	16	8	16	18	8	9	2	-
American Indian 1/, both sexes	2	-	4	2	4	7	6	2	3	2	-	-
Male	2	-	2	1	4	4	1	2	1	-	-	-
Female	-	-	2	1	-	3	5	-	2	2	-	-
Asian or Pacific Islander 2/, both sexes	4	4	7	4	14	18	18	10	7	2	2	-
Male	2	4	4	2	4	6	8	5	2	1	1	-
Female	2	-	3	2	10	12	10	5	5	1	1	-
Connecticut												
All races, both sexes	218	356	392	462	652	1,110	1,709	2,079	1,536	733	183	1
Male	169	241	260	302	394	570	791	833	475	131	17	1
Female	49	115	132	160	258	540	918	1,246	1,061	602	166	-
White, both sexes	178	301	334	395	581	1,029	1,627	2,026	1,492	717	174	1
Male	142	206	225	262	350	540	757	812	468	129	15	1
Female	36	95	109	133	231	489	870	1,214	1,024	588	159	-
Black, both sexes	35	49	52	58	64	70	74	46	37	14	8	-
Male	23	31	30	34	40	26	31	17	5	2	2	-
Female	12	18	22	24	24	44	43	29	32	12	6	-
American Indian 1/, both sexes	-	1	2	1	1	1	2	1	2	-	-	-
Male	-	1	1	1	1	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	5	4	8	6	10	6	6	5	2	1	-
Male	4	3	4	5	3	4	2	4	2	-	-	-
Female	1	2	-	3	3	6	4	2	3	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Delaware												
All races, both sexes	94	99	141	154	220	274	397	433	284	101	28	1
Male	62	67	88	96	131	154	167	169	81	22	3	1
Female	32	32	53	58	89	120	230	264	203	79	25	-
White, both sexes	68	77	112	117	164	246	351	395	252	88	24	1
Male	45	53	70	75	97	138	147	153	74	20	3	1
Female	23	24	42	42	67	108	204	242	178	68	21	-
Black, both sexes	25	22	28	35	54	27	46	37	31	12	4	-
Male	17	14	17	21	32	16	20	16	7	2	-	-
Female	8	8	11	14	22	11	26	21	24	10	4	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	1	1	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	112	98	120	174	172	210	323	267	205	89	27	-
Male	71	62	71	103	93	113	138	94	66	23	3	-
Female	41	36	49	71	79	97	185	173	139	66	24	-
White, both sexes	6	16	23	23	22	51	83	82	74	36	8	-
Male	5	13	12	14	13	36	38	44	28	13	-	-
Female	1	3	11	9	9	15	45	38	46	23	8	-
Black, both sexes	106	81	97	150	149	157	238	182	128	50	19	-
Male	66	48	59	88	80	76	98	48	37	8	3	-
Female	40	33	38	62	69	81	140	134	91	42	16	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	2	1	-	-
Male	-	-	-	-	-	-	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	2	2	1	2	-	-
Male	-	1	-	1	-	1	2	1	-	1	-	-
Female	-	-	-	-	1	1	-	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Florida												
All races, both sexes	1,661	2,264	2,810	3,408	4,682	7,298	10,236	10,840	7,998	3,259	687	2
Male	1,184	1,560	1,887	2,211	2,808	4,009	5,133	4,679	2,911	861	159	1
Female	477	704	923	1,197	1,874	3,289	5,103	6,161	5,087	2,398	528	1
White, both sexes	1,230	1,796	2,301	2,830	3,985	6,567	9,420	10,182	7,504	3,048	620	1
Male	909	1,294	1,580	1,872	2,432	3,667	4,778	4,469	2,780	810	150	-
Female	321	502	721	958	1,553	2,900	4,642	5,713	4,724	2,238	470	1
Black, both sexes	413	451	482	549	660	685	767	612	473	202	65	1
Male	259	257	293	320	357	317	332	193	127	47	8	1
Female	154	194	189	229	303	368	435	419	346	155	57	-
American Indian 1/, both sexes	4	4	6	5	5	4	6	5	4	1	-	-
Male	4	1	4	4	3	1	2	-	-	-	-	-
Female	-	3	2	1	2	3	4	5	4	1	-	-
Asian or Pacific Islander 2/, both sexes	14	13	21	24	32	42	43	41	17	8	2	-
Male	12	8	10	15	16	24	21	17	4	4	1	-
Female	2	5	11	9	16	18	22	24	13	4	1	-
Georgia												
All races, both sexes	1,096	1,406	1,619	1,734	2,028	2,583	3,373	3,320	2,254	898	164	1
Male	711	979	1,057	1,048	1,168	1,259	1,519	1,194	596	184	18	1
Female	385	427	562	686	860	1,324	1,854	2,126	1,658	714	146	-
White, both sexes	575	813	1,035	1,164	1,394	1,892	2,678	2,658	1,824	711	110	1
Male	406	603	688	731	828	952	1,260	987	483	147	11	1
Female	169	210	347	433	566	940	1,418	1,671	1,341	564	99	-
Black, both sexes	509	580	570	550	605	650	670	637	418	184	52	-
Male	298	366	365	307	324	290	249	195	111	36	6	-
Female	211	214	205	243	281	360	421	442	307	148	46	-
American Indian 1/, both sexes	2	2	4	1	3	6	3	3	2	1	-	-
Male	1	2	1	1	2	5	2	2	-	1	-	-
Female	1	-	3	-	1	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	10	11	10	19	26	35	22	22	10	2	2	-
Male	6	8	3	9	14	12	8	10	2	-	1	-
Female	4	3	7	10	12	23	14	12	8	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Hawaii												
All races, both sexes	121	158	178	216	253	377	521	545	420	146	43	-
Male	87	114	125	143	147	216	292	260	193	64	14	-
Female	34	44	53	73	106	161	229	285	227	82	29	-
White, both sexes	35	46	44	48	58	111	131	138	99	37	7	-
Male	29	32	35	32	38	63	75	66	44	10	2	-
Female	6	14	9	16	20	48	56	72	55	27	5	-
Black, both sexes	-	2	1	2	4	1	2	3	-	-	1	-
Male	-	1	1	2	3	1	1	-	-	-	-	-
Female	-	1	-	-	1	-	1	3	-	-	1	-
American Indian 1/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	86	109	133	166	191	264	387	404	321	109	35	-
Male	58	80	89	109	106	152	216	194	149	54	12	-
Female	28	29	44	57	85	112	171	210	172	55	23	-
Idaho												
All races, both sexes	95	146	156	203	285	430	576	620	506	182	31	-
Male	62	103	113	136	170	248	289	269	170	50	9	-
Female	33	43	43	67	115	182	287	351	336	132	22	-
White, both sexes	89	141	154	199	282	423	566	610	504	180	30	-
Male	59	99	112	134	168	245	283	263	170	49	9	-
Female	30	42	42	65	114	178	283	347	334	131	21	-
Black, both sexes	3	1	-	-	-	-	2	-	-	-	-	-
Male	2	1	-	-	-	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	2	2	3	1	5	4	3	2	1	-	-
Male	1	2	1	2	-	2	2	1	-	-	-	-
Female	2	-	1	1	1	3	2	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	2	2	4	7	-	1	1	-
Male	-	1	-	-	2	1	2	5	-	1	-	-
Female	-	1	-	1	-	1	2	2	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Illinois												
All races, both sexes	1,219	1,602	1,776	2,111	2,830	4,226	6,001	6,332	4,984	2,072	476	1
Male	863	1,093	1,172	1,317	1,616	2,178	2,713	2,452	1,429	431	71	-
Female	356	509	604	794	1,214	2,048	3,288	3,880	3,555	1,641	405	1
White, both sexes	832	1,128	1,321	1,578	2,215	3,574	5,211	5,791	4,632	1,897	426	-
Male	606	793	878	998	1,293	1,866	2,376	2,245	1,342	393	66	-
Female	226	335	443	580	922	1,708	2,835	3,546	3,290	1,504	360	-
Black, both sexes	366	452	429	483	567	606	727	470	323	158	47	1
Male	244	285	280	286	296	292	305	176	80	34	5	-
Female	122	167	149	197	271	314	422	294	243	124	42	1
American Indian 1/, both sexes	2	1	2	3	-	5	2	2	-	-	-	-
Male	1	1	1	2	-	1	-	2	-	-	-	-
Female	1	-	1	1	-	4	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	21	24	47	48	41	61	69	29	17	3	-
Male	12	14	13	31	27	19	32	29	7	4	-	-
Female	7	7	11	16	21	22	29	40	22	13	3	-
Indiana												
All races, both sexes	656	864	1,003	1,277	1,652	2,379	3,220	3,285	2,434	969	240	-
Male	447	583	660	818	962	1,283	1,457	1,245	699	196	31	-
Female	209	281	343	459	690	1,096	1,763	2,040	1,735	773	209	-
White, both sexes	558	719	879	1,122	1,503	2,199	3,017	3,143	2,319	927	223	-
Male	379	487	575	722	872	1,201	1,364	1,202	669	185	29	-
Female	179	232	304	400	631	998	1,653	1,941	1,650	742	194	-
Black, both sexes	97	142	120	148	139	174	199	136	115	42	17	-
Male	68	94	81	90	83	81	91	42	30	11	2	-
Female	29	48	39	58	56	93	108	94	85	31	15	-
American Indian 1/, both sexes	-	1	1	1	2	1	-	-	-	-	-	-
Male	-	1	1	-	2	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	6	8	5	4	6	-	-	-	-
Male	-	1	3	6	5	-	2	1	-	-	-	-
Female	1	1	-	-	3	5	2	5	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Iowa												
All races, both sexes	258	356	398	499	649	1,070	1,643	1,931	1,592	743	187	-
Male	182	245	248	325	387	570	803	790	498	163	18	-
Female	76	111	150	174	262	500	840	1,141	1,094	580	169	-
White, both sexes	239	341	387	485	629	1,054	1,624	1,917	1,589	739	186	-
Male	169	237	242	317	379	558	797	783	498	162	18	-
Female	70	104	145	168	250	496	827	1,134	1,091	577	168	-
Black, both sexes	18	13	9	12	15	13	15	13	3	3	1	-
Male	12	8	4	6	6	10	4	6	-	-	-	-
Female	6	5	5	6	9	3	11	7	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	2	5	3	2	1	-	1	-	-
Male	1	-	2	2	2	2	2	1	-	1	-	-
Female	-	2	-	-	3	1	-	-	-	-	-	-
Kansas												
All races, both sexes	233	316	340	416	612	927	1,363	1,592	1,271	594	137	-
Male	176	234	218	257	345	487	636	627	376	129	17	-
Female	57	82	122	159	267	440	727	965	895	465	120	-
White, both sexes	205	279	303	378	562	848	1,304	1,532	1,245	575	133	-
Male	155	209	195	237	318	449	611	603	372	123	17	-
Female	50	70	108	141	244	399	693	929	873	452	116	-
Black, both sexes	24	31	28	28	37	63	46	53	22	18	4	-
Male	17	20	16	14	20	29	17	22	3	6	-	-
Female	7	11	12	14	17	34	29	31	19	12	4	-
American Indian 1/, both sexes	3	2	5	8	7	11	8	4	3	-	-	-
Male	3	2	4	4	5	8	5	2	-	-	-	-
Female	-	-	1	4	2	3	3	2	3	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	4	2	6	5	5	3	1	1	-	-
Male	1	3	3	2	2	1	3	-	1	-	-	-
Female	-	1	1	-	4	4	2	3	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Kentucky												
All races, both sexes	578	664	805	1,054	1,233	1,685	2,218	2,151	1,533	549	118	-
Male	415	441	507	635	710	894	954	777	402	116	17	-
Female	163	223	298	419	523	791	1,264	1,374	1,131	433	101	-
White, both sexes	507	599	740	959	1,149	1,581	2,117	2,043	1,476	517	111	-
Male	367	407	471	577	658	846	915	743	383	112	14	-
Female	140	192	269	382	491	735	1,202	1,300	1,093	405	97	-
Black, both sexes	69	63	59	90	79	101	96	104	56	32	7	-
Male	47	34	32	56	50	47	36	34	18	4	3	-
Female	22	29	27	34	29	54	60	70	38	28	4	-
American Indian 1/, both sexes	2	-	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	6	4	4	3	5	4	1	-	-	-
Male	-	-	4	1	2	1	3	-	1	-	-	-
Female	-	2	2	3	2	2	2	4	-	-	-	-
Louisiana												
All races, both sexes	648	806	915	991	1,258	1,658	2,053	1,950	1,279	496	93	-
Male	436	506	591	605	732	850	936	697	367	100	20	-
Female	212	300	324	386	526	808	1,117	1,253	912	396	73	-
White, both sexes	344	458	572	615	862	1,213	1,594	1,542	1,024	382	61	-
Male	242	295	383	399	521	636	747	565	296	74	13	-
Female	102	163	189	216	341	577	847	977	728	308	48	-
Black, both sexes	299	341	337	367	384	440	453	400	252	113	32	-
Male	189	205	203	200	206	211	187	128	70	26	7	-
Female	110	136	134	167	178	229	266	272	182	87	25	-
American Indian 1/, both sexes	3	2	2	3	3	4	3	-	-	-	-	-
Male	3	2	2	1	2	2	2	-	-	-	-	-
Female	-	-	-	2	1	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	4	6	9	1	3	8	3	1	-	-
Male	2	4	3	5	3	1	-	4	1	-	-	-
Female	-	1	1	1	6	-	3	4	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Maine												
All races, both sexes	109	161	197	216	312	453	643	695	519	220	51	-
Male	82	124	137	133	184	259	323	267	153	54	6	-
Female	27	37	60	83	128	194	320	428	366	166	45	-
White, both sexes	103	157	197	216	312	451	640	694	519	219	51	-
Male	76	120	137	133	184	258	323	267	153	53	6	-
Female	27	37	60	83	128	193	317	427	366	166	45	-
Black, both sexes	4	2	-	-	-	2	-	-	-	-	-	-
Male	4	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	3	1	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	564	742	852	943	1,222	1,774	2,390	2,514	1,793	713	163	2
Male	360	478	544	553	693	918	1,046	978	506	157	23	2
Female	204	264	308	390	529	856	1,344	1,536	1,287	556	140	-
White, both sexes	307	406	495	569	777	1,287	1,834	2,098	1,490	597	127	2
Male	205	267	328	343	451	674	831	848	434	138	18	2
Female	102	139	167	226	326	613	1,003	1,250	1,056	459	109	-
Black, both sexes	250	323	336	347	420	455	510	384	279	105	34	-
Male	151	200	202	190	224	224	195	118	65	17	5	-
Female	99	123	134	157	196	231	315	266	214	88	29	-
American Indian 1/, both sexes	-	2	3	2	-	1	2	1	2	-	-	-
Male	-	1	3	-	-	1	-	-	1	-	-	-
Female	-	1	-	2	-	-	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	11	18	25	25	31	44	31	22	11	2	-
Male	4	10	11	20	18	19	20	12	6	2	-	-
Female	3	1	7	5	7	12	24	19	16	9	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Massachusetts												
All races, both sexes	406	569	712	811	1,168	1,980	3,093	3,547	2,738	1,169	307	-
Male	308	419	456	522	676	1,067	1,438	1,284	791	227	36	-
Female	98	150	256	289	492	913	1,655	2,263	1,947	942	271	-
White, both sexes	368	532	654	744	1,090	1,872	2,976	3,448	2,668	1,136	297	-
Male	280	389	418	480	641	1,010	1,389	1,243	777	219	35	-
Female	88	143	236	264	449	862	1,587	2,205	1,891	917	262	-
Black, both sexes	28	29	49	50	55	74	85	75	54	25	5	-
Male	21	23	33	30	22	42	42	30	11	6	1	-
Female	7	6	16	20	33	32	43	45	43	19	4	-
American Indian 1/, both sexes	2	-	1	-	2	6	4	-	-	1	-	-
Male	2	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	6	3	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	8	8	8	17	21	28	28	24	16	7	5	-
Male	5	7	4	12	12	15	6	11	3	2	-	-
Female	3	1	4	5	9	13	22	13	13	5	5	-
Michigan												
All races, both sexes	1,115	1,506	1,606	1,789	2,515	3,908	5,371	5,741	4,167	1,629	349	-
Male	770	1,033	1,041	1,082	1,408	2,037	2,487	2,274	1,231	351	59	-
Female	345	473	565	707	1,107	1,871	2,884	3,467	2,936	1,278	290	-
White, both sexes	751	1,065	1,208	1,409	2,032	3,332	4,688	5,222	3,843	1,489	301	-
Male	550	750	802	866	1,161	1,779	2,198	2,064	1,138	321	51	-
Female	201	315	406	543	871	1,553	2,490	3,158	2,705	1,168	250	-
Black, both sexes	345	424	380	355	446	538	644	483	291	137	45	-
Male	208	272	232	204	227	240	276	193	84	29	7	-
Female	137	152	148	151	219	298	368	290	207	108	38	-
American Indian 1/, both sexes	11	11	9	18	11	20	20	14	17	1	1	-
Male	6	7	5	9	5	10	7	6	4	-	-	-
Female	5	4	4	9	6	10	13	8	13	1	1	-
Asian or Pacific Islander 2/, both sexes	8	6	9	7	26	18	19	22	16	2	2	-
Male	6	4	2	3	15	8	6	11	5	1	1	-
Female	2	2	7	4	11	10	13	11	11	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Minnesota												
All races, both sexes	303	380	436	497	752	1,196	1,803	2,155	1,788	768	184	-
Male	238	280	289	323	450	646	926	923	571	179	32	-
Female	65	100	147	174	302	550	877	1,232	1,217	589	152	-
White, both sexes	270	340	394	458	713	1,162	1,771	2,123	1,777	761	182	-
Male	217	251	266	300	424	632	912	910	570	177	32	-
Female	53	89	128	158	289	530	859	1,213	1,207	584	150	-
Black, both sexes	18	23	22	13	18	10	14	16	6	5	2	-
Male	12	15	11	3	12	5	6	6	-	2	-	-
Female	6	8	11	10	6	5	8	10	6	3	2	-
American Indian 1/, both sexes	10	11	13	13	10	13	6	8	1	1	-	-
Male	5	8	7	9	7	5	5	1	-	-	-	-
Female	5	3	6	4	3	8	1	7	1	1	-	-
Asian or Pacific Islander 2/, both sexes	5	6	7	13	11	11	12	8	4	1	-	-
Male	4	6	5	11	7	4	3	6	1	-	-	-
Female	1	-	2	2	4	7	9	2	3	1	-	-
Mississippi												
All races, both sexes	507	619	649	814	964	1,340	1,483	1,505	1,109	492	102	-
Male	341	396	415	509	529	670	596	503	298	95	15	-
Female	166	223	234	305	435	670	887	1,002	811	397	87	-
White, both sexes	245	328	384	523	627	972	1,090	1,152	835	342	53	-
Male	168	208	248	345	362	501	464	381	215	65	8	-
Female	77	120	136	178	265	471	626	771	620	277	45	-
Black, both sexes	262	283	263	287	335	364	390	350	274	150	49	-
Male	173	182	165	162	167	166	130	122	83	30	7	-
Female	89	101	98	125	168	198	260	228	191	120	42	-
American Indian 1/, both sexes	-	4	1	1	1	1	1	1	-	-	-	-
Male	-	3	1	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	1	3	1	3	2	2	-	-	-	-
Male	-	3	1	1	-	3	1	-	-	-	-	-
Female	-	1	-	2	1	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Missouri												
All races, both sexes	707	873	978	1,238	1,598	2,269	3,226	3,445	2,615	1,092	241	-
Male	474	600	601	777	912	1,177	1,444	1,240	763	223	42	-
Female	233	273	377	461	686	1,092	1,782	2,205	1,852	869	199	-
White, both sexes	550	696	825	1,054	1,385	2,032	2,958	3,234	2,478	1,028	226	-
Male	375	489	517	680	800	1,066	1,355	1,184	724	214	40	-
Female	175	207	308	374	585	966	1,603	2,050	1,754	814	186	-
Black, both sexes	153	169	144	178	201	221	257	200	129	63	14	-
Male	96	107	79	95	105	102	88	54	34	9	2	-
Female	57	62	65	83	96	119	169	146	95	54	12	-
American Indian 1/, both sexes	2	2	5	-	4	5	6	6	2	-	-	-
Male	2	-	3	-	4	4	1	-	1	-	-	-
Female	-	2	2	-	-	1	5	6	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	6	4	6	8	11	5	5	6	1	1	-
Male	1	4	2	2	3	5	-	2	4	-	-	-
Female	1	2	2	4	5	6	5	3	2	1	1	-
Montana												
All races, both sexes	86	97	118	134	184	287	409	495	394	156	19	-
Male	63	66	84	88	109	164	215	209	143	44	2	-
Female	23	31	34	46	75	123	194	286	251	112	17	-
White, both sexes	76	92	109	125	170	278	401	488	386	156	18	-
Male	55	62	79	82	103	161	209	207	142	44	2	-
Female	21	30	30	43	67	117	192	281	244	112	16	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	4	9	9	13	9	8	7	7	-	1	-
Male	7	3	5	6	6	3	6	2	1	-	-	-
Female	2	1	4	3	7	6	2	5	6	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Nebraska												
All races, both sexes	134	152	181	213	336	532	832	904	773	431	88	-
Male	99	113	117	130	187	288	416	313	236	100	13	-
Female	35	39	64	83	149	244	416	591	537	331	75	-
White, both sexes	117	141	159	200	327	513	814	887	758	427	86	-
Male	88	106	100	123	183	281	408	308	233	100	13	-
Female	29	35	59	77	144	232	406	579	525	327	73	-
Black, both sexes	16	10	15	8	8	16	13	12	14	3	2	-
Male	10	7	12	5	3	7	7	3	3	-	-	-
Female	6	3	3	3	5	9	6	9	11	3	2	-
American Indian 1/, both sexes	1	1	6	2	1	3	1	-	-	1	-	-
Male	1	-	4	1	1	-	-	-	-	-	-	-
Female	-	1	2	1	-	3	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	-	-	4	5	1	-	-	-
Male	-	-	1	1	-	-	1	2	-	-	-	-
Female	-	-	-	2	-	-	3	3	1	-	-	-
Nevada												
All races, both sexes	304	432	503	580	702	775	1,001	857	511	165	28	-
Male	218	309	340	393	429	446	498	367	162	36	3	-
Female	86	123	163	187	273	329	503	490	349	129	25	-
White, both sexes	246	364	432	490	605	681	918	802	485	152	23	-
Male	181	267	298	343	372	396	463	341	157	33	3	-
Female	65	97	134	147	233	285	455	461	328	119	20	-
Black, both sexes	44	46	43	52	65	60	47	31	15	7	4	-
Male	25	28	25	35	40	28	20	14	2	1	-	-
Female	19	18	18	17	25	32	27	17	13	6	4	-
American Indian 1/, both sexes	3	7	7	8	6	3	6	6	4	3	1	-
Male	2	2	3	4	3	1	3	3	2	-	-	-
Female	1	5	4	4	3	2	3	3	2	3	1	-
Asian or Pacific Islander 2/, both sexes	11	15	21	30	26	31	30	18	7	3	-	-
Male	10	12	14	11	14	21	12	9	1	2	-	-
Female	1	3	7	19	12	10	18	9	6	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
New Hampshire												
All races, both sexes	93	147	139	153	251	394	540	624	491	215	45	-
Male	72	109	97	113	151	219	238	247	142	46	6	-
Female	21	38	42	40	100	175	302	377	349	169	39	-
White, both sexes	90	144	137	151	250	392	538	620	491	215	45	-
Male	70	107	96	112	150	218	238	245	142	46	6	-
Female	20	37	41	39	100	174	300	375	349	169	39	-
Black, both sexes	-	2	1	1	1	-	2	-	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	1	-	2	-	3	-	-	-	-
Male	2	1	1	-	-	1	-	1	-	-	-	-
Female	1	-	-	1	-	1	-	2	-	-	-	-
New Jersey												
All races, both sexes	689	902	1,091	1,296	1,878	3,056	4,416	4,824	3,647	1,505	320	2
Male	489	620	719	818	1,095	1,568	2,019	1,906	1,112	323	39	2
Female	200	282	372	478	783	1,488	2,397	2,918	2,535	1,182	281	-
White, both sexes	496	666	814	1,003	1,501	2,684	3,947	4,483	3,401	1,409	293	2
Male	372	467	554	645	891	1,393	1,826	1,785	1,044	306	37	2
Female	124	199	260	358	610	1,291	2,121	2,698	2,357	1,103	256	-
Black, both sexes	174	207	237	241	331	315	395	281	200	86	23	-
Male	105	131	136	140	179	150	164	99	52	16	1	-
Female	69	76	101	101	152	165	231	182	148	70	22	-
American Indian 1/, both sexes	1	3	1	2	2	3	6	2	3	-	-	-
Male	-	3	-	-	-	2	1	1	1	-	-	-
Female	1	-	1	2	2	1	5	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	18	26	39	50	44	54	68	58	43	10	4	-
Male	12	19	29	33	25	23	28	21	15	1	1	-
Female	6	7	10	17	19	31	40	37	28	9	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
New Mexico												
All races, both sexes	148	204	223	279	398	576	714	769	600	262	48	-
Male	103	140	148	179	246	320	348	310	191	51	8	-
Female	45	64	75	100	152	256	366	459	409	211	40	-
White, both sexes	132	186	204	251	361	548	686	740	575	249	42	-
Male	94	129	136	161	228	308	334	303	180	48	6	-
Female	38	57	68	90	133	240	352	437	395	201	36	-
Black, both sexes	5	8	3	12	14	12	10	9	6	7	1	-
Male	3	4	1	9	6	5	3	1	2	1	-	-
Female	2	4	2	3	8	7	7	8	4	6	1	-
American Indian 1/, both sexes	9	7	16	13	20	14	13	19	18	6	5	-
Male	5	5	11	8	11	7	8	5	8	2	2	-
Female	4	2	5	5	9	7	5	14	10	4	3	-
Asian or Pacific Islander 2/, both sexes	2	3	-	3	3	2	5	1	1	-	-	-
Male	1	2	-	1	1	-	3	1	1	-	-	-
Female	1	1	-	2	2	2	2	-	-	-	-	-
New York												
All races, both sexes	1,529	2,278	2,772	3,327	4,649	7,336	10,324	11,739	9,111	4,070	1,001	1
Male	1,062	1,563	1,765	2,044	2,583	3,811	4,692	4,515	2,698	867	143	1
Female	467	715	1,007	1,283	2,066	3,525	5,632	7,224	6,413	3,203	858	-
White, both sexes	1,042	1,679	2,037	2,451	3,642	6,144	8,930	10,515	8,345	3,693	887	1
Male	767	1,205	1,327	1,547	2,092	3,278	4,152	4,095	2,491	797	131	1
Female	275	474	710	904	1,550	2,866	4,778	6,420	5,854	2,896	756	-
Black, both sexes	437	532	649	772	874	1,015	1,146	1,012	624	320	106	-
Male	256	306	380	435	416	442	434	318	144	55	11	-
Female	181	226	269	337	458	573	712	694	480	265	95	-
American Indian 1/, both sexes	1	2	6	8	8	9	8	7	3	4	-	-
Male	1	2	5	2	2	2	2	2	-	-	-	-
Female	-	-	1	6	6	7	6	5	3	4	-	-
Asian or Pacific Islander 2/, both sexes	49	65	80	96	125	168	240	205	139	53	8	-
Male	38	50	53	60	73	89	104	100	63	15	1	-
Female	11	15	27	36	52	79	136	105	76	38	7	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
North Carolina												
All races, both sexes	962	1,254	1,513	1,681	2,165	3,003	3,886	3,887	2,687	937	222	1
Male	675	881	1,005	1,063	1,294	1,558	1,793	1,426	721	179	28	1
Female	287	373	508	618	871	1,445	2,093	2,461	1,966	758	194	-
White, both sexes	591	837	1,052	1,170	1,634	2,385	3,155	3,296	2,293	785	165	1
Male	433	606	727	739	998	1,271	1,510	1,238	636	154	18	1
Female	158	231	325	431	636	1,114	1,645	2,058	1,657	631	147	-
Black, both sexes	350	389	434	479	494	586	683	564	376	145	56	-
Male	222	251	261	298	276	273	259	184	76	25	10	-
Female	128	138	173	181	218	313	424	380	300	120	46	-
American Indian 1/, both sexes	16	23	18	24	22	24	34	17	12	6	-	-
Male	15	20	13	20	13	10	14	1	6	-	-	-
Female	1	3	5	4	9	14	20	16	6	6	-	-
Asian or Pacific Islander 2/, both sexes	5	5	9	8	15	8	14	10	6	1	1	-
Male	5	4	4	6	7	4	10	3	3	-	-	-
Female	-	1	5	2	8	4	4	7	3	1	1	-
North Dakota												
All races, both sexes	62	59	75	110	145	217	351	429	378	164	40	-
Male	45	48	57	72	87	127	188	183	113	44	5	-
Female	17	11	18	38	58	90	163	246	265	120	35	-
White, both sexes	55	54	68	102	134	210	345	425	375	164	40	-
Male	40	44	54	67	81	123	186	181	113	44	5	-
Female	15	10	14	35	53	87	159	244	262	120	35	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	7	5	6	8	11	7	5	4	2	-	-	-
Male	5	4	2	5	6	4	1	2	-	-	-	-
Female	2	1	4	3	5	3	4	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Ohio												
All races, both sexes	1,265	1,711	1,788	2,244	3,025	4,699	6,434	6,749	4,765	1,785	345	-
Male	899	1,176	1,163	1,356	1,690	2,460	2,916	2,577	1,383	365	48	-
Female	366	535	625	888	1,335	2,239	3,518	4,172	3,382	1,420	297	-
White, both sexes	974	1,369	1,509	1,900	2,598	4,165	5,882	6,322	4,475	1,663	303	-
Male	702	954	1,004	1,151	1,450	2,206	2,671	2,427	1,304	338	41	-
Female	272	415	505	749	1,148	1,959	3,211	3,895	3,171	1,325	262	-
Black, both sexes	284	330	267	329	415	519	532	416	280	117	42	-
Male	195	212	152	195	236	248	235	145	74	25	7	-
Female	89	118	115	134	179	271	297	271	206	92	35	-
American Indian 1/, both sexes	2	3	3	5	-	3	3	2	-	-	-	-
Male	1	2	2	2	-	1	1	1	-	-	-	-
Female	1	1	1	3	-	2	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	9	9	10	12	12	17	9	10	5	-	-
Male	1	8	5	8	4	5	9	4	5	2	-	-
Female	4	1	4	2	8	7	8	5	5	3	-	-
Oklahoma												
All races, both sexes	492	603	786	841	1,179	1,528	2,085	2,097	1,564	654	139	1
Male	335	420	498	527	681	835	980	796	464	133	25	1
Female	157	183	288	314	498	693	1,105	1,301	1,100	521	114	-
White, both sexes	386	481	667	692	1,007	1,365	1,895	1,947	1,437	606	126	1
Male	269	338	427	431	587	752	894	740	432	123	23	1
Female	117	143	240	261	420	613	1,001	1,207	1,005	483	103	-
Black, both sexes	68	72	63	74	84	86	96	69	60	36	7	-
Male	41	45	35	44	47	42	42	22	19	9	2	-
Female	27	27	28	30	37	44	54	47	41	27	5	-
American Indian 1/, both sexes	34	44	52	67	80	70	89	79	63	11	4	-
Male	23	33	35	45	45	39	42	33	13	1	-	-
Female	11	11	17	22	35	31	47	46	50	10	4	-
Asian or Pacific Islander 2/, both sexes	4	6	4	8	8	7	5	2	4	1	2	-
Male	2	4	1	7	2	2	2	1	-	-	-	-
Female	2	2	3	1	6	5	3	1	4	1	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Oregon												
All races, both sexes	291	372	414	531	729	1,108	1,669	1,885	1,420	580	110	-
Male	209	270	289	350	449	606	812	828	484	168	21	-
Female	82	102	125	181	280	502	857	1,057	936	412	89	-
White, both sexes	274	351	385	506	701	1,073	1,628	1,842	1,400	574	108	-
Male	198	260	275	335	432	591	799	806	478	166	20	-
Female	76	91	110	171	269	482	829	1,036	922	408	88	-
Black, both sexes	10	14	13	2	10	11	14	12	4	3	-	-
Male	7	9	6	1	7	7	5	5	2	1	-	-
Female	3	5	7	1	3	4	9	7	2	2	-	-
American Indian 1/, both sexes	5	1	10	11	5	7	4	8	5	-	-	-
Male	3	-	4	4	4	3	1	6	2	-	-	-
Female	2	1	6	7	1	4	3	2	3	-	-	-
Asian or Pacific Islander 2/, both sexes	2	6	6	12	13	17	23	23	11	3	2	-
Male	1	1	4	10	6	5	7	11	2	1	1	-
Female	1	5	2	2	7	12	16	12	9	2	1	-
Pennsylvania												
All races, both sexes	1,144	1,637	2,032	2,354	3,421	5,591	7,948	8,737	6,264	2,473	460	3
Male	791	1,134	1,373	1,479	1,931	2,900	3,692	3,302	1,822	517	58	2
Female	353	503	659	875	1,490	2,691	4,256	5,435	4,442	1,956	402	1
White, both sexes	899	1,331	1,716	2,013	3,023	5,081	7,426	8,307	5,983	2,356	424	2
Male	646	930	1,173	1,285	1,720	2,659	3,482	3,154	1,745	495	55	1
Female	253	401	543	728	1,303	2,422	3,944	5,153	4,238	1,861	369	1
Black, both sexes	235	294	304	314	381	489	501	402	270	112	35	1
Male	140	197	191	177	199	229	201	138	76	21	3	1
Female	95	97	113	137	182	260	300	264	194	91	32	-
American Indian 1/, both sexes	-	2	1	1	2	2	2	-	-	1	1	-
Male	-	2	1	1	1	2	2	-	-	1	-	-
Female	-	-	-	-	1	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	10	10	11	26	15	19	19	28	11	4	-	-
Male	5	5	8	16	11	10	7	10	1	-	-	-
Female	5	5	3	10	4	9	12	18	10	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Rhode Island												
All races, both sexes	85	90	121	167	211	410	588	708	572	241	57	-
Male	65	67	74	108	121	217	282	269	161	44	11	-
Female	20	23	47	59	90	193	306	439	411	197	46	-
White, both sexes	78	84	115	154	200	393	574	703	565	236	56	-
Male	61	64	71	101	116	208	277	267	160	43	11	-
Female	17	20	44	53	84	185	297	436	405	193	45	-
Black, both sexes	3	5	3	9	9	10	11	5	4	3	1	-
Male	2	2	1	4	4	6	4	2	-	-	-	-
Female	1	3	2	5	5	4	7	3	4	3	1	-
American Indian 1/, both sexes	3	-	-	3	1	2	1	-	2	1	-	-
Male	2	-	-	2	-	1	-	-	-	-	-	-
Female	1	-	-	1	1	1	1	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	1	1	5	2	-	1	1	-	-
Male	-	1	2	1	1	2	1	-	1	1	-	-
Female	1	-	1	-	-	3	1	-	-	-	-	-
South Carolina												
All races, both sexes	566	801	893	966	1,174	1,529	1,933	1,817	1,113	398	88	5
Male	391	533	573	612	668	851	878	684	335	81	8	5
Female	175	268	320	354	506	678	1,055	1,133	778	317	80	-
White, both sexes	333	479	563	647	816	1,125	1,460	1,424	871	310	47	2
Male	235	324	372	433	480	660	683	562	270	61	4	2
Female	98	155	191	214	336	465	777	862	601	249	43	-
Black, both sexes	230	314	327	313	353	395	464	386	239	87	41	3
Male	156	202	199	176	185	188	194	118	64	19	4	3
Female	74	112	128	137	168	207	270	268	175	68	37	-
American Indian 1/, both sexes	-	5	2	3	2	4	6	1	2	1	-	-
Male	-	4	1	-	2	2	1	-	1	1	-	-
Female	-	1	1	3	-	2	5	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	1	3	3	5	3	6	1	-	-	-
Male	-	3	1	3	1	1	-	4	-	-	-	-
Female	3	-	-	-	2	4	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
South Dakota												
All races, both sexes	58	81	99	112	145	272	390	450	418	211	58	-
Male	41	57	70	75	85	160	204	195	135	49	13	-
Female	17	24	29	37	60	112	186	255	283	162	45	-
White, both sexes	50	68	91	102	132	259	377	445	414	208	57	-
Male	33	49	68	69	81	149	199	191	135	49	12	-
Female	17	19	23	33	51	110	178	254	279	159	45	-
Black, both sexes	-	-	-	1	-	3	2	1	1	-	-	-
Male	-	-	-	1	-	3	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	8	13	7	9	13	10	11	4	3	3	-	-
Male	8	8	1	5	4	8	4	3	-	-	-	-
Female	-	5	6	4	9	2	7	1	3	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	803	1,114	1,311	1,385	1,836	2,369	3,139	2,968	2,117	815	169	1
Male	533	770	856	853	1,061	1,210	1,353	1,137	617	165	28	1
Female	270	344	455	532	775	1,159	1,786	1,831	1,500	650	141	-
White, both sexes	563	858	1,054	1,123	1,542	2,033	2,775	2,677	1,885	724	129	1
Male	386	620	691	699	905	1,054	1,215	1,037	554	148	25	1
Female	177	238	363	424	637	979	1,560	1,640	1,331	576	104	-
Black, both sexes	237	249	253	255	286	332	359	284	222	91	40	-
Male	144	147	162	150	151	154	134	97	61	17	3	-
Female	93	102	91	105	135	178	225	187	161	74	37	-
American Indian 1/, both sexes	-	2	1	2	1	2	1	-	2	-	-	-
Male	-	1	1	2	1	1	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	3	5	7	2	4	7	8	-	-	-
Male	3	2	2	2	4	1	3	3	2	-	-	-
Female	-	3	1	3	3	1	1	4	6	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Texas												
All races, both sexes	2,147	2,805	3,104	3,557	4,556	6,573	8,359	8,215	5,941	2,442	484	2
Male	1,491	1,869	1,993	2,139	2,609	3,444	3,800	3,146	1,843	541	85	-
Female	656	936	1,111	1,418	1,947	3,129	4,559	5,069	4,098	1,901	399	2
White, both sexes	1,615	2,195	2,532	2,933	3,870	5,735	7,501	7,534	5,440	2,196	410	2
Male	1,150	1,488	1,668	1,784	2,220	3,050	3,438	2,911	1,697	489	67	-
Female	465	707	864	1,149	1,650	2,685	4,063	4,623	3,743	1,707	343	2
Black, both sexes	497	550	528	567	628	752	787	620	464	235	73	-
Male	316	346	296	320	362	345	342	209	128	49	18	-
Female	181	204	232	247	266	407	445	411	336	186	55	-
American Indian 1/, both sexes	6	11	6	10	5	6	11	8	4	3	-	-
Male	4	5	4	7	3	2	3	3	1	-	-	-
Female	2	6	2	3	2	4	8	5	3	3	-	-
Asian or Pacific Islander 2/, both sexes	29	49	38	47	53	80	60	53	33	8	1	-
Male	21	30	25	28	24	47	17	23	17	3	-	-
Female	8	19	13	19	29	33	43	30	16	5	1	-
Utah												
All races, both sexes	112	150	187	234	316	470	712	812	531	201	21	1
Male	85	102	126	154	176	260	368	341	176	60	3	1
Female	27	48	61	80	140	210	344	471	355	141	18	-
White, both sexes	106	141	184	225	300	453	702	801	527	201	19	1
Male	80	99	124	149	168	250	363	339	174	60	3	1
Female	26	42	60	76	132	203	339	462	353	141	16	-
Black, both sexes	3	2	-	1	2	5	5	-	1	-	1	-
Male	2	2	-	-	1	4	3	-	-	-	-	-
Female	1	-	-	1	1	1	2	-	1	-	1	-
American Indian 1/, both sexes	1	2	1	3	2	4	-	1	-	-	1	-
Male	1	-	1	2	-	1	-	-	-	-	-	-
Female	-	2	-	1	2	3	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	5	2	5	12	8	5	10	3	-	-	-
Male	2	1	1	3	7	5	2	2	2	-	-	-
Female	-	4	1	2	5	3	3	8	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Vermont												
All races, both sexes	40	69	64	88	127	215	269	316	241	99	28	-
Male	25	51	41	63	72	110	135	126	77	22	5	-
Female	15	18	23	25	55	105	134	190	164	77	23	-
White, both sexes	40	68	63	88	126	215	268	316	241	99	28	-
Male	25	50	41	63	71	110	135	126	77	22	5	-
Female	15	18	22	25	55	105	133	190	164	77	23	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	694	942	1,070	1,202	1,715	2,372	3,093	3,268	2,262	897	186	-
Male	477	643	710	765	962	1,312	1,377	1,228	668	166	21	-
Female	217	299	360	437	753	1,060	1,716	2,040	1,594	731	165	-
White, both sexes	464	625	768	844	1,272	1,901	2,519	2,752	1,957	762	139	-
Male	326	438	523	540	734	1,072	1,151	1,038	582	143	13	-
Female	138	187	245	304	538	829	1,368	1,714	1,375	619	126	-
Black, both sexes	220	299	285	333	416	431	537	482	292	127	46	-
Male	143	189	173	207	215	219	205	179	82	21	8	-
Female	77	110	112	126	201	212	332	303	210	106	38	-
American Indian 1/, both sexes	-	1	1	1	1	1	4	3	2	1	-	-
Male	-	1	1	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	3	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	10	17	16	24	26	39	33	31	11	7	1	-
Male	8	15	13	17	13	20	20	10	4	2	-	-
Female	2	2	3	7	13	19	13	21	7	5	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Washington												
All races, both sexes	436	578	725	852	1,113	1,701	2,495	2,755	2,212	888	186	-
Male	314	399	477	534	666	920	1,180	1,145	731	252	29	-
Female	122	179	248	318	447	781	1,315	1,610	1,481	636	157	-
White, both sexes	379	507	631	754	1,021	1,569	2,352	2,646	2,136	863	183	-
Male	280	354	416	477	616	855	1,123	1,093	699	243	28	-
Female	99	153	215	277	405	714	1,229	1,553	1,437	620	155	-
Black, both sexes	26	37	43	40	29	43	42	25	20	7	-	-
Male	19	23	29	26	14	20	15	9	8	2	-	-
Female	7	14	14	14	15	23	27	16	12	5	-	-
American Indian 1/, both sexes	9	11	12	12	22	18	25	13	12	-	1	-
Male	4	5	5	6	13	8	8	6	4	-	1	-
Female	5	6	7	6	9	10	17	7	8	-	-	-
Asian or Pacific Islander 2/, both sexes	22	23	39	46	41	71	76	71	44	18	2	-
Male	11	17	27	25	23	37	34	37	20	7	-	-
Female	11	6	12	21	18	34	42	34	24	11	2	-
West Virginia												
All races, both sexes	251	320	358	516	647	955	1,218	1,127	830	333	68	-
Male	188	222	224	332	366	499	547	429	230	56	6	-
Female	63	98	134	184	281	456	671	698	600	277	62	-
White, both sexes	237	304	344	506	623	925	1,175	1,104	807	322	62	-
Male	177	209	216	324	352	482	530	424	226	53	5	-
Female	60	95	128	182	271	443	645	680	581	269	57	-
Black, both sexes	13	15	14	9	24	30	41	23	23	10	6	-
Male	10	12	8	7	14	17	16	5	4	3	1	-
Female	3	3	6	2	10	13	25	18	19	7	5	-
American Indian 1/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Wisconsin												
All races, both sexes	429	556	641	787	1,182	1,806	2,643	3,012	2,466	1,044	239	-
Male	300	394	436	528	703	998	1,271	1,289	753	231	30	-
Female	129	162	205	259	479	808	1,372	1,723	1,713	813	209	-
White, both sexes	366	479	579	726	1,107	1,721	2,569	2,956	2,437	1,030	238	-
Male	267	350	403	496	663	962	1,243	1,270	747	228	30	-
Female	99	129	176	230	444	759	1,326	1,686	1,690	802	208	-
Black, both sexes	55	61	53	46	59	70	52	39	24	11	-	-
Male	28	31	29	22	31	31	24	13	5	2	-	-
Female	27	30	24	24	28	39	28	26	19	9	-	-
American Indian 1/, both sexes	7	11	6	5	10	7	16	5	2	2	-	-
Male	4	8	2	4	6	3	2	1	-	1	-	-
Female	3	3	4	1	4	4	14	4	2	1	-	-
Asian or Pacific Islander 2/, both sexes	1	5	3	10	6	8	6	12	3	1	1	-
Male	1	5	2	6	3	2	2	5	1	-	-	-
Female	-	-	1	4	3	6	4	7	2	1	1	-
Wyoming												
All races, both sexes	41	61	74	101	113	163	209	220	174	72	12	-
Male	30	40	47	63	65	101	105	88	58	11	3	-
Female	11	21	27	38	48	62	104	132	116	61	9	-
White, both sexes	40	57	72	100	108	162	207	219	173	71	12	-
Male	30	38	45	62	62	100	104	88	58	10	3	-
Female	10	19	27	38	46	62	103	131	115	61	9	-
Black, both sexes	1	1	1	-	2	-	1	1	-	-	-	-
Male	-	-	1	-	2	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	3	1	1	2	1	1	-	1	1	-	-
Male	-	2	1	1	1	1	1	-	-	1	-	-
Female	-	1	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
United States												
All races, both sexes	3,257	3	-	1	6	3	4	9	22	21	45	84
Male	1,008	2	-	1	1	2	3	4	13	11	17	41
Female	2,249	1	-	-	5	1	1	5	9	10	28	43
White, both sexes	2,903	3	-	-	2	1	2	7	13	13	29	61
Male	887	2	-	-	-	1	2	3	8	8	14	32
Female	2,016	1	-	-	2	-	-	4	5	5	15	29
Black, both sexes	249	-	-	-	3	2	-	1	6	7	13	22
Male	87	-	-	-	1	1	-	1	3	3	3	9
Female	162	-	-	-	2	1	-	-	3	4	10	13
American Indian 1/, both sexes	20	-	-	1	-	-	2	-	-	-	-	-
Male	6	-	-	1	-	-	1	-	-	-	-	-
Female	14	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	85	-	-	-	1	-	-	1	3	1	3	1
Male	28	-	-	-	-	-	-	-	2	-	-	-
Female	57	-	-	-	1	-	-	1	1	1	3	1
Alabama												
All races, both sexes	50	-	-	-	-	-	-	-	1	1	1	4
Male	15	-	-	-	-	-	-	-	-	1	-	2
Female	35	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	41	-	-	-	-	-	-	-	1	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	2
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Alaska												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	59	-	-	-	-	-	-	1	-	-	-	1
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	-	-	-	-	-	-	1	-	-	-	1
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Arkansas												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	394	1	-	1	-	-	-	1	4	1	9	14
Male	137	-	-	1	-	-	-	-	1	1	3	3
Female	257	1	-	-	-	-	-	1	3	-	6	11
White, both sexes	333	1	-	-	-	-	-	1	2	1	6	12
Male	116	-	-	-	-	-	-	-	-	1	2	3
Female	217	1	-	-	-	-	-	1	2	-	4	9
Black, both sexes	20	-	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	1	-	1	1
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Colorado												
All races, both sexes	60	-	-	-	-	-	1	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	56	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	5
Male	26	-	-	-	-	-	-	-	-	-	-	2
Female	24	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	3
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Florida												
All races, both sexes	208	-	-	-	1	-	1	-	2	1	2	7
Male	74	-	-	-	-	-	1	-	2	-	1	4
Female	134	-	-	-	1	-	-	-	-	1	1	3
White, both sexes	185	-	-	-	-	-	1	-	1	1	1	6
Male	67	-	-	-	-	-	1	-	1	-	1	4
Female	118	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	21	-	-	-	1	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	55	-	-	-	-	1	-	1	-	2	2	1
Male	21	-	-	-	-	-	-	1	-	-	1	1
Female	34	-	-	-	-	1	-	-	-	2	1	-
White, both sexes	40	-	-	-	-	-	-	1	-	1	2	1
Male	16	-	-	-	-	-	-	1	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	14	-	-	-	-	1	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hawaii												
All races, both sexes	22	-	-	-	-	-	-	1	1	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	1	-	2	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	1	1	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	1	-	2	-
Idaho												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Illinois												
All races, both sexes	94	-	-	-	-	-	1	-	1	2	-	3
Male	35	-	-	-	-	-	1	-	1	1	-	2
Female	59	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	85	-	-	-	-	-	1	-	1	1	-	2
Male	32	-	-	-	-	-	1	-	1	1	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Iowa												
All races, both sexes	69	-	-	-	-	-	-	-	-	2	1	1
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	56	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	68	-	-	-	-	-	-	-	-	2	1	1
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	55	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	25	1	-	-	-	-	-	1	2	-	-	-
Male	7	1	-	-	-	-	-	1	1	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	21	1	-	-	-	-	-	-	2	-	-	-
Male	4	1	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Kentucky												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	30	-	-	-	-	1	-	-	3	-	2	1
Male	11	-	-	-	-	1	-	-	2	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	19	-	-	-	-	-	-	-	1	-	1	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	11	-	-	-	-	1	-	-	2	-	1	-
Male	5	-	-	-	-	1	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Maine												
All races, both sexes	25	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	24	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	50	-	-	-	-	-	-	-	-	1	1	5
Male	13	-	-	-	-	-	-	-	-	1	-	2
Female	37	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	1	1	3
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Massachusetts												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	95	-	-	-	-	-	-	-	-	1	-	1
Male	35	-	-	-	-	-	-	-	-	1	-	1
Female	60	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	84	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Minnesota												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	21	-	-	-	-	-	-	-	1	-	-	2
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	13	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Missouri												
All races, both sexes	71	-	-	-	-	-	-	-	-	1	1	1
Male	25	-	-	-	-	-	-	-	-	1	1	1
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	1	1	1
Male	25	-	-	-	-	-	-	-	-	1	1	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Nebraska												
All races, both sexes	29	-	-	-	-	-	1	-	1	-	-	-
Male	5	-	-	-	-	-	1	-	1	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	20	-	-	-	1	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Hampshire												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	121	-	-	-	-	-	-	-	1	1	1	1
Male	31	-	-	-	-	-	-	-	1	-	-	1
Female	90	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	109	-	-	-	-	-	-	-	1	-	1	1
Male	29	-	-	-	-	-	-	-	1	-	-	1
Female	80	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Mexico												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	180	-	-	-	1	1	-	1	1	-	5	5
Male	58	-	-	-	1	1	-	-	1	-	3	5
Female	122	-	-	-	-	-	-	1	-	-	2	-
White, both sexes	158	-	-	-	-	1	-	1	-	-	5	1
Male	46	-	-	-	-	1	-	-	-	-	3	1
Female	112	-	-	-	-	-	-	1	-	-	2	-
Black, both sexes	18	-	-	-	1	-	-	-	-	-	-	4
Male	10	-	-	-	1	-	-	-	-	-	-	4
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
North Carolina												
All races, both sexes	97	-	-	-	1	-	-	-	-	2	1	3
Male	30	-	-	-	-	-	-	-	-	1	-	-
Female	67	-	-	-	1	-	-	-	-	1	1	3
White, both sexes	78	-	-	-	-	-	-	-	-	2	-	2
Male	23	-	-	-	-	-	-	-	-	1	-	-
Female	55	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	18	-	-	-	1	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Ohio												
All races, both sexes	160	-	-	-	-	-	-	-	1	1	2	7
Male	44	-	-	-	-	-	-	-	1	-	2	4
Female	116	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	146	-	-	-	-	-	-	-	-	-	2	5
Male	39	-	-	-	-	-	-	-	-	-	2	3
Female	107	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	14	-	-	-	-	-	-	-	1	1	-	2
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	9	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	38	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Oregon												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	75	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	171	-	-	-	-	-	-	-	1	-	1	2
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	131	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	162	-	-	-	-	-	-	-	-	-	1	1
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	124	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Rhode Island												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	3	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
South Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	73	-	-	-	1	-	-	-	1	1	2	3
Male	32	-	-	-	-	-	-	-	1	1	2	3
Female	41	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	1	-	-	-	1	1	-	3
Male	30	-	-	-	-	-	-	-	1	1	-	3
Female	38	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Texas												
All races, both sexes	118	1	-	-	-	-	-	1	-	2	-	2
Male	26	1	-	-	-	-	-	-	-	-	-	1
Female	92	-	-	-	-	-	-	1	-	2	-	1
White, both sexes	106	1	-	-	-	-	-	1	-	1	-	1
Male	23	1	-	-	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Utah												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Vermont												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	68	-	-	-	1	-	-	1	-	2	4	2
Male	20	-	-	-	-	-	-	-	-	2	-	1
Female	48	-	-	-	1	-	-	1	-	-	4	1
White, both sexes	56	-	-	-	1	-	-	1	-	2	1	1
Male	20	-	-	-	-	-	-	-	-	2	-	1
Female	36	-	-	-	1	-	-	1	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	3	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Washington												
All races, both sexes	103	-	-	-	-	-	-	-	-	-	2	1
Male	38	-	-	-	-	-	-	-	-	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	98	-	-	-	-	-	-	-	-	-	2	1
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	62	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Wisconsin												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	63	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
United States												
All races, both sexes	92	157	195	257	351	502	621	512	258	96	18	-
Male	40	61	74	92	116	159	168	134	51	17	1	-
Female	52	96	121	165	235	343	453	378	207	79	17	-
White, both sexes	63	125	152	219	309	460	586	500	251	89	18	-
Male	27	46	59	79	101	146	163	131	49	15	1	-
Female	36	79	93	140	208	314	423	369	202	74	17	-
Black, both sexes	17	22	32	28	25	28	26	6	7	4	-	-
Male	7	10	13	10	9	10	4	-	2	1	-	-
Female	10	12	19	18	16	18	22	6	5	3	-	-
American Indian 1/, both sexes	3	3	3	1	1	4	2	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	3	1	1	4	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	7	8	9	16	10	7	6	-	3	-	-
Male	4	3	2	3	6	3	1	3	-	1	-	-
Female	5	4	6	6	10	7	6	3	-	2	-	-
Alabama												
All races, both sexes	4	2	5	5	7	9	7	2	2	-	-	-
Male	2	1	-	1	2	5	1	-	-	-	-	-
Female	2	1	5	4	5	4	6	2	2	-	-	-
White, both sexes	2	2	5	4	6	7	7	2	2	-	-	-
Male	1	1	-	-	2	4	1	-	-	-	-	-
Female	1	1	5	4	4	3	6	2	2	-	-	-
Black, both sexes	1	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Alaska												
All races, both sexes	2	1	-	1	-	2	1	-	1	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	1	-	1	-	-	-
White, both sexes	1	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	4	4	7	8	15	8	6	3	-	-	-
Male	2	1	2	2	2	3	1	1	-	-	-	-
Female	-	3	2	5	6	12	7	5	3	-	-	-
White, both sexes	2	4	4	6	7	15	7	6	3	-	-	-
Male	2	1	2	2	2	3	1	1	-	-	-	-
Female	-	3	2	4	5	12	6	5	3	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Arkansas												
All races, both sexes	1	-	1	1	3	2	5	5	2	-	-	-
Male	1	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	1	1	3	2	4	2	2	-	-	-
White, both sexes	-	-	1	1	3	2	5	5	2	-	-	-
Male	-	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	1	1	3	2	4	2	2	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12	13	27	37	41	55	79	61	29	9	-	-
Male	4	6	11	16	15	16	27	22	9	2	-	-
Female	8	7	16	21	26	39	52	39	20	7	-	-
White, both sexes	6	9	20	26	32	47	76	58	29	7	-	-
Male	3	4	7	14	12	13	27	19	9	2	-	-
Female	3	5	13	12	20	34	49	39	20	5	-	-
Black, both sexes	1	-	2	5	3	3	1	-	-	1	-	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	1	-	-	4	2	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	4	5	6	6	4	2	3	-	1	-	-
Male	1	2	2	1	2	1	-	3	-	-	-	-
Female	4	2	3	5	4	3	2	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Colorado												
All races, both sexes	1	3	4	7	10	11	10	6	4	2	-	-
Male	1	1	1	-	2	5	3	2	-	1	-	-
Female	-	2	3	7	8	6	7	4	4	1	-	-
White, both sexes	1	3	3	7	9	11	9	6	4	2	-	-
Male	1	1	1	-	2	5	2	2	-	1	-	-
Female	-	2	2	7	7	6	7	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	2	1	3	5	11	6	9	7	-	-	-
Male	1	1	-	3	3	7	2	3	4	-	-	-
Female	-	1	1	-	2	4	4	6	3	-	-	-
White, both sexes	1	1	1	3	4	9	5	9	6	-	-	-
Male	1	1	-	3	3	6	2	3	3	-	-	-
Female	-	-	1	-	1	3	3	6	3	-	-	-
Black, both sexes	-	1	-	-	-	2	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Delaware												
All races, both sexes	1	1	-	1	1	2	2	1	-	-	-	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	2	2	-	-	-	-	-
White, both sexes	1	1	-	-	1	1	2	1	-	-	-	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	2	-	-	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Florida												
All races, both sexes	4	10	17	19	21	34	43	30	12	3	1	-
Male	1	4	9	7	13	10	13	7	1	1	-	-
Female	3	6	8	12	8	24	30	23	11	2	1	-
White, both sexes	2	7	16	17	18	31	41	27	12	3	1	-
Male	1	2	8	7	11	9	13	7	1	1	-	-
Female	1	5	8	10	7	22	28	20	11	2	1	-
Black, both sexes	2	3	1	2	3	2	2	2	-	-	-	-
Male	-	2	1	-	2	1	-	-	-	-	-	-
Female	2	1	-	2	1	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Georgia												
All races, both sexes	1	2	7	11	4	7	7	3	2	3	1	-
Male	-	1	2	7	-	2	1	2	2	1	-	-
Female	1	1	5	4	4	5	6	1	-	2	1	-
White, both sexes	-	2	3	8	3	6	5	3	2	2	1	-
Male	-	1	-	5	-	2	1	2	2	-	-	-
Female	-	1	3	3	3	4	4	1	-	2	1	-
Black, both sexes	1	-	4	3	1	1	1	-	-	1	-	-
Male	-	-	2	2	-	-	-	-	-	1	-	-
Female	1	-	2	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hawaii												
All races, both sexes	3	2	-	-	2	6	1	3	-	1	-	-
Male	2	1	-	-	1	2	-	1	-	1	-	-
Female	1	1	-	-	1	4	1	2	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	-	-	2	5	1	-	-	1	-	-
Male	2	1	-	-	1	2	-	-	-	1	-	-
Female	1	1	-	-	1	3	1	-	-	-	-	-
Idaho												
All races, both sexes	1	3	1	1	2	7	3	4	5	1	-	-
Male	-	1	1	-	-	1	-	1	1	-	-	-
Female	1	2	-	1	2	6	3	3	4	1	-	-
White, both sexes	1	3	1	1	2	7	3	4	5	1	-	-
Male	-	1	1	-	-	1	-	1	1	-	-	-
Female	1	2	-	1	2	6	3	3	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Illinois												
All races, both sexes	1	4	3	7	12	13	21	17	8	1	-	-
Male	-	3	2	2	6	5	6	4	1	1	-	-
Female	1	1	1	5	6	8	15	13	7	-	-	-
White, both sexes	1	4	2	6	11	13	18	16	8	1	-	-
Male	-	3	2	2	5	5	5	4	1	1	-	-
Female	1	1	-	4	6	8	13	12	7	-	-	-
Black, both sexes	-	-	-	-	1	-	3	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	2	6	3	9	6	8	10	7	2	-	-
Male	1	-	3	2	1	-	2	3	3	-	-	-
Female	1	2	3	1	8	6	6	7	4	2	-	-
White, both sexes	2	1	5	2	8	6	8	10	7	2	-	-
Male	1	-	2	1	1	-	2	3	3	-	-	-
Female	1	1	3	1	7	6	6	7	4	2	-	-
Black, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Iowa												
All races, both sexes	1	2	4	3	4	7	14	15	8	6	1	-
Male	-	-	-	1	3	4	2	1	1	-	-	-
Female	1	2	4	2	1	3	12	14	7	6	1	-
White, both sexes	1	2	3	3	4	7	14	15	8	6	1	-
Male	-	-	-	1	3	4	2	1	1	-	-	-
Female	1	2	3	2	1	3	12	14	7	6	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	1	2	1	3	7	3	2	-	1	-
Male	-	1	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	2	1	3	5	3	2	-	1	-
White, both sexes	-	-	1	2	1	2	6	3	2	-	1	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	1	2	5	3	2	-	1	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Kentucky												
All races, both sexes	-	2	5	8	5	9	10	6	2	1	-	-
Male	-	1	2	4	3	3	5	-	1	-	-	-
Female	-	1	3	4	2	6	5	6	1	1	-	-
White, both sexes	-	2	4	8	5	8	10	6	1	1	-	-
Male	-	1	2	4	3	3	5	-	1	-	-	-
Female	-	1	2	4	2	5	5	6	-	1	-	-
Black, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	2	3	3	3	4	2	2	2	-	-	-
Male	-	2	1	-	1	2	-	1	1	-	-	-
Female	2	-	2	3	2	2	2	1	1	-	-	-
White, both sexes	2	1	1	2	3	2	1	2	2	-	-	-
Male	-	1	-	-	1	1	-	1	1	-	-	-
Female	2	-	1	2	2	1	1	1	1	-	-	-
Black, both sexes	-	1	2	1	-	2	1	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Maine												
All races, both sexes	2	1	2	5	-	2	6	2	1	-	1	-
Male	-	-	-	1	-	2	2	-	1	-	1	-
Female	2	1	2	4	-	-	4	2	-	-	-	-
White, both sexes	2	1	2	5	-	2	5	2	1	-	1	-
Male	-	-	-	1	-	2	2	-	1	-	1	-
Female	2	1	2	4	-	-	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	4	6	3	10	10	6	-	2	1	-
Male	-	-	1	1	-	5	2	1	-	-	-	-
Female	-	1	3	5	3	5	8	5	-	2	1	-
White, both sexes	-	1	-	5	3	10	10	6	-	2	1	-
Male	-	-	-	1	-	5	2	1	-	-	-	-
Female	-	1	-	4	3	5	8	5	-	2	1	-
Black, both sexes	-	-	4	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	3	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Massachusetts												
All races, both sexes	-	1	4	7	6	14	16	15	9	2	-	-
Male	-	1	2	1	5	6	4	3	-	-	-	-
Female	-	-	2	6	1	8	12	12	9	2	-	-
White, both sexes	-	1	3	7	6	13	16	15	9	2	-	-
Male	-	1	2	1	5	6	4	3	-	-	-	-
Female	-	-	1	6	1	7	12	12	9	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	4	9	5	12	20	18	18	7	-	-	-
Male	-	2	3	3	2	7	7	6	3	-	-	-
Female	-	2	6	2	10	13	11	12	4	-	-	-
White, both sexes	-	3	6	3	12	18	16	18	7	-	-	-
Male	-	1	2	1	2	6	6	6	3	-	-	-
Female	-	2	4	2	10	12	10	12	4	-	-	-
Black, both sexes	-	1	3	1	-	2	1	-	-	-	-	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	-	-	2	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Minnesota												
All races, both sexes	2	4	2	9	6	13	16	13	6	1	-	-
Male	-	2	-	3	1	6	4	4	-	-	-	-
Female	2	2	2	6	5	7	12	9	6	1	-	-
White, both sexes	2	4	2	9	6	13	16	13	6	1	-	-
Male	-	2	-	3	1	6	4	4	-	-	-	-
Female	2	2	2	6	5	7	12	9	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	4	-	2	1	4	2	1	2	-	-	-
Male	1	2	-	-	-	2	1	1	-	-	-	-
Female	1	2	-	2	1	2	1	-	2	-	-	-
White, both sexes	1	1	-	2	1	4	-	1	2	-	-	-
Male	1	1	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	2	1	2	-	-	2	-	-	-
Black, both sexes	1	3	-	-	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	2	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Missouri												
All races, both sexes	1	1	2	7	12	11	13	10	10	1	-	-
Male	-	-	1	2	6	9	1	2	1	-	-	-
Female	1	1	1	5	6	2	12	8	9	1	-	-
White, both sexes	1	1	2	6	11	11	12	10	10	1	-	-
Male	-	-	1	2	6	9	1	2	1	-	-	-
Female	1	1	1	4	5	2	11	8	9	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	1	1	3	2	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	3	2	-	1	-	-
White, both sexes	-	-	1	-	1	1	3	2	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Nebraska												
All races, both sexes	-	1	-	1	3	8	6	4	1	2	1	-
Male	-	-	-	-	-	3	-	-	-	-	-	-
Female	-	1	-	1	3	5	6	4	1	2	1	-
White, both sexes	-	1	-	1	3	8	6	4	1	2	1	-
Male	-	-	-	-	-	3	-	-	-	-	-	-
Female	-	1	-	1	3	5	6	4	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	1	3	3	3	2	3	2	-	-	-
Male	-	1	-	1	2	2	-	-	1	-	-	-
Female	-	-	1	2	1	1	2	3	1	-	-	-
White, both sexes	-	1	1	3	3	3	1	3	2	-	-	-
Male	-	1	-	1	2	2	-	-	1	-	-	-
Female	-	-	1	2	1	1	1	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Hampshire												
All races, both sexes	1	1	-	-	1	1	3	4	3	1	2	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	-	1	-	-	1	1	3	3	3	-	2	-
White, both sexes	1	1	-	-	1	1	3	4	3	1	2	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	-	1	-	-	1	1	3	3	3	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	2	6	9	11	21	30	17	13	5	1	-
Male	-	-	4	4	2	4	9	5	1	-	-	-
Female	2	2	2	5	9	17	21	12	12	5	1	-
White, both sexes	2	2	4	9	8	20	29	15	11	5	1	-
Male	-	-	2	4	2	4	9	5	1	-	-	-
Female	2	2	2	5	6	16	20	10	10	5	1	-
Black, both sexes	-	-	2	-	3	1	1	2	2	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	3	1	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Mexico												
All races, both sexes	-	2	2	2	2	1	7	3	1	1	-	-
Male	-	1	1	1	-	-	2	1	-	-	-	-
Female	-	1	1	1	2	1	5	2	1	1	-	-
White, both sexes	-	2	2	1	2	1	6	3	1	1	-	-
Male	-	1	1	1	-	-	2	1	-	-	-	-
Female	-	1	1	-	2	1	4	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	11	11	9	17	30	31	21	20	9	1	-
Male	4	4	4	3	6	9	9	6	1	1	-	-
Female	2	7	7	6	11	21	22	15	19	8	1	-
White, both sexes	3	11	8	7	14	28	30	21	20	7	1	-
Male	2	4	3	2	4	9	9	6	1	1	-	-
Female	1	7	5	5	10	19	21	15	19	6	1	-
Black, both sexes	3	-	2	2	2	2	1	-	-	1	-	-
Male	2	-	1	1	1	-	-	-	-	-	-	-
Female	1	-	1	1	1	2	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
North Carolina												
All races, both sexes	3	12	5	7	6	12	16	19	10	-	-	-
Male	1	6	2	2	1	2	7	3	5	-	-	-
Female	2	6	3	5	5	10	9	16	5	-	-	-
White, both sexes	3	7	4	4	5	10	14	19	8	-	-	-
Male	1	3	2	1	1	-	7	3	4	-	-	-
Female	2	4	2	3	4	10	7	16	4	-	-	-
Black, both sexes	-	4	1	3	1	2	2	-	2	-	-	-
Male	-	2	-	1	-	2	-	-	1	-	-	-
Female	-	2	1	2	1	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	2	4	3	3	2	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	2	4	1	2	2	-	-	-
White, both sexes	-	-	-	1	2	4	3	3	2	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	2	4	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Ohio												
All races, both sexes	8	11	2	7	17	36	26	24	11	6	1	-
Male	2	6	1	1	7	10	4	5	1	-	-	-
Female	6	5	1	6	10	26	22	19	10	6	1	-
White, both sexes	6	8	2	7	16	34	25	24	11	5	1	-
Male	1	5	1	1	7	9	4	5	1	-	-	-
Female	5	3	1	6	9	25	21	19	10	5	1	-
Black, both sexes	2	3	-	-	1	2	1	-	-	1	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	1	2	-	-	1	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	5	5	2	4	6	9	3	1	1	-	-
Male	1	1	1	-	-	1	4	2	-	-	-	-
Female	-	4	4	2	4	5	5	1	1	1	-	-
White, both sexes	1	2	5	1	4	5	9	3	1	1	-	-
Male	1	1	1	-	-	1	4	2	-	-	-	-
Female	-	1	4	1	4	4	5	1	1	1	-	-
Black, both sexes	-	2	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Oregon												
All races, both sexes	-	3	3	1	8	8	16	16	14	7	-	-
Male	-	1	1	-	2	1	4	3	1	2	-	-
Female	-	2	2	1	6	7	12	13	13	5	-	-
White, both sexes	-	3	3	1	6	8	16	16	14	7	-	-
Male	-	1	1	-	1	1	4	3	1	2	-	-
Female	-	2	2	1	5	7	12	13	13	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	5	8	7	14	26	42	37	16	6	2	-
Male	1	2	4	2	4	6	9	8	3	1	-	-
Female	3	3	4	5	10	20	33	29	13	5	2	-
White, both sexes	3	5	6	7	13	25	41	36	16	6	2	-
Male	1	2	3	2	3	6	9	8	3	1	-	-
Female	2	3	3	5	10	19	32	28	13	5	2	-
Black, both sexes	1	-	2	-	1	1	1	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Rhode Island												
All races, both sexes	-	1	-	2	2	1	3	3	1	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	1	-	1	2	1	3	2	-	-	-	-
White, both sexes	-	1	-	1	2	1	3	3	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	1	-	1	2	1	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	3	4	8	12	3	6	3	1	3	-	-
Male	1	2	1	4	8	2	-	1	-	-	-	-
Female	2	1	3	4	4	1	6	2	1	3	-	-
White, both sexes	2	2	4	6	7	3	4	3	1	3	-	-
Male	-	1	1	3	4	2	-	1	-	-	-	-
Female	2	1	3	3	3	1	4	2	1	3	-	-
Black, both sexes	1	1	-	2	4	-	2	-	-	-	-	-
Male	1	1	-	1	3	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
South Dakota												
All races, both sexes	-	-	2	2	2	2	3	2	1	1	-	-
Male	-	-	-	1	-	-	2	-	-	1	-	-
Female	-	-	2	1	2	2	1	2	1	-	-	-
White, both sexes	-	-	-	2	2	2	3	2	1	1	-	-
Male	-	-	-	1	-	-	2	-	-	1	-	-
Female	-	-	-	1	2	2	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	5	7	5	6	6	18	8	5	1	-	-
Male	3	1	1	1	3	3	8	2	2	1	-	-
Female	1	4	6	4	3	3	10	6	3	-	-	-
White, both sexes	4	4	6	5	6	6	18	8	4	1	-	-
Male	3	1	1	1	3	3	8	2	2	1	-	-
Female	1	3	5	4	3	3	10	6	2	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Texas												
All races, both sexes	7	8	9	9	18	16	16	16	10	2	1	-
Male	4	1	3	3	4	3	2	2	1	1	-	-
Female	3	7	6	6	14	13	14	14	9	1	1	-
White, both sexes	5	8	6	7	17	15	15	16	10	2	1	-
Male	3	1	3	2	4	3	2	2	1	1	-	-
Female	2	7	3	5	13	12	13	14	9	1	1	-
Black, both sexes	2	-	2	2	1	1	1	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	2	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	3	3	4	6	4	8	5	2	-	-	-
Male	-	-	1	1	1	1	2	1	-	-	-	-
Female	-	3	2	3	5	3	6	4	2	-	-	-
White, both sexes	-	3	3	4	6	4	8	5	2	-	-	-
Male	-	-	1	1	1	1	2	1	-	-	-	-
Female	-	3	2	3	5	3	6	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Vermont												
All races, both sexes	-	-	-	1	2	2	2	4	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	2	1	1	3	-	-	-	-
White, both sexes	-	-	-	1	2	2	2	4	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	2	1	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	2	7	3	7	11	13	12	1	1	-	-
Male	-	-	3	1	-	3	5	5	-	-	-	-
Female	1	2	4	2	7	8	8	7	1	1	-	-
White, both sexes	1	2	5	3	5	9	11	12	1	1	-	-
Male	-	-	3	1	-	3	5	5	-	-	-	-
Female	1	2	2	2	5	6	6	7	1	1	-	-
Black, both sexes	-	-	2	-	2	2	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	2	2	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Washington												
All races, both sexes	4	1	2	4	12	11	16	35	5	8	2	-
Male	3	1	1	2	8	3	3	12	2	1	-	-
Female	1	-	1	2	4	8	13	23	3	7	2	-
White, both sexes	1	1	2	4	12	11	16	33	5	8	2	-
Male	1	1	1	2	8	3	3	12	2	1	-	-
Female	-	-	1	2	4	8	13	21	3	7	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
West Virginia												
All races, both sexes	-	1	2	4	2	3	5	2	-	-	-	-
Male	-	1	-	1	-	1	3	2	-	-	-	-
Female	-	-	2	3	2	2	2	-	-	-	-	-
White, both sexes	-	1	2	4	2	3	5	2	-	-	-	-
Male	-	1	-	1	-	1	3	2	-	-	-	-
Female	-	-	2	3	2	2	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Wisconsin												
All races, both sexes	2	6	3	4	12	8	20	15	6	6	1	-
Male	1	-	3	3	1	2	4	2	3	1	-	-
Female	1	6	-	1	11	6	16	13	3	5	1	-
White, both sexes	2	5	3	4	12	8	20	15	6	6	1	-
Male	1	-	3	3	1	2	4	2	3	1	-	-
Female	1	5	-	1	11	6	16	13	3	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	2	1	1	2	2	1	-	-	-
Male	-	-	-	2	1	-	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	1	-	-	-
White, both sexes	-	1	-	2	1	1	2	2	1	-	-	-
Male	-	-	-	2	1	-	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
United States												
All races, both sexes	25	-	-	-	2	-	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	2	-	-	-	1	-	1	-
White, both sexes	18	-	-	-	1	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	1	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	1	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Florida												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Massachusetts												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
North Carolina												
All races, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Texas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Wisconsin												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
United States												
All races, both sexes	2	1	4	1	1	3	5	1	1	1	-	-
Male	2	-	2	-	-	1	1	-	1	-	-	-
Female	-	1	2	1	1	2	4	1	-	1	-	-
White, both sexes	2	1	1	-	1	3	5	1	1	1	-	-
Male	2	-	1	-	-	1	1	-	1	-	-	-
Female	-	1	-	-	1	2	4	1	-	1	-	-
Black, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	1	-	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Colorado												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Delaware												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Florida												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Texas												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Washington												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Wisconsin												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
United States												
All races, both sexes	3,232	3	-	1	4	3	4	9	21	21	44	83
Male	1,000	2	-	1	1	2	3	4	13	11	17	40
Female	2,232	1	-	-	3	1	1	5	8	10	27	43
White, both sexes	2,885	3	-	-	1	1	2	7	13	13	28	61
Male	881	2	-	-	-	1	2	3	8	8	14	32
Female	2,004	1	-	-	1	-	-	4	5	5	14	29
Black, both sexes	243	-	-	-	2	2	-	1	5	7	13	21
Male	86	-	-	-	1	1	-	1	3	3	3	8
Female	157	-	-	-	1	1	-	-	2	4	10	13
American Indian 1/, both sexes	20	-	-	1	-	-	2	-	-	-	-	-
Male	6	-	-	1	-	-	1	-	-	-	-	-
Female	14	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	84	-	-	-	1	-	-	1	3	1	3	1
Male	27	-	-	-	-	-	-	-	2	-	-	-
Female	57	-	-	-	1	-	-	1	1	1	3	1
Alabama												
All races, both sexes	50	-	-	-	-	-	-	-	1	1	1	4
Male	15	-	-	-	-	-	-	-	-	1	-	2
Female	35	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	41	-	-	-	-	-	-	-	1	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	2
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Alaska												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	59	-	-	-	-	-	-	1	-	-	-	1
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	-	-	-	-	-	-	1	-	-	-	1
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Arkansas												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	391	1	-	1	-	-	-	1	4	1	9	14
Male	135	-	-	1	-	-	-	-	1	1	3	3
Female	256	1	-	-	-	-	-	1	3	-	6	11
White, both sexes	331	1	-	-	-	-	-	1	2	1	6	12
Male	115	-	-	-	-	-	-	-	-	1	2	3
Female	216	1	-	-	-	-	-	1	2	-	4	9
Black, both sexes	20	-	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	-	-	-	-	-	-	1	-	1	1
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Colorado												
All races, both sexes	59	-	-	-	-	-	1	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	5
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	24	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	3
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Delaware												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Florida												
All races, both sexes	207	-	-	-	1	-	1	-	2	1	2	7
Male	74	-	-	-	-	-	1	-	2	-	1	4
Female	133	-	-	-	1	-	-	-	-	1	1	3
White, both sexes	184	-	-	-	-	-	1	-	1	1	1	6
Male	67	-	-	-	-	-	1	-	1	-	1	4
Female	117	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	21	-	-	-	1	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	55	-	-	-	-	1	-	1	-	2	2	1
Male	21	-	-	-	-	-	-	1	-	-	1	1
Female	34	-	-	-	-	1	-	-	-	2	1	-
White, both sexes	40	-	-	-	-	-	-	1	-	1	2	1
Male	16	-	-	-	-	-	-	1	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	14	-	-	-	-	1	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Hawaii												
All races, both sexes	22	-	-	-	-	-	-	1	1	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	1	-	2	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	1	1	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	1	-	2	-
Idaho												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Illinois												
All races, both sexes	93	-	-	-	-	-	1	-	1	2	-	2
Male	34	-	-	-	-	-	1	-	1	1	-	1
Female	59	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	85	-	-	-	-	-	1	-	1	1	-	2
Male	32	-	-	-	-	-	1	-	1	1	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Iowa												
All races, both sexes	69	-	-	-	-	-	-	-	-	2	1	1
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	56	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	68	-	-	-	-	-	-	-	-	2	1	1
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	55	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	25	1	-	-	-	-	-	1	2	-	-	-
Male	7	1	-	-	-	-	-	1	1	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	21	1	-	-	-	-	-	-	2	-	-	-
Male	4	1	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Kentucky												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	29	-	-	-	-	1	-	-	2	-	2	1
Male	11	-	-	-	-	1	-	-	2	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	19	-	-	-	-	-	-	-	1	-	1	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	10	-	-	-	-	1	-	-	1	-	1	-
Male	5	-	-	-	-	1	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Maine												
All races, both sexes	25	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	24	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	1	5
Male	13	-	-	-	-	-	-	-	-	1	-	2
Female	36	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	1	3
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Massachusetts												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	71	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	94	-	-	-	-	-	-	-	-	1	-	1
Male	35	-	-	-	-	-	-	-	-	1	-	1
Female	59	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Minnesota												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	21	-	-	-	-	-	-	-	1	-	-	2
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	13	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Missouri												
All races, both sexes	71	-	-	-	-	-	-	-	-	1	1	1
Male	25	-	-	-	-	-	-	-	-	1	1	1
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	1	1	1
Male	25	-	-	-	-	-	-	-	-	1	1	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Nebraska												
All races, both sexes	28	-	-	-	-	-	1	-	1	-	-	-
Male	4	-	-	-	-	-	1	-	1	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	20	-	-	-	1	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
New Hampshire												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	121	-	-	-	-	-	-	-	1	1	1	1
Male	31	-	-	-	-	-	-	-	1	-	-	1
Female	90	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	109	-	-	-	-	-	-	-	1	-	1	1
Male	29	-	-	-	-	-	-	-	1	-	-	1
Female	80	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
New Mexico												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	178	-	-	-	1	1	-	1	1	-	4	5
Male	58	-	-	-	1	1	-	-	1	-	3	5
Female	120	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	157	-	-	-	-	1	-	1	-	-	4	1
Male	46	-	-	-	-	1	-	-	-	-	3	1
Female	111	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	17	-	-	-	1	-	-	-	-	-	-	4
Male	10	-	-	-	1	-	-	-	-	-	-	4
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
North Carolina												
All races, both sexes	94	-	-	-	-	-	-	-	-	2	1	3
Male	30	-	-	-	-	-	-	-	-	1	-	-
Female	64	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	76	-	-	-	-	-	-	-	-	2	-	2
Male	23	-	-	-	-	-	-	-	-	1	-	-
Female	53	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Ohio												
All races, both sexes	160	-	-	-	-	-	-	-	1	1	2	7
Male	44	-	-	-	-	-	-	-	1	-	2	4
Female	116	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	146	-	-	-	-	-	-	-	-	-	2	5
Male	39	-	-	-	-	-	-	-	-	-	2	3
Female	107	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	14	-	-	-	-	-	-	-	1	1	-	2
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	9	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	38	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Oregon												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	75	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	170	-	-	-	-	-	-	-	1	-	1	2
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	130	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	161	-	-	-	-	-	-	-	-	-	1	1
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	123	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Rhode Island												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	3	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
South Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	72	-	-	-	1	-	-	-	1	1	2	3
Male	31	-	-	-	-	-	-	-	1	1	2	3
Female	41	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	67	-	-	-	1	-	-	-	1	1	-	3
Male	29	-	-	-	-	-	-	-	1	1	-	3
Female	38	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Texas												
All races, both sexes	117	1	-	-	-	-	-	1	-	2	-	2
Male	26	1	-	-	-	-	-	-	-	-	-	1
Female	91	-	-	-	-	-	-	1	-	2	-	1
White, both sexes	105	1	-	-	-	-	-	1	-	1	-	1
Male	23	1	-	-	-	-	-	-	-	-	-	-
Female	82	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Utah												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Vermont												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	67	-	-	-	-	-	-	1	-	2	4	2
Male	20	-	-	-	-	-	-	-	-	2	-	1
Female	47	-	-	-	-	-	-	1	-	-	4	1
White, both sexes	55	-	-	-	-	-	-	1	-	2	1	1
Male	20	-	-	-	-	-	-	-	-	2	-	1
Female	35	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	3	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Washington												
All races, both sexes	102	-	-	-	-	-	-	-	-	-	2	1
Male	38	-	-	-	-	-	-	-	-	-	1	1
Female	64	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	97	-	-	-	-	-	-	-	-	-	2	1
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	61	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Wisconsin												
All races, both sexes	81	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	80	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
United States												
All races, both sexes	90	156	191	256	350	499	616	511	257	95	18	-
Male	38	61	72	92	116	158	167	134	50	17	1	-
Female	52	95	119	164	234	341	449	377	207	78	17	-
White, both sexes	61	124	151	219	308	457	581	499	250	88	18	-
Male	25	46	58	79	101	145	162	131	48	15	1	-
Female	36	78	93	140	207	312	419	368	202	73	17	-
Black, both sexes	17	22	30	27	25	28	26	6	7	4	-	-
Male	7	10	13	10	9	10	4	-	2	1	-	-
Female	10	12	17	17	16	18	22	6	5	3	-	-
American Indian 1/, both sexes	3	3	3	1	1	4	2	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	3	1	1	4	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	7	7	9	16	10	7	6	-	3	-	-
Male	4	3	1	3	6	3	1	3	-	1	-	-
Female	5	4	6	6	10	7	6	3	-	2	-	-
Alabama												
All races, both sexes	4	2	5	5	7	9	7	2	2	-	-	-
Male	2	1	-	1	2	5	1	-	-	-	-	-
Female	2	1	5	4	5	4	6	2	2	-	-	-
White, both sexes	2	2	5	4	6	7	7	2	2	-	-	-
Male	1	1	-	-	2	4	1	-	-	-	-	-
Female	1	1	5	4	4	3	6	2	2	-	-	-
Black, both sexes	1	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Alaska												
All races, both sexes	2	1	-	1	-	2	1	-	1	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	1	-	1	-	-	-
White, both sexes	1	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	4	4	7	8	15	8	6	3	-	-	-
Male	2	1	2	2	2	3	1	1	-	-	-	-
Female	-	3	2	5	6	12	7	5	3	-	-	-
White, both sexes	2	4	4	6	7	15	7	6	3	-	-	-
Male	2	1	2	2	2	3	1	1	-	-	-	-
Female	-	3	2	4	5	12	6	5	3	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Arkansas												
All races, both sexes	1	-	1	1	3	2	5	5	2	-	-	-
Male	1	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	1	1	3	2	4	2	2	-	-	-
White, both sexes	-	-	1	1	3	2	5	5	2	-	-	-
Male	-	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	1	1	3	2	4	2	2	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12	13	26	37	41	55	78	61	29	8	-	-
Male	4	6	10	16	15	16	26	22	9	2	-	-
Female	8	7	16	21	26	39	52	39	20	6	-	-
White, both sexes	6	9	20	26	32	47	75	58	29	6	-	-
Male	3	4	7	14	12	13	26	19	9	2	-	-
Female	3	5	13	12	20	34	49	39	20	4	-	-
Black, both sexes	1	-	2	5	3	3	1	-	-	1	-	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	1	-	-	4	2	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	4	4	6	6	4	2	3	-	1	-	-
Male	1	2	1	1	2	1	-	3	-	-	-	-
Female	4	2	3	5	4	3	2	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Colorado												
All races, both sexes	1	3	3	7	10	11	10	6	4	2	-	-
Male	1	1	-	-	2	5	3	2	-	1	-	-
Female	-	2	3	7	8	6	7	4	4	1	-	-
White, both sexes	1	3	2	7	9	11	9	6	4	2	-	-
Male	1	1	-	-	2	5	2	2	-	1	-	-
Female	-	2	2	7	7	6	7	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	2	1	3	5	11	6	9	6	-	-	-
Male	1	1	-	3	3	7	2	3	3	-	-	-
Female	-	1	1	-	2	4	4	6	3	-	-	-
White, both sexes	1	1	1	3	4	9	5	9	5	-	-	-
Male	1	1	-	3	3	6	2	3	2	-	-	-
Female	-	-	1	-	1	3	3	6	3	-	-	-
Black, both sexes	-	1	-	-	-	2	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Delaware												
All races, both sexes	1	1	-	-	1	2	2	1	-	-	-	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	2	2	-	-	-	-	-
White, both sexes	1	1	-	-	1	1	2	1	-	-	-	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Florida												
All races, both sexes	4	10	17	19	21	33	43	30	12	3	1	-
Male	1	4	9	7	13	10	13	7	1	1	-	-
Female	3	6	8	12	8	23	30	23	11	2	1	-
White, both sexes	2	7	16	17	18	30	41	27	12	3	1	-
Male	1	2	8	7	11	9	13	7	1	1	-	-
Female	1	5	8	10	7	21	28	20	11	2	1	-
Black, both sexes	2	3	1	2	3	2	2	2	-	-	-	-
Male	-	2	1	-	2	1	-	-	-	-	-	-
Female	2	1	-	2	1	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Georgia												
All races, both sexes	1	2	7	11	4	7	7	3	2	3	1	-
Male	-	1	2	7	-	2	1	2	2	1	-	-
Female	1	1	5	4	4	5	6	1	-	2	1	-
White, both sexes	-	2	3	8	3	6	5	3	2	2	1	-
Male	-	1	-	5	-	2	1	2	2	-	-	-
Female	-	1	3	3	3	4	4	1	-	2	1	-
Black, both sexes	1	-	4	3	1	1	1	-	-	1	-	-
Male	-	-	2	2	-	-	-	-	-	1	-	-
Female	1	-	2	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Hawaii												
All races, both sexes	3	2	-	-	2	6	1	3	-	1	-	-
Male	2	1	-	-	1	2	-	1	-	1	-	-
Female	1	1	-	-	1	4	1	2	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	-	-	2	5	1	-	-	1	-	-
Male	2	1	-	-	1	2	-	-	-	1	-	-
Female	1	1	-	-	1	3	1	-	-	-	-	-
Idaho												
All races, both sexes	1	3	1	1	2	7	3	4	5	1	-	-
Male	-	1	1	-	-	1	-	1	1	-	-	-
Female	1	2	-	1	2	6	3	3	4	1	-	-
White, both sexes	1	3	1	1	2	7	3	4	5	1	-	-
Male	-	1	1	-	-	1	-	1	1	-	-	-
Female	1	2	-	1	2	6	3	3	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Illinois												
All races, both sexes	1	4	3	7	12	13	21	17	8	1	-	-
Male	-	3	2	2	6	5	6	4	1	1	-	-
Female	1	1	1	5	6	8	15	13	7	-	-	-
White, both sexes	1	4	2	6	11	13	18	16	8	1	-	-
Male	-	3	2	2	5	5	5	4	1	1	-	-
Female	1	1	-	4	6	8	13	12	7	-	-	-
Black, both sexes	-	-	-	-	1	-	3	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	2	6	3	9	6	8	10	7	2	-	-
Male	1	-	3	2	1	-	2	3	3	-	-	-
Female	1	2	3	1	8	6	6	7	4	2	-	-
White, both sexes	2	1	5	2	8	6	8	10	7	2	-	-
Male	1	-	2	1	1	-	2	3	3	-	-	-
Female	1	1	3	1	7	6	6	7	4	2	-	-
Black, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Iowa												
All races, both sexes	1	2	4	3	4	7	14	15	8	6	1	-
Male	-	-	-	1	3	4	2	1	1	-	-	-
Female	1	2	4	2	1	3	12	14	7	6	1	-
White, both sexes	1	2	3	3	4	7	14	15	8	6	1	-
Male	-	-	-	1	3	4	2	1	1	-	-	-
Female	1	2	3	2	1	3	12	14	7	6	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	1	2	1	3	7	3	2	-	1	-
Male	-	1	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	2	1	3	5	3	2	-	1	-
White, both sexes	-	-	1	2	1	2	6	3	2	-	1	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	1	2	5	3	2	-	1	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Kentucky												
All races, both sexes	-	2	5	8	5	9	10	6	2	1	-	-
Male	-	1	2	4	3	3	5	-	1	-	-	-
Female	-	1	3	4	2	6	5	6	1	1	-	-
White, both sexes	-	2	4	8	5	8	10	6	1	1	-	-
Male	-	1	2	4	3	3	5	-	1	-	-	-
Female	-	1	2	4	2	5	5	6	-	1	-	-
Black, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	2	3	3	3	4	2	2	2	-	-	-
Male	-	2	1	-	1	2	-	1	1	-	-	-
Female	2	-	2	3	2	2	2	1	1	-	-	-
White, both sexes	2	1	1	2	3	2	1	2	2	-	-	-
Male	-	1	-	-	1	1	-	1	1	-	-	-
Female	2	-	1	2	2	1	1	1	1	-	-	-
Black, both sexes	-	1	2	1	-	2	1	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Maine												
All races, both sexes	2	1	2	5	-	2	6	2	1	-	1	-
Male	-	-	-	1	-	2	2	-	1	-	1	-
Female	2	1	2	4	-	-	4	2	-	-	-	-
White, both sexes	2	1	2	5	-	2	5	2	1	-	1	-
Male	-	-	-	1	-	2	2	-	1	-	1	-
Female	2	1	2	4	-	-	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	3	6	3	10	10	6	-	2	1	-
Male	-	-	1	1	-	5	2	1	-	-	-	-
Female	-	1	2	5	3	5	8	5	-	2	1	-
White, both sexes	-	1	-	5	3	10	10	6	-	2	1	-
Male	-	-	-	1	-	5	2	1	-	-	-	-
Female	-	1	-	4	3	5	8	5	-	2	1	-
Black, both sexes	-	-	3	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Massachusetts												
All races, both sexes	-	1	4	7	6	14	15	15	9	2	-	-
Male	-	1	2	1	5	6	4	3	-	-	-	-
Female	-	-	2	6	1	8	11	12	9	2	-	-
White, both sexes	-	1	3	7	6	13	15	15	9	2	-	-
Male	-	1	2	1	5	6	4	3	-	-	-	-
Female	-	-	1	6	1	7	11	12	9	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	4	9	5	12	20	17	18	7	-	-	-
Male	-	2	3	3	2	7	7	6	3	-	-	-
Female	-	2	6	2	10	13	10	12	4	-	-	-
White, both sexes	-	3	6	3	12	18	15	18	7	-	-	-
Male	-	1	2	1	2	6	6	6	3	-	-	-
Female	-	2	4	2	10	12	9	12	4	-	-	-
Black, both sexes	-	1	3	1	-	2	1	-	-	-	-	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	-	-	2	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Minnesota												
All races, both sexes	2	4	2	9	6	13	16	13	6	1	-	-
Male	-	2	-	3	1	6	4	4	-	-	-	-
Female	2	2	2	6	5	7	12	9	6	1	-	-
White, both sexes	2	4	2	9	6	13	16	13	6	1	-	-
Male	-	2	-	3	1	6	4	4	-	-	-	-
Female	2	2	2	6	5	7	12	9	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	4	-	2	1	4	2	1	2	-	-	-
Male	1	2	-	-	-	2	1	1	-	-	-	-
Female	1	2	-	2	1	2	1	-	2	-	-	-
White, both sexes	1	1	-	2	1	4	-	1	2	-	-	-
Male	1	1	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	2	1	2	-	-	2	-	-	-
Black, both sexes	1	3	-	-	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	2	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Missouri												
All races, both sexes	1	1	2	7	12	11	13	10	10	1	-	-
Male	-	-	1	2	6	9	1	2	1	-	-	-
Female	1	1	1	5	6	2	12	8	9	1	-	-
White, both sexes	1	1	2	6	11	11	12	10	10	1	-	-
Male	-	-	1	2	6	9	1	2	1	-	-	-
Female	1	1	1	4	5	2	11	8	9	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	1	1	3	2	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	3	2	-	1	-	-
White, both sexes	-	-	1	-	1	1	3	2	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Nebraska												
All races, both sexes	-	1	-	1	3	7	6	4	1	2	1	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	-	1	3	5	6	4	1	2	1	-
White, both sexes	-	1	-	1	3	7	6	4	1	2	1	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	-	1	3	5	6	4	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	1	3	3	3	2	3	2	-	-	-
Male	-	1	-	1	2	2	-	-	1	-	-	-
Female	-	-	1	2	1	1	2	3	1	-	-	-
White, both sexes	-	1	1	3	3	3	1	3	2	-	-	-
Male	-	1	-	1	2	2	-	-	1	-	-	-
Female	-	-	1	2	1	1	1	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
New Hampshire												
All races, both sexes	1	1	-	-	1	1	3	4	3	1	2	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	-	1	-	-	1	1	3	3	3	-	2	-
White, both sexes	1	1	-	-	1	1	3	4	3	1	2	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	-	1	-	-	1	1	3	3	3	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	2	6	9	11	21	30	17	13	5	1	-
Male	-	-	4	4	2	4	9	5	1	-	-	-
Female	2	2	2	5	9	17	21	12	12	5	1	-
White, both sexes	2	2	4	9	8	20	29	15	11	5	1	-
Male	-	-	2	4	2	4	9	5	1	-	-	-
Female	2	2	2	5	6	16	20	10	10	5	1	-
Black, both sexes	-	-	2	-	3	1	1	2	2	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	3	1	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
New Mexico												
All races, both sexes	-	2	2	2	2	1	7	3	1	1	-	-
Male	-	1	1	1	-	-	2	1	-	-	-	-
Female	-	1	1	1	2	1	5	2	1	1	-	-
White, both sexes	-	2	2	1	2	1	6	3	1	1	-	-
Male	-	1	1	1	-	-	2	1	-	-	-	-
Female	-	1	1	-	2	1	4	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	11	10	9	17	30	31	21	20	9	1	-
Male	4	4	4	3	6	9	9	6	1	1	-	-
Female	2	7	6	6	11	21	22	15	19	8	1	-
White, both sexes	3	11	8	7	14	28	30	21	20	7	1	-
Male	2	4	3	2	4	9	9	6	1	1	-	-
Female	1	7	5	5	10	19	21	15	19	6	1	-
Black, both sexes	3	-	1	2	2	2	1	-	-	1	-	-
Male	2	-	1	1	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	2	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
North Carolina												
All races, both sexes	3	12	5	7	6	12	15	18	10	-	-	-
Male	1	6	2	2	1	2	7	3	5	-	-	-
Female	2	6	3	5	5	10	8	15	5	-	-	-
White, both sexes	3	7	4	4	5	10	13	18	8	-	-	-
Male	1	3	2	1	1	-	7	3	4	-	-	-
Female	2	4	2	3	4	10	6	15	4	-	-	-
Black, both sexes	-	4	1	3	1	2	2	-	2	-	-	-
Male	-	2	-	1	-	2	-	-	1	-	-	-
Female	-	2	1	2	1	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	2	4	3	3	2	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	2	4	1	2	2	-	-	-
White, both sexes	-	-	-	1	2	4	3	3	2	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	2	4	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Ohio												
All races, both sexes	8	11	2	7	17	36	26	24	11	6	1	-
Male	2	6	1	1	7	10	4	5	1	-	-	-
Female	6	5	1	6	10	26	22	19	10	6	1	-
White, both sexes	6	8	2	7	16	34	25	24	11	5	1	-
Male	1	5	1	1	7	9	4	5	1	-	-	-
Female	5	3	1	6	9	25	21	19	10	5	1	-
Black, both sexes	2	3	-	-	1	2	1	-	-	1	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	1	2	-	-	1	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	5	5	2	4	6	9	3	1	1	-	-
Male	1	1	1	-	-	1	4	2	-	-	-	-
Female	-	4	4	2	4	5	5	1	1	1	-	-
White, both sexes	1	2	5	1	4	5	9	3	1	1	-	-
Male	1	1	1	-	-	1	4	2	-	-	-	-
Female	-	1	4	1	4	4	5	1	1	1	-	-
Black, both sexes	-	2	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Oregon												
All races, both sexes	-	3	3	1	8	8	16	16	14	7	-	-
Male	-	1	1	-	2	1	4	3	1	2	-	-
Female	-	2	2	1	6	7	12	13	13	5	-	-
White, both sexes	-	3	3	1	6	8	16	16	14	7	-	-
Male	-	1	1	-	1	1	4	3	1	2	-	-
Female	-	2	2	1	5	7	12	13	13	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	5	8	7	14	26	41	37	16	6	2	-
Male	1	2	4	2	4	6	9	8	3	1	-	-
Female	3	3	4	5	10	20	32	29	13	5	2	-
White, both sexes	3	5	6	7	13	25	40	36	16	6	2	-
Male	1	2	3	2	3	6	9	8	3	1	-	-
Female	2	3	3	5	10	19	31	28	13	5	2	-
Black, both sexes	1	-	2	-	1	1	1	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Rhode Island												
All races, both sexes	-	1	-	2	2	1	3	3	1	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	1	-	1	2	1	3	2	-	-	-	-
White, both sexes	-	1	-	1	2	1	3	3	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	1	-	1	2	1	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	3	4	8	12	3	6	3	1	3	-	-
Male	1	2	1	4	8	2	-	1	-	-	-	-
Female	2	1	3	4	4	1	6	2	1	3	-	-
White, both sexes	2	2	4	6	7	3	4	3	1	3	-	-
Male	-	1	1	3	4	2	-	1	-	-	-	-
Female	2	1	3	3	3	1	4	2	1	3	-	-
Black, both sexes	1	1	-	2	4	-	2	-	-	-	-	-
Male	1	1	-	1	3	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
South Dakota												
All races, both sexes	-	-	2	2	2	2	3	2	1	1	-	-
Male	-	-	-	1	-	-	2	-	-	1	-	-
Female	-	-	2	1	2	2	1	2	1	-	-	-
White, both sexes	-	-	-	2	2	2	3	2	1	1	-	-
Male	-	-	-	1	-	-	2	-	-	1	-	-
Female	-	-	-	1	2	2	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	5	7	5	6	6	18	8	5	1	-	-
Male	2	1	1	1	3	3	8	2	2	1	-	-
Female	1	4	6	4	3	3	10	6	3	-	-	-
White, both sexes	3	4	6	5	6	6	18	8	4	1	-	-
Male	2	1	1	1	3	3	8	2	2	1	-	-
Female	1	3	5	4	3	3	10	6	2	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Texas												
All races, both sexes	7	8	9	9	18	15	16	16	10	2	1	-
Male	4	1	3	3	4	3	2	2	1	1	-	-
Female	3	7	6	6	14	12	14	14	9	1	1	-
White, both sexes	5	8	6	7	17	14	15	16	10	2	1	-
Male	3	1	3	2	4	3	2	2	1	1	-	-
Female	2	7	3	5	13	11	13	14	9	1	1	-
Black, both sexes	2	-	2	2	1	1	1	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	2	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	3	3	4	6	4	8	5	2	-	-	-
Male	-	-	1	1	1	1	2	1	-	-	-	-
Female	-	3	2	3	5	3	6	4	2	-	-	-
White, both sexes	-	3	3	4	6	4	8	5	2	-	-	-
Male	-	-	1	1	1	1	2	1	-	-	-	-
Female	-	3	2	3	5	3	6	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Vermont												
All races, both sexes	-	-	-	1	2	2	2	4	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	2	1	1	3	-	-	-	-
White, both sexes	-	-	-	1	2	2	2	4	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	2	1	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	2	7	3	7	11	13	12	1	1	-	-
Male	-	-	3	1	-	3	5	5	-	-	-	-
Female	1	2	4	2	7	8	8	7	1	1	-	-
White, both sexes	1	2	5	3	5	9	11	12	1	1	-	-
Male	-	-	3	1	-	3	5	5	-	-	-	-
Female	1	2	2	2	5	6	6	7	1	1	-	-
Black, both sexes	-	-	2	-	2	2	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	2	2	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Washington												
All races, both sexes	4	1	2	4	11	11	16	35	5	8	2	-
Male	3	1	1	2	8	3	3	12	2	1	-	-
Female	1	-	1	2	3	8	13	23	3	7	2	-
White, both sexes	1	1	2	4	11	11	16	33	5	8	2	-
Male	1	1	1	2	8	3	3	12	2	1	-	-
Female	-	-	1	2	3	8	13	21	3	7	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
West Virginia												
All races, both sexes	-	1	2	4	2	3	5	2	-	-	-	-
Male	-	1	-	1	-	1	3	2	-	-	-	-
Female	-	-	2	3	2	2	2	-	-	-	-	-
White, both sexes	-	1	2	4	2	3	5	2	-	-	-	-
Male	-	1	-	1	-	1	3	2	-	-	-	-
Female	-	-	2	3	2	2	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Wisconsin												
All races, both sexes	1	5	3	4	12	8	20	15	6	6	1	-
Male	-	-	3	3	1	2	4	2	3	1	-	-
Female	1	5	-	1	11	6	16	13	3	5	1	-
White, both sexes	1	4	3	4	12	8	20	15	6	6	1	-
Male	-	-	3	3	1	2	4	2	3	1	-	-
Female	1	4	-	1	11	6	16	13	3	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	2	1	1	2	2	1	-	-	-
Male	-	-	-	2	1	-	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	1	-	-	-
White, both sexes	-	1	-	2	1	1	2	2	1	-	-	-
Male	-	-	-	2	1	-	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
United States												
All races, both sexes	1,503	2	-	-	2	2	2	4	13	13	28	43
Male	469	1	-	-	-	1	2	2	8	6	11	17
Female	1,034	1	-	-	2	1	-	2	5	7	17	26
White, both sexes	1,329	2	-	-	-	1	2	2	7	9	16	31
Male	413	1	-	-	-	1	2	1	4	5	9	12
Female	916	1	-	-	-	-	-	1	3	4	7	19
Black, both sexes	125	-	-	-	1	1	-	1	5	3	9	11
Male	41	-	-	-	-	-	-	1	3	1	2	5
Female	84	-	-	-	1	1	-	-	2	2	7	6
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	1	-	-	1	1	1	3	1
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	27	-	-	-	1	-	-	1	-	1	3	1
Alabama												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Arkansas												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	193	1	-	-	-	-	-	-	2	1	5	6
Male	64	-	-	-	-	-	-	-	-	1	2	1
Female	129	1	-	-	-	-	-	-	2	-	3	5
White, both sexes	162	1	-	-	-	-	-	-	1	1	3	4
Male	55	-	-	-	-	-	-	-	-	1	1	1
Female	107	1	-	-	-	-	-	-	1	-	2	3
Black, both sexes	10	-	-	-	-	-	-	-	1	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Colorado												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	3
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Delaware												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Florida												
All races, both sexes	88	-	-	-	1	-	1	-	1	1	2	1
Male	30	-	-	-	-	-	1	-	1	-	1	-
Female	58	-	-	-	1	-	-	-	-	1	1	1
White, both sexes	79	-	-	-	-	-	1	-	-	1	1	1
Male	27	-	-	-	-	-	1	-	-	-	1	-
Female	52	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	8	-	-	-	1	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	23	-	-	-	-	1	-	1	-	1	2	-
Male	8	-	-	-	-	-	-	1	-	-	1	-
Female	15	-	-	-	-	1	-	-	-	1	1	-
White, both sexes	18	-	-	-	-	-	-	1	-	1	2	-
Male	6	-	-	-	-	-	-	1	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Hawaii												
All races, both sexes	10	-	-	-	-	-	-	1	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	2	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	1	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	2	-
Idaho												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Illinois												
All races, both sexes	42	-	-	-	-	-	1	-	1	2	-	2
Male	17	-	-	-	-	-	1	-	1	1	-	1
Female	25	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	39	-	-	-	-	-	1	-	1	1	-	2
Male	17	-	-	-	-	-	1	-	1	1	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Iowa												
All races, both sexes	28	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	12	-	-	-	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Kentucky												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	18	-	-	-	-	-	-	-	2	-	1	1
Male	8	-	-	-	-	-	-	-	2	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Maine												
All races, both sexes	19	-	-	-	-	-	-	-	1	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	18	-	-	-	-	-	-	-	1	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	1	4
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	19	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Massachusetts												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	48	-	-	-	-	-	-	-	-	1	-	-
Male	22	-	-	-	-	-	-	-	-	1	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Minnesota												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	11	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Missouri												
All races, both sexes	28	-	-	-	-	-	-	-	-	1	1	-
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	1	1	-
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Nebraska												
All races, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	14	-	-	-	1	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
New Hampshire												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
New Mexico												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	81	-	-	-	-	1	-	-	1	-	2	3
Male	29	-	-	-	-	1	-	-	1	-	2	3
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	1	-	-	-	-	2	-
Male	23	-	-	-	-	1	-	-	-	-	2	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	3
Male	5	-	-	-	-	-	-	-	-	-	-	3
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
North Carolina												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	3
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Ohio												
All races, both sexes	75	-	-	-	-	-	-	-	1	1	1	2
Male	27	-	-	-	-	-	-	-	1	-	1	1
Female	48	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	65	-	-	-	-	-	-	-	-	-	1	1
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	42	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	1	1	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Oregon												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	72	-	-	-	-	-	-	-	1	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	68	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Rhode Island												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	2	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
South Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	38	-	-	-	-	-	-	-	-	1	1	1
Male	17	-	-	-	-	-	-	-	-	1	1	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Texas												
All races, both sexes	56	1	-	-	-	-	-	1	-	2	-	1
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	1	-	2	-	1
White, both sexes	53	1	-	-	-	-	-	1	-	1	-	1
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Utah												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	39	-	-	-	-	-	-	-	-	1	3	1
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	29	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	32	-	-	-	-	-	-	-	-	1	1	1
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Washington												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Wisconsin												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
United States												
All races, both sexes	43	86	105	134	168	242	276	217	85	33	5	-
Male	17	33	40	43	62	84	74	52	12	4	-	-
Female	26	53	65	91	106	158	202	165	73	29	5	-
White, both sexes	31	70	86	118	146	222	259	212	82	28	5	-
Male	12	26	35	39	54	77	72	49	12	2	-	-
Female	19	44	51	79	92	145	187	163	70	26	5	-
Black, both sexes	9	10	17	11	16	11	12	2	3	3	-	-
Male	4	4	5	3	6	4	2	-	-	1	-	-
Female	5	6	12	8	10	7	10	2	3	2	-	-
American Indian 1/, both sexes	2	3	-	-	1	3	1	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	3	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	2	5	5	6	4	3	-	2	-	-
Male	-	1	-	1	2	3	-	3	-	1	-	-
Female	1	2	2	4	3	3	4	-	-	1	-	-
Alabama												
All races, both sexes	3	1	3	4	4	4	3	1	1	-	-	-
Male	1	1	-	1	1	2	1	-	-	-	-	-
Female	2	-	3	3	3	2	2	1	1	-	-	-
White, both sexes	1	1	3	3	3	3	3	1	1	-	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	1	-	3	3	2	2	2	1	1	-	-	-
Black, both sexes	1	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Alaska												
All races, both sexes	1	1	-	1	-	1	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	4	4	5	3	4	4	2	-	-	-
Male	-	-	2	2	2	1	-	1	-	-	-	-
Female	-	1	2	2	3	2	4	3	2	-	-	-
White, both sexes	-	1	4	3	4	3	3	4	2	-	-	-
Male	-	-	2	2	2	1	-	1	-	-	-	-
Female	-	1	2	1	2	2	3	3	2	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Arkansas												
All races, both sexes	1	-	1	-	2	-	4	5	-	-	-	-
Male	1	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	1	-	2	-	3	2	-	-	-	-
White, both sexes	-	-	1	-	2	-	4	5	-	-	-	-
Male	-	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	1	-	2	-	3	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	5	14	19	24	29	40	31	9	2	-	-
Male	1	1	6	5	14	9	13	10	1	-	-	-
Female	4	4	8	14	10	20	27	21	8	2	-	-
White, both sexes	3	4	12	13	18	25	39	28	9	1	-	-
Male	1	1	5	5	11	8	13	7	1	-	-	-
Female	2	3	7	8	7	17	26	21	8	1	-	-
Black, both sexes	1	-	1	2	3	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	2	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	4	3	3	1	3	-	1	-	-
Male	-	-	-	-	2	1	-	3	-	-	-	-
Female	1	1	1	4	1	2	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Colorado												
All races, both sexes	-	1	2	4	6	4	6	4	1	-	-	-
Male	-	1	-	-	1	2	1	1	-	-	-	-
Female	-	-	2	4	5	2	5	3	1	-	-	-
White, both sexes	-	1	2	4	5	4	5	4	1	-	-	-
Male	-	1	-	-	1	2	-	1	-	-	-	-
Female	-	-	2	4	4	2	5	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	-	1	1	6	1	4	2	-	-	-
Male	1	1	-	1	-	4	1	1	1	-	-	-
Female	-	-	-	-	1	2	-	3	1	-	-	-
White, both sexes	1	1	-	1	1	4	1	4	2	-	-	-
Male	1	1	-	1	-	3	1	1	1	-	-	-
Female	-	-	-	-	1	1	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Delaware												
All races, both sexes	1	-	-	-	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Florida												
All races, both sexes	3	5	11	10	9	11	17	10	3	2	-	-
Male	1	3	6	4	6	2	4	1	-	-	-	-
Female	2	2	5	6	3	9	13	9	3	2	-	-
White, both sexes	2	4	11	9	8	10	17	9	3	2	-	-
Male	1	2	6	4	5	2	4	1	-	-	-	-
Female	1	2	5	5	3	8	13	8	3	2	-	-
Black, both sexes	1	1	-	1	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	4	5	1	2	4	-	-	1	-	-
Male	-	-	1	3	-	-	1	-	-	1	-	-
Female	-	1	3	2	1	2	3	-	-	-	-	-
White, both sexes	-	1	1	5	1	2	4	-	-	-	-	-
Male	-	-	-	3	-	-	1	-	-	-	-	-
Female	-	1	1	2	1	2	3	-	-	-	-	-
Black, both sexes	-	-	3	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Hawaii												
All races, both sexes	-	1	-	-	-	3	1	1	-	1	-	-
Male	-	1	-	-	-	2	-	-	-	1	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	1	-	-	1	-	-
Male	-	1	-	-	-	2	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	-	1	-	-	-	5	2	2	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	4	2	2	-	1	-	-
White, both sexes	-	1	-	-	-	5	2	2	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	4	2	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Illinois												
All races, both sexes	-	2	2	4	5	7	6	6	4	-	-	-
Male	-	2	1	1	2	1	3	3	-	-	-	-
Female	-	-	1	3	3	6	3	3	4	-	-	-
White, both sexes	-	2	1	4	5	7	5	6	4	-	-	-
Male	-	2	1	1	2	1	3	3	-	-	-	-
Female	-	-	-	3	3	6	2	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	3	1	3	2	7	4	-	1	-	-
Male	1	-	1	1	-	-	2	1	-	-	-	-
Female	-	1	2	-	3	2	5	3	-	1	-	-
White, both sexes	1	-	3	-	3	2	7	4	-	1	-	-
Male	1	-	1	-	-	-	2	1	-	-	-	-
Female	-	-	2	-	3	2	5	3	-	1	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Iowa												
All races, both sexes	1	2	3	1	1	4	6	5	1	2	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	2	3	1	1	2	6	5	1	2	-	-
White, both sexes	1	2	2	1	1	4	6	5	1	2	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	2	2	1	1	2	6	5	1	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	2	1	2	2	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	1	2	2	1	1	-	1	-
White, both sexes	-	-	-	2	1	1	2	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	1	1	2	1	1	-	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Kentucky												
All races, both sexes	-	2	3	5	5	7	3	2	2	-	-	-
Male	-	1	1	2	3	3	2	-	1	-	-	-
Female	-	1	2	3	2	4	1	2	1	-	-	-
White, both sexes	-	2	2	5	5	7	3	2	1	-	-	-
Male	-	1	1	2	3	3	2	-	1	-	-	-
Female	-	1	1	3	2	4	1	2	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	2	1	1	3	2	1	1	1	-	-	-
Male	-	2	-	-	1	1	-	1	1	-	-	-
Female	2	-	1	1	2	1	1	-	-	-	-	-
White, both sexes	2	1	1	1	3	1	-	1	1	-	-	-
Male	-	1	-	-	1	1	-	1	1	-	-	-
Female	2	-	1	1	2	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Maine												
All races, both sexes	1	1	2	4	-	1	5	2	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	1	1	2	3	-	-	3	2	-	-	-	-
White, both sexes	1	1	2	4	-	1	4	2	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	1	1	2	3	-	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	2	4	2	5	3	2	-	-	1	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	1	4	2	3	3	2	-	-	1	-
White, both sexes	-	-	-	3	2	5	3	2	-	-	1	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	3	2	3	3	2	-	-	1	-
Black, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Massachusetts												
All races, both sexes	-	-	2	5	2	5	7	7	3	-	-	-
Male	-	-	1	1	1	2	2	2	-	-	-	-
Female	-	-	1	4	1	3	5	5	3	-	-	-
White, both sexes	-	-	1	5	2	4	7	7	3	-	-	-
Male	-	-	1	1	1	2	2	2	-	-	-	-
Female	-	-	-	4	1	2	5	5	3	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	5	5	5	15	7	6	3	-	-	-
Male	-	1	3	3	1	6	4	2	1	-	-	-
Female	-	-	2	2	4	9	3	4	2	-	-	-
White, both sexes	-	1	3	3	5	14	6	6	3	-	-	-
Male	-	1	2	1	1	6	4	2	1	-	-	-
Female	-	-	1	2	4	8	2	4	2	-	-	-
Black, both sexes	-	-	2	1	-	1	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Minnesota												
All races, both sexes	1	2	2	1	3	6	7	4	2	-	-	-
Male	-	1	-	-	1	2	3	2	-	-	-	-
Female	1	1	2	1	2	4	4	2	2	-	-	-
White, both sexes	1	2	2	1	3	6	7	4	2	-	-	-
Male	-	1	-	-	1	2	3	2	-	-	-	-
Female	1	1	2	1	2	4	4	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	3	-	1	-	2	1	1	1	-	-	-
Male	-	1	-	-	-	1	1	1	-	-	-	-
Female	-	2	-	1	-	1	-	-	1	-	-	-
White, both sexes	-	1	-	1	-	2	-	1	1	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Black, both sexes	-	2	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Missouri												
All races, both sexes	-	1	1	3	7	4	4	3	3	-	-	-
Male	-	-	-	-	3	4	-	1	-	-	-	-
Female	-	1	1	3	4	-	4	2	3	-	-	-
White, both sexes	-	1	1	2	6	4	3	3	3	-	-	-
Male	-	-	-	-	3	4	-	1	-	-	-	-
Female	-	1	1	2	3	-	3	2	3	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Nebraska												
All races, both sexes	-	-	-	-	3	5	3	2	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	3	3	3	2	1	-	-	-
White, both sexes	-	-	-	-	3	5	3	2	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	3	3	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	3	3	2	2	2	1	-	-	-
Male	-	-	-	1	2	1	-	-	1	-	-	-
Female	-	-	-	2	1	1	2	2	-	-	-	-
White, both sexes	-	-	-	3	3	2	1	2	1	-	-	-
Male	-	-	-	1	2	1	-	-	1	-	-	-
Female	-	-	-	2	1	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
New Hampshire												
All races, both sexes	-	1	-	-	1	1	1	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	2	-	1	-
White, both sexes	-	1	-	-	1	1	1	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	1	1	5	6	10	13	5	9	2	-	-
Male	-	-	1	2	-	2	2	-	-	-	-	-
Female	-	1	-	3	6	8	11	5	9	2	-	-
White, both sexes	-	1	-	5	3	10	12	4	7	2	-	-
Male	-	-	-	2	-	2	2	-	-	-	-	-
Female	-	1	-	3	3	8	10	4	7	2	-	-
Black, both sexes	-	-	1	-	3	-	1	1	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	3	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
New Mexico												
All races, both sexes	-	1	-	-	-	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	-	1	-	-
White, both sexes	-	1	-	-	-	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	7	5	3	8	16	9	8	7	4	1	-
Male	4	2	3	-	3	6	1	2	1	-	-	-
Female	2	5	2	3	5	10	8	6	6	4	1	-
White, both sexes	3	7	5	2	7	15	9	8	7	3	1	-
Male	2	2	3	-	3	6	1	2	1	-	-	-
Female	1	5	2	2	4	9	8	6	6	3	1	-
Black, both sexes	3	-	-	1	1	1	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
North Carolina												
All races, both sexes	1	6	4	5	4	6	5	9	3	-	-	-
Male	1	3	2	1	1	1	3	1	3	-	-	-
Female	-	3	2	4	3	5	2	8	-	-	-	-
White, both sexes	1	4	3	4	4	5	5	9	3	-	-	-
Male	1	1	2	1	1	-	3	1	3	-	-	-
Female	-	3	1	3	3	5	2	8	-	-	-	-
Black, both sexes	-	1	1	1	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	1	2	-	2	2	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	2	-	-	-
White, both sexes	-	-	-	1	1	2	-	2	2	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Ohio												
All races, both sexes	5	7	1	4	6	16	13	11	4	3	-	-
Male	1	4	1	1	4	7	2	4	-	-	-	-
Female	4	3	-	3	2	9	11	7	4	3	-	-
White, both sexes	3	5	1	4	6	14	13	11	4	2	-	-
Male	-	4	1	1	4	6	2	4	-	-	-	-
Female	3	1	-	3	2	8	11	7	4	2	-	-
Black, both sexes	2	2	-	-	-	2	-	-	-	1	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	2	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	1	1	2	3	4	2	-	-	-	-
Male	-	1	-	-	-	1	1	2	-	-	-	-
Female	-	1	1	1	2	2	3	-	-	-	-	-
White, both sexes	-	1	1	1	2	2	4	2	-	-	-	-
Male	-	1	-	-	-	1	1	2	-	-	-	-
Female	-	-	1	1	2	1	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Oregon												
All races, both sexes	-	2	2	1	3	4	8	5	3	1	-	-
Male	-	1	-	-	1	1	1	1	-	-	-	-
Female	-	1	2	1	2	3	7	4	3	1	-	-
White, both sexes	-	2	2	1	3	4	8	5	3	1	-	-
Male	-	1	-	-	1	1	1	1	-	-	-	-
Female	-	1	2	1	2	3	7	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	2	3	2	8	11	16	19	5	2	-	-
Male	1	-	2	1	4	4	5	2	1	-	-	-
Female	1	2	1	1	4	7	11	17	4	2	-	-
White, both sexes	2	2	3	2	7	11	15	19	5	2	-	-
Male	1	-	2	1	3	4	5	2	1	-	-	-
Female	1	2	1	1	4	7	10	17	4	2	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Rhode Island												
All races, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	2	1	5	6	2	3	-	-	1	-	-
Male	-	2	-	3	4	2	-	-	-	-	-	-
Female	2	-	1	2	2	-	3	-	-	1	-	-
White, both sexes	2	1	1	5	2	2	1	-	-	1	-	-
Male	-	1	-	3	1	2	-	-	-	-	-	-
Female	2	-	1	2	1	-	1	-	-	1	-	-
Black, both sexes	-	1	-	-	4	-	2	-	-	-	-	-
Male	-	1	-	-	3	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
South Dakota												
All races, both sexes	-	-	-	-	2	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	1	-	-
Female	-	-	-	-	2	1	1	1	-	-	-	-
White, both sexes	-	-	-	-	2	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	1	-	-
Female	-	-	-	-	2	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	4	4	3	4	5	10	2	1	1	-	-
Male	1	1	1	1	2	2	4	1	-	1	-	-
Female	-	3	3	2	2	3	6	1	1	-	-	-
White, both sexes	1	3	3	3	4	5	10	2	1	1	-	-
Male	1	1	1	1	2	2	4	1	-	1	-	-
Female	-	2	2	2	2	3	6	1	1	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Texas												
All races, both sexes	2	6	7	6	5	6	8	7	3	1	-	-
Male	1	1	3	1	2	2	-	-	-	-	-	-
Female	1	5	4	5	3	4	8	7	3	1	-	-
White, both sexes	2	6	6	6	5	6	7	7	3	1	-	-
Male	1	1	3	1	2	2	-	-	-	-	-	-
Female	1	5	3	5	3	4	7	7	3	1	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	2	2	-	1	1	3	3	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	2	2	-	1	-	2	2	-	-	-	-
White, both sexes	-	2	2	-	1	1	3	3	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	2	2	-	1	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Vermont												
All races, both sexes	-	-	-	1	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	2	5	2	4	5	8	8	-	-	-	-
Male	-	-	2	1	-	-	3	2	-	-	-	-
Female	-	2	3	1	4	5	5	6	-	-	-	-
White, both sexes	-	2	3	2	3	4	7	8	-	-	-	-
Male	-	-	2	1	-	-	3	2	-	-	-	-
Female	-	2	1	1	3	4	4	6	-	-	-	-
Black, both sexes	-	-	2	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Washington												
All races, both sexes	2	-	-	1	3	5	8	10	2	4	1	-
Male	1	-	-	-	2	2	2	3	-	-	-	-
Female	1	-	-	1	1	3	6	7	2	4	1	-
White, both sexes	1	-	-	1	3	5	8	10	2	4	1	-
Male	1	-	-	-	2	2	2	3	-	-	-	-
Female	-	-	-	1	1	3	6	7	2	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	2	1	1	2	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	2	1	1	2	-	-	-	-	-	-
White, both sexes	-	1	2	1	1	2	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	2	1	1	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Wisconsin												
All races, both sexes	1	4	1	3	5	4	9	9	2	2	-	-
Male	-	-	1	2	1	1	4	1	-	-	-	-
Female	1	4	-	1	4	3	5	8	2	2	-	-
White, both sexes	1	3	1	3	5	4	9	9	2	2	-	-
Male	-	-	1	2	1	1	4	1	-	-	-	-
Female	1	3	-	1	4	3	5	8	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	1	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
United States												
All races, both sexes	93	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	91	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Colorado												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Florida												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Illinois												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Iowa												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Missouri												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Ohio												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Oregon												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Texas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Washington												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
United States												
All races, both sexes	-	5	1	8	10	7	23	21	10	7	-	-
Male	-	3	1	6	4	3	10	7	2	3	-	-
Female	-	2	-	2	6	4	13	14	8	4	-	-
White, both sexes	-	5	1	8	10	6	23	20	10	7	-	-
Male	-	3	1	6	4	2	10	7	2	3	-	-
Female	-	2	-	2	6	4	13	13	8	4	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	2	-	1	2	1	1	-	-	-
Male	-	-	-	2	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	2	-	1	2	1	1	-	-	-
Male	-	-	-	2	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Colorado												
All races, both sexes	-	-	-	-	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Delaware												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Florida												
All races, both sexes	-	-	-	-	-	1	4	2	2	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	2	-	-	-
White, both sexes	-	-	-	-	-	-	4	2	2	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	-	-	-
White, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Illinois												
All races, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Iowa												
All races, both sexes	-	-	-	1	1	-	2	2	1	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	-	1	1	-	2	2	1	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Missouri												
All races, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	2	1	-	3	2	1	1	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	1	1	-	2	1	1	-	-	-
White, both sexes	-	-	-	2	1	-	3	2	1	1	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	1	1	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
North Carolina												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Ohio												
All races, both sexes	-	-	-	-	-	1	2	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
White, both sexes	-	-	-	-	-	1	2	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Oregon												
All races, both sexes	-	-	-	-	1	-	1	2	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	1	-	-	2	1	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	1	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	-	-	-	3	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	3	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
South Dakota												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Texas												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	2	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
United States												
All races, both sexes	690	1	-	-	-	-	2	2	4	3	5	10
Male	240	1	-	-	-	-	1	1	4	2	3	7
Female	450	-	-	-	-	-	1	1	-	1	2	3
White, both sexes	632	1	-	-	-	-	-	2	4	1	4	10
Male	218	1	-	-	-	-	-	1	4	1	3	7
Female	414	-	-	-	-	-	-	1	-	-	1	3
Black, both sexes	42	-	-	-	-	-	-	-	-	2	1	-
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	26	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Arkansas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	94	-	-	-	-	-	-	1	-	-	2	2
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	58	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	81	-	-	-	-	-	-	1	-	-	2	2
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	50	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Colorado												
All races, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Delaware												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Florida												
All races, both sexes	46	-	-	-	-	-	-	-	1	-	-	2
Male	21	-	-	-	-	-	-	-	1	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	1	-	-	2
Male	19	-	-	-	-	-	-	-	1	-	-	2
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Hawaii												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Illinois												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Iowa												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	12	1	-	-	-	-	-	-	1	-	-	-
Male	5	1	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	-	1	-	-	-
Male	4	1	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Kentucky												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Massachusetts												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Minnesota												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Missouri												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Nebraska												
All races, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
New Hampshire												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	36	-	-	-	-	-	-	-	1	1	-	-
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	23	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	33	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
New Mexico												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
North Carolina												
All races, both sexes	25	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	20	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Ohio												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Oregon												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Rhode Island												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Texas												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Washington												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Wisconsin												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
United States												
All races, both sexes	12	21	29	32	71	102	137	129	90	32	8	-
Male	7	10	14	12	21	37	49	41	22	8	-	-
Female	5	11	15	20	50	65	88	88	68	24	8	-
White, both sexes	9	17	23	22	63	90	130	128	89	31	8	-
Male	5	8	10	8	18	34	47	41	22	8	-	-
Female	4	9	13	14	45	56	83	87	67	23	8	-
Black, both sexes	2	4	4	7	4	10	5	1	1	1	-	-
Male	1	2	3	3	2	3	1	-	-	-	-	-
Female	1	2	1	4	2	7	4	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	3	4	2	2	-	-	-	-	-
Male	1	-	1	1	1	-	1	-	-	-	-	-
Female	-	-	1	2	3	2	1	-	-	-	-	-
Alabama												
All races, both sexes	-	-	1	1	1	4	2	1	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	1	1	2	2	1	1	-	-	-
White, both sexes	-	-	1	1	1	3	2	1	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	1	1	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	2	-	1	-	5	3	1	-	-	-	-
Male	1	1	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	1	-	4	2	1	-	-	-	-
White, both sexes	1	2	-	1	-	5	3	1	-	-	-	-
Male	1	1	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	1	-	4	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Arkansas												
All races, both sexes	-	-	-	1	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	2	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	3	5	7	8	10	20	16	15	4	-	-
Male	1	2	2	4	1	4	6	6	6	2	-	-
Female	-	1	3	3	7	6	14	10	9	2	-	-
White, both sexes	1	3	3	3	6	7	19	16	15	3	-	-
Male	1	2	1	2	1	2	6	6	6	2	-	-
Female	-	1	2	1	5	5	13	10	9	1	-	-
Black, both sexes	-	-	-	2	-	2	1	-	-	1	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	2	1	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	2	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Colorado												
All races, both sexes	-	-	-	2	-	3	1	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	2	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	2	-	3	1	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	2	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	1	3	2	1	2	-	-	-	-
Male	-	-	-	1	2	1	1	2	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	1	1	2	2	1	2	-	-	-	-
Male	-	-	-	1	2	1	1	2	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Delaware												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Florida												
All races, both sexes	-	1	2	3	7	10	7	8	5	-	-	-
Male	-	-	1	-	4	4	4	4	1	-	-	-
Female	-	1	1	3	3	6	3	4	4	-	-	-
White, both sexes	-	1	1	3	6	9	7	8	5	-	-	-
Male	-	-	-	-	3	4	4	4	1	-	-	-
Female	-	1	1	3	3	5	3	4	4	-	-	-
Black, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	2	2	2	2	2	2	2	2	1	-
Male	-	-	-	2	-	1	-	1	2	-	-	-
Female	1	-	2	-	2	1	2	1	-	2	1	-
White, both sexes	-	-	2	1	2	2	-	2	2	2	1	-
Male	-	-	-	1	-	1	-	1	2	-	-	-
Female	-	-	2	-	2	1	-	1	-	2	1	-
Black, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Hawaii												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	1	-	-	-	-	1	-	3	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	3	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Illinois												
All races, both sexes	1	1	-	1	1	4	8	8	-	-	-	-
Male	-	1	-	-	1	4	3	-	-	-	-	-
Female	1	-	-	1	-	-	5	8	-	-	-	-
White, both sexes	1	1	-	-	-	4	7	7	-	-	-	-
Male	-	1	-	-	-	4	2	-	-	-	-	-
Female	1	-	-	-	-	-	5	7	-	-	-	-
Black, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	1	3	1	1	3	6	1	-	-
Male	-	-	-	1	-	-	-	1	2	-	-	-
Female	-	-	-	-	3	1	1	2	4	1	-	-
White, both sexes	-	-	-	1	3	1	1	3	6	1	-	-
Male	-	-	-	1	-	-	-	1	2	-	-	-
Female	-	-	-	-	3	1	1	2	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Iowa												
All races, both sexes	-	-	-	-	1	1	3	3	1	1	1	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	1	1	1	-
White, both sexes	-	-	-	-	1	1	3	3	1	1	1	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	1	-	-	1	5	2	1	-	-	-
Male	-	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	3	2	1	-	-	-
White, both sexes	-	-	1	-	-	1	4	2	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Kentucky												
All races, both sexes	-	-	-	1	-	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	3	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	1	-	-
White, both sexes	-	-	-	1	-	-	3	1	-	1	-	-
Male	-	-	-	-	-	-	3	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	1	1	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Maine												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	3	1	-	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	3	1	-	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Massachusetts												
All races, both sexes	-	-	-	-	1	1	3	3	5	1	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	5	1	-	-
White, both sexes	-	-	-	-	1	1	3	3	5	1	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	2	1	-	5	2	3	6	-	-	-	-
Male	-	1	-	-	1	1	1	1	-	-	-	-
Female	-	1	1	-	4	1	2	5	-	-	-	-
White, both sexes	-	1	1	-	5	1	3	6	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	1	1	-	4	1	2	5	-	-	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Minnesota												
All races, both sexes	-	-	-	1	1	2	4	4	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	1	4	3	1	-	-	-
White, both sexes	-	-	-	1	1	2	4	4	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	1	4	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	-	-	1	2	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Missouri												
All races, both sexes	-	-	-	-	3	2	3	3	5	-	-	-
Male	-	-	-	-	1	2	1	-	1	-	-	-
Female	-	-	-	-	2	-	2	3	4	-	-	-
White, both sexes	-	-	-	-	3	2	3	3	5	-	-	-
Male	-	-	-	-	1	2	1	-	1	-	-	-
Female	-	-	-	-	2	-	2	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Nebraska												
All races, both sexes	-	1	-	1	-	1	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	1	1	-
White, both sexes	-	1	-	1	-	1	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	3	1	4	6	8	5	3	2	-	-
Male	-	-	3	-	1	1	5	2	-	-	-	-
Female	2	-	-	1	3	5	3	3	3	2	-	-
White, both sexes	2	-	2	1	4	5	8	5	3	2	-	-
Male	-	-	2	-	1	1	5	2	-	-	-	-
Female	2	-	-	1	3	4	3	3	3	2	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
New Mexico												
All races, both sexes	-	-	1	1	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	1	9	7	3	6	2	-	-
Male	-	-	-	-	-	3	4	1	-	-	-	-
Female	-	-	-	-	1	6	3	2	6	2	-	-
White, both sexes	-	-	-	-	1	8	6	3	6	2	-	-
Male	-	-	-	-	-	3	4	1	-	-	-	-
Female	-	-	-	-	1	5	2	2	6	2	-	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
North Carolina												
All races, both sexes	-	3	-	1	1	3	3	6	6	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	2	-	1	1	3	3	6	5	-	-	-
White, both sexes	-	2	-	-	-	3	3	6	5	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	3	3	6	4	-	-	-
Black, both sexes	-	1	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	1	2	2	1	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	2	1	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Ohio												
All races, both sexes	1	1	1	-	2	2	8	6	4	2	1	-
Male	-	-	-	-	1	1	2	1	1	-	-	-
Female	1	1	1	-	1	1	6	5	3	2	1	-
White, both sexes	1	1	1	-	2	2	7	6	4	2	1	-
Male	-	-	-	-	1	1	2	1	1	-	-	-
Female	1	1	1	-	1	1	5	5	3	2	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	2	1	-	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	1	2	1	-	1	1	1	-	1	-	-
White, both sexes	-	-	2	-	-	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	2	-	-	1	1	1	-	1	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Oregon												
All races, both sexes	-	-	-	-	-	1	2	4	6	1	-	-
Male	-	-	-	-	-	-	-	1	1	1	-	-
Female	-	-	-	-	-	1	2	3	5	-	-	-
White, both sexes	-	-	-	-	-	1	2	4	6	1	-	-
Male	-	-	-	-	-	-	-	1	1	1	-	-
Female	-	-	-	-	-	1	2	3	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	2	1	4	5	12	8	7	4	1	-
Male	-	1	1	1	-	1	3	3	2	1	-	-
Female	-	-	1	-	4	4	9	5	5	3	1	-
White, both sexes	-	1	1	1	4	5	12	8	7	4	1	-
Male	-	1	1	1	-	1	3	3	2	1	-	-
Female	-	-	-	-	4	4	9	5	5	3	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Rhode Island												
All races, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	2	3	1	1	1	1	-	-	-
Male	-	-	1	1	2	-	-	1	-	-	-	-
Female	-	-	-	1	1	1	1	-	1	-	-	-
White, both sexes	-	-	1	-	2	1	1	1	1	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	-	-	1	-	3	-	2	-	-	-
Male	-	-	-	-	1	-	2	-	2	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	1	-	3	-	2	-	-	-
Male	-	-	-	-	1	-	2	-	2	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Texas												
All races, both sexes	2	-	-	-	6	7	3	7	3	1	1	-
Male	2	-	-	-	2	1	1	2	1	1	-	-
Female	-	-	-	-	4	6	2	5	2	-	1	-
White, both sexes	2	-	-	-	6	6	3	7	3	1	1	-
Male	2	-	-	-	2	1	1	2	1	1	-	-
Female	-	-	-	-	4	5	2	5	2	-	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	1	1	2	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	1	1	1	2	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Vermont												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	2	1	1	3	1	-	-	-
Male	-	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	-	-	2	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	1	1	3	1	-	-	-
Male	-	-	-	-	-	-	1	3	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Washington												
All races, both sexes	2	-	2	-	2	1	3	11	-	2	1	-
Male	2	-	1	-	2	-	1	7	-	1	-	-
Female	-	-	1	-	-	1	2	4	-	1	1	-
White, both sexes	-	-	2	-	2	1	3	11	-	2	1	-
Male	-	-	1	-	2	-	1	7	-	1	-	-
Female	-	-	1	-	-	1	2	4	-	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Wisconsin												
All races, both sexes	-	-	2	-	4	1	5	1	2	3	1	-
Male	-	-	2	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	4	1	5	1	1	2	1	-
White, both sexes	-	-	2	-	4	1	5	1	2	3	1	-
Male	-	-	2	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	4	1	5	1	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
United States												
All races, both sexes	946	-	-	1	2	1	-	3	4	5	11	29
Male	251	-	-	1	1	1	-	1	1	3	3	15
Female	695	-	-	-	1	-	-	2	3	2	8	14
White, both sexes	833	-	-	-	1	-	-	3	2	3	8	19
Male	211	-	-	-	-	-	-	1	-	2	2	12
Female	622	-	-	-	1	-	-	2	2	1	6	7
Black, both sexes	75	-	-	-	1	1	-	-	-	2	3	10
Male	28	-	-	-	1	1	-	-	-	1	1	3
Female	47	-	-	-	-	-	-	-	-	1	2	7
American Indian 1/, both sexes	8	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	2	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	-	1	-	-	-
Alabama												
All races, both sexes	12	-	-	-	-	-	-	-	1	1	1	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	10	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Arkansas												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	97	-	-	1	-	-	-	-	2	-	2	6
Male	30	-	-	1	-	-	-	-	1	-	-	1
Female	67	-	-	-	-	-	-	-	1	-	2	5
White, both sexes	81	-	-	-	-	-	-	-	1	-	1	6
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	57	-	-	-	-	-	-	-	1	-	1	5
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Colorado												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Florida												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	-	4
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	44	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	3
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Hawaii												
All races, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
Idaho												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Illinois												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Iowa												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	23	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Kentucky												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	7	-	-	-	-	1	-	-	-	-	1	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Maine												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Massachusetts												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Minnesota												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Missouri												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Nebraska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Mexico												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	59	-	-	-	1	-	-	1	-	-	2	2
Male	17	-	-	-	1	-	-	-	-	-	1	2
Female	42	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	51	-	-	-	-	-	-	1	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	40	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	5	-	-	-	1	-	-	-	-	-	-	1
Male	5	-	-	-	1	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
North Carolina												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Ohio												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	3
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	41	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Oregon												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Rhode Island												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	25	-	-	-	1	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	18	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	1	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Texas												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	19	-	-	-	-	-	-	1	-	1	1	1
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	15	-	-	-	-	-	-	1	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Washington												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Wisconsin												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
United States												
All races, both sexes	35	44	56	82	101	148	180	144	72	23	5	-
Male	14	15	17	31	29	34	34	34	14	2	1	-
Female	21	29	39	51	72	114	146	110	58	21	4	-
White, both sexes	21	32	41	71	89	139	169	139	69	22	5	-
Male	8	9	12	26	25	32	33	34	12	2	1	-
Female	13	23	29	45	64	107	136	105	57	20	4	-
Black, both sexes	6	8	9	9	5	6	9	3	3	-	-	-
Male	2	4	5	4	1	2	1	-	2	-	-	-
Female	4	4	4	5	4	4	8	3	1	-	-	-
American Indian 1/, both sexes	1	-	3	1	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	3	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	4	3	1	7	2	1	2	-	1	-	-
Male	3	2	-	1	3	-	-	-	-	-	-	-
Female	4	2	3	-	4	2	1	2	-	1	-	-
Alabama												
All races, both sexes	1	1	1	-	2	1	2	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	1	-	1	-	2	-	-	-	-	-
White, both sexes	1	1	1	-	2	1	2	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	1	-	1	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Alaska												
All races, both sexes	1	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	1	-	2	2	6	1	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	2	2	5	1	-	1	-	-	-
White, both sexes	1	1	-	2	2	6	1	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	2	2	5	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Arkansas												
All races, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	5	7	9	9	15	16	13	4	2	-	-
Male	2	3	2	5	-	3	5	6	1	-	-	-
Female	4	2	5	4	9	12	11	7	3	2	-	-
White, both sexes	2	2	5	8	8	14	15	13	4	2	-	-
Male	1	1	1	5	-	3	5	6	1	-	-	-
Female	1	1	4	3	8	11	10	7	3	2	-	-
Black, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	1	-	1	-	1	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	3	1	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Colorado												
All races, both sexes	1	2	1	1	4	2	3	2	3	1	-	-
Male	1	-	-	-	1	-	2	1	-	-	-	-
Female	-	2	1	1	3	2	1	1	3	1	-	-
White, both sexes	1	2	-	1	4	2	3	2	3	1	-	-
Male	1	-	-	-	1	-	2	1	-	-	-	-
Female	-	2	-	1	3	2	1	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	-	1	3	4	3	4	-	-	-
Male	-	-	-	-	1	2	-	-	2	-	-	-
Female	-	1	-	-	-	1	4	3	2	-	-	-
White, both sexes	-	-	-	-	1	3	3	3	3	-	-	-
Male	-	-	-	-	1	2	-	-	1	-	-	-
Female	-	-	-	-	-	1	3	3	2	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Delaware												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Florida												
All races, both sexes	1	4	4	6	5	11	15	10	2	1	1	-
Male	-	1	2	3	3	3	3	2	-	1	-	-
Female	1	3	2	3	2	8	12	8	2	-	1	-
White, both sexes	-	2	4	5	4	11	13	8	2	1	1	-
Male	-	-	2	3	3	3	3	2	-	1	-	-
Female	-	2	2	2	1	8	10	6	2	-	1	-
Black, both sexes	1	2	-	1	1	-	2	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	-	1	1	4	1	3	1	1	-	-	-	-
Male	-	1	1	2	-	1	-	1	-	-	-	-
Female	-	-	-	2	1	2	1	-	-	-	-	-
White, both sexes	-	1	-	2	-	2	1	1	-	-	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Black, both sexes	-	-	1	2	1	1	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Hawaii												
All races, both sexes	3	1	-	-	2	2	-	1	-	-	-	-
Male	2	-	-	-	1	-	-	1	-	-	-	-
Female	1	1	-	-	1	2	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	2	2	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	1	2	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	1	-	2	2	-	2	1	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	2	2	-	1	1	-	-	-
White, both sexes	1	-	1	-	2	2	-	2	1	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	2	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Illinois												
All races, both sexes	-	1	1	2	6	1	7	1	4	1	-	-
Male	-	-	1	1	3	-	-	-	1	1	-	-
Female	-	1	-	1	3	1	7	1	3	-	-	-
White, both sexes	-	1	1	2	6	1	6	1	4	1	-	-
Male	-	-	1	1	3	-	-	-	1	1	-	-
Female	-	1	-	1	3	1	6	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	3	1	1	3	-	3	1	-	-	-
Male	-	-	2	-	-	-	-	1	1	-	-	-
Female	1	1	1	1	1	3	-	2	-	-	-	-
White, both sexes	1	1	2	1	-	3	-	3	1	-	-	-
Male	-	-	1	-	-	-	-	1	1	-	-	-
Female	1	1	1	1	-	3	-	2	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Iowa												
All races, both sexes	-	-	1	1	1	2	3	5	5	3	-	-
Male	-	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	1	1	-	1	3	4	4	3	-	-
White, both sexes	-	-	1	1	1	2	3	5	5	3	-	-
Male	-	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	1	1	-	1	3	4	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Kentucky												
All races, both sexes	-	-	2	2	-	1	4	3	-	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	4	3	-	-	-	-
White, both sexes	-	-	2	2	-	1	4	3	-	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	4	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	1	1	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Maine												
All races, both sexes	1	-	-	1	-	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	1	2	1	2	6	3	-	1	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	1	1	1	1	1	4	3	-	1	-	-
White, both sexes	-	1	-	2	1	2	6	3	-	1	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	1	-	1	1	1	4	3	-	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Massachusetts												
All races, both sexes	-	1	2	2	3	8	5	5	1	1	-	-
Male	-	1	1	-	3	3	1	1	-	-	-	-
Female	-	-	1	2	-	5	4	4	1	1	-	-
White, both sexes	-	1	2	2	3	8	5	5	1	1	-	-
Male	-	1	1	-	3	3	1	1	-	-	-	-
Female	-	-	1	2	-	5	4	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	3	-	2	3	7	5	4	-	-	-
Male	-	-	-	-	-	-	2	2	2	-	-	-
Female	-	-	3	-	2	3	5	3	2	-	-	-
White, both sexes	-	-	2	-	2	3	6	5	4	-	-	-
Male	-	-	-	-	-	-	1	2	2	-	-	-
Female	-	-	2	-	2	3	5	3	2	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Minnesota												
All races, both sexes	1	2	-	7	2	5	5	5	3	1	-	-
Male	-	1	-	3	-	3	1	1	-	-	-	-
Female	1	1	-	4	2	2	4	4	3	1	-	-
White, both sexes	1	2	-	7	2	5	5	5	3	1	-	-
Male	-	1	-	3	-	3	1	1	-	-	-	-
Female	1	1	-	4	2	2	4	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	-	1	-	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	-	1	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Missouri												
All races, both sexes	1	-	1	4	1	5	5	4	2	1	-	-
Male	-	-	1	2	1	3	-	1	-	-	-	-
Female	1	-	-	2	-	2	5	3	2	1	-	-
White, both sexes	1	-	1	4	1	5	5	4	2	1	-	-
Male	-	-	1	2	1	3	-	1	-	-	-	-
Female	1	-	-	2	-	2	5	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	1	-	1	-	-
White, both sexes	-	-	-	-	-	1	3	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	1	3	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	1	-
White, both sexes	-	-	-	-	-	-	1	3	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	1	2	3	1	5	9	6	-	-	1	-
Male	-	-	-	2	1	1	2	2	-	-	-	-
Female	-	1	2	1	-	4	7	4	-	-	1	-
White, both sexes	-	1	2	3	1	5	9	5	-	-	1	-
Male	-	-	-	2	1	1	2	2	-	-	-	-
Female	-	1	2	1	-	4	7	3	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Mexico												
All races, both sexes	-	1	1	1	2	-	3	1	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	3	1	1	-	-	-
White, both sexes	-	1	1	-	2	-	2	1	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	4	5	4	7	5	12	8	6	2	-	-
Male	-	2	1	2	3	-	3	2	-	-	-	-
Female	-	2	4	2	4	5	9	6	6	2	-	-
White, both sexes	-	4	3	3	5	5	12	8	6	1	-	-
Male	-	2	-	1	1	-	3	2	-	-	-	-
Female	-	2	3	2	4	5	9	6	6	1	-	-
Black, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
North Carolina												
All races, both sexes	2	2	1	1	1	3	7	3	1	-	-	-
Male	-	1	-	1	-	1	4	2	1	-	-	-
Female	2	1	1	-	1	2	3	1	-	-	-	-
White, both sexes	2	-	1	-	1	2	5	3	-	-	-	-
Male	-	-	-	-	-	-	4	2	-	-	-	-
Female	2	-	1	-	1	2	1	1	-	-	-	-
Black, both sexes	-	2	-	1	-	1	2	-	1	-	-	-
Male	-	1	-	1	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Ohio												
All races, both sexes	2	3	-	3	9	17	3	6	2	1	-	-
Male	1	2	-	-	2	1	-	-	-	-	-	-
Female	1	1	-	3	7	16	3	6	2	1	-	-
White, both sexes	2	2	-	3	8	17	3	6	2	1	-	-
Male	1	1	-	-	2	1	-	-	-	-	-	-
Female	1	1	-	3	6	16	3	6	2	1	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	2	1	-	2	2	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	-	2	2	1	-	1	-	-	-
White, both sexes	1	1	1	-	2	2	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	2	2	1	-	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Oregon												
All races, both sexes	-	1	1	-	4	3	5	5	4	4	-	-
Male	-	-	1	-	1	-	2	1	-	-	-	-
Female	-	1	-	-	3	3	3	4	4	4	-	-
White, both sexes	-	1	1	-	2	3	5	5	4	4	-	-
Male	-	-	1	-	-	-	2	1	-	-	-	-
Female	-	1	-	-	2	3	3	4	4	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	1	3	4	2	10	10	8	4	-	1	-
Male	-	-	1	-	-	1	1	1	-	-	-	-
Female	2	1	2	4	2	9	9	7	4	-	1	-
White, both sexes	1	1	2	4	2	9	10	7	4	-	1	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	1	1	2	4	2	8	9	6	4	-	1	-
Black, both sexes	1	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Rhode Island												
All races, both sexes	-	1	-	2	-	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	2	1	-	-	-	-
White, both sexes	-	1	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	2	1	2	-	2	2	-	2	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	2	1	1	-	2	2	-	2	-	-
White, both sexes	-	1	2	1	2	-	2	2	-	2	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	2	1	1	-	2	2	-	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
South Dakota												
All races, both sexes	-	-	2	1	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	3	2	1	1	5	6	2	-	-	-
Male	1	-	-	-	-	1	2	1	-	-	-	-
Female	1	-	3	2	1	-	3	5	2	-	-	-
White, both sexes	2	-	3	2	1	1	5	6	1	-	-	-
Male	1	-	-	-	-	1	2	1	-	-	-	-
Female	1	-	3	2	1	-	3	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Texas												
All races, both sexes	3	2	2	3	7	2	4	2	4	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	2	2	2	1	7	2	4	2	4	-	-	-
White, both sexes	1	2	-	1	6	2	4	2	4	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	2	-	-	6	2	4	2	4	-	-	-
Black, both sexes	2	-	1	2	1	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	3	2	3	5	2	2	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	3	1	3	4	2	2	-	-	-
White, both sexes	-	-	-	3	2	3	5	2	2	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	3	1	3	4	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Vermont												
All races, both sexes	-	-	-	-	1	-	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	2	1	1	5	4	-	-	1	-	-
Male	-	-	1	-	-	3	1	-	-	-	-	-
Female	1	-	1	1	1	2	3	-	-	1	-	-
White, both sexes	1	-	2	1	1	4	3	-	-	1	-	-
Male	-	-	1	-	-	3	1	-	-	-	-	-
Female	1	-	1	1	1	1	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Washington												
All races, both sexes	-	1	-	3	6	5	5	13	3	-	-	-
Male	-	1	-	2	4	1	-	2	2	-	-	-
Female	-	-	-	1	2	4	5	11	1	-	-	-
White, both sexes	-	1	-	3	6	5	5	12	3	-	-	-
Male	-	1	-	2	4	1	-	2	2	-	-	-
Female	-	-	-	1	2	4	5	10	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	3	1	1	3	-	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	2	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	3	1	1	3	-	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	2	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Wisconsin												
All races, both sexes	-	1	-	1	3	3	6	5	2	-	-	-
Male	-	-	-	1	-	1	-	1	2	-	-	-
Female	-	1	-	-	3	2	6	4	-	-	-	-
White, both sexes	-	1	-	1	3	3	6	5	2	-	-	-
Male	-	-	-	1	-	1	-	1	2	-	-	-
Female	-	1	-	-	3	2	6	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
United States												
All races, both sexes	56,561	1	1	2	3	12	67	138	291	606	1,218	2,104
Male	24,382	1	-	1	2	5	51	101	200	435	789	1,435
Female	32,179	-	1	1	1	7	16	37	91	171	429	669
White, both sexes	42,469	-	1	1	2	5	39	62	135	336	669	1,223
Male	17,581	-	-	1	1	3	29	46	101	259	472	891
Female	24,888	-	1	-	1	2	10	16	34	77	197	332
Black, both sexes	12,569	1	-	1	1	6	25	69	150	259	508	837
Male	6,089	1	-	-	1	2	20	49	94	169	285	512
Female	6,480	-	-	1	-	4	5	20	56	90	223	325
American Indian 1/, both sexes	256	-	-	-	-	-	1	1	4	3	17	21
Male	140	-	-	-	-	-	1	-	3	3	13	15
Female	116	-	-	-	-	-	-	1	1	-	4	6
Asian or Pacific Islander 2/, both sexes	1,267	-	-	-	-	1	2	6	2	8	24	23
Male	572	-	-	-	-	-	1	6	2	4	19	17
Female	695	-	-	-	-	1	1	-	-	4	5	6
Alabama												
All races, both sexes	894	-	-	-	-	-	1	1	12	10	24	37
Male	379	-	-	-	-	-	-	1	9	6	9	19
Female	515	-	-	-	-	-	1	-	3	4	15	18
White, both sexes	506	-	-	-	-	-	1	-	3	7	3	13
Male	203	-	-	-	-	-	-	-	1	6	1	7
Female	303	-	-	-	-	-	1	-	2	1	2	6
Black, both sexes	386	-	-	-	-	-	-	1	9	3	21	24
Male	174	-	-	-	-	-	-	1	8	-	8	12
Female	212	-	-	-	-	-	-	-	1	3	13	12
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Alaska												
All races, both sexes	65	-	-	-	-	-	-	-	1	3	1	3
Male	33	-	-	-	-	-	-	-	-	2	-	3
Female	32	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	38	-	-	-	-	-	-	-	-	2	-	2
Male	19	-	-	-	-	-	-	-	-	1	-	2
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	965	-	-	-	-	2	1	3	2	8	18	27
Male	442	-	-	-	-	2	1	2	1	6	8	21
Female	523	-	-	-	-	-	-	1	1	2	10	6
White, both sexes	868	-	-	-	-	-	-	3	2	7	13	21
Male	386	-	-	-	-	-	-	2	1	5	6	17
Female	482	-	-	-	-	-	-	1	1	2	7	4
Black, both sexes	49	-	-	-	-	2	-	-	-	1	2	3
Male	30	-	-	-	-	2	-	-	-	1	1	1
Female	19	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	32	-	-	-	-	-	1	-	-	-	3	3
Male	18	-	-	-	-	-	1	-	-	-	1	3
Female	14	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Arkansas												
All races, both sexes	582	-	-	-	-	-	1	1	5	5	21	22
Male	250	-	-	-	-	-	1	-	3	3	13	17
Female	332	-	-	-	-	-	-	1	2	2	8	5
White, both sexes	428	-	-	-	-	-	-	-	2	2	13	10
Male	167	-	-	-	-	-	-	-	2	2	7	8
Female	261	-	-	-	-	-	-	-	-	-	6	2
Black, both sexes	152	-	-	-	-	-	1	1	3	3	8	11
Male	81	-	-	-	-	-	1	-	1	1	6	8
Female	71	-	-	-	-	-	-	1	2	2	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,593	-	1	-	-	1	7	12	23	80	138	235
Male	3,318	-	-	-	-	-	6	9	15	59	92	164
Female	4,275	-	1	-	-	1	1	3	8	21	46	71
White, both sexes	5,880	-	1	-	-	1	4	8	17	52	96	142
Male	2,558	-	-	-	-	-	4	5	12	39	66	105
Female	3,322	-	1	-	-	1	-	3	5	13	30	37
Black, both sexes	1,029	-	-	-	-	-	2	2	5	21	30	81
Male	466	-	-	-	-	-	2	2	2	17	17	49
Female	563	-	-	-	-	-	-	-	3	4	13	32
American Indian 1/, both sexes	44	-	-	-	-	-	-	-	-	1	4	3
Male	32	-	-	-	-	-	-	-	-	1	3	3
Female	12	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	640	-	-	-	-	-	1	2	1	6	8	9
Male	262	-	-	-	-	-	-	2	1	2	6	7
Female	378	-	-	-	-	-	1	-	-	4	2	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Colorado												
All races, both sexes	488	-	-	-	-	-	-	-	3	3	8	11
Male	199	-	-	-	-	-	-	-	1	1	6	10
Female	289	-	-	-	-	-	-	-	2	2	2	1
White, both sexes	443	-	-	-	-	-	-	-	2	3	6	11
Male	180	-	-	-	-	-	-	-	1	1	5	10
Female	263	-	-	-	-	-	-	-	1	2	1	1
Black, both sexes	37	-	-	-	-	-	-	-	1	-	2	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	546	-	-	-	-	-	-	-	2	4	6	18
Male	218	-	-	-	-	-	-	-	2	3	6	13
Female	328	-	-	-	-	-	-	-	-	1	-	5
White, both sexes	491	-	-	-	-	-	-	-	2	3	6	15
Male	189	-	-	-	-	-	-	-	2	2	6	11
Female	302	-	-	-	-	-	-	-	-	1	-	4
Black, both sexes	51	-	-	-	-	-	-	-	-	1	-	2
Male	26	-	-	-	-	-	-	-	-	1	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Delaware												
All races, both sexes	135	-	-	-	-	-	-	-	-	4	4	8
Male	65	-	-	-	-	-	-	-	-	3	2	5
Female	70	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	95	-	-	-	-	-	-	-	-	2	3	4
Male	40	-	-	-	-	-	-	-	-	1	2	3
Female	55	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	40	-	-	-	-	-	-	-	-	2	1	4
Male	25	-	-	-	-	-	-	-	-	2	-	2
Female	15	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	269	-	-	-	-	-	2	-	4	3	6	18
Male	129	-	-	-	-	-	2	-	3	2	3	9
Female	140	-	-	-	-	-	-	-	1	1	3	9
White, both sexes	41	-	-	-	-	-	1	-	-	-	1	1
Male	20	-	-	-	-	-	1	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	225	-	-	-	-	-	1	-	4	3	5	17
Male	108	-	-	-	-	-	1	-	3	2	2	9
Female	117	-	-	-	-	-	-	-	1	1	3	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Florida												
All races, both sexes	4,228	-	-	1	-	2	6	14	23	45	91	149
Male	1,976	-	-	1	-	-	5	10	21	31	55	94
Female	2,252	-	-	-	-	2	1	4	2	14	36	55
White, both sexes	3,377	-	-	1	-	-	3	6	12	31	53	91
Male	1,565	-	-	1	-	-	2	4	11	23	36	64
Female	1,812	-	-	-	-	-	1	2	1	8	17	27
Black, both sexes	828	-	-	-	-	2	3	7	10	14	37	57
Male	398	-	-	-	-	-	3	5	9	8	18	29
Female	430	-	-	-	-	2	-	2	1	6	19	28
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	1	-	-	1	1
Male	11	-	-	-	-	-	-	1	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,151	-	-	-	1	1	4	12	13	29	77	109
Male	993	-	-	-	1	1	4	9	8	20	49	71
Female	1,158	-	-	-	-	-	-	3	5	9	28	38
White, both sexes	1,065	-	-	-	-	1	3	1	3	11	30	48
Male	467	-	-	-	-	1	3	1	2	8	21	35
Female	598	-	-	-	-	-	-	-	1	3	9	13
Black, both sexes	1,065	-	-	-	1	-	1	11	10	18	47	59
Male	514	-	-	-	1	-	1	8	6	12	28	34
Female	551	-	-	-	-	-	-	3	4	6	19	25
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Hawaii												
All races, both sexes	190	-	-	-	-	-	-	2	-	-	4	2
Male	94	-	-	-	-	-	-	1	-	-	2	2
Female	96	-	-	-	-	-	-	1	-	-	2	-
White, both sexes	34	-	-	-	-	-	-	1	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	153	-	-	-	-	-	-	1	-	-	3	2
Male	73	-	-	-	-	-	-	1	-	-	1	2
Female	80	-	-	-	-	-	-	-	-	-	2	-
Idaho												
All races, both sexes	217	-	-	-	-	-	1	-	1	1	2	8
Male	91	-	-	-	-	-	-	-	1	1	1	6
Female	126	-	-	-	-	-	1	-	-	-	1	2
White, both sexes	212	-	-	-	-	-	1	-	1	1	1	7
Male	87	-	-	-	-	-	-	-	1	1	-	5
Female	125	-	-	-	-	-	1	-	-	-	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Illinois												
All races, both sexes	2,363	-	-	-	-	-	3	10	13	34	57	92
Male	1,035	-	-	-	-	-	3	8	9	22	40	70
Female	1,328	-	-	-	-	-	-	2	4	12	17	22
White, both sexes	1,599	-	-	-	-	-	1	5	4	14	23	38
Male	619	-	-	-	-	-	1	3	4	10	19	32
Female	980	-	-	-	-	-	-	2	-	4	4	6
Black, both sexes	738	-	-	-	-	-	2	5	9	20	33	53
Male	407	-	-	-	-	-	2	5	5	12	20	38
Female	331	-	-	-	-	-	-	-	4	8	13	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Indiana												
All races, both sexes	1,053	-	-	-	-	-	1	4	7	12	20	42
Male	446	-	-	-	-	-	1	2	6	8	16	29
Female	607	-	-	-	-	-	-	2	1	4	4	13
White, both sexes	869	-	-	-	-	-	1	3	1	7	10	32
Male	344	-	-	-	-	-	1	2	1	5	7	20
Female	525	-	-	-	-	-	-	1	-	2	3	12
Black, both sexes	181	-	-	-	-	-	-	1	6	5	9	10
Male	100	-	-	-	-	-	-	-	5	3	8	9
Female	81	-	-	-	-	-	-	1	1	2	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Iowa												
All races, both sexes	565	-	-	-	-	-	-	2	-	-	7	4
Male	207	-	-	-	-	-	-	2	-	-	3	1
Female	358	-	-	-	-	-	-	-	-	-	4	3
White, both sexes	555	-	-	-	-	-	-	2	-	-	7	4
Male	204	-	-	-	-	-	-	2	-	-	3	1
Female	351	-	-	-	-	-	-	-	-	-	4	3
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	381	-	-	-	-	-	-	-	-	2	1	10
Male	118	-	-	-	-	-	-	-	-	1	-	6
Female	263	-	-	-	-	-	-	-	-	1	1	4
White, both sexes	331	-	-	-	-	-	-	-	-	1	1	7
Male	95	-	-	-	-	-	-	-	-	1	-	5
Female	236	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	45	-	-	-	-	-	-	-	-	1	-	3
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Kentucky												
All races, both sexes	732	-	-	-	-	-	-	4	6	9	21	42
Male	350	-	-	-	-	-	-	3	6	6	14	36
Female	382	-	-	-	-	-	-	1	-	3	7	6
White, both sexes	649	-	-	-	-	-	-	4	5	6	15	38
Male	311	-	-	-	-	-	-	3	5	4	10	32
Female	338	-	-	-	-	-	-	1	-	2	5	6
Black, both sexes	82	-	-	-	-	-	-	-	1	3	6	4
Male	39	-	-	-	-	-	-	-	1	2	4	4
Female	43	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,166	-	-	1	-	-	3	-	10	18	24	59
Male	531	-	-	-	-	-	2	-	3	16	14	37
Female	635	-	-	1	-	-	1	-	7	2	10	22
White, both sexes	665	-	-	-	-	-	2	-	1	7	7	25
Male	281	-	-	-	-	-	1	-	-	6	5	18
Female	384	-	-	-	-	-	1	-	1	1	2	7
Black, both sexes	496	-	-	1	-	-	1	-	9	11	17	34
Male	245	-	-	-	-	-	1	-	3	10	9	19
Female	251	-	-	1	-	-	-	-	6	1	8	15
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Maine												
All races, both sexes	158	-	-	-	-	-	-	-	-	3	2	8
Male	68	-	-	-	-	-	-	-	-	3	1	5
Female	90	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	154	-	-	-	-	-	-	-	-	2	2	6
Male	65	-	-	-	-	-	-	-	-	2	1	4
Female	89	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,280	-	-	-	-	-	2	6	5	17	56	91
Male	647	-	-	-	-	-	2	5	5	15	39	64
Female	633	-	-	-	-	-	-	1	-	2	17	27
White, both sexes	745	-	-	-	-	-	-	1	2	9	30	45
Male	368	-	-	-	-	-	-	1	2	9	24	35
Female	377	-	-	-	-	-	-	-	-	-	6	10
Black, both sexes	509	-	-	-	-	-	2	5	3	8	25	46
Male	263	-	-	-	-	-	2	4	3	6	14	29
Female	246	-	-	-	-	-	-	1	-	2	11	17
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Massachusetts												
All races, both sexes	986	-	-	-	-	1	1	3	3	9	28	40
Male	419	-	-	-	-	1	1	2	3	6	24	33
Female	567	-	-	-	-	-	-	1	-	3	4	7
White, both sexes	899	-	-	-	-	1	1	2	2	9	25	36
Male	370	-	-	-	-	1	1	1	2	6	21	29
Female	529	-	-	-	-	-	-	1	-	3	4	7
Black, both sexes	74	-	-	-	-	-	-	1	1	-	2	4
Male	42	-	-	-	-	-	-	1	1	-	2	4
Female	32	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,173	-	-	-	1	-	1	5	11	24	36	72
Male	909	-	-	-	-	-	1	4	6	17	17	47
Female	1,264	-	-	-	1	-	-	1	5	7	19	25
White, both sexes	1,640	-	-	-	1	-	1	-	3	9	18	41
Male	647	-	-	-	-	-	1	-	2	8	8	27
Female	993	-	-	-	1	-	-	-	1	1	10	14
Black, both sexes	507	-	-	-	-	-	-	4	8	15	18	29
Male	254	-	-	-	-	-	-	3	4	9	9	20
Female	253	-	-	-	-	-	-	1	4	6	9	9
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Minnesota												
All races, both sexes	663	-	-	-	-	-	2	-	1	5	12	8
Male	256	-	-	-	-	-	1	-	1	5	8	5
Female	407	-	-	-	-	-	1	-	-	-	4	3
White, both sexes	627	-	-	-	-	-	1	-	-	5	7	6
Male	240	-	-	-	-	-	-	-	-	5	6	4
Female	387	-	-	-	-	-	1	-	-	-	1	2
Black, both sexes	24	-	-	-	-	-	1	-	-	-	4	2
Male	11	-	-	-	-	-	1	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,132	-	-	-	-	-	5	5	10	8	19	40
Male	475	-	-	-	-	-	4	4	6	5	13	24
Female	657	-	-	-	-	-	1	1	4	3	6	16
White, both sexes	558	-	-	-	-	-	2	1	2	1	5	13
Male	216	-	-	-	-	-	2	1	1	1	3	9
Female	342	-	-	-	-	-	-	-	1	-	2	4
Black, both sexes	571	-	-	-	-	-	3	4	8	7	14	27
Male	257	-	-	-	-	-	2	3	5	4	10	15
Female	314	-	-	-	-	-	1	1	3	3	4	12
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Missouri												
All races, both sexes	958	-	-	-	-	-	2	2	8	12	33	56
Male	395	-	-	-	-	-	2	2	2	9	23	39
Female	563	-	-	-	-	-	-	-	6	3	10	17
White, both sexes	683	-	-	-	-	-	1	-	5	3	16	29
Male	254	-	-	-	-	-	1	-	1	3	12	23
Female	429	-	-	-	-	-	-	-	4	-	4	6
Black, both sexes	269	-	-	-	-	-	1	2	3	9	17	27
Male	138	-	-	-	-	-	1	2	1	6	11	16
Female	131	-	-	-	-	-	-	-	2	3	6	11
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	116	-	-	-	-	-	-	-	1	1	3	2
Male	52	-	-	-	-	-	-	-	1	1	1	1
Female	64	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	113	-	-	-	-	-	-	-	-	1	3	1
Male	51	-	-	-	-	-	-	-	-	1	1	1
Female	62	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Nebraska												
All races, both sexes	291	-	-	-	-	-	-	2	2	2	6	9
Male	113	-	-	-	-	-	-	1	1	2	4	7
Female	178	-	-	-	-	-	-	1	1	-	2	2
White, both sexes	274	-	-	-	-	-	-	2	2	1	6	9
Male	107	-	-	-	-	-	-	1	1	1	4	7
Female	167	-	-	-	-	-	-	1	1	-	2	2
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	565	-	-	-	-	1	-	-	2	3	11	27
Male	307	-	-	-	-	1	-	-	1	2	9	21
Female	258	-	-	-	-	-	-	-	1	1	2	6
White, both sexes	478	-	-	-	-	1	-	-	2	2	7	25
Male	264	-	-	-	-	1	-	-	1	1	6	21
Female	214	-	-	-	-	-	-	-	1	1	1	4
Black, both sexes	61	-	-	-	-	-	-	-	-	-	3	2
Male	33	-	-	-	-	-	-	-	-	-	2	-
Female	28	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
New Hampshire												
All races, both sexes	206	-	-	-	-	-	-	-	-	-	2	2
Male	78	-	-	-	-	-	-	-	-	-	2	2
Female	128	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	204	-	-	-	-	-	-	-	-	-	2	2
Male	78	-	-	-	-	-	-	-	-	-	2	2
Female	126	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,403	-	-	-	-	-	1	2	4	9	21	49
Male	598	-	-	-	-	-	-	-	3	5	15	34
Female	805	-	-	-	-	-	1	2	1	4	6	15
White, both sexes	1,089	-	-	-	-	-	1	1	2	4	12	28
Male	446	-	-	-	-	-	-	-	1	3	10	17
Female	643	-	-	-	-	-	1	1	1	1	2	11
Black, both sexes	286	-	-	-	-	-	-	1	2	5	9	21
Male	142	-	-	-	-	-	-	-	2	2	5	17
Female	144	-	-	-	-	-	-	1	-	3	4	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
New Mexico												
All races, both sexes	268	-	-	-	-	-	-	2	1	8	9	14
Male	122	-	-	-	-	-	-	2	1	7	8	11
Female	146	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	238	-	-	-	-	-	-	2	1	6	6	11
Male	110	-	-	-	-	-	-	2	1	5	5	9
Female	128	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	1	2	2
Male	7	-	-	-	-	-	-	-	-	1	2	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,429	-	-	-	-	-	6	10	17	41	87	162
Male	1,891	-	-	-	-	-	4	7	12	28	62	114
Female	2,538	-	-	-	-	-	2	3	5	13	25	48
White, both sexes	3,038	-	-	-	-	-	2	5	7	16	37	78
Male	1,220	-	-	-	-	-	1	4	6	12	32	58
Female	1,818	-	-	-	-	-	1	1	1	4	5	20
Black, both sexes	1,277	-	-	-	-	-	3	5	9	24	47	81
Male	603	-	-	-	-	-	2	3	5	15	28	54
Female	674	-	-	-	-	-	1	2	4	9	19	27
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	114	-	-	-	-	-	1	-	1	1	3	3
Male	68	-	-	-	-	-	1	-	1	1	2	2
Female	46	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
North Carolina												
All races, both sexes	1,407	1	-	-	-	1	2	5	4	23	39	77
Male	608	1	-	-	-	-	2	3	2	16	23	42
Female	799	-	-	-	-	1	-	2	2	7	16	35
White, both sexes	846	-	-	-	-	-	1	1	2	8	15	33
Male	344	-	-	-	-	-	1	1	1	6	8	21
Female	502	-	-	-	-	-	-	-	1	2	7	12
Black, both sexes	543	1	-	-	-	1	1	4	2	15	23	41
Male	254	1	-	-	-	-	1	2	1	10	14	19
Female	289	-	-	-	-	1	-	2	1	5	9	22
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	1	3
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	119	-	-	-	-	-	-	1	-	1	1	1
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	76	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	112	-	-	-	-	-	-	-	-	1	1	1
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	70	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Ohio												
All races, both sexes	2,916	-	-	-	-	-	3	8	14	33	53	112
Male	1,209	-	-	-	-	-	1	8	11	21	36	73
Female	1,707	-	-	-	-	-	2	-	3	12	17	39
White, both sexes	2,347	-	-	-	-	-	3	3	8	19	37	69
Male	922	-	-	-	-	-	1	3	6	13	29	50
Female	1,425	-	-	-	-	-	2	-	2	6	8	19
Black, both sexes	560	-	-	-	-	-	-	5	6	14	15	42
Male	283	-	-	-	-	-	-	5	5	8	6	22
Female	277	-	-	-	-	-	-	-	1	6	9	20
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	540	-	-	-	-	-	-	2	5	6	5	18
Male	233	-	-	-	-	-	-	2	5	4	3	15
Female	307	-	-	-	-	-	-	-	-	2	2	3
White, both sexes	437	-	-	-	-	-	-	2	4	2	2	14
Male	178	-	-	-	-	-	-	2	4	2	1	11
Female	259	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	74	-	-	-	-	-	-	-	1	4	2	1
Male	38	-	-	-	-	-	-	-	1	2	1	1
Female	36	-	-	-	-	-	-	-	-	2	1	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	1	3
Male	17	-	-	-	-	-	-	-	-	-	1	3
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Oregon												
All races, both sexes	657	-	-	-	-	-	-	1	2	3	5	7
Male	242	-	-	-	-	-	-	1	2	2	3	7
Female	415	-	-	-	-	-	-	-	-	1	2	-
White, both sexes	641	-	-	-	-	-	-	1	1	3	5	7
Male	236	-	-	-	-	-	-	1	1	2	3	7
Female	405	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,860	-	-	-	-	1	1	2	7	8	29	56
Male	784	-	-	-	-	-	1	1	3	5	19	39
Female	1,076	-	-	-	-	1	-	1	4	3	10	17
White, both sexes	1,571	-	-	-	-	-	1	1	3	5	17	34
Male	654	-	-	-	-	-	1	-	2	4	11	23
Female	917	-	-	-	-	-	-	1	1	1	6	11
Black, both sexes	278	-	-	-	-	1	-	1	4	3	11	21
Male	123	-	-	-	-	-	-	1	1	1	7	15
Female	155	-	-	-	-	1	-	-	3	2	4	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Rhode Island												
All races, both sexes	109	-	-	-	-	-	-	-	-	1	4	3
Male	49	-	-	-	-	-	-	-	-	1	2	3
Female	60	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	102	-	-	-	-	-	-	-	-	1	3	3
Male	45	-	-	-	-	-	-	-	-	1	1	3
Female	57	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	812	-	-	-	-	1	1	1	4	5	18	29
Male	355	-	-	-	-	-	1	1	1	3	12	18
Female	457	-	-	-	-	1	-	-	3	2	6	11
White, both sexes	434	-	-	-	-	-	1	-	-	2	5	13
Male	186	-	-	-	-	-	1	-	-	1	4	10
Female	248	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	373	-	-	-	-	-	-	1	4	3	13	16
Male	167	-	-	-	-	-	-	1	1	2	8	8
Female	206	-	-	-	-	-	-	-	3	1	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
South Dakota												
All races, both sexes	141	-	-	-	-	-	-	-	-	1	-	-
Male	47	-	-	-	-	-	-	-	-	1	-	-
Female	94	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	135	-	-	-	-	-	-	-	-	1	-	-
Male	42	-	-	-	-	-	-	-	-	1	-	-
Female	93	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,468	-	-	-	-	-	4	6	13	26	41	53
Male	603	-	-	-	-	-	3	5	5	20	23	30
Female	865	-	-	-	-	-	1	1	8	6	18	23
White, both sexes	1,022	-	-	-	-	-	3	4	4	13	17	28
Male	401	-	-	-	-	-	3	4	1	12	9	16
Female	621	-	-	-	-	-	-	-	3	1	8	12
Black, both sexes	438	-	-	-	-	-	1	2	9	13	23	25
Male	199	-	-	-	-	-	-	1	4	8	13	14
Female	239	-	-	-	-	-	1	1	5	5	10	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Texas												
All races, both sexes	3,507	-	-	-	1	-	3	7	33	49	109	158
Male	1,481	-	-	-	1	-	1	4	25	36	72	103
Female	2,026	-	-	-	-	-	2	3	8	13	37	55
White, both sexes	2,688	-	-	-	1	-	2	2	15	33	67	101
Male	1,093	-	-	-	1	-	1	2	12	24	48	67
Female	1,595	-	-	-	-	-	1	-	3	9	19	34
Black, both sexes	776	-	-	-	-	-	1	5	18	16	40	57
Male	365	-	-	-	-	-	-	2	13	12	22	36
Female	411	-	-	-	-	-	1	3	5	4	18	21
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	2	-
Male	18	-	-	-	-	-	-	-	-	-	2	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	234	-	-	-	-	-	-	-	-	1	2	4
Male	83	-	-	-	-	-	-	-	-	1	2	3
Female	151	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	227	-	-	-	-	-	-	-	-	1	2	4
Male	80	-	-	-	-	-	-	-	-	1	2	3
Female	147	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Vermont												
All races, both sexes	100	-	-	-	-	1	-	-	1	1	1	2
Male	42	-	-	-	-	-	-	-	1	1	1	1
Female	58	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	98	-	-	-	-	1	-	-	1	1	1	2
Male	42	-	-	-	-	-	-	-	1	1	1	1
Female	56	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	961	-	-	-	-	-	2	1	8	18	21	45
Male	418	-	-	-	-	-	1	-	6	14	12	37
Female	543	-	-	-	-	-	1	1	2	4	9	8
White, both sexes	626	-	-	-	-	-	1	-	4	7	8	27
Male	253	-	-	-	-	-	1	-	3	6	6	23
Female	373	-	-	-	-	-	-	-	1	1	2	4
Black, both sexes	323	-	-	-	-	-	1	1	4	11	12	18
Male	159	-	-	-	-	-	-	-	3	8	6	14
Female	164	-	-	-	-	-	1	1	1	3	6	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Washington												
All races, both sexes	1,247	-	-	-	-	-	-	1	4	12	20	27
Male	522	-	-	-	-	-	-	1	3	11	14	25
Female	725	-	-	-	-	-	-	-	1	1	6	2
White, both sexes	1,122	-	-	-	-	-	-	-	4	10	14	22
Male	456	-	-	-	-	-	-	-	3	9	11	22
Female	666	-	-	-	-	-	-	-	1	1	3	-
Black, both sexes	67	-	-	-	-	-	-	-	-	2	5	2
Male	38	-	-	-	-	-	-	-	-	2	2	2
Female	29	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	48	-	-	-	-	-	-	1	-	-	1	1
Male	23	-	-	-	-	-	-	1	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
West Virginia												
All races, both sexes	358	-	-	-	-	-	-	-	3	3	3	8
Male	132	-	-	-	-	-	-	-	3	3	2	3
Female	226	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	336	-	-	-	-	-	-	-	3	3	3	7
Male	122	-	-	-	-	-	-	-	3	3	2	2
Female	214	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Wisconsin												
All races, both sexes	855	-	-	-	-	-	1	1	3	3	10	25
Male	301	-	-	-	-	-	1	1	3	1	4	13
Female	554	-	-	-	-	-	-	-	-	2	6	12
White, both sexes	780	-	-	-	-	-	1	1	3	2	5	16
Male	269	-	-	-	-	-	1	1	3	1	3	8
Female	511	-	-	-	-	-	-	-	-	1	2	8
Black, both sexes	66	-	-	-	-	-	-	-	-	1	5	8
Male	29	-	-	-	-	-	-	-	-	-	1	5
Female	37	-	-	-	-	-	-	-	-	1	4	3
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	2	3
Male	26	-	-	-	-	-	-	-	-	-	2	2
Female	33	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	59	-	-	-	-	-	-	-	-	-	2	3
Male	26	-	-	-	-	-	-	-	-	-	2	2
Female	33	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
United States												
All races, both sexes	2,960	3,271	3,296	3,418	3,975	5,617	8,075	9,332	7,589	3,669	914	2
Male	1,965	2,113	2,059	1,969	2,040	2,544	3,166	2,950	1,814	635	105	2
Female	995	1,158	1,237	1,449	1,935	3,073	4,909	6,382	5,775	3,034	809	-
White, both sexes	1,716	2,038	2,131	2,232	2,715	4,197	6,533	7,947	6,559	3,181	745	2
Male	1,220	1,360	1,375	1,286	1,388	1,908	2,552	2,516	1,542	542	87	2
Female	496	678	756	946	1,327	2,289	3,981	5,431	5,017	2,639	658	-
Black, both sexes	1,158	1,154	1,072	1,068	1,143	1,237	1,337	1,148	826	420	149	-
Male	691	701	632	608	588	558	525	355	211	74	13	-
Female	467	453	440	460	555	679	812	793	615	346	136	-
American Indian 1/, both sexes	24	19	16	31	17	28	21	21	20	8	4	-
Male	14	13	8	22	10	15	7	7	6	2	1	-
Female	10	6	8	9	7	13	14	14	14	6	3	-
Asian or Pacific Islander 2/, both sexes	62	60	77	87	100	155	184	216	184	60	16	-
Male	40	39	44	53	54	63	82	72	55	17	4	-
Female	22	21	33	34	46	92	102	144	129	43	12	-
Alabama												
All races, both sexes	46	49	62	66	78	95	133	126	99	43	12	-
Male	34	30	36	33	33	46	50	35	27	10	1	-
Female	12	19	26	33	45	49	83	91	72	33	11	-
White, both sexes	18	15	34	35	49	52	77	93	69	31	6	-
Male	15	10	20	20	23	24	27	23	18	7	1	-
Female	3	5	14	15	26	28	50	70	51	24	5	-
Black, both sexes	27	34	27	31	29	43	56	33	30	12	6	-
Male	18	20	15	13	10	22	23	12	9	3	-	-
Female	9	14	12	18	19	21	33	21	21	9	6	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Alaska												
All races, both sexes	5	7	3	7	4	7	8	7	5	1	3	-
Male	4	2	2	3	3	6	3	2	3	-	-	-
Female	1	5	1	4	1	1	5	5	2	1	3	-
White, both sexes	4	2	1	5	3	5	4	6	2	1	1	-
Male	3	-	1	2	2	4	1	2	1	-	-	-
Female	1	2	-	3	1	1	3	4	1	1	1	-
Black, both sexes	-	1	-	-	1	-	1	-	-	-	1	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	2	2	1	-	2	3	-	3	-	1	-
Male	1	1	1	-	-	2	1	-	2	-	-	-
Female	-	1	1	1	-	-	2	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	51	65	49	57	82	109	139	164	124	49	15	-
Male	39	40	31	36	47	42	52	68	32	12	2	-
Female	12	25	18	21	35	67	87	96	92	37	13	-
White, both sexes	39	51	39	50	68	104	133	158	122	44	14	-
Male	31	32	25	33	37	41	47	66	31	11	1	-
Female	8	19	14	17	31	63	86	92	91	33	13	-
Black, both sexes	8	9	5	1	7	2	3	2	1	3	-	-
Male	6	6	3	-	6	-	3	1	-	-	-	-
Female	2	3	2	1	1	2	-	1	1	3	-	-
American Indian 1/, both sexes	3	2	4	5	5	1	1	2	-	1	1	-
Male	2	1	3	3	3	-	-	-	-	-	1	-
Female	1	1	1	2	2	1	1	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	1	2	2	2	2	1	1	-	-
Male	-	1	-	-	1	1	2	1	1	1	-	-
Female	1	2	1	1	1	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Arkansas												
All races, both sexes	40	32	28	33	38	47	72	100	77	45	15	-
Male	26	23	21	21	19	20	31	27	13	10	2	-
Female	14	9	7	12	19	27	41	73	64	35	13	-
White, both sexes	20	21	20	25	27	38	61	82	63	33	11	-
Male	15	14	14	16	14	14	24	19	9	8	1	-
Female	5	7	6	9	13	24	37	63	54	25	10	-
Black, both sexes	20	10	8	8	11	9	11	18	14	12	4	-
Male	11	8	7	5	5	6	7	8	4	2	1	-
Female	9	2	1	3	6	3	4	10	10	10	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	342	419	442	431	481	749	1,123	1,386	1,068	518	136	1
Male	217	267	286	247	269	349	445	470	301	106	15	1
Female	125	152	156	184	212	400	678	916	767	412	121	-
White, both sexes	227	305	321	302	363	557	910	1,128	880	451	114	1
Male	149	197	219	174	208	266	364	390	253	92	14	1
Female	78	108	102	128	155	291	546	738	627	359	100	-
Black, both sexes	89	87	86	81	72	109	110	127	78	34	15	-
Male	52	52	47	41	38	47	34	40	21	5	-	-
Female	37	35	39	40	34	62	76	87	57	29	15	-
American Indian 1/, both sexes	3	6	1	9	1	7	2	6	-	1	-	-
Male	1	5	1	7	1	5	2	3	-	-	-	-
Female	2	1	-	2	-	2	-	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	23	21	34	39	45	76	101	125	110	32	7	-
Male	15	13	19	25	22	31	45	37	27	9	1	-
Female	8	8	15	14	23	45	56	88	83	23	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Colorado												
All races, both sexes	23	35	26	28	31	63	52	89	65	42	9	-
Male	13	27	16	17	15	25	24	23	12	8	1	-
Female	10	8	10	11	16	38	28	66	53	34	8	-
White, both sexes	19	29	21	25	23	60	47	84	64	41	8	-
Male	11	23	13	16	11	24	22	22	12	8	1	-
Female	8	6	8	9	12	36	25	62	52	33	7	-
Black, both sexes	4	5	4	3	6	2	3	4	1	1	1	-
Male	2	3	2	1	3	1	1	1	-	-	-	-
Female	2	2	2	2	3	1	2	3	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	2	1	2	1	-	-	-	-
Male	-	1	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-
Connecticut												
All races, both sexes	18	39	20	26	22	51	81	99	100	48	12	-
Male	14	25	8	19	14	19	38	29	21	7	-	-
Female	4	14	12	7	8	32	43	70	79	41	12	-
White, both sexes	13	32	13	20	17	42	79	96	96	46	11	-
Male	11	21	3	14	11	16	36	28	21	7	-	-
Female	2	11	10	6	6	26	43	68	75	39	11	-
Black, both sexes	5	6	6	5	5	9	2	3	4	2	1	-
Male	3	4	4	4	3	3	2	1	-	-	-	-
Female	2	2	2	1	2	6	-	2	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Delaware												
All races, both sexes	11	9	6	9	12	15	17	21	15	2	2	-
Male	6	7	4	7	6	9	7	5	4	-	-	-
Female	5	2	2	2	6	6	10	16	11	2	2	-
White, both sexes	7	3	5	5	7	13	13	19	11	1	2	-
Male	3	2	3	4	3	7	5	3	4	-	-	-
Female	4	1	2	1	4	6	8	16	7	1	2	-
Black, both sexes	4	6	1	4	5	2	4	2	4	1	-	-
Male	3	5	1	3	3	2	2	2	-	-	-	-
Female	1	1	-	1	2	-	2	-	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	27	20	18	19	22	23	42	31	19	11	4	-
Male	18	12	12	12	10	12	19	8	5	2	-	-
Female	9	8	6	7	12	11	23	23	14	9	4	-
White, both sexes	-	4	-	1	1	3	8	7	8	5	1	-
Male	-	2	-	1	-	1	7	4	2	1	-	-
Female	-	2	-	-	1	2	1	3	6	4	1	-
Black, both sexes	27	16	18	18	21	20	34	23	11	4	3	-
Male	18	10	12	11	10	11	12	4	3	-	-	-
Female	9	6	6	7	11	9	22	19	8	4	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Florida												
All races, both sexes	210	217	269	260	317	427	633	691	574	231	68	-
Male	142	152	175	153	156	209	274	258	168	58	14	-
Female	68	65	94	107	161	218	359	433	406	173	54	-
White, both sexes	142	151	185	185	238	349	546	620	508	200	56	-
Male	103	113	122	110	117	170	241	235	150	52	11	-
Female	39	38	63	75	121	179	305	385	358	148	45	-
Black, both sexes	67	65	81	72	77	75	87	68	63	31	12	-
Male	38	38	51	42	38	38	33	22	17	6	3	-
Female	29	27	30	30	39	37	54	46	46	25	9	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	2	2	3	-	3	2	-	-	-
Male	1	1	2	-	1	1	-	1	1	-	-	-
Female	-	-	1	2	1	2	-	2	1	-	-	-
Georgia												
All races, both sexes	164	194	174	175	197	202	246	266	168	97	22	-
Male	113	119	109	105	96	83	83	74	35	13	-	-
Female	51	75	65	70	101	119	163	192	133	84	22	-
White, both sexes	54	81	65	69	87	111	152	164	116	58	11	-
Male	45	49	41	40	49	46	53	49	20	4	-	-
Female	9	32	24	29	38	65	99	115	96	54	11	-
Black, both sexes	110	110	108	103	104	89	94	99	52	38	11	-
Male	68	68	68	64	43	36	30	24	15	8	-	-
Female	42	42	40	39	61	53	64	75	37	30	11	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	3	6	2	-	2	-	-	-	-
Male	-	2	-	1	4	1	-	1	-	-	-	-
Female	-	1	1	2	2	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Hawaii												
All races, both sexes	7	7	11	9	14	14	23	40	38	12	7	-
Male	6	5	7	6	9	6	10	17	17	4	2	-
Female	1	2	4	3	5	8	13	23	21	8	5	-
White, both sexes	1	3	1	2	4	1	6	7	6	1	-	-
Male	1	1	1	1	1	1	2	5	5	-	-	-
Female	-	2	-	1	3	-	4	2	1	1	-	-
Black, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	3	10	7	10	13	16	32	32	11	7	-
Male	5	3	6	5	8	5	7	12	12	4	2	-
Female	1	-	4	2	2	8	9	20	20	7	5	-
Idaho												
All races, both sexes	6	11	13	7	18	22	32	30	41	19	5	-
Male	6	6	10	4	6	10	18	9	9	3	1	-
Female	-	5	3	3	12	12	14	21	32	16	4	-
White, both sexes	5	11	13	7	18	22	32	29	41	19	4	-
Male	5	6	10	4	6	10	18	8	9	3	1	-
Female	-	5	3	3	12	12	14	21	32	16	3	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Illinois												
All races, both sexes	144	157	135	162	193	226	325	336	275	162	39	-
Male	97	101	85	86	102	106	117	109	57	18	5	-
Female	47	56	50	76	91	120	208	227	218	144	34	-
White, both sexes	65	74	67	87	114	161	253	281	235	140	37	-
Male	48	49	37	44	59	73	86	88	49	12	5	-
Female	17	25	30	43	55	88	167	193	186	128	32	-
Black, both sexes	77	83	67	71	75	63	66	52	39	21	2	-
Male	48	52	47	39	42	33	29	21	8	6	-	-
Female	29	31	20	32	33	30	37	31	31	15	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	4	4	1	6	3	1	1	-	-
Male	1	-	1	3	1	-	2	-	-	-	-	-
Female	1	-	-	1	3	1	4	3	1	1	-	-
Indiana												
All races, both sexes	56	52	48	53	66	111	178	175	140	72	16	-
Male	40	37	28	30	33	57	65	60	25	8	1	-
Female	16	15	20	23	33	54	113	115	115	64	15	-
White, both sexes	39	37	37	42	54	84	151	163	129	66	13	-
Male	28	25	21	23	27	46	52	55	24	7	-	-
Female	11	12	16	19	27	38	99	108	105	59	13	-
Black, both sexes	17	15	11	11	11	26	27	12	11	6	3	-
Male	12	12	7	7	5	11	13	5	1	1	1	-
Female	5	3	4	4	6	15	14	7	10	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Iowa												
All races, both sexes	12	16	16	20	28	50	88	123	122	60	17	-
Male	8	8	10	8	18	21	37	50	32	9	-	-
Female	4	8	6	12	10	29	51	73	90	51	17	-
White, both sexes	12	13	16	19	27	49	86	122	122	59	17	-
Male	8	6	10	8	18	21	37	49	32	9	-	-
Female	4	7	6	11	9	28	49	73	90	50	17	-
Black, both sexes	-	3	-	1	1	1	2	1	-	1	-	-
Male	-	2	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	1	1	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	7	16	11	15	18	31	57	100	58	45	10	-
Male	4	8	6	8	8	15	14	28	11	9	-	-
Female	3	8	5	7	10	16	43	72	47	36	10	-
White, both sexes	4	10	10	13	15	24	49	91	54	42	10	-
Male	3	4	5	6	7	9	12	25	10	8	-	-
Female	1	6	5	7	8	15	37	66	44	34	10	-
Black, both sexes	3	6	1	2	2	5	7	9	3	3	-	-
Male	1	4	1	2	-	4	2	3	1	1	-	-
Female	2	2	-	-	2	1	5	6	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	2	1	-	1	-	-	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Kentucky												
All races, both sexes	48	39	40	51	44	74	113	102	88	39	12	-
Male	29	31	23	30	31	35	45	30	22	6	3	-
Female	19	8	17	21	13	39	68	72	66	33	9	-
White, both sexes	42	36	34	42	40	65	103	93	81	34	11	-
Male	27	28	21	26	27	31	41	27	21	5	3	-
Female	15	8	13	16	13	34	62	66	60	29	8	-
Black, both sexes	6	3	6	8	4	9	10	9	7	5	1	-
Male	2	3	2	4	4	4	4	3	1	1	-	-
Female	4	-	4	4	-	5	6	6	6	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	82	91	72	79	91	138	155	160	124	48	11	-
Male	52	58	51	45	47	58	60	51	22	12	3	-
Female	30	33	21	34	44	80	95	109	102	36	8	-
White, both sexes	38	43	34	40	48	78	105	115	83	34	5	-
Male	23	28	26	24	24	28	41	40	11	5	1	-
Female	15	15	8	16	24	50	64	75	72	29	4	-
Black, both sexes	42	48	38	38	43	58	50	45	41	14	6	-
Male	27	30	25	20	23	28	19	11	11	7	2	-
Female	15	18	13	18	20	30	31	34	30	7	4	-
American Indian 1/, both sexes	2	-	-	-	-	2	-	-	-	-	-	-
Male	2	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Maine												
All races, both sexes	6	8	10	9	12	15	22	19	23	17	4	-
Male	5	7	5	7	5	8	9	6	5	2	-	-
Female	1	1	5	2	7	7	13	13	18	15	4	-
White, both sexes	6	8	10	9	12	14	22	19	23	17	4	-
Male	5	7	5	7	5	7	9	6	5	2	-	-
Female	1	1	5	2	7	7	13	13	18	15	4	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	92	90	105	101	94	105	143	165	130	58	20	-
Male	59	56	74	61	52	49	60	68	30	6	2	-
Female	33	34	31	40	42	56	83	97	100	52	18	-
White, both sexes	44	43	53	50	42	57	93	127	94	41	14	-
Male	32	31	36	34	21	24	38	54	22	4	1	-
Female	12	12	17	16	21	33	55	73	72	37	13	-
Black, both sexes	46	45	50	48	50	44	48	35	33	15	6	-
Male	26	23	36	26	29	22	21	12	7	2	1	-
Female	20	22	14	22	21	22	27	23	26	13	5	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	2	2	4	2	3	2	2	-	-
Male	1	1	2	1	2	3	1	2	-	-	-	-
Female	1	-	-	1	-	1	1	1	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Massachusetts												
All races, both sexes	52	54	66	56	51	88	130	176	134	77	17	-
Male	36	36	42	34	24	40	52	41	34	9	1	-
Female	16	18	24	22	27	48	78	135	100	68	16	-
White, both sexes	44	50	55	47	46	73	118	173	127	73	17	-
Male	30	36	35	28	20	32	48	41	30	8	1	-
Female	14	14	20	19	26	41	70	132	97	65	16	-
Black, both sexes	7	4	11	9	2	12	10	2	5	4	-	-
Male	5	-	7	6	1	8	4	-	2	1	-	-
Female	2	4	4	3	1	4	6	2	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	1	1	1	2	-	-	-
Male	1	-	-	-	2	-	-	-	2	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Michigan												
All races, both sexes	111	122	128	107	151	241	308	373	316	138	28	-
Male	77	71	80	58	70	107	129	127	70	24	4	-
Female	34	51	48	49	81	134	179	246	246	114	24	-
White, both sexes	58	72	82	60	104	188	250	329	277	124	23	-
Male	47	42	55	33	49	79	103	111	58	21	3	-
Female	11	30	27	27	55	109	147	218	219	103	20	-
Black, both sexes	50	50	44	45	45	51	56	41	34	12	5	-
Male	29	29	25	24	21	28	25	14	11	2	1	-
Female	21	21	19	21	24	23	31	27	23	10	4	-
American Indian 1/, both sexes	2	-	1	2	-	-	-	1	2	-	-	-
Male	1	-	-	1	-	-	-	1	1	-	-	-
Female	1	-	1	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	2	2	2	2	3	2	-	-
Male	-	-	-	-	-	-	1	1	-	1	-	-
Female	1	-	1	-	2	2	1	1	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Minnesota												
All races, both sexes	12	20	31	32	49	62	84	139	117	67	22	-
Male	9	14	20	24	22	22	37	49	25	10	4	-
Female	3	6	11	8	27	40	47	90	92	57	18	-
White, both sexes	12	20	24	29	46	58	82	134	116	65	22	-
Male	9	14	16	22	20	21	37	48	25	9	4	-
Female	3	6	8	7	26	37	45	86	91	56	18	-
Black, both sexes	-	-	4	3	2	1	1	4	1	1	-	-
Male	-	-	2	2	2	-	-	1	-	1	-	-
Female	-	-	2	1	-	1	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	2	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	3	1	-	-	1	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	1	-	-
Mississippi												
All races, both sexes	63	76	63	90	103	134	149	163	115	74	15	-
Male	38	46	40	49	51	63	42	40	34	15	1	-
Female	25	30	23	41	52	71	107	123	81	59	14	-
White, both sexes	24	22	24	37	48	75	84	102	73	43	2	-
Male	16	12	16	21	25	31	27	22	19	9	1	-
Female	8	10	8	16	23	44	57	80	54	34	1	-
Black, both sexes	39	53	39	52	55	58	65	61	42	31	13	-
Male	22	33	24	28	26	31	15	18	15	6	-	-
Female	17	20	15	24	29	27	50	43	27	25	13	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Missouri												
All races, both sexes	58	53	46	50	63	88	139	147	124	65	12	-
Male	36	32	26	27	32	40	48	39	25	9	4	-
Female	22	21	20	23	31	48	91	108	99	56	8	-
White, both sexes	25	26	27	33	38	56	118	125	113	58	10	-
Male	17	16	15	17	18	24	42	34	19	9	3	-
Female	8	10	12	16	20	32	76	91	94	49	7	-
Black, both sexes	33	26	18	17	24	30	21	22	10	7	2	-
Male	19	15	11	10	14	15	6	5	5	-	1	-
Female	14	11	7	7	10	15	15	17	5	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	2	-	-	1	-	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Montana												
All races, both sexes	5	2	4	7	5	6	17	28	22	11	2	-
Male	4	1	3	4	4	4	10	7	7	4	-	-
Female	1	1	1	3	1	2	7	21	15	7	2	-
White, both sexes	4	2	4	7	5	6	17	28	22	11	2	-
Male	4	1	3	4	4	4	10	7	7	4	-	-
Female	-	1	1	3	1	2	7	21	15	7	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Nebraska												
All races, both sexes	10	3	11	12	17	25	37	64	53	29	9	-
Male	7	2	6	9	6	10	18	20	14	5	1	-
Female	3	1	5	3	11	15	19	44	39	24	8	-
White, both sexes	7	3	9	11	16	22	36	63	49	29	9	-
Male	6	2	4	8	5	10	18	20	14	5	1	-
Female	1	1	5	3	11	12	18	43	35	24	8	-
Black, both sexes	3	-	2	1	1	2	1	1	3	-	-	-
Male	1	-	2	1	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	2	1	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Nevada												
All races, both sexes	38	47	52	56	52	70	68	74	46	16	2	-
Male	25	39	41	39	31	32	26	28	11	1	-	-
Female	13	8	11	17	21	38	42	46	35	15	2	-
White, both sexes	25	40	46	46	40	57	60	67	44	14	2	-
Male	18	33	37	33	24	28	23	26	11	1	-	-
Female	7	7	9	13	16	29	37	41	33	13	2	-
Black, both sexes	11	6	3	7	9	10	4	3	1	2	-	-
Male	6	5	2	6	5	4	2	1	-	-	-	-
Female	5	1	1	1	4	6	2	2	1	2	-	-
American Indian 1/, both sexes	2	-	-	1	1	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	2	2	2	4	3	1	-	-	-
Male	-	1	2	-	2	-	1	1	-	-	-	-
Female	-	-	1	2	-	2	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
New Hampshire												
All races, both sexes	9	18	12	14	13	16	29	30	36	20	5	-
Male	8	13	6	6	10	9	8	8	4	2	-	-
Female	1	5	6	8	3	7	21	22	32	18	5	-
White, both sexes	9	18	12	13	13	16	29	29	36	20	5	-
Male	8	13	6	6	10	9	8	8	4	2	-	-
Female	1	5	6	7	3	7	21	21	32	18	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
New Jersey												
All races, both sexes	85	80	75	76	94	126	208	239	227	85	22	-
Male	55	59	50	39	42	60	84	79	53	15	5	-
Female	30	21	25	37	52	66	124	160	174	70	17	-
White, both sexes	46	47	48	47	66	103	170	214	207	74	19	-
Male	31	38	36	22	28	50	70	72	49	14	5	-
Female	15	9	12	25	38	53	100	142	158	60	14	-
Black, both sexes	38	30	24	26	25	22	31	22	17	10	3	-
Male	23	18	14	14	13	10	13	7	3	1	-	-
Female	15	12	10	12	12	12	18	15	14	9	3	-
American Indian 1/, both sexes	-	1	-	-	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	3	3	1	5	3	3	1	-	-
Male	1	2	-	3	1	-	1	-	1	-	-	-
Female	-	-	3	-	2	1	4	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
New Mexico												
All races, both sexes	21	13	10	18	21	33	27	31	32	25	3	-
Male	12	7	6	10	11	16	12	8	6	4	1	-
Female	9	6	4	8	10	17	15	23	26	21	2	-
White, both sexes	17	12	9	16	19	28	26	30	30	23	2	-
Male	12	7	6	8	10	16	12	7	5	4	1	-
Female	5	5	3	8	9	12	14	23	25	19	1	-
Black, both sexes	1	1	-	1	1	2	1	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	2	1	-	-	1	-	-
American Indian 1/, both sexes	3	-	1	1	1	2	-	1	2	1	1	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	3	-	1	-	1	2	-	-	1	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	222	262	317	284	329	467	587	737	531	285	85	-
Male	148	152	175	165	174	204	252	235	104	45	10	-
Female	74	110	142	119	155	263	335	502	427	240	75	-
White, both sexes	106	136	174	164	189	318	436	609	445	246	70	-
Male	77	85	98	99	100	143	182	191	83	40	9	-
Female	29	51	76	65	89	175	254	418	362	206	61	-
Black, both sexes	106	122	137	114	131	131	128	111	77	36	15	-
Male	63	63	72	62	69	51	58	35	17	5	1	-
Female	43	59	65	52	62	80	70	76	60	31	14	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	4	6	6	9	18	23	17	9	3	-	-
Male	8	4	5	4	5	10	12	9	4	-	-	-
Female	2	-	1	2	4	8	11	8	5	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
North Carolina												
All races, both sexes	80	85	87	80	110	148	208	200	164	72	20	1
Male	57	61	54	48	61	58	84	52	34	9	-	1
Female	23	24	33	32	49	90	124	148	130	63	20	-
White, both sexes	29	47	45	34	66	90	144	145	122	51	12	1
Male	23	37	34	17	38	36	54	33	25	8	-	1
Female	6	10	11	17	28	54	90	112	97	43	12	-
Black, both sexes	49	37	41	46	42	54	62	54	42	20	8	-
Male	32	24	20	31	21	20	29	19	9	1	-	-
Female	17	13	21	15	21	34	33	35	33	19	8	-
American Indian 1/, both sexes	2	1	-	-	2	2	-	1	-	1	-	-
Male	2	-	-	-	2	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
North Dakota												
All races, both sexes	1	4	4	1	3	5	21	24	29	15	8	-
Male	1	2	2	-	2	3	12	10	6	4	-	-
Female	-	2	2	1	1	2	9	14	23	11	8	-
White, both sexes	-	3	2	1	3	5	20	24	28	15	8	-
Male	-	2	2	-	2	3	12	10	6	4	-	-
Female	-	1	-	1	1	2	8	14	22	11	8	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	1	2	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	2	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Ohio												
All races, both sexes	129	163	148	185	189	273	442	505	438	178	43	-
Male	94	114	96	105	102	117	161	139	96	31	4	-
Female	35	49	52	80	87	156	281	366	342	147	39	-
White, both sexes	78	113	108	134	135	209	369	454	409	158	41	-
Male	57	80	70	76	73	89	128	126	90	27	4	-
Female	21	33	38	58	62	120	241	328	319	131	37	-
Black, both sexes	51	49	40	51	54	61	73	48	29	20	2	-
Male	37	34	26	29	29	27	33	12	6	4	-	-
Female	14	15	14	22	25	34	40	36	23	16	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	3	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	2	-	1	-	-	-	-
Oklahoma												
All races, both sexes	39	27	40	42	38	35	55	95	75	47	11	-
Male	28	16	18	26	27	18	27	16	16	8	4	-
Female	11	11	22	16	11	17	28	79	59	39	7	-
White, both sexes	30	20	30	33	27	31	42	87	62	42	9	-
Male	22	11	13	20	19	15	18	16	13	7	4	-
Female	8	9	17	13	8	16	24	71	49	35	5	-
Black, both sexes	7	5	9	5	10	3	10	5	7	4	1	-
Male	5	3	4	2	8	2	6	-	2	1	-	-
Female	2	2	5	3	2	1	4	5	5	3	1	-
American Indian 1/, both sexes	1	2	1	4	1	1	3	3	6	1	1	-
Male	1	2	1	4	-	1	3	-	1	-	-	-
Female	-	-	-	-	1	-	-	3	5	1	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Oregon												
All races, both sexes	23	21	23	25	41	65	106	130	128	65	12	-
Male	17	18	15	10	21	28	39	42	26	9	2	-
Female	6	3	8	15	20	37	67	88	102	56	10	-
White, both sexes	20	19	23	24	40	64	101	129	127	65	12	-
Male	16	17	15	10	20	28	37	42	26	9	2	-
Female	4	2	8	14	20	36	64	87	101	56	10	-
Black, both sexes	1	2	-	-	-	-	2	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	1	-	2	-	1	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	1	-	-	1	-	-	1	-	1	-	-	-
Pennsylvania												
All races, both sexes	82	78	100	89	127	199	310	339	273	142	17	-
Male	57	55	61	55	52	98	124	119	71	23	1	-
Female	25	23	39	34	75	101	186	220	202	119	16	-
White, both sexes	57	51	73	70	101	170	280	307	252	133	16	-
Male	42	39	47	43	42	85	115	111	65	23	1	-
Female	15	12	26	27	59	85	165	196	187	110	15	-
Black, both sexes	23	26	27	16	25	28	30	31	21	9	1	-
Male	14	15	14	10	10	12	9	8	6	-	-	-
Female	9	11	13	6	15	16	21	23	15	9	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	3	1	-	-	1	-	-	-	-
Male	1	1	-	2	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Rhode Island												
All races, both sexes	4	3	3	6	5	12	15	17	26	9	1	-
Male	3	2	1	3	5	5	9	8	5	2	-	-
Female	1	1	2	3	-	7	6	9	21	7	1	-
White, both sexes	4	3	3	4	4	11	14	17	26	9	-	-
Male	3	2	1	2	4	4	9	8	5	2	-	-
Female	1	1	2	2	-	7	5	9	21	7	-	-
Black, both sexes	-	-	-	2	-	1	1	-	-	-	1	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	39	70	49	61	59	85	132	134	81	33	10	-
Male	25	46	31	34	29	34	46	47	19	7	1	-
Female	14	24	18	27	30	51	86	87	62	26	9	-
White, both sexes	17	29	18	33	32	42	85	82	49	22	4	-
Male	10	18	14	17	14	22	30	31	10	4	-	-
Female	7	11	4	16	18	20	55	51	39	18	4	-
Black, both sexes	21	41	30	28	27	42	47	51	32	11	6	-
Male	15	28	16	17	15	12	16	15	9	3	1	-
Female	6	13	14	11	12	30	31	36	23	8	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
South Dakota												
All races, both sexes	2	-	2	4	2	13	16	29	42	20	10	-
Male	1	-	1	4	2	7	4	8	11	6	2	-
Female	1	-	1	-	-	6	12	21	31	14	8	-
White, both sexes	2	-	2	3	2	11	16	28	42	19	9	-
Male	1	-	1	3	2	5	4	7	11	6	1	-
Female	1	-	1	-	-	6	12	21	31	13	8	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	1	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	77	95	92	82	87	146	231	233	182	80	20	-
Male	40	58	58	45	47	57	86	58	51	16	1	-
Female	37	37	34	37	40	89	145	175	131	64	19	-
White, both sexes	34	54	59	53	45	104	180	188	156	66	14	-
Male	20	37	38	28	25	40	65	45	45	12	1	-
Female	14	17	21	25	20	64	115	143	111	54	13	-
Black, both sexes	42	38	33	29	42	42	51	44	24	14	6	-
Male	19	20	20	17	22	17	21	13	6	4	-	-
Female	23	18	13	12	20	25	30	31	18	10	6	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	-	-	-	-	1	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Texas												
All races, both sexes	244	226	203	228	263	345	483	492	433	196	34	-
Male	155	131	116	125	113	158	175	138	98	27	3	-
Female	89	95	87	103	150	187	308	354	335	169	31	-
White, both sexes	156	150	147	148	190	267	407	432	373	171	26	-
Male	107	85	90	76	76	128	146	118	85	25	2	-
Female	49	65	57	72	114	139	261	314	288	146	24	-
Black, both sexes	82	72	53	73	70	72	72	57	56	24	8	-
Male	45	45	24	43	36	27	28	19	10	2	1	-
Female	37	27	29	30	34	45	44	38	46	22	7	-
American Indian 1/, both sexes	3	-	-	1	-	-	3	1	-	-	-	-
Male	2	-	-	1	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	3	6	3	6	1	2	4	1	-	-
Male	1	1	2	5	1	3	-	-	3	-	-	-
Female	2	3	1	1	2	3	1	2	1	1	-	-
Utah												
All races, both sexes	11	6	7	9	11	21	35	57	45	22	3	-
Male	6	3	4	6	3	10	12	18	10	4	1	-
Female	5	3	3	3	8	11	23	39	35	18	2	-
White, both sexes	11	4	7	8	10	20	33	57	45	22	3	-
Male	6	3	4	5	3	9	11	18	10	4	1	-
Female	5	1	3	3	7	11	22	39	35	18	2	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Vermont												
All races, both sexes	6	10	4	5	7	6	12	21	14	7	2	-
Male	2	8	1	4	4	3	6	4	4	2	-	-
Female	4	2	3	1	3	3	6	17	10	5	2	-
White, both sexes	6	10	3	5	7	6	11	21	14	7	2	-
Male	2	8	1	4	4	3	6	4	4	2	-	-
Female	4	2	2	1	3	3	5	17	10	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	51	47	48	50	83	87	138	168	111	66	17	-
Male	26	25	29	26	42	44	55	56	33	11	1	-
Female	25	22	19	24	41	43	83	112	78	55	16	-
White, both sexes	23	21	25	24	40	64	98	131	87	54	12	-
Male	13	7	18	9	17	33	38	45	24	9	1	-
Female	10	14	7	15	23	31	60	86	63	45	11	-
Black, both sexes	27	26	22	25	43	22	39	35	22	10	5	-
Male	12	18	10	16	25	11	16	10	9	1	-	-
Female	15	8	12	9	18	11	23	25	13	9	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	1	1	1	-	2	-	-
Male	1	-	1	1	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Washington												
All races, both sexes	41	66	50	69	57	118	212	231	215	98	26	-
Male	32	50	34	36	30	56	85	67	57	19	2	-
Female	9	16	16	33	27	62	127	164	158	79	24	-
White, both sexes	34	53	40	58	54	102	191	218	200	96	26	-
Male	28	40	28	31	27	50	76	59	52	18	2	-
Female	6	13	12	27	27	52	115	159	148	78	24	-
Black, both sexes	5	8	8	7	2	7	9	8	4	-	-	-
Male	3	5	6	3	2	3	3	5	2	-	-	-
Female	2	3	2	4	-	4	6	3	2	-	-	-
American Indian 1/, both sexes	-	1	-	3	1	1	1	-	1	-	-	-
Male	-	1	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	2	1	-	8	11	5	10	2	-	-
Male	1	4	-	-	-	3	6	3	3	1	-	-
Female	1	-	2	1	-	5	5	2	7	1	-	-
West Virginia												
All races, both sexes	15	18	23	24	24	43	51	44	63	28	8	-
Male	11	10	17	15	14	16	17	10	9	2	-	-
Female	4	8	6	9	10	27	34	34	54	26	8	-
White, both sexes	14	17	22	22	22	39	48	43	58	28	7	-
Male	10	10	16	14	13	14	16	9	8	2	-	-
Female	4	7	6	8	9	25	32	34	50	26	7	-
Black, both sexes	1	1	1	2	2	4	3	1	5	-	1	-
Male	1	-	1	1	1	2	1	1	1	-	-	-
Female	-	1	-	1	1	2	2	-	4	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Wisconsin												
All races, both sexes	29	27	36	36	57	81	134	174	154	73	11	-
Male	19	19	25	22	28	38	43	49	28	7	-	-
Female	10	8	11	14	29	43	91	125	126	66	11	-
White, both sexes	21	22	27	32	48	68	129	169	153	72	11	-
Male	15	15	21	20	24	32	42	48	28	7	-	-
Female	6	7	6	12	24	36	87	121	125	65	11	-
Black, both sexes	8	4	9	4	7	13	3	2	1	1	-	-
Male	4	3	4	2	3	6	1	-	-	-	-	-
Female	4	1	5	2	4	7	2	2	1	1	-	-
American Indian 1/, both sexes	-	1	-	-	2	-	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Wyoming												
All races, both sexes	4	2	4	3	2	5	9	8	10	7	-	-
Male	4	2	2	3	2	2	2	1	2	2	-	-
Female	-	-	2	-	-	3	7	7	8	5	-	-
White, both sexes	4	2	4	3	2	5	9	8	10	7	-	-
Male	4	2	2	3	2	2	2	1	2	2	-	-
Female	-	-	2	-	-	3	7	7	8	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
United States												
All races, both sexes	14,024	-	-	1	-	1	3	11	34	53	127	275
Male	5,096	-	-	1	-	-	2	8	23	29	86	172
Female	8,928	-	-	-	-	1	1	3	11	24	41	103
White, both sexes	11,303	-	-	1	-	-	2	6	16	29	72	174
Male	3,977	-	-	1	-	-	1	5	11	16	52	118
Female	7,326	-	-	-	-	-	1	1	5	13	20	56
Black, both sexes	2,338	-	-	-	-	1	1	5	18	24	53	93
Male	962	-	-	-	-	-	1	3	12	13	32	48
Female	1,376	-	-	-	-	1	-	2	6	11	21	45
American Indian 1/, both sexes	50	-	-	-	-	-	-	-	-	-	-	5
Male	23	-	-	-	-	-	-	-	-	-	-	5
Female	27	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	333	-	-	-	-	-	-	-	-	-	2	3
Male	134	-	-	-	-	-	-	-	-	-	2	1
Female	199	-	-	-	-	-	-	-	-	-	-	2
Alabama												
All races, both sexes	294	-	-	-	-	-	-	-	3	-	6	16
Male	122	-	-	-	-	-	-	-	1	-	1	10
Female	172	-	-	-	-	-	-	-	2	-	5	6
White, both sexes	172	-	-	-	-	-	-	-	2	-	1	5
Male	61	-	-	-	-	-	-	-	-	-	-	4
Female	111	-	-	-	-	-	-	-	2	-	1	1
Black, both sexes	120	-	-	-	-	-	-	-	1	-	5	11
Male	59	-	-	-	-	-	-	-	1	-	1	6
Female	61	-	-	-	-	-	-	-	-	-	4	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Alaska												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	191	-	-	-	-	-	-	1	-	-	-	2
Male	63	-	-	-	-	-	-	1	-	-	-	1
Female	128	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	174	-	-	-	-	-	-	1	-	-	-	1
Male	56	-	-	-	-	-	-	1	-	-	-	-
Female	118	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Arkansas												
All races, both sexes	146	-	-	-	-	-	-	-	-	1	3	5
Male	62	-	-	-	-	-	-	-	-	-	3	4
Female	84	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	112	-	-	-	-	-	-	-	-	-	2	2
Male	42	-	-	-	-	-	-	-	-	-	2	2
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	34	-	-	-	-	-	-	-	-	1	1	3
Male	20	-	-	-	-	-	-	-	-	-	1	2
Female	14	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,910	-	-	-	-	-	1	-	2	5	11	33
Male	708	-	-	-	-	-	1	-	2	3	8	17
Female	1,202	-	-	-	-	-	-	-	-	2	3	16
White, both sexes	1,546	-	-	-	-	-	1	-	2	4	7	20
Male	575	-	-	-	-	-	1	-	2	2	5	14
Female	971	-	-	-	-	-	-	-	-	2	2	6
Black, both sexes	192	-	-	-	-	-	-	-	-	1	3	13
Male	70	-	-	-	-	-	-	-	-	1	2	3
Female	122	-	-	-	-	-	-	-	-	-	1	10
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	164	-	-	-	-	-	-	-	-	-	1	-
Male	59	-	-	-	-	-	-	-	-	-	1	-
Female	105	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Colorado												
All races, both sexes	132	-	-	-	-	-	-	-	-	-	-	1
Male	56	-	-	-	-	-	-	-	-	-	-	1
Female	76	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	125	-	-	-	-	-	-	-	-	-	-	1
Male	52	-	-	-	-	-	-	-	-	-	-	1
Female	73	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	227	-	-	-	-	-	-	-	1	-	2	2
Male	79	-	-	-	-	-	-	-	1	-	2	1
Female	148	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	206	-	-	-	-	-	-	-	1	-	2	1
Male	69	-	-	-	-	-	-	-	1	-	2	1
Female	137	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Delaware												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Florida												
All races, both sexes	981	-	-	1	-	1	-	1	3	5	8	21
Male	369	-	-	1	-	-	-	1	3	1	5	10
Female	612	-	-	-	-	1	-	-	-	4	3	11
White, both sexes	818	-	-	1	-	-	-	-	2	4	6	15
Male	304	-	-	1	-	-	-	-	2	1	3	8
Female	514	-	-	-	-	-	-	-	-	3	3	7
Black, both sexes	160	-	-	-	-	1	-	1	1	1	2	5
Male	63	-	-	-	-	-	-	1	1	-	2	1
Female	97	-	-	-	-	1	-	-	-	1	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	629	-	-	-	-	-	-	2	3	7	14	19
Male	240	-	-	-	-	-	-	1	2	5	12	8
Female	389	-	-	-	-	-	-	1	1	2	2	11
White, both sexes	384	-	-	-	-	-	-	-	1	4	9	13
Male	136	-	-	-	-	-	-	-	1	2	7	6
Female	248	-	-	-	-	-	-	-	-	2	2	7
Black, both sexes	239	-	-	-	-	-	-	2	2	3	5	6
Male	102	-	-	-	-	-	-	1	1	3	5	2
Female	137	-	-	-	-	-	-	1	1	-	-	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Hawaii												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	1	-
Male	34	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	65	-	-	-	-	-	-	-	-	-	1	-
Male	31	-	-	-	-	-	-	-	-	-	1	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Illinois												
All races, both sexes	550	-	-	-	-	-	-	-	-	2	2	9
Male	169	-	-	-	-	-	-	-	-	-	1	7
Female	381	-	-	-	-	-	-	-	-	2	1	2
White, both sexes	426	-	-	-	-	-	-	-	-	1	1	5
Male	122	-	-	-	-	-	-	-	-	-	1	5
Female	304	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	113	-	-	-	-	-	-	-	-	1	1	4
Male	46	-	-	-	-	-	-	-	-	-	-	2
Female	67	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	358	-	-	-	-	-	-	1	3	4	5	11
Male	141	-	-	-	-	-	-	1	3	2	3	6
Female	217	-	-	-	-	-	-	-	-	2	2	5
White, both sexes	311	-	-	-	-	-	-	1	1	1	3	9
Male	110	-	-	-	-	-	-	1	1	-	1	4
Female	201	-	-	-	-	-	-	-	-	1	2	5
Black, both sexes	47	-	-	-	-	-	-	-	2	3	2	2
Male	31	-	-	-	-	-	-	-	2	2	2	2
Female	16	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Iowa												
All races, both sexes	111	-	-	-	-	-	-	1	-	-	-	-
Male	33	-	-	-	-	-	-	1	-	-	-	-
Female	78	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	109	-	-	-	-	-	-	1	-	-	-	-
Male	32	-	-	-	-	-	-	1	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	123	-	-	-	-	-	-	-	-	2	-	-
Male	38	-	-	-	-	-	-	-	-	1	-	-
Female	85	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	107	-	-	-	-	-	-	-	-	1	-	-
Male	30	-	-	-	-	-	-	-	-	1	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Kentucky												
All races, both sexes	194	-	-	-	-	-	-	-	-	-	4	7
Male	71	-	-	-	-	-	-	-	-	-	2	5
Female	123	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	168	-	-	-	-	-	-	-	-	-	3	7
Male	63	-	-	-	-	-	-	-	-	-	2	5
Female	105	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	224	-	-	-	-	-	-	-	3	-	3	8
Male	90	-	-	-	-	-	-	-	2	-	1	7
Female	134	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	128	-	-	-	-	-	-	-	-	-	-	3
Male	44	-	-	-	-	-	-	-	-	-	-	3
Female	84	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	96	-	-	-	-	-	-	-	3	-	3	5
Male	46	-	-	-	-	-	-	-	2	-	1	4
Female	50	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Maine												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	1	3
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	47	-	-	-	-	-	-	-	-	-	1	2
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	186	-	-	-	-	-	-	-	-	-	2	1
Male	60	-	-	-	-	-	-	-	-	-	1	1
Female	126	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	127	-	-	-	-	-	-	-	-	-	1	-
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	56	-	-	-	-	-	-	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	35	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Massachusetts												
All races, both sexes	307	-	-	-	-	-	-	1	1	1	2	-
Male	115	-	-	-	-	-	-	-	1	1	1	-
Female	192	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	286	-	-	-	-	-	-	1	1	1	2	-
Male	107	-	-	-	-	-	-	-	1	1	1	-
Female	179	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	432	-	-	-	-	-	-	-	-	2	3	2
Male	148	-	-	-	-	-	-	-	-	1	2	2
Female	284	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	358	-	-	-	-	-	-	-	-	1	1	1
Male	116	-	-	-	-	-	-	-	-	-	1	1
Female	242	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	69	-	-	-	-	-	-	-	-	1	2	1
Male	30	-	-	-	-	-	-	-	-	1	1	1
Female	39	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Minnesota												
All races, both sexes	308	-	-	-	-	-	-	-	-	1	3	3
Male	107	-	-	-	-	-	-	-	-	1	2	1
Female	201	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	296	-	-	-	-	-	-	-	-	1	2	3
Male	101	-	-	-	-	-	-	-	-	1	2	1
Female	195	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	247	-	-	-	-	-	-	-	-	-	5	9
Male	93	-	-	-	-	-	-	-	-	-	3	7
Female	154	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	126	-	-	-	-	-	-	-	-	-	2	2
Male	46	-	-	-	-	-	-	-	-	-	1	2
Female	80	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	120	-	-	-	-	-	-	-	-	-	3	7
Male	46	-	-	-	-	-	-	-	-	-	2	5
Female	74	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Missouri												
All races, both sexes	233	-	-	-	-	-	-	-	1	1	5	7
Male	77	-	-	-	-	-	-	-	1	-	4	7
Female	156	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	204	-	-	-	-	-	-	-	-	-	2	7
Male	64	-	-	-	-	-	-	-	-	-	2	7
Female	140	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	28	-	-	-	-	-	-	-	1	1	3	-
Male	12	-	-	-	-	-	-	-	1	-	2	-
Female	16	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Nebraska												
All races, both sexes	95	-	-	-	-	-	-	-	-	-	-	1
Male	37	-	-	-	-	-	-	-	-	-	-	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	89	-	-	-	-	-	-	-	-	-	-	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	70	-	-	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	62	-	-	-	-	-	-	-	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
New Hampshire												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	371	-	-	-	-	-	-	-	-	1	1	8
Male	142	-	-	-	-	-	-	-	-	-	-	7
Female	229	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	308	-	-	-	-	-	-	-	-	-	-	6
Male	118	-	-	-	-	-	-	-	-	-	-	5
Female	190	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	58	-	-	-	-	-	-	-	-	1	1	2
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	34	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
New Mexico												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	947	-	-	-	-	-	1	-	-	4	6	13
Male	332	-	-	-	-	-	1	-	-	3	6	9
Female	615	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	724	-	-	-	-	-	-	-	-	1	3	7
Male	253	-	-	-	-	-	-	-	-	1	3	5
Female	471	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	200	-	-	-	-	-	1	-	-	3	3	6
Male	65	-	-	-	-	-	1	-	-	2	3	4
Female	135	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
North Carolina												
All races, both sexes	413	-	-	-	-	-	-	-	3	4	4	9
Male	147	-	-	-	-	-	-	-	1	3	2	4
Female	266	-	-	-	-	-	-	-	2	1	2	5
White, both sexes	274	-	-	-	-	-	-	-	1	2	3	3
Male	88	-	-	-	-	-	-	-	-	1	2	1
Female	186	-	-	-	-	-	-	-	1	1	1	2
Black, both sexes	132	-	-	-	-	-	-	-	2	2	1	6
Male	56	-	-	-	-	-	-	-	1	2	-	3
Female	76	-	-	-	-	-	-	-	1	-	1	3
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Ohio												
All races, both sexes	663	-	-	-	-	-	-	1	2	1	3	12
Male	233	-	-	-	-	-	-	1	1	-	3	8
Female	430	-	-	-	-	-	-	-	1	1	-	4
White, both sexes	579	-	-	-	-	-	-	-	1	1	3	10
Male	198	-	-	-	-	-	-	-	-	-	3	7
Female	381	-	-	-	-	-	-	-	1	1	-	3
Black, both sexes	83	-	-	-	-	-	-	1	1	-	-	2
Male	35	-	-	-	-	-	-	1	1	-	-	1
Female	48	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	139	-	-	-	-	-	-	1	-	-	-	5
Male	48	-	-	-	-	-	-	1	-	-	-	4
Female	91	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	116	-	-	-	-	-	-	1	-	-	-	2
Male	35	-	-	-	-	-	-	1	-	-	-	1
Female	81	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	3
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Oregon												
All races, both sexes	248	-	-	-	-	-	-	-	-	1	-	4
Male	96	-	-	-	-	-	-	-	-	1	-	4
Female	152	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	244	-	-	-	-	-	-	-	-	1	-	4
Male	95	-	-	-	-	-	-	-	-	1	-	4
Female	149	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	619	-	-	-	-	-	-	-	1	1	6	12
Male	243	-	-	-	-	-	-	-	-	-	4	4
Female	376	-	-	-	-	-	-	-	1	1	2	8
White, both sexes	548	-	-	-	-	-	-	-	-	-	4	10
Male	215	-	-	-	-	-	-	-	-	-	3	3
Female	333	-	-	-	-	-	-	-	-	-	1	7
Black, both sexes	68	-	-	-	-	-	-	-	1	1	2	2
Male	27	-	-	-	-	-	-	-	-	-	1	1
Female	41	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Rhode Island												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	198	-	-	-	-	-	-	-	2	-	5	8
Male	92	-	-	-	-	-	-	-	-	-	3	5
Female	106	-	-	-	-	-	-	-	2	-	2	3
White, both sexes	109	-	-	-	-	-	-	-	-	-	-	5
Male	47	-	-	-	-	-	-	-	-	-	-	4
Female	62	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	88	-	-	-	-	-	-	-	2	-	5	3
Male	44	-	-	-	-	-	-	-	-	-	3	1
Female	44	-	-	-	-	-	-	-	2	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
South Dakota												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	269	-	-	-	-	-	-	1	1	3	6	6
Male	91	-	-	-	-	-	-	-	1	2	3	3
Female	178	-	-	-	-	-	-	1	-	1	3	3
White, both sexes	206	-	-	-	-	-	-	-	-	1	3	3
Male	71	-	-	-	-	-	-	-	-	1	1	2
Female	135	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	61	-	-	-	-	-	-	1	1	2	3	3
Male	19	-	-	-	-	-	-	-	1	1	2	1
Female	42	-	-	-	-	-	-	1	-	1	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Texas												
All races, both sexes	708	-	-	-	-	-	1	-	3	3	6	11
Male	230	-	-	-	-	-	-	-	2	2	5	6
Female	478	-	-	-	-	-	1	-	1	1	1	5
White, both sexes	591	-	-	-	-	-	1	-	2	2	4	5
Male	192	-	-	-	-	-	-	-	1	2	3	3
Female	399	-	-	-	-	-	1	-	1	-	1	2
Black, both sexes	109	-	-	-	-	-	-	-	1	1	2	6
Male	35	-	-	-	-	-	-	-	1	-	2	3
Female	74	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	81	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Vermont												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	301	-	-	-	-	-	-	-	-	2	3	12
Male	118	-	-	-	-	-	-	-	-	2	1	12
Female	183	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	212	-	-	-	-	-	-	-	-	1	2	10
Male	80	-	-	-	-	-	-	-	-	1	1	10
Female	132	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	84	-	-	-	-	-	-	-	-	1	1	2
Male	35	-	-	-	-	-	-	-	-	1	-	2
Female	49	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Washington												
All races, both sexes	248	-	-	-	-	-	-	-	-	-	4	4
Male	83	-	-	-	-	-	-	-	-	-	4	3
Female	165	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	226	-	-	-	-	-	-	-	-	-	3	2
Male	74	-	-	-	-	-	-	-	-	-	3	2
Female	152	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
West Virginia												
All races, both sexes	160	-	-	-	-	-	-	-	1	1	1	3
Male	73	-	-	-	-	-	-	-	1	1	1	1
Female	87	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	151	-	-	-	-	-	-	-	1	1	1	3
Male	69	-	-	-	-	-	-	-	1	1	1	1
Female	82	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Wisconsin												
All races, both sexes	215	-	-	-	-	-	-	1	1	1	1	2
Male	62	-	-	-	-	-	-	1	1	-	1	-
Female	153	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	208	-	-	-	-	-	-	1	1	1	1	1
Male	60	-	-	-	-	-	-	1	1	-	1	-
Female	148	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
United States												
All races, both sexes	487	578	698	794	1,002	1,466	2,174	2,569	2,180	1,219	352	-
Male	322	361	423	445	506	592	771	708	440	172	35	-
Female	165	217	275	349	496	874	1,403	1,861	1,740	1,047	317	-
White, both sexes	310	389	508	571	778	1,177	1,805	2,243	1,885	1,053	284	-
Male	215	248	324	324	384	479	630	628	370	143	28	-
Female	95	141	184	247	394	698	1,175	1,615	1,515	910	256	-
Black, both sexes	165	172	171	197	195	242	311	252	237	141	60	-
Male	98	103	90	105	112	94	117	55	52	23	4	-
Female	67	69	81	92	83	148	194	197	185	118	56	-
American Indian 1/, both sexes	1	2	2	7	3	6	5	7	6	5	1	-
Male	1	-	1	5	1	2	3	2	2	1	-	-
Female	-	2	1	2	2	4	2	5	4	4	1	-
Asian or Pacific Islander 2/, both sexes	11	15	17	19	26	41	53	67	52	20	7	-
Male	8	10	8	11	9	17	21	23	16	5	3	-
Female	3	5	9	8	17	24	32	44	36	15	4	-
Alabama												
All races, both sexes	18	21	16	31	28	27	39	46	27	11	5	-
Male	13	16	8	15	13	15	16	9	3	2	-	-
Female	5	5	8	16	15	12	23	37	24	9	5	-
White, both sexes	6	8	8	18	19	16	24	38	17	8	2	-
Male	4	8	4	9	8	8	9	5	1	1	-	-
Female	2	-	4	9	11	8	15	33	16	7	2	-
Black, both sexes	11	13	7	13	9	11	15	8	10	3	3	-
Male	8	8	3	6	5	7	7	4	2	1	-	-
Female	3	5	4	7	4	4	8	4	8	2	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Alaska												
All races, both sexes	-	-	-	1	1	2	1	1	2	-	1	-
Male	-	-	-	-	1	1	-	1	2	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	1	-
White, both sexes	-	-	-	1	-	2	1	1	-	-	1	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	11	5	10	21	25	26	34	36	10	6	-
Male	4	5	4	6	9	8	5	10	6	3	1	-
Female	-	6	1	4	12	17	21	24	30	7	5	-
White, both sexes	4	6	5	6	19	25	25	31	36	9	6	-
Male	4	3	4	4	8	8	4	10	6	3	1	-
Female	-	3	1	2	11	17	21	21	30	6	5	-
Black, both sexes	-	4	-	1	1	-	1	1	-	1	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	3	-	1	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	3	1	-	-	1	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Arkansas												
All races, both sexes	9	11	6	11	6	15	22	19	20	13	5	-
Male	4	9	5	6	4	4	11	8	3	1	-	-
Female	5	2	1	5	2	11	11	11	17	12	5	-
White, both sexes	2	6	4	9	6	13	18	15	19	11	5	-
Male	1	4	3	5	4	3	8	6	3	1	-	-
Female	1	2	1	4	2	10	10	9	16	10	5	-
Black, both sexes	7	5	2	2	-	2	4	4	1	2	-	-
Male	3	5	2	1	-	1	3	2	-	-	-	-
Female	4	-	-	1	-	1	1	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	51	69	84	97	124	211	320	393	308	161	40	-
Male	29	41	52	61	67	76	112	128	81	25	5	-
Female	22	28	32	36	57	135	208	265	227	136	35	-
White, both sexes	37	49	68	73	98	168	260	331	257	139	32	-
Male	23	29	45	47	54	61	90	108	70	20	4	-
Female	14	20	23	26	44	107	170	223	187	119	28	-
Black, both sexes	11	17	12	15	14	19	30	18	22	11	6	-
Male	4	11	5	8	9	5	8	6	6	2	-	-
Female	7	6	7	7	5	14	22	12	16	9	6	-
American Indian 1/, both sexes	-	-	-	2	-	1	1	3	-	1	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	3	4	7	12	23	29	41	29	10	2	-
Male	2	1	2	5	4	9	13	13	5	3	1	-
Female	1	2	2	2	8	14	16	28	24	7	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Colorado												
All races, both sexes	4	5	9	5	9	19	23	28	13	14	2	-
Male	2	4	6	5	6	9	8	10	-	4	1	-
Female	2	1	3	-	3	10	15	18	13	10	1	-
White, both sexes	4	4	6	5	9	18	22	28	13	14	1	-
Male	2	3	4	5	6	8	8	10	-	4	1	-
Female	2	1	2	-	3	10	14	18	13	10	-	-
Black, both sexes	-	-	2	-	-	1	-	-	-	-	1	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	6	16	10	11	10	28	38	39	39	20	5	-
Male	5	10	4	9	5	9	12	14	5	2	-	-
Female	1	6	6	2	5	19	26	25	34	18	5	-
White, both sexes	4	12	8	9	9	24	37	37	37	20	5	-
Male	4	8	2	7	5	8	11	13	5	2	-	-
Female	-	4	6	2	4	16	26	24	32	18	5	-
Black, both sexes	2	3	1	1	1	4	1	2	2	-	-	-
Male	1	2	1	1	-	1	1	1	-	-	-	-
Female	1	1	-	-	1	3	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Delaware												
All races, both sexes	3	1	1	1	-	2	3	4	3	-	2	-
Male	2	1	1	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	3	4	3	-	2	-
White, both sexes	1	-	1	1	-	2	3	4	2	-	2	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	3	4	2	-	2	-
Black, both sexes	2	1	-	-	-	-	-	-	1	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	3	1	1	8	4	4	2	-	-
Male	-	-	-	1	1	1	5	1	-	1	-	-
Female	-	-	1	2	-	-	3	3	4	1	-	-
White, both sexes	-	-	-	1	-	-	-	2	1	1	-	-
Male	-	-	-	1	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	1	2	1	1	8	2	3	1	-	-
Male	-	-	-	-	1	1	5	-	-	-	-	-
Female	-	-	1	2	-	-	3	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Florida												
All races, both sexes	38	24	62	52	70	95	156	179	160	75	30	-
Male	27	16	38	30	30	33	57	61	36	14	6	-
Female	11	8	24	22	40	62	99	118	124	61	24	-
White, both sexes	27	16	43	37	57	82	136	164	142	62	24	-
Male	21	11	27	21	26	27	48	57	33	13	5	-
Female	6	5	16	16	31	55	88	107	109	49	19	-
Black, both sexes	11	7	19	15	13	13	20	15	17	13	6	-
Male	6	4	11	9	4	6	9	4	3	1	1	-
Female	5	3	8	6	9	7	11	11	14	12	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Georgia												
All races, both sexes	39	44	40	49	53	62	81	104	62	37	13	-
Male	34	26	23	23	28	24	18	22	9	5	-	-
Female	5	18	17	26	25	38	63	82	53	32	13	-
White, both sexes	17	25	19	22	29	40	58	71	47	24	5	-
Male	16	13	10	8	16	18	12	20	5	2	-	-
Female	1	12	9	14	13	22	46	51	42	22	5	-
Black, both sexes	22	17	20	26	24	22	23	32	15	12	8	-
Male	18	12	13	15	12	6	6	2	4	2	-	-
Female	4	5	7	11	12	16	17	30	11	10	8	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Hawaii												
All races, both sexes	2	3	6	3	7	5	9	14	13	5	5	-
Male	2	3	4	2	4	2	3	5	6	-	2	-
Female	-	-	2	1	3	3	6	9	7	5	3	-
White, both sexes	-	1	1	-	1	1	3	-	-	1	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	5	3	6	4	6	14	13	4	5	-
Male	2	2	3	2	4	1	3	5	6	-	2	-
Female	-	-	2	1	2	3	3	9	7	4	3	-
Idaho												
All races, both sexes	-	3	3	1	6	6	8	7	13	7	-	-
Male	-	2	2	-	3	3	2	2	1	-	-	-
Female	-	1	1	1	3	3	6	5	12	7	-	-
White, both sexes	-	3	3	1	6	6	8	7	13	7	-	-
Male	-	2	2	-	3	3	2	2	1	-	-	-
Female	-	1	1	1	3	3	6	5	12	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Illinois												
All races, both sexes	18	23	19	29	48	58	81	90	84	66	21	-
Male	13	14	10	12	27	21	25	20	9	7	3	-
Female	5	9	9	17	21	37	56	70	75	59	18	-
White, both sexes	9	14	12	17	33	42	67	78	71	55	21	-
Male	8	10	6	7	17	16	21	18	6	4	3	-
Female	1	4	6	10	16	26	46	60	65	51	18	-
Black, both sexes	9	9	7	10	12	15	12	11	12	10	-	-
Male	5	4	4	4	10	5	4	2	3	3	-	-
Female	4	5	3	6	2	10	8	9	9	7	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	1	2	1	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	3	1	2	1	1	1	-	-
Indiana												
All races, both sexes	16	21	20	17	20	37	60	58	51	29	5	-
Male	10	16	11	11	9	16	20	15	12	6	-	-
Female	6	5	9	6	11	21	40	43	39	23	5	-
White, both sexes	12	15	16	15	19	32	51	56	49	27	4	-
Male	8	10	8	10	9	14	14	13	12	5	-	-
Female	4	5	8	5	10	18	37	43	37	22	4	-
Black, both sexes	4	6	4	2	1	5	9	2	2	2	1	-
Male	2	6	3	1	-	2	6	2	-	1	-	-
Female	2	-	1	1	1	3	3	-	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Iowa												
All races, both sexes	1	1	3	5	9	11	18	15	28	12	7	-
Male	-	1	2	1	5	2	9	3	8	1	-	-
Female	1	-	1	4	4	9	9	12	20	11	7	-
White, both sexes	1	-	3	5	9	10	18	15	28	12	7	-
Male	-	-	2	1	5	2	9	3	8	1	-	-
Female	1	-	1	4	4	8	9	12	20	11	7	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	8	4	7	6	12	20	28	17	15	3	-
Male	1	5	2	3	3	7	5	6	4	1	-	-
Female	-	3	2	4	3	5	15	22	13	14	3	-
White, both sexes	-	5	4	7	6	9	15	28	16	13	3	-
Male	-	2	2	3	3	4	4	6	4	1	-	-
Female	-	3	2	4	3	5	11	22	12	12	3	-
Black, both sexes	1	3	-	-	-	3	4	-	1	2	-	-
Male	1	3	-	-	-	3	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Kentucky												
All races, both sexes	8	6	7	17	9	20	30	33	27	19	7	-
Male	3	4	3	8	5	10	14	10	5	2	-	-
Female	5	2	4	9	4	10	16	23	22	17	7	-
White, both sexes	6	6	5	13	8	16	26	30	24	18	6	-
Male	3	4	3	7	4	8	12	9	4	2	-	-
Female	3	2	2	6	4	8	14	21	20	16	6	-
Black, both sexes	2	-	2	3	1	4	4	3	3	1	1	-
Male	-	-	-	1	1	2	2	1	1	-	-	-
Female	2	-	2	2	-	2	2	2	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	11	10	10	18	20	26	33	40	25	12	5	-
Male	7	5	5	11	11	9	10	13	3	5	1	-
Female	4	5	5	7	9	17	23	27	22	7	4	-
White, both sexes	4	4	7	10	11	14	23	27	16	7	2	-
Male	2	3	4	8	4	1	8	9	1	1	-	-
Female	2	1	3	2	7	13	15	18	15	6	2	-
Black, both sexes	7	6	3	8	9	12	10	13	9	5	3	-
Male	5	2	1	3	7	8	2	4	2	4	1	-
Female	2	4	2	5	2	4	8	9	7	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Maine												
All races, both sexes	1	1	1	5	4	6	8	5	8	4	2	-
Male	1	1	-	3	2	2	4	2	2	-	-	-
Female	-	-	1	2	2	4	4	3	6	4	2	-
White, both sexes	1	1	1	5	4	5	8	5	8	4	2	-
Male	1	1	-	3	2	1	4	2	2	-	-	-
Female	-	-	1	2	2	4	4	3	6	4	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7	8	9	10	14	18	26	37	29	21	4	-
Male	3	4	5	7	7	7	7	9	6	2	1	-
Female	4	4	4	3	7	11	19	28	23	19	3	-
White, both sexes	4	3	6	5	6	12	18	31	24	13	4	-
Male	2	3	3	4	3	4	5	8	6	-	1	-
Female	2	-	3	1	3	8	13	23	18	13	3	-
Black, both sexes	3	5	3	5	8	6	7	6	4	7	-	-
Male	1	1	2	3	4	3	2	1	-	2	-	-
Female	2	4	1	2	4	3	5	5	4	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Massachusetts												
All races, both sexes	8	13	15	22	21	37	47	64	41	28	6	-
Male	4	9	12	13	10	17	14	16	10	6	1	-
Female	4	4	3	9	11	20	33	48	31	22	5	-
White, both sexes	7	12	13	19	20	34	43	62	37	28	6	-
Male	3	9	10	13	9	16	13	16	8	6	1	-
Female	4	3	3	6	11	18	30	46	29	22	5	-
Black, both sexes	1	1	2	3	1	2	4	2	3	-	-	-
Male	1	-	2	-	1	1	1	-	1	-	-	-
Female	-	1	-	3	-	1	3	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	14	11	14	14	29	60	57	83	91	40	12	-
Male	8	5	11	6	9	24	25	33	15	7	-	-
Female	6	6	3	8	20	36	32	50	76	33	12	-
White, both sexes	8	7	12	8	23	49	54	73	79	31	11	-
Male	6	3	9	2	7	16	24	29	13	5	-	-
Female	2	4	3	6	16	33	30	44	66	26	11	-
Black, both sexes	5	4	2	6	6	11	3	9	11	7	1	-
Male	2	2	2	4	2	8	1	3	2	1	-	-
Female	3	2	-	2	4	3	2	6	9	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Minnesota												
All races, both sexes	2	8	18	21	35	26	34	54	56	37	10	-
Male	1	6	11	17	16	7	16	16	5	6	2	-
Female	1	2	7	4	19	19	18	38	51	31	8	-
White, both sexes	2	8	16	19	32	23	34	53	56	37	10	-
Male	1	6	10	15	14	6	16	16	5	6	2	-
Female	1	2	6	4	18	17	18	37	51	31	8	-
Black, both sexes	-	-	1	2	2	1	-	1	-	-	-	-
Male	-	-	1	2	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	15	20	11	17	27	20	28	41	28	18	8	-
Male	11	11	8	10	13	6	6	7	5	5	1	-
Female	4	9	3	7	14	14	22	34	23	13	7	-
White, both sexes	7	7	6	9	14	11	10	31	16	10	1	-
Male	4	3	4	6	6	5	3	5	3	3	1	-
Female	3	4	2	3	8	6	7	26	13	7	-	-
Black, both sexes	8	12	5	8	13	9	18	10	12	8	7	-
Male	7	7	4	4	7	1	3	2	2	2	-	-
Female	1	5	1	4	6	8	15	8	10	6	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Missouri												
All races, both sexes	6	7	10	9	16	18	53	33	35	27	5	-
Male	3	6	3	7	9	9	17	6	3	1	1	-
Female	3	1	7	2	7	9	36	27	32	26	4	-
White, both sexes	5	5	8	7	14	14	50	29	33	25	5	-
Male	3	4	3	5	7	6	17	6	2	1	1	-
Female	2	1	5	2	7	8	33	23	31	24	4	-
Black, both sexes	1	2	2	2	2	3	3	4	2	2	-	-
Male	-	2	-	2	2	2	-	-	1	-	-	-
Female	1	-	2	-	-	1	3	4	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	1	2	1	4	7	10	7	1	1	-
Male	1	-	1	1	-	2	5	3	2	-	-	-
Female	-	-	-	1	1	2	2	7	5	1	1	-
White, both sexes	1	-	1	2	1	4	7	10	7	1	1	-
Male	1	-	1	1	-	2	5	3	2	-	-	-
Female	-	-	-	1	1	2	2	7	5	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Nebraska												
All races, both sexes	-	1	3	4	7	6	15	23	24	10	1	-
Male	-	1	1	3	1	5	7	9	6	3	-	-
Female	-	-	2	1	6	1	8	14	18	7	1	-
White, both sexes	-	1	2	4	7	6	14	23	20	10	1	-
Male	-	1	-	3	1	5	7	9	6	3	-	-
Female	-	-	2	1	6	1	7	14	14	7	1	-
Black, both sexes	-	-	1	-	-	-	1	-	3	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Nevada												
All races, both sexes	1	2	2	3	4	6	17	16	9	6	2	-
Male	1	1	1	2	4	3	5	5	1	-	-	-
Female	-	1	1	1	-	3	12	11	8	6	2	-
White, both sexes	1	2	1	1	3	6	14	15	9	6	2	-
Male	1	1	1	1	3	3	4	5	1	-	-	-
Female	-	1	-	-	-	3	10	10	8	6	2	-
Black, both sexes	-	-	-	1	1	-	2	1	-	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
New Hampshire												
All races, both sexes	2	4	3	3	5	6	6	5	9	6	1	-
Male	2	3	1	2	3	3	3	2	-	1	-	-
Female	-	1	2	1	2	3	3	3	9	5	1	-
White, both sexes	2	4	3	3	5	6	6	5	9	6	1	-
Male	2	3	1	2	3	3	3	2	-	1	-	-
Female	-	1	2	1	2	3	3	3	9	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	14	10	23	20	22	37	69	55	79	27	5	-
Male	8	6	14	9	5	18	27	18	24	5	1	-
Female	6	4	9	11	17	19	42	37	55	22	4	-
White, both sexes	8	5	15	12	19	32	59	53	72	22	5	-
Male	5	3	11	6	5	16	23	17	22	4	1	-
Female	3	2	4	6	14	16	36	36	50	18	4	-
Black, both sexes	6	5	7	8	1	5	9	2	6	5	-	-
Male	3	3	3	3	-	2	4	1	2	1	-	-
Female	3	2	4	5	1	3	5	1	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
New Mexico												
All races, both sexes	2	5	-	2	4	9	5	11	6	7	2	-
Male	1	2	-	1	3	6	1	2	-	2	1	-
Female	1	3	-	1	1	3	4	9	6	5	1	-
White, both sexes	2	4	-	2	4	7	5	11	6	7	1	-
Male	1	2	-	1	3	6	1	2	-	2	1	-
Female	1	2	-	1	1	1	4	9	6	5	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	20	36	51	45	67	92	147	207	134	89	35	-
Male	15	20	24	27	35	38	60	50	30	12	2	-
Female	5	16	27	18	32	54	87	157	104	77	33	-
White, both sexes	13	22	31	34	49	71	113	172	104	76	28	-
Male	10	13	19	22	28	32	44	42	22	11	1	-
Female	3	9	12	12	21	39	69	130	82	65	27	-
Black, both sexes	6	13	20	11	15	16	28	30	28	13	7	-
Male	4	6	5	5	6	3	13	5	6	1	1	-
Female	2	7	15	6	9	13	15	25	22	12	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	3	5	6	5	2	-	-	-
Male	1	1	-	-	1	3	3	3	2	-	-	-
Female	-	-	-	-	2	2	3	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
North Carolina												
All races, both sexes	18	14	22	20	37	38	65	76	69	28	6	-
Male	11	10	13	11	24	14	23	17	12	2	-	-
Female	7	4	9	9	13	24	42	59	57	26	6	-
White, both sexes	4	10	10	9	25	25	47	58	52	21	4	-
Male	2	7	5	3	16	10	17	12	10	2	-	-
Female	2	3	5	6	9	15	30	46	42	19	4	-
Black, both sexes	13	3	12	11	11	11	18	17	17	6	2	-
Male	8	3	8	8	7	3	6	5	2	-	-	-
Female	5	-	4	3	4	8	12	12	15	6	2	-
American Indian 1/, both sexes	1	1	-	-	1	1	-	1	-	1	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	1	2	3	7	5	3	5	-
Male	-	-	-	-	-	2	2	1	-	-	-	-
Female	-	-	-	1	1	-	1	6	5	3	5	-
White, both sexes	-	-	-	1	1	2	3	7	5	3	5	-
Male	-	-	-	-	-	2	2	1	-	-	-	-
Female	-	-	-	1	1	-	1	6	5	3	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Ohio												
All races, both sexes	20	17	41	34	39	78	113	125	105	55	17	-
Male	15	10	28	18	22	29	42	24	24	7	1	-
Female	5	7	13	16	17	49	71	101	81	48	16	-
White, both sexes	9	16	38	29	30	65	94	115	102	49	17	-
Male	7	10	26	17	16	27	31	23	23	7	1	-
Female	2	6	12	12	14	38	63	92	79	42	16	-
Black, both sexes	11	1	3	5	9	13	19	9	3	6	-	-
Male	8	-	2	1	6	2	11	1	1	-	-	-
Female	3	1	1	4	3	11	8	8	2	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	7	2	13	9	10	7	14	25	26	17	3	-
Male	6	-	5	6	9	3	6	3	2	2	1	-
Female	1	2	8	3	1	4	8	22	24	15	2	-
White, both sexes	6	2	9	8	8	6	8	24	24	15	3	-
Male	5	-	4	5	7	2	2	3	2	2	1	-
Female	1	2	5	3	1	4	6	21	22	13	2	-
Black, both sexes	1	-	4	-	2	1	4	1	1	1	-	-
Male	1	-	1	-	2	1	2	-	-	-	-	-
Female	-	-	3	-	-	-	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	2	-	1	1	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Oregon												
All races, both sexes	10	9	14	15	19	25	37	47	42	22	3	-
Male	8	6	10	5	12	11	15	15	7	2	-	-
Female	2	3	4	10	7	14	22	32	35	20	3	-
White, both sexes	9	8	14	15	19	24	36	47	42	22	3	-
Male	7	6	10	5	12	11	15	15	7	2	-	-
Female	2	2	4	10	7	13	21	32	35	20	3	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	25	19	34	35	49	68	95	116	96	56	6	-
Male	14	16	24	20	19	33	45	37	19	7	1	-
Female	11	3	10	15	30	35	50	79	77	49	5	-
White, both sexes	21	14	26	32	42	60	83	107	90	54	5	-
Male	13	12	20	18	17	29	40	35	17	7	1	-
Female	8	2	6	14	25	31	43	72	73	47	4	-
Black, both sexes	3	4	8	2	7	8	12	9	6	2	1	-
Male	1	3	4	2	2	4	5	2	2	-	-	-
Female	2	1	4	-	5	4	7	7	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Rhode Island												
All races, both sexes	-	1	2	2	2	2	2	3	9	4	-	-
Male	-	-	-	2	2	1	1	1	-	-	-	-
Female	-	1	2	-	-	1	1	2	9	4	-	-
White, both sexes	-	1	2	2	2	1	2	3	9	4	-	-
Male	-	-	-	2	2	-	1	1	-	-	-	-
Female	-	1	2	-	-	1	1	2	9	4	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	7	23	8	23	15	13	33	28	22	8	3	-
Male	6	15	5	13	7	4	13	11	6	3	1	-
Female	1	8	3	10	8	9	20	17	16	5	2	-
White, both sexes	5	14	3	14	9	5	23	14	12	4	1	-
Male	4	9	2	6	3	2	7	6	2	2	-	-
Female	1	5	1	8	6	3	16	8	10	2	1	-
Black, both sexes	2	9	5	9	6	8	10	13	10	4	2	-
Male	2	6	3	7	4	2	6	4	4	1	1	-
Female	-	3	2	2	2	6	4	9	6	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
South Dakota												
All races, both sexes	1	-	-	1	-	2	3	10	11	7	3	-
Male	-	-	-	1	-	1	1	3	5	-	-	-
Female	1	-	-	-	-	1	2	7	6	7	3	-
White, both sexes	1	-	-	-	-	2	3	10	11	6	3	-
Male	-	-	-	-	-	1	1	3	5	-	-	-
Female	1	-	-	-	-	1	2	7	6	6	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	14	11	19	10	16	29	41	47	37	20	8	-
Male	8	2	14	5	10	10	10	10	8	4	1	-
Female	6	9	5	5	6	19	31	37	29	16	7	-
White, both sexes	7	6	16	8	10	21	35	42	31	18	5	-
Male	5	1	12	4	6	8	10	10	7	3	1	-
Female	2	5	4	4	4	13	25	32	24	15	4	-
Black, both sexes	6	4	3	2	6	8	6	5	6	2	3	-
Male	2	1	2	1	4	2	-	-	1	1	-	-
Female	4	3	1	1	2	6	6	5	5	1	3	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Texas												
All races, both sexes	28	29	21	38	54	73	116	134	103	70	18	-
Male	19	16	11	19	22	31	39	33	21	4	-	-
Female	9	13	10	19	32	42	77	101	82	66	18	-
White, both sexes	23	23	19	26	42	62	101	119	90	59	13	-
Male	17	13	11	12	16	28	34	30	18	4	-	-
Female	6	10	8	14	26	34	67	89	72	55	13	-
Black, both sexes	5	6	1	11	12	10	14	13	12	10	5	-
Male	2	3	-	6	6	3	5	2	2	-	-	-
Female	3	3	1	5	6	7	9	11	10	10	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	1	-	1	1	1	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	-	1	-	1	-	-
Utah												
All races, both sexes	4	2	2	2	3	12	13	21	14	9	1	-
Male	2	1	1	1	1	5	4	4	3	-	-	-
Female	2	1	1	1	2	7	9	17	11	9	1	-
White, both sexes	4	1	2	2	3	12	12	21	14	9	1	-
Male	2	1	1	1	1	5	4	4	3	-	-	-
Female	2	-	1	1	2	7	8	17	11	9	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Vermont												
All races, both sexes	-	1	-	-	1	1	4	6	3	2	1	-
Male	-	1	-	-	1	-	3	1	-	1	-	-
Female	-	-	-	-	-	1	1	5	3	1	1	-
White, both sexes	-	1	-	-	1	1	4	6	3	2	1	-
Male	-	1	-	-	1	-	3	1	-	1	-	-
Female	-	-	-	-	-	1	1	5	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	15	17	26	22	17	30	42	43	35	29	8	-
Male	6	10	19	11	9	15	11	12	8	1	1	-
Female	9	7	7	11	8	15	31	31	27	28	7	-
White, both sexes	10	9	15	11	12	23	32	34	25	23	5	-
Male	5	4	13	4	5	11	9	11	5	-	1	-
Female	5	5	2	7	7	12	23	23	20	23	4	-
Black, both sexes	5	8	10	10	5	7	10	8	9	5	3	-
Male	1	6	5	6	4	4	2	1	3	-	-	-
Female	4	2	5	4	1	3	8	7	6	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	1	-	1	-	-
Male	-	-	1	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Washington												
All races, both sexes	2	12	8	15	14	23	43	39	49	27	8	-
Male	2	7	4	5	8	14	11	7	11	7	-	-
Female	-	5	4	10	6	9	32	32	38	20	8	-
White, both sexes	2	11	5	14	14	18	39	38	45	27	8	-
Male	2	7	3	5	8	12	9	7	9	7	-	-
Female	-	4	2	9	6	6	30	31	36	20	8	-
Black, both sexes	-	1	2	1	-	1	1	1	2	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	1	1	1	-	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	3	3	-	1	-	-	-
Male	-	-	-	-	-	2	2	-	1	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
West Virginia												
All races, both sexes	9	9	16	15	12	24	23	12	22	10	2	-
Male	7	6	12	10	9	8	8	3	5	1	-	-
Female	2	3	4	5	3	16	15	9	17	9	2	-
White, both sexes	9	9	16	13	11	22	22	12	19	10	2	-
Male	7	6	12	9	8	7	8	3	4	1	-	-
Female	2	3	4	4	3	15	14	9	15	9	2	-
Black, both sexes	-	-	-	2	1	2	1	-	3	-	-	-
Male	-	-	-	1	1	1	-	-	1	-	-	-
Female	-	-	-	1	-	1	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Wisconsin												
All races, both sexes	4	9	5	5	9	31	30	50	42	20	4	-
Male	2	8	5	3	3	13	8	10	6	1	-	-
Female	2	1	-	2	6	18	22	40	36	19	4	-
White, both sexes	4	9	5	5	9	27	29	50	41	20	4	-
Male	2	8	5	3	3	11	8	10	6	1	-	-
Female	2	1	-	2	6	16	21	40	35	19	4	-
Black, both sexes	-	-	-	-	-	4	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	2	-	1	2	-	2	3	-	-
Male	1	-	-	2	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	1	2	-	1	2	-	-
White, both sexes	1	-	-	2	-	1	2	-	2	3	-	-
Male	1	-	-	2	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	1	2	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
United States												
All races, both sexes	29,788	1	-	1	1	7	45	104	184	456	902	1,542
Male	13,677	1	-	-	1	4	38	78	133	337	588	1,100
Female	16,111	-	-	1	-	3	7	26	51	119	314	442
White, both sexes	22,220	-	-	-	-	4	30	48	92	270	544	938
Male	9,755	-	-	-	-	3	25	36	72	213	387	709
Female	12,465	-	-	-	-	1	5	12	20	57	157	229
Black, both sexes	6,845	1	-	1	1	2	15	51	88	177	330	574
Male	3,559	1	-	-	1	1	13	38	58	118	176	367
Female	3,286	-	-	1	-	1	2	13	30	59	154	207
American Indian 1/, both sexes	133	-	-	-	-	-	-	1	3	3	12	13
Male	80	-	-	-	-	-	-	-	2	3	11	9
Female	53	-	-	-	-	-	-	1	1	-	1	4
Asian or Pacific Islander 2/, both sexes	590	-	-	-	-	1	-	4	1	6	16	17
Male	283	-	-	-	-	-	-	4	1	3	14	15
Female	307	-	-	-	-	1	-	-	-	3	2	2
Alabama												
All races, both sexes	390	-	-	-	-	-	1	1	6	6	11	11
Male	165	-	-	-	-	-	-	1	5	4	5	5
Female	225	-	-	-	-	-	1	-	1	2	6	6
White, both sexes	244	-	-	-	-	-	1	-	1	5	2	5
Male	105	-	-	-	-	-	-	-	1	4	1	2
Female	139	-	-	-	-	-	1	-	-	1	1	3
Black, both sexes	146	-	-	-	-	-	-	1	5	1	9	6
Male	60	-	-	-	-	-	-	1	4	-	4	3
Female	86	-	-	-	-	-	-	-	1	1	5	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Alaska												
All races, both sexes	43	-	-	-	-	-	-	-	1	3	1	3
Male	21	-	-	-	-	-	-	-	-	2	-	3
Female	22	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	26	-	-	-	-	-	-	-	-	2	-	2
Male	14	-	-	-	-	-	-	-	-	1	-	2
Female	12	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	518	-	-	-	-	1	-	2	1	8	13	24
Male	263	-	-	-	-	1	-	1	1	6	6	20
Female	255	-	-	-	-	-	-	1	-	2	7	4
White, both sexes	466	-	-	-	-	-	-	2	1	7	11	19
Male	231	-	-	-	-	-	-	1	1	5	5	17
Female	235	-	-	-	-	-	-	1	-	2	6	2
Black, both sexes	32	-	-	-	-	1	-	-	-	1	2	3
Male	21	-	-	-	-	1	-	-	-	1	1	1
Female	11	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Arkansas												
All races, both sexes	324	-	-	-	-	-	1	1	4	3	16	15
Male	141	-	-	-	-	-	1	-	3	2	8	12
Female	183	-	-	-	-	-	-	1	1	1	8	3
White, both sexes	250	-	-	-	-	-	-	-	2	2	11	6
Male	102	-	-	-	-	-	-	-	2	2	5	5
Female	148	-	-	-	-	-	-	-	-	-	6	1
Black, both sexes	72	-	-	-	-	-	1	1	2	1	5	8
Male	37	-	-	-	-	-	1	-	1	-	3	6
Female	35	-	-	-	-	-	-	1	1	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,014	-	-	-	-	-	4	10	16	64	104	170
Male	1,852	-	-	-	-	-	4	7	11	48	69	128
Female	2,162	-	-	-	-	-	-	3	5	16	35	42
White, both sexes	3,142	-	-	-	-	-	3	7	11	44	76	106
Male	1,423	-	-	-	-	-	3	4	9	34	53	83
Female	1,719	-	-	-	-	-	-	3	2	10	23	23
Black, both sexes	558	-	-	-	-	-	1	2	5	14	20	52
Male	276	-	-	-	-	-	1	2	2	11	10	35
Female	282	-	-	-	-	-	-	-	3	3	10	17
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	1	3	3
Male	22	-	-	-	-	-	-	-	-	1	2	3
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	286	-	-	-	-	-	-	1	-	5	5	9
Male	131	-	-	-	-	-	-	1	-	2	4	7
Female	155	-	-	-	-	-	-	-	-	3	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Colorado												
All races, both sexes	254	-	-	-	-	-	-	-	3	2	5	10
Male	106	-	-	-	-	-	-	-	1	1	4	9
Female	148	-	-	-	-	-	-	-	2	1	1	1
White, both sexes	231	-	-	-	-	-	-	-	2	2	5	10
Male	97	-	-	-	-	-	-	-	1	1	4	9
Female	134	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	213	-	-	-	-	-	-	-	-	3	4	14
Male	94	-	-	-	-	-	-	-	-	2	4	10
Female	119	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	192	-	-	-	-	-	-	-	-	3	4	13
Male	80	-	-	-	-	-	-	-	-	2	4	9
Female	112	-	-	-	-	-	-	-	-	1	-	4
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Delaware												
All races, both sexes	83	-	-	-	-	-	-	-	-	4	3	5
Male	43	-	-	-	-	-	-	-	-	3	2	3
Female	40	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	61	-	-	-	-	-	-	-	-	2	3	4
Male	28	-	-	-	-	-	-	-	-	1	2	3
Female	33	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	22	-	-	-	-	-	-	-	-	2	-	1
Male	15	-	-	-	-	-	-	-	-	2	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	181	-	-	-	-	-	2	-	3	2	6	14
Male	88	-	-	-	-	-	2	-	2	2	3	7
Female	93	-	-	-	-	-	-	-	1	-	3	7
White, both sexes	26	-	-	-	-	-	1	-	-	-	1	1
Male	11	-	-	-	-	-	1	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	153	-	-	-	-	-	1	-	3	2	5	13
Male	77	-	-	-	-	-	1	-	2	2	2	7
Female	76	-	-	-	-	-	-	-	1	-	3	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Florida												
All races, both sexes	2,211	-	-	-	-	-	4	10	13	35	71	100
Male	1,088	-	-	-	-	-	4	7	13	25	45	70
Female	1,123	-	-	-	-	-	-	3	-	10	26	30
White, both sexes	1,793	-	-	-	-	-	2	4	8	24	44	64
Male	875	-	-	-	-	-	2	3	8	19	31	49
Female	918	-	-	-	-	-	-	1	-	5	13	15
Black, both sexes	401	-	-	-	-	-	2	5	5	11	26	36
Male	205	-	-	-	-	-	2	3	5	6	13	21
Female	196	-	-	-	-	-	-	2	-	5	13	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	1	-	-	1	-
Male	7	-	-	-	-	-	-	1	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,119	-	-	-	1	1	3	7	7	15	49	75
Male	585	-	-	-	1	1	3	6	5	11	28	53
Female	534	-	-	-	-	-	-	1	2	4	21	22
White, both sexes	549	-	-	-	-	1	3	1	2	5	19	32
Male	274	-	-	-	-	1	3	1	1	5	12	27
Female	275	-	-	-	-	-	-	-	1	-	7	5
Black, both sexes	557	-	-	-	1	-	-	6	5	10	30	41
Male	302	-	-	-	1	-	-	5	4	6	16	24
Female	255	-	-	-	-	-	-	1	1	4	14	17
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Hawaii												
All races, both sexes	67	-	-	-	-	-	-	2	-	-	2	2
Male	34	-	-	-	-	-	-	1	-	-	1	2
Female	33	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	15	-	-	-	-	-	-	1	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	-	-	-	-	-	1	-	-	1	2
Male	22	-	-	-	-	-	-	1	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	100	-	-	-	-	-	1	-	1	1	2	7
Male	47	-	-	-	-	-	-	-	1	1	1	6
Female	53	-	-	-	-	-	1	-	-	-	1	1
White, both sexes	96	-	-	-	-	-	1	-	1	1	1	6
Male	44	-	-	-	-	-	-	-	1	1	-	5
Female	52	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Illinois												
All races, both sexes	1,369	-	-	-	-	-	3	10	10	29	48	72
Male	676	-	-	-	-	-	3	8	8	22	34	57
Female	693	-	-	-	-	-	-	2	2	7	14	15
White, both sexes	870	-	-	-	-	-	1	5	4	11	21	30
Male	375	-	-	-	-	-	1	3	4	10	17	25
Female	495	-	-	-	-	-	-	2	-	1	4	5
Black, both sexes	488	-	-	-	-	-	2	5	6	18	26	42
Male	295	-	-	-	-	-	2	5	4	12	16	32
Female	193	-	-	-	-	-	-	-	2	6	10	10
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	467	-	-	-	-	-	1	3	2	7	13	27
Male	199	-	-	-	-	-	1	1	2	6	11	20
Female	268	-	-	-	-	-	-	2	-	1	2	7
White, both sexes	382	-	-	-	-	-	1	2	-	6	7	21
Male	154	-	-	-	-	-	1	1	-	5	6	15
Female	228	-	-	-	-	-	-	1	-	1	1	6
Black, both sexes	83	-	-	-	-	-	-	1	2	1	5	6
Male	43	-	-	-	-	-	-	-	2	1	4	5
Female	40	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Iowa												
All races, both sexes	267	-	-	-	-	-	-	1	-	-	7	2
Male	98	-	-	-	-	-	-	1	-	-	3	1
Female	169	-	-	-	-	-	-	-	-	-	4	1
White, both sexes	266	-	-	-	-	-	-	1	-	-	7	2
Male	98	-	-	-	-	-	-	1	-	-	3	1
Female	168	-	-	-	-	-	-	-	-	-	4	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	158	-	-	-	-	-	-	-	-	-	1	6
Male	35	-	-	-	-	-	-	-	-	-	-	4
Female	123	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	144	-	-	-	-	-	-	-	-	-	1	4
Male	31	-	-	-	-	-	-	-	-	-	-	3
Female	113	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Kentucky												
All races, both sexes	368	-	-	-	-	-	-	3	5	7	15	34
Male	197	-	-	-	-	-	-	3	5	5	11	30
Female	171	-	-	-	-	-	-	-	-	2	4	4
White, both sexes	329	-	-	-	-	-	-	3	4	5	11	30
Male	173	-	-	-	-	-	-	3	4	3	8	26
Female	156	-	-	-	-	-	-	-	-	2	3	4
Black, both sexes	39	-	-	-	-	-	-	-	1	2	4	4
Male	24	-	-	-	-	-	-	-	1	2	3	4
Female	15	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	724	-	-	1	-	-	1	-	6	14	16	45
Male	353	-	-	-	-	-	1	-	1	12	10	27
Female	371	-	-	1	-	-	-	-	5	2	6	18
White, both sexes	436	-	-	-	-	-	1	-	1	6	5	22
Male	202	-	-	-	-	-	1	-	-	5	4	15
Female	234	-	-	-	-	-	-	-	1	1	1	7
Black, both sexes	284	-	-	1	-	-	-	-	5	8	11	23
Male	147	-	-	-	-	-	-	-	1	7	6	12
Female	137	-	-	1	-	-	-	-	4	1	5	11
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Maine												
All races, both sexes	65	-	-	-	-	-	-	-	-	3	1	5
Male	28	-	-	-	-	-	-	-	-	3	-	4
Female	37	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	63	-	-	-	-	-	-	-	-	2	1	4
Male	26	-	-	-	-	-	-	-	-	2	-	3
Female	37	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	835	-	-	-	-	-	1	5	5	17	47	77
Male	465	-	-	-	-	-	1	4	5	15	34	56
Female	370	-	-	-	-	-	-	1	-	2	13	21
White, both sexes	486	-	-	-	-	-	-	1	2	9	29	43
Male	269	-	-	-	-	-	-	1	2	9	24	33
Female	217	-	-	-	-	-	-	-	-	-	5	10
Black, both sexes	330	-	-	-	-	-	1	4	3	8	17	34
Male	183	-	-	-	-	-	1	3	3	6	9	23
Female	147	-	-	-	-	-	-	1	-	2	8	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Massachusetts												
All races, both sexes	524	-	-	-	-	1	1	2	1	7	25	39
Male	240	-	-	-	-	1	1	2	1	4	22	33
Female	284	-	-	-	-	-	-	-	-	3	3	6
White, both sexes	479	-	-	-	-	1	1	1	1	7	22	35
Male	212	-	-	-	-	1	1	1	1	4	19	29
Female	267	-	-	-	-	-	-	-	-	3	3	6
Black, both sexes	41	-	-	-	-	-	-	1	-	-	2	4
Male	25	-	-	-	-	-	-	1	-	-	2	4
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,278	-	-	-	-	-	1	2	9	17	26	63
Male	552	-	-	-	-	-	1	2	4	12	11	41
Female	726	-	-	-	-	-	-	-	5	5	15	22
White, both sexes	959	-	-	-	-	-	1	-	2	7	14	36
Male	393	-	-	-	-	-	1	-	1	7	5	24
Female	566	-	-	-	-	-	-	-	1	-	9	12
Black, both sexes	305	-	-	-	-	-	-	2	7	10	12	25
Male	155	-	-	-	-	-	-	2	3	5	6	17
Female	150	-	-	-	-	-	-	-	4	5	6	8
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Minnesota												
All races, both sexes	177	-	-	-	-	-	1	-	1	2	9	5
Male	83	-	-	-	-	-	1	-	1	2	6	4
Female	94	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	166	-	-	-	-	-	-	-	-	2	5	3
Male	78	-	-	-	-	-	-	-	-	2	4	3
Female	88	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	1	-	-	-	3	2
Male	3	-	-	-	-	-	1	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	688	-	-	-	-	-	4	4	4	8	10	24
Male	302	-	-	-	-	-	4	3	3	5	6	11
Female	386	-	-	-	-	-	-	1	1	3	4	13
White, both sexes	359	-	-	-	-	-	2	1	1	1	3	9
Male	141	-	-	-	-	-	2	1	1	1	2	5
Female	218	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	328	-	-	-	-	-	2	3	3	7	7	15
Male	160	-	-	-	-	-	2	2	2	4	4	6
Female	168	-	-	-	-	-	-	1	1	3	3	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Missouri												
All races, both sexes	474	-	-	-	-	-	2	2	5	8	26	40
Male	218	-	-	-	-	-	2	2	1	7	17	25
Female	256	-	-	-	-	-	-	-	4	1	9	15
White, both sexes	303	-	-	-	-	-	1	-	4	3	14	19
Male	123	-	-	-	-	-	1	-	1	3	10	13
Female	180	-	-	-	-	-	-	-	3	-	4	6
Black, both sexes	170	-	-	-	-	-	1	2	1	5	12	21
Male	95	-	-	-	-	-	1	2	-	4	7	12
Female	75	-	-	-	-	-	-	-	1	1	5	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	58	-	-	-	-	-	-	-	1	1	2	2
Male	27	-	-	-	-	-	-	-	1	1	1	1
Female	31	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	55	-	-	-	-	-	-	-	-	1	2	1
Male	26	-	-	-	-	-	-	-	-	1	1	1
Female	29	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Nebraska												
All races, both sexes	117	-	-	-	-	-	-	2	2	2	6	7
Male	45	-	-	-	-	-	-	1	1	2	4	6
Female	72	-	-	-	-	-	-	1	1	-	2	1
White, both sexes	111	-	-	-	-	-	-	2	2	1	6	7
Male	42	-	-	-	-	-	-	1	1	1	4	6
Female	69	-	-	-	-	-	-	1	1	-	2	1
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	439	-	-	-	-	1	-	-	2	2	8	23
Male	255	-	-	-	-	1	-	-	1	1	7	17
Female	184	-	-	-	-	-	-	-	1	1	1	6
White, both sexes	369	-	-	-	-	1	-	-	2	2	6	21
Male	219	-	-	-	-	1	-	-	1	1	5	17
Female	150	-	-	-	-	-	-	-	1	1	1	4
Black, both sexes	49	-	-	-	-	-	-	-	-	-	1	2
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
New Hampshire												
All races, both sexes	114	-	-	-	-	-	-	-	-	-	2	2
Male	41	-	-	-	-	-	-	-	-	-	2	2
Female	73	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	113	-	-	-	-	-	-	-	-	-	2	2
Male	41	-	-	-	-	-	-	-	-	-	2	2
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	751	-	-	-	-	-	1	1	2	8	14	35
Male	323	-	-	-	-	-	-	-	1	5	12	24
Female	428	-	-	-	-	-	1	1	1	3	2	11
White, both sexes	592	-	-	-	-	-	1	-	1	4	10	20
Male	241	-	-	-	-	-	-	-	-	3	8	12
Female	351	-	-	-	-	-	1	-	1	1	2	8
Black, both sexes	142	-	-	-	-	-	-	1	1	4	4	15
Male	76	-	-	-	-	-	-	-	1	2	4	12
Female	66	-	-	-	-	-	-	1	-	2	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
New Mexico												
All races, both sexes	159	-	-	-	-	-	-	2	-	6	8	14
Male	79	-	-	-	-	-	-	2	-	5	7	11
Female	80	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	146	-	-	-	-	-	-	2	-	5	6	11
Male	74	-	-	-	-	-	-	2	-	4	5	9
Female	72	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	1	2	2
Male	4	-	-	-	-	-	-	-	-	1	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,596	-	-	-	-	-	4	7	13	32	59	122
Male	1,156	-	-	-	-	-	2	5	10	22	42	90
Female	1,440	-	-	-	-	-	2	2	3	10	17	32
White, both sexes	1,739	-	-	-	-	-	2	4	6	11	28	59
Male	722	-	-	-	-	-	1	3	5	9	25	44
Female	1,017	-	-	-	-	-	1	1	1	2	3	15
Black, both sexes	791	-	-	-	-	-	2	3	6	20	29	62
Male	397	-	-	-	-	-	1	2	4	12	15	45
Female	394	-	-	-	-	-	1	1	2	8	14	17
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	66	-	-	-	-	-	-	-	1	1	2	1
Male	37	-	-	-	-	-	-	-	1	1	2	1
Female	29	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
North Carolina												
All races, both sexes	632	1	-	-	-	1	1	3	1	15	28	48
Male	314	1	-	-	-	-	1	2	1	11	16	28
Female	318	-	-	-	-	1	-	1	-	4	12	20
White, both sexes	390	-	-	-	-	-	1	1	1	5	12	23
Male	189	-	-	-	-	-	1	1	1	4	6	16
Female	201	-	-	-	-	-	-	-	-	1	6	7
Black, both sexes	237	1	-	-	-	1	-	2	-	10	15	24
Male	122	1	-	-	-	-	-	1	-	7	9	11
Female	115	-	-	-	-	1	-	1	-	3	6	13
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	52	-	-	-	-	-	-	1	-	1	1	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	47	-	-	-	-	-	-	-	-	1	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Ohio												
All races, both sexes	1,587	-	-	-	-	-	1	7	9	26	44	87
Male	705	-	-	-	-	-	-	7	7	16	30	56
Female	882	-	-	-	-	-	1	-	2	10	14	31
White, both sexes	1,266	-	-	-	-	-	1	3	5	16	31	54
Male	530	-	-	-	-	-	-	3	4	11	24	40
Female	736	-	-	-	-	-	1	-	1	5	7	14
Black, both sexes	316	-	-	-	-	-	-	4	4	10	12	32
Male	172	-	-	-	-	-	-	4	3	5	5	15
Female	144	-	-	-	-	-	-	-	1	5	7	17
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	276	-	-	-	-	-	-	-	4	4	4	12
Male	131	-	-	-	-	-	-	-	4	3	2	10
Female	145	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	223	-	-	-	-	-	-	-	3	2	2	12
Male	103	-	-	-	-	-	-	-	3	2	1	10
Female	120	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	40	-	-	-	-	-	-	-	1	2	1	-
Male	21	-	-	-	-	-	-	-	1	1	-	-
Female	19	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Oregon												
All races, both sexes	251	-	-	-	-	-	-	-	2	2	3	3
Male	89	-	-	-	-	-	-	-	2	1	3	3
Female	162	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	246	-	-	-	-	-	-	-	1	2	3	3
Male	87	-	-	-	-	-	-	-	1	1	3	3
Female	159	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	787	-	-	-	-	-	1	2	4	6	20	37
Male	348	-	-	-	-	-	1	1	3	5	12	30
Female	439	-	-	-	-	-	-	1	1	1	8	7
White, both sexes	649	-	-	-	-	-	1	1	2	5	12	23
Male	282	-	-	-	-	-	1	-	2	4	7	19
Female	367	-	-	-	-	-	-	1	-	1	5	4
Black, both sexes	133	-	-	-	-	-	-	1	2	1	7	13
Male	61	-	-	-	-	-	-	1	1	1	4	10
Female	72	-	-	-	-	-	-	-	1	-	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Rhode Island												
All races, both sexes	45	-	-	-	-	-	-	-	-	1	4	3
Male	27	-	-	-	-	-	-	-	-	1	2	3
Female	18	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	41	-	-	-	-	-	-	-	-	1	3	3
Male	24	-	-	-	-	-	-	-	-	1	1	3
Female	17	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	384	-	-	-	-	1	1	1	-	5	9	16
Male	157	-	-	-	-	-	1	1	-	3	6	11
Female	227	-	-	-	-	1	-	-	-	2	3	5
White, both sexes	223	-	-	-	-	-	1	-	-	2	5	6
Male	89	-	-	-	-	-	1	-	-	1	4	5
Female	134	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	158	-	-	-	-	-	-	1	-	3	4	10
Male	67	-	-	-	-	-	-	1	-	2	2	6
Female	91	-	-	-	-	-	-	-	-	1	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
South Dakota												
All races, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	824	-	-	-	-	-	3	5	8	14	29	39
Male	347	-	-	-	-	-	3	5	2	12	16	21
Female	477	-	-	-	-	-	-	-	6	2	13	18
White, both sexes	571	-	-	-	-	-	3	4	3	10	13	24
Male	226	-	-	-	-	-	3	4	-	9	8	13
Female	345	-	-	-	-	-	-	-	3	1	5	11
Black, both sexes	247	-	-	-	-	-	-	1	5	4	15	15
Male	119	-	-	-	-	-	-	1	2	3	7	8
Female	128	-	-	-	-	-	-	-	3	1	8	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Texas												
All races, both sexes	1,888	-	-	-	-	-	1	7	23	37	86	129
Male	855	-	-	-	-	-	-	4	18	25	57	87
Female	1,033	-	-	-	-	-	1	3	5	12	29	42
White, both sexes	1,445	-	-	-	-	-	-	2	11	27	56	87
Male	627	-	-	-	-	-	-	2	9	18	41	58
Female	818	-	-	-	-	-	-	-	2	9	15	29
Black, both sexes	419	-	-	-	-	-	1	5	12	10	29	42
Male	213	-	-	-	-	-	-	2	9	7	15	29
Female	206	-	-	-	-	-	1	3	3	3	14	13
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	116	-	-	-	-	-	-	-	-	1	2	2
Male	46	-	-	-	-	-	-	-	-	1	2	2
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	112	-	-	-	-	-	-	-	-	1	2	2
Male	43	-	-	-	-	-	-	-	-	1	2	2
Female	69	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Vermont												
All races, both sexes	61	-	-	-	-	1	-	-	1	1	1	2
Male	27	-	-	-	-	-	-	-	1	1	1	1
Female	34	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	60	-	-	-	-	1	-	-	1	1	1	2
Male	27	-	-	-	-	-	-	-	1	1	1	1
Female	33	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	403	-	-	-	-	-	-	-	5	13	13	27
Male	182	-	-	-	-	-	-	-	4	9	9	21
Female	221	-	-	-	-	-	-	-	1	4	4	6
White, both sexes	271	-	-	-	-	-	-	-	3	5	6	16
Male	110	-	-	-	-	-	-	-	2	4	5	13
Female	161	-	-	-	-	-	-	-	1	1	1	3
Black, both sexes	129	-	-	-	-	-	-	-	2	8	7	11
Male	71	-	-	-	-	-	-	-	2	5	4	8
Female	58	-	-	-	-	-	-	-	-	3	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Washington												
All races, both sexes	724	-	-	-	-	-	-	1	2	11	16	21
Male	324	-	-	-	-	-	-	1	2	10	10	21
Female	400	-	-	-	-	-	-	-	-	1	6	-
White, both sexes	656	-	-	-	-	-	-	-	2	9	11	20
Male	285	-	-	-	-	-	-	-	2	8	8	20
Female	371	-	-	-	-	-	-	-	-	1	3	-
Black, both sexes	37	-	-	-	-	-	-	-	-	2	4	1
Male	24	-	-	-	-	-	-	-	-	2	1	1
Female	13	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	1	-	-	1	-
Male	11	-	-	-	-	-	-	1	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	110	-	-	-	-	-	-	-	-	2	2	3
Male	26	-	-	-	-	-	-	-	-	2	1	2
Female	84	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	105	-	-	-	-	-	-	-	-	2	2	2
Male	23	-	-	-	-	-	-	-	-	2	1	1
Female	82	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Wisconsin												
All races, both sexes	415	-	-	-	-	-	1	-	2	1	8	17
Male	150	-	-	-	-	-	1	-	2	1	3	11
Female	265	-	-	-	-	-	-	-	-	-	5	6
White, both sexes	369	-	-	-	-	-	1	-	2	1	4	12
Male	127	-	-	-	-	-	1	-	2	1	2	7
Female	242	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	44	-	-	-	-	-	-	-	-	-	4	5
Male	22	-	-	-	-	-	-	-	-	-	1	4
Female	22	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	2	2
Male	17	-	-	-	-	-	-	-	-	-	2	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	36	-	-	-	-	-	-	-	-	-	2	2
Male	17	-	-	-	-	-	-	-	-	-	2	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
United States												
All races, both sexes	2,044	2,143	1,987	1,891	1,969	2,595	3,637	4,322	3,698	1,818	439	2
Male	1,386	1,437	1,300	1,160	1,066	1,255	1,376	1,261	807	301	46	2
Female	658	706	687	731	903	1,340	2,261	3,061	2,891	1,517	393	-
White, both sexes	1,245	1,380	1,317	1,258	1,319	1,939	2,938	3,689	3,234	1,611	362	2
Male	909	966	872	763	725	928	1,100	1,068	676	261	40	2
Female	336	414	445	495	594	1,011	1,838	2,621	2,558	1,350	322	-
Black, both sexes	741	720	626	572	599	578	614	527	383	175	70	-
Male	441	441	401	359	308	290	241	160	108	31	6	-
Female	300	279	225	213	291	288	373	367	275	144	64	-
American Indian 1/, both sexes	20	14	6	16	7	15	8	5	9	-	1	-
Male	11	10	3	12	4	8	2	3	2	-	-	-
Female	9	4	3	4	3	7	6	2	7	-	1	-
Asian or Pacific Islander 2/, both sexes	38	29	38	45	44	63	77	101	72	32	6	-
Male	25	20	24	26	29	29	33	30	21	9	-	-
Female	13	9	14	19	15	34	44	71	51	23	6	-
Alabama												
All races, both sexes	21	14	31	21	33	39	53	60	56	22	4	-
Male	15	6	21	14	15	18	18	16	16	6	-	-
Female	6	8	10	7	18	21	35	44	40	16	4	-
White, both sexes	10	3	21	10	22	25	32	45	42	18	2	-
Male	9	-	13	8	13	13	11	12	13	5	-	-
Female	1	3	8	2	9	12	21	33	29	13	2	-
Black, both sexes	11	11	10	11	11	14	21	15	14	4	2	-
Male	6	6	8	6	2	5	7	4	3	1	-	-
Female	5	5	2	5	9	9	14	11	11	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Alaska												
All races, both sexes	4	7	2	6	3	3	4	2	2	1	1	-
Male	3	2	1	3	2	3	2	-	-	-	-	-
Female	1	5	1	3	1	-	2	2	2	1	1	-
White, both sexes	4	2	1	4	3	2	2	2	1	1	-	-
Male	3	-	1	2	2	2	1	-	-	-	-	-
Female	1	2	-	2	1	-	1	2	1	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	1	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	2	1	1	-	1	2	-	1	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	1	1	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	44	39	36	37	37	49	54	78	60	30	5	-
Male	33	27	22	28	25	23	19	31	14	6	-	-
Female	11	12	14	9	12	26	35	47	46	24	5	-
White, both sexes	33	34	28	35	29	46	52	77	59	28	5	-
Male	25	24	17	27	19	22	18	30	14	6	-	-
Female	8	10	11	8	10	24	34	47	45	22	5	-
Black, both sexes	7	3	5	-	5	1	-	1	1	2	-	-
Male	6	3	3	-	4	-	-	1	-	-	-	-
Female	1	-	2	-	1	1	-	-	1	2	-	-
American Indian 1/, both sexes	3	1	2	1	3	1	1	-	-	-	-	-
Male	2	-	2	1	2	-	-	-	-	-	-	-
Female	1	1	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	1	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Arkansas												
All races, both sexes	24	14	19	13	18	25	36	60	47	21	7	-
Male	20	12	14	10	9	12	11	13	7	7	-	-
Female	4	2	5	3	9	13	25	47	40	14	7	-
White, both sexes	16	12	14	9	14	20	33	52	39	15	5	-
Male	14	10	10	7	8	8	9	11	5	6	-	-
Female	2	2	4	2	6	12	24	41	34	9	5	-
Black, both sexes	8	1	5	4	4	5	3	8	8	6	2	-
Male	6	1	4	3	1	4	2	2	2	1	-	-
Female	2	-	1	1	3	1	1	6	6	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	241	269	263	219	240	353	501	688	513	277	81	1
Male	162	180	185	128	145	192	184	226	119	56	7	1
Female	79	89	78	91	95	161	317	462	394	221	74	-
White, both sexes	164	200	192	157	186	267	412	557	442	247	70	1
Male	112	138	140	90	114	146	153	187	99	50	7	1
Female	52	62	52	67	72	121	259	370	343	197	63	-
Black, both sexes	61	50	55	38	36	53	50	71	31	12	7	-
Male	39	26	33	23	19	29	14	22	9	1	-	-
Female	22	24	22	15	17	24	36	49	22	11	7	-
American Indian 1/, both sexes	3	6	1	3	1	6	-	1	-	-	-	-
Male	1	5	1	3	1	4	-	1	-	-	-	-
Female	2	1	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	13	15	21	17	27	39	59	40	18	4	-
Male	10	11	11	12	11	13	17	16	11	5	-	-
Female	3	2	4	9	6	14	22	43	29	13	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Colorado												
All races, both sexes	17	25	16	17	14	28	13	41	35	24	4	-
Male	11	19	10	10	5	10	10	6	7	3	-	-
Female	6	6	6	7	9	18	3	35	28	21	4	-
White, both sexes	14	20	14	15	10	27	12	39	34	23	4	-
Male	9	16	9	9	4	10	9	6	7	3	-	-
Female	5	4	5	6	6	17	3	33	27	20	4	-
Black, both sexes	3	5	2	2	3	1	-	2	1	1	-	-
Male	2	3	1	1	1	-	-	-	-	-	-	-
Female	1	2	1	1	2	1	-	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	8	22	8	11	10	13	25	30	39	20	6	-
Male	7	15	4	8	8	7	13	4	9	3	-	-
Female	1	7	4	3	2	6	12	26	30	17	6	-
White, both sexes	7	19	4	8	7	12	24	30	37	19	5	-
Male	6	13	1	5	6	6	12	4	9	3	-	-
Female	1	6	3	3	1	6	12	26	28	16	5	-
Black, both sexes	1	3	4	3	3	1	1	-	2	1	1	-
Male	1	2	3	3	2	1	1	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Delaware												
All races, both sexes	7	6	2	5	9	10	12	11	7	2	-	-
Male	3	5	1	4	4	6	6	4	2	-	-	-
Female	4	1	1	1	5	4	6	7	5	2	-	-
White, both sexes	6	2	2	3	6	8	9	9	6	1	-	-
Male	3	2	1	2	2	4	4	2	2	-	-	-
Female	3	-	1	1	4	4	5	7	4	1	-	-
Black, both sexes	1	4	-	2	3	2	3	2	1	1	-	-
Male	-	3	-	2	2	2	2	2	-	-	-	-
Female	1	1	-	-	1	-	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	20	13	13	13	11	16	25	19	13	7	4	-
Male	15	8	10	10	5	7	9	5	3	-	-	-
Female	5	5	3	3	6	9	16	14	10	7	4	-
White, both sexes	-	3	-	-	-	3	5	2	5	4	1	-
Male	-	2	-	-	-	1	4	2	-	-	-	-
Female	-	1	-	-	-	2	1	-	5	4	1	-
Black, both sexes	20	10	13	13	11	13	20	16	8	2	3	-
Male	15	6	10	10	5	6	5	3	3	-	-	-
Female	5	4	3	3	6	7	15	13	5	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Florida												
All races, both sexes	148	158	153	148	149	190	295	317	277	112	31	-
Male	99	110	109	95	76	101	121	108	73	26	6	-
Female	49	48	44	53	73	89	174	209	204	86	25	-
White, both sexes	105	115	114	111	113	167	262	284	246	103	27	-
Male	75	86	82	73	54	89	112	101	62	24	5	-
Female	30	29	32	38	59	78	150	183	184	79	22	-
Black, both sexes	42	43	37	34	35	20	33	30	29	9	4	-
Male	23	24	26	21	22	11	9	6	10	2	1	-
Female	19	19	11	13	13	9	24	24	19	7	3	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	2	1	3	-	3	1	-	-	-
Male	1	-	1	-	-	1	-	1	1	-	-	-
Female	-	-	1	2	1	2	-	2	-	-	-	-
Georgia												
All races, both sexes	103	125	108	93	98	95	106	105	76	44	8	-
Male	63	82	74	64	50	44	44	33	16	7	-	-
Female	40	43	34	29	48	51	62	72	60	37	8	-
White, both sexes	36	51	44	38	44	53	66	68	52	29	5	-
Male	28	34	29	27	25	22	28	20	9	2	-	-
Female	8	17	15	11	19	31	38	48	43	27	5	-
Black, both sexes	67	74	64	53	49	40	40	35	24	15	3	-
Male	35	48	45	36	21	21	16	12	7	5	-	-
Female	32	26	19	17	28	19	24	23	17	10	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	5	2	-	2	-	-	-	-
Male	-	-	-	1	4	1	-	1	-	-	-	-
Female	-	-	-	1	1	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Hawaii												
All races, both sexes	4	4	2	4	4	4	6	16	11	5	1	-
Male	3	2	1	3	3	2	2	7	4	3	-	-
Female	1	2	1	1	1	2	4	9	7	2	1	-
White, both sexes	1	2	-	1	2	-	1	4	2	-	-	-
Male	1	-	-	1	1	-	1	3	2	-	-	-
Female	-	2	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	2	3	2	4	4	11	9	5	1	-
Male	2	1	1	2	2	2	-	4	2	3	-	-
Female	1	-	1	1	-	2	4	7	7	2	1	-
Idaho												
All races, both sexes	6	7	7	2	4	4	11	11	22	10	4	-
Male	6	4	5	1	1	1	7	3	7	2	1	-
Female	-	3	2	1	3	3	4	8	15	8	3	-
White, both sexes	5	7	7	2	4	4	11	11	22	10	3	-
Male	5	4	5	1	1	1	7	3	7	2	1	-
Female	-	3	2	1	3	3	4	8	15	8	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Illinois												
All races, both sexes	106	121	89	103	101	117	166	168	137	75	14	-
Male	73	78	57	59	52	63	64	57	31	8	2	-
Female	33	43	32	44	49	54	102	111	106	67	12	-
White, both sexes	52	54	44	55	56	82	123	136	116	67	13	-
Male	37	35	23	29	30	41	41	45	27	5	2	-
Female	15	19	21	26	26	41	82	91	89	62	11	-
Black, both sexes	53	67	44	47	44	34	40	30	21	8	1	-
Male	35	43	33	29	21	22	22	12	4	3	-	-
Female	18	24	11	18	23	12	18	18	17	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	-	3	2	-	-	-	-
Male	1	-	1	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	-	-	-	-
Indiana												
All races, both sexes	35	21	24	27	31	41	65	68	62	30	10	-
Male	25	14	14	14	19	23	22	18	7	1	1	-
Female	10	7	10	13	12	18	43	50	55	29	9	-
White, both sexes	24	18	19	20	24	29	57	62	57	27	8	-
Male	17	13	12	10	13	17	20	16	7	1	-	-
Female	7	5	7	10	11	12	37	46	50	26	8	-
Black, both sexes	11	3	5	7	6	12	8	6	5	3	2	-
Male	8	1	2	4	5	6	2	2	-	-	1	-
Female	3	2	3	3	1	6	6	4	5	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Iowa												
All races, both sexes	10	11	11	9	11	19	31	65	52	31	7	-
Male	8	6	8	6	9	10	10	20	12	4	-	-
Female	2	5	3	3	2	9	21	45	40	27	7	-
White, both sexes	10	10	11	9	11	19	31	65	52	31	7	-
Male	8	6	8	6	9	10	10	20	12	4	-	-
Female	2	4	3	3	2	9	21	45	40	27	7	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	6	2	3	4	11	21	41	33	22	3	-
Male	2	3	2	2	2	3	3	7	3	4	-	-
Female	3	3	-	1	2	8	18	34	30	18	3	-
White, both sexes	3	4	2	3	4	9	21	37	31	22	3	-
Male	2	2	2	2	2	2	3	7	2	4	-	-
Female	1	2	-	1	2	7	18	30	29	18	3	-
Black, both sexes	2	2	-	-	-	1	-	4	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	2	1	-	-	-	1	-	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Kentucky												
All races, both sexes	35	28	28	23	21	29	43	44	40	10	3	-
Male	24	22	18	15	16	12	12	11	8	2	3	-
Female	11	6	10	8	5	17	31	33	32	8	-	-
White, both sexes	31	26	25	21	18	26	39	41	38	8	3	-
Male	22	20	17	14	13	11	10	10	8	1	3	-
Female	9	6	8	7	5	15	29	31	30	7	-	-
Black, both sexes	4	2	3	2	3	3	4	3	2	2	-	-
Male	2	2	1	1	3	1	2	1	-	1	-	-
Female	2	-	2	1	-	2	2	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	60	70	49	50	56	72	90	84	72	33	5	-
Male	39	47	38	29	29	33	37	29	13	6	2	-
Female	21	23	11	21	27	39	53	55	59	27	3	-
White, both sexes	29	37	24	27	30	45	63	67	52	25	2	-
Male	18	24	20	13	18	19	27	25	8	4	1	-
Female	11	13	4	14	12	26	36	42	44	21	1	-
Black, both sexes	29	33	25	22	26	26	27	17	20	8	3	-
Male	19	23	18	15	11	13	10	4	5	2	1	-
Female	10	10	7	7	15	13	17	13	15	6	2	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Maine												
All races, both sexes	4	6	5	2	4	5	4	4	11	9	2	-
Male	3	5	2	2	2	2	2	-	1	2	-	-
Female	1	1	3	-	2	3	2	4	10	7	2	-
White, both sexes	4	6	5	2	4	5	4	4	11	9	2	-
Male	3	5	2	2	2	2	2	-	1	2	-	-
Female	1	1	3	-	2	3	2	4	10	7	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	73	73	85	70	58	57	80	68	78	28	13	-
Male	47	45	65	43	33	31	38	28	17	2	1	-
Female	26	28	20	27	25	26	42	40	61	26	12	-
White, both sexes	39	38	38	38	30	30	53	52	55	21	8	-
Male	29	26	30	28	16	15	22	22	10	2	-	-
Female	10	12	8	10	14	15	31	30	45	19	8	-
Black, both sexes	32	34	45	29	26	24	26	14	22	6	5	-
Male	17	18	33	14	15	13	15	5	7	-	1	-
Female	15	16	12	15	11	11	11	9	15	6	4	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	2	2	3	1	2	1	1	-	-
Male	1	-	2	1	2	3	1	1	-	-	-	-
Female	1	-	-	1	-	-	-	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Massachusetts												
All races, both sexes	42	39	45	29	25	34	51	75	63	36	9	-
Male	30	26	27	18	11	15	22	10	16	1	-	-
Female	12	13	18	11	14	19	29	65	47	35	9	-
White, both sexes	36	36	39	26	23	27	47	74	62	32	9	-
Male	26	26	23	15	10	10	20	10	16	-	-	-
Female	10	10	16	11	13	17	27	64	46	32	9	-
Black, both sexes	5	3	6	3	1	7	4	-	1	4	-	-
Male	3	-	4	3	-	5	2	-	-	1	-	-
Female	2	3	2	-	1	2	2	-	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	84	91	89	69	88	117	161	200	180	71	10	-
Male	60	55	56	41	48	54	62	50	42	11	2	-
Female	24	36	33	28	40	63	99	150	138	60	8	-
White, both sexes	45	58	62	41	58	90	127	181	162	67	8	-
Male	37	36	43	27	34	41	46	44	35	10	2	-
Female	8	22	19	14	24	49	81	137	127	57	6	-
Black, both sexes	37	33	26	28	28	27	32	18	14	4	2	-
Male	22	19	13	14	14	13	15	5	6	1	-	-
Female	15	14	13	14	14	14	17	13	8	3	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	-	2	-	3	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	2	-	1	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Minnesota												
All races, both sexes	9	8	6	5	4	15	21	37	29	15	10	-
Male	7	6	4	5	2	7	9	17	8	2	2	-
Female	2	2	2	-	2	8	12	20	21	13	8	-
White, both sexes	9	8	5	5	4	15	20	36	29	15	10	-
Male	7	6	4	5	2	7	9	17	8	2	2	-
Female	2	2	1	-	2	8	11	19	21	13	8	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	34	45	40	56	61	80	95	92	76	49	6	-
Male	23	29	27	33	32	41	26	25	26	8	-	-
Female	11	16	13	23	29	39	69	67	50	41	6	-
White, both sexes	14	14	14	24	28	47	61	60	51	28	1	-
Male	11	8	11	12	16	20	19	13	14	5	-	-
Female	3	6	3	12	12	27	42	47	37	23	1	-
Black, both sexes	20	31	26	32	33	32	34	32	25	21	5	-
Male	12	21	16	21	16	20	7	12	12	3	-	-
Female	8	10	10	11	17	12	27	20	13	18	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Missouri												
All races, both sexes	43	38	26	31	28	44	46	50	52	27	6	-
Male	29	22	17	17	15	21	15	14	10	2	2	-
Female	14	16	9	14	13	23	31	36	42	25	4	-
White, both sexes	16	19	13	19	15	26	37	42	47	24	4	-
Male	13	11	7	10	7	10	13	13	8	2	1	-
Female	3	8	6	9	8	16	24	29	39	22	3	-
Black, both sexes	27	19	12	12	13	18	9	8	5	3	2	-
Male	16	11	10	7	8	11	2	1	2	-	1	-
Female	11	8	2	5	5	7	7	7	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	2	2	4	3	2	6	9	11	8	1	-
Male	3	1	1	3	3	2	2	-	4	4	-	-
Female	1	1	1	1	-	-	4	9	7	4	1	-
White, both sexes	3	2	2	4	3	2	6	9	11	8	1	-
Male	3	1	1	3	3	2	2	-	4	4	-	-
Female	-	1	1	1	-	-	4	9	7	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Nebraska												
All races, both sexes	10	1	6	7	5	7	11	20	14	12	5	-
Male	7	1	4	5	2	2	4	4	1	1	-	-
Female	3	-	2	2	3	5	7	16	13	11	5	-
White, both sexes	7	1	6	6	5	7	11	19	14	12	5	-
Male	6	1	4	4	2	2	4	4	1	1	-	-
Female	1	-	2	2	3	5	7	15	13	11	5	-
Black, both sexes	3	-	-	1	-	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	31	44	48	50	40	57	43	53	29	8	-	-
Male	21	37	39	35	26	25	18	20	6	1	-	-
Female	10	7	9	15	14	32	25	33	23	7	-	-
White, both sexes	18	38	43	43	30	45	38	48	27	7	-	-
Male	14	32	35	31	20	21	16	18	6	1	-	-
Female	4	6	8	12	10	24	22	30	21	6	-	-
Black, both sexes	11	5	3	5	7	9	2	2	1	1	-	-
Male	6	4	2	4	4	4	1	1	-	-	-	-
Female	5	1	1	1	3	5	1	1	1	1	-	-
American Indian 1/, both sexes	2	-	-	1	1	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	2	2	3	3	1	-	-	-
Male	-	1	2	-	2	-	1	1	-	-	-	-
Female	-	-	-	1	-	2	2	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
New Hampshire												
All races, both sexes	5	14	7	6	6	8	17	15	16	12	4	-
Male	4	10	4	3	5	5	2	2	2	-	-	-
Female	1	4	3	3	1	3	15	13	14	12	4	-
White, both sexes	5	14	7	6	6	8	17	14	16	12	4	-
Male	4	10	4	3	5	5	2	2	2	-	-	-
Female	1	4	3	3	1	3	15	12	14	12	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
New Jersey												
All races, both sexes	61	53	39	42	55	62	79	128	113	43	15	-
Male	41	40	28	25	29	30	26	35	16	8	3	-
Female	20	13	11	17	26	32	53	93	97	35	12	-
White, both sexes	35	38	30	28	38	53	64	116	101	40	13	-
Male	25	32	22	15	20	25	22	32	14	8	3	-
Female	10	6	8	13	18	28	42	84	87	32	10	-
Black, both sexes	25	13	7	13	16	8	11	10	10	2	2	-
Male	15	6	6	9	8	5	4	3	1	-	-	-
Female	10	7	1	4	8	3	7	7	9	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	1	1	1	2	2	2	1	-	-
Male	1	2	-	1	1	-	-	-	1	-	-	-
Female	-	-	2	-	-	1	2	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
New Mexico												
All races, both sexes	17	7	7	11	8	18	13	17	16	14	1	-
Male	10	5	5	6	4	7	8	5	3	1	-	-
Female	7	2	2	5	4	11	5	12	13	13	1	-
White, both sexes	13	7	7	11	8	16	13	17	15	14	1	-
Male	10	5	5	6	4	7	8	5	3	1	-	-
Female	3	2	2	5	4	9	5	12	12	13	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	160	167	205	186	187	266	321	369	305	151	42	-
Male	106	105	112	108	99	120	134	119	54	22	6	-
Female	54	62	93	78	88	146	187	250	251	129	36	-
White, both sexes	84	85	113	106	102	176	232	306	260	130	35	-
Male	60	60	58	62	54	81	96	98	42	18	6	-
Female	24	25	55	44	48	95	136	208	218	112	29	-
Black, both sexes	69	81	88	76	80	81	75	55	39	18	7	-
Male	41	44	51	44	42	34	31	17	10	4	-	-
Female	28	37	37	32	38	47	44	38	29	14	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	1	4	4	5	9	14	8	6	3	-	-
Male	5	1	3	2	3	5	7	4	2	-	-	-
Female	2	-	1	2	2	4	7	4	4	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
North Carolina												
All races, both sexes	47	54	47	32	39	54	81	74	63	30	12	1
Male	34	44	32	22	24	22	34	20	14	7	-	1
Female	13	10	15	10	15	32	47	54	49	23	12	-
White, both sexes	24	32	29	18	24	34	57	56	45	21	6	1
Male	20	27	24	10	15	14	22	13	8	6	-	1
Female	4	5	5	8	9	20	35	43	37	15	6	-
Black, both sexes	23	22	17	14	15	18	24	18	18	9	6	-
Male	14	17	8	12	9	7	12	7	6	1	-	-
Female	9	5	9	2	6	11	12	11	12	8	6	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	2	2	-	2	1	13	9	10	7	2	-
Male	1	-	1	-	2	1	8	5	2	3	-	-
Female	-	2	1	-	-	-	5	4	8	4	2	-
White, both sexes	-	1	1	-	2	1	12	9	10	7	2	-
Male	-	-	1	-	2	1	8	5	2	3	-	-
Female	-	1	-	-	-	-	4	4	8	4	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Ohio												
All races, both sexes	98	124	89	121	101	112	196	235	221	94	22	-
Male	71	91	54	69	63	49	71	62	43	14	2	-
Female	27	33	35	52	38	63	125	173	178	80	20	-
White, both sexes	66	85	59	89	74	86	169	214	207	87	20	-
Male	47	64	36	47	47	36	60	56	40	13	2	-
Female	19	21	23	42	27	50	109	158	167	74	18	-
Black, both sexes	32	38	30	32	27	25	27	20	14	7	2	-
Male	24	27	18	22	16	13	11	5	3	1	-	-
Female	8	11	12	10	11	12	16	15	11	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	28	13	20	26	19	15	22	48	33	22	6	-
Male	21	12	11	17	13	7	10	7	9	4	1	-
Female	7	1	9	9	6	8	12	41	24	18	5	-
White, both sexes	21	10	16	21	12	14	19	42	26	19	4	-
Male	17	9	8	14	8	6	8	7	6	3	1	-
Female	4	1	8	7	4	8	11	35	20	16	3	-
Black, both sexes	5	2	4	2	6	1	3	4	5	3	1	-
Male	3	2	3	-	5	1	2	-	2	1	-	-
Female	2	-	1	2	1	-	1	4	3	2	1	-
American Indian 1/, both sexes	1	1	-	3	1	-	-	2	2	-	1	-
Male	1	1	-	3	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	2	1	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Oregon												
All races, both sexes	12	9	7	7	9	23	43	43	47	35	6	-
Male	8	9	5	4	4	11	13	12	10	4	-	-
Female	4	-	2	3	5	12	30	31	37	31	6	-
White, both sexes	10	9	7	7	9	23	41	43	47	35	6	-
Male	8	9	5	4	4	11	12	12	10	4	-	-
Female	2	-	2	3	5	12	29	31	37	31	6	-
Black, both sexes	1	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	46	47	53	43	57	75	121	107	101	56	11	-
Male	35	31	30	29	25	42	41	29	25	9	-	-
Female	11	16	23	14	32	33	80	78	76	47	11	-
White, both sexes	32	32	38	30	41	64	109	98	96	54	11	-
Male	26	24	22	21	19	37	40	27	24	9	-	-
Female	6	8	16	9	22	27	69	71	72	45	11	-
Black, both sexes	13	15	15	12	16	10	12	9	5	2	-	-
Male	8	7	8	7	6	4	1	2	1	-	-	-
Female	5	8	7	5	10	6	11	7	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Rhode Island												
All races, both sexes	4	2	1	2	2	3	5	6	8	4	-	-
Male	3	2	1	1	2	2	3	3	3	1	-	-
Female	1	-	-	1	-	1	2	3	5	3	-	-
White, both sexes	4	2	1	-	1	3	5	6	8	4	-	-
Male	3	2	1	-	1	2	3	3	3	1	-	-
Female	1	-	-	-	-	1	2	3	5	3	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	25	30	26	19	30	48	53	58	37	19	6	-
Male	15	17	16	12	14	20	14	17	8	2	-	-
Female	10	13	10	7	16	28	39	41	29	17	6	-
White, both sexes	11	12	13	12	17	29	36	38	21	17	3	-
Male	6	7	10	6	8	15	11	10	3	2	-	-
Female	5	5	3	6	9	14	25	28	18	15	3	-
Black, both sexes	14	18	12	7	13	18	17	20	16	2	3	-
Male	9	10	5	6	6	5	3	7	5	-	-	-
Female	5	8	7	1	7	13	14	13	11	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
South Dakota												
All races, both sexes	-	-	1	2	1	5	2	7	22	8	4	-
Male	-	-	-	2	1	1	-	-	3	3	-	-
Female	-	-	1	-	-	4	2	7	19	5	4	-
White, both sexes	-	-	1	2	1	5	2	7	22	8	4	-
Male	-	-	-	2	1	1	-	-	3	3	-	-
Female	-	-	1	-	-	4	2	7	19	5	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	49	67	54	45	44	70	128	122	97	44	6	-
Male	26	46	31	27	23	30	46	25	26	8	-	-
Female	23	21	23	18	21	40	82	97	71	36	6	-
White, both sexes	20	38	36	31	22	49	100	92	85	36	5	-
Male	11	30	21	18	12	19	31	18	23	6	-	-
Female	9	8	15	13	10	30	69	74	62	30	5	-
Black, both sexes	29	27	18	14	22	21	28	29	10	8	1	-
Male	15	15	10	9	11	11	15	7	3	2	-	-
Female	14	12	8	5	11	10	13	22	7	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Texas												
All races, both sexes	163	151	129	127	120	157	215	223	212	95	13	-
Male	106	87	83	71	52	72	77	56	45	13	2	-
Female	57	64	46	56	68	85	138	167	167	82	11	-
White, both sexes	108	102	98	86	89	119	179	198	186	87	10	-
Male	77	56	63	46	38	56	62	47	40	13	1	-
Female	31	46	35	40	51	63	117	151	146	74	9	-
Black, both sexes	50	46	29	37	30	35	35	24	23	8	3	-
Male	26	30	18	21	14	15	14	9	3	-	1	-
Female	24	16	11	16	16	20	21	15	20	8	2	-
American Indian 1/, both sexes	2	-	-	1	-	-	1	-	-	-	-	-
Male	2	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	2	3	1	3	-	1	3	-	-	-
Male	1	1	2	3	-	1	-	-	2	-	-	-
Female	2	2	-	-	1	2	-	1	1	-	-	-
Utah												
All races, both sexes	7	3	4	5	7	5	16	27	25	10	2	-
Male	4	2	2	3	2	2	6	10	5	4	1	-
Female	3	1	2	2	5	3	10	17	20	6	1	-
White, both sexes	7	3	4	4	6	4	15	27	25	10	2	-
Male	4	2	2	2	2	1	5	10	5	4	1	-
Female	3	1	2	2	4	3	10	17	20	6	1	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Vermont												
All races, both sexes	5	8	4	4	6	3	5	10	4	5	1	-
Male	2	6	1	3	3	1	2	2	2	1	-	-
Female	3	2	3	1	3	2	3	8	2	4	1	-
White, both sexes	5	8	3	4	6	3	5	10	4	5	1	-
Male	2	6	1	3	3	1	2	2	2	1	-	-
Female	3	2	2	1	3	2	3	8	2	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	22	21	12	17	38	34	45	74	49	25	8	-
Male	11	12	6	9	19	21	18	25	11	7	-	-
Female	11	9	6	8	19	13	27	49	38	18	8	-
White, both sexes	12	10	8	8	17	27	33	55	43	22	6	-
Male	7	3	4	3	7	17	12	18	9	6	-	-
Female	5	7	4	5	10	10	21	37	34	16	6	-
Black, both sexes	10	11	4	9	21	7	12	18	5	2	2	-
Male	4	9	2	6	12	4	6	6	2	1	-	-
Female	6	2	2	3	9	3	6	12	3	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Washington												
All races, both sexes	35	44	35	40	31	60	115	136	113	52	12	-
Male	28	36	24	27	14	25	48	38	30	8	2	-
Female	7	8	11	13	17	35	67	98	83	44	10	-
White, both sexes	28	38	30	33	29	54	103	130	107	50	12	-
Male	24	30	21	23	12	24	42	35	27	7	2	-
Female	4	8	9	10	17	30	61	95	80	43	10	-
Black, both sexes	5	4	4	3	1	3	6	3	1	-	-	-
Male	3	4	3	2	1	1	3	2	1	-	-	-
Female	2	-	1	1	-	2	3	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	3	1	-	-	-	-	-	-	-
Male	-	1	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	1	-	3	6	3	5	2	-	-
Male	1	1	-	-	-	-	3	1	2	1	-	-
Female	1	-	1	1	-	3	3	2	3	1	-	-
West Virginia												
All races, both sexes	4	3	2	5	5	7	15	22	24	13	3	-
Male	3	2	1	2	2	2	6	3	-	-	-	-
Female	1	1	1	3	3	5	9	19	24	13	3	-
White, both sexes	3	3	2	5	5	6	14	22	23	13	3	-
Male	2	2	1	2	2	2	5	3	-	-	-	-
Female	1	1	1	3	3	4	9	19	23	13	3	-
Black, both sexes	1	-	-	-	-	1	1	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Wisconsin												
All races, both sexes	20	15	25	23	30	30	53	72	79	36	3	-
Male	13	9	16	14	15	13	16	19	14	3	-	-
Female	7	6	9	9	15	17	37	53	65	33	3	-
White, both sexes	13	10	18	20	25	24	50	72	79	35	3	-
Male	10	5	12	12	13	11	15	19	14	3	-	-
Female	3	5	6	8	12	13	35	53	65	32	3	-
Black, both sexes	7	4	7	3	5	6	2	-	-	1	-	-
Male	3	3	4	2	2	2	1	-	-	-	-	-
Female	4	1	3	1	3	4	1	-	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	2	3	1	2	3	4	4	6	4	-	-
Male	3	2	1	1	2	2	1	1	-	1	-	-
Female	-	-	2	-	-	1	3	3	6	3	-	-
White, both sexes	3	2	3	1	2	3	4	4	6	4	-	-
Male	3	2	1	1	2	2	1	1	-	1	-	-
Female	-	-	2	-	-	1	3	3	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
United States												
All races, both sexes	9,831	-	1	-	2	3	14	15	50	62	143	212
Male	4,319	-	-	-	1	1	8	10	25	46	81	125
Female	5,512	-	1	-	1	2	6	5	25	16	62	87
White, both sexes	6,961	-	1	-	2	1	7	5	17	24	39	84
Male	3,013	-	-	-	1	-	3	3	9	20	21	51
Female	3,948	-	1	-	1	1	4	2	8	4	18	33
Black, both sexes	2,562	-	-	-	-	2	6	9	32	37	95	123
Male	1,164	-	-	-	-	1	4	6	15	26	56	72
Female	1,398	-	-	-	-	1	2	3	17	11	39	51
American Indian 1/, both sexes	56	-	-	-	-	-	1	-	-	-	4	2
Male	29	-	-	-	-	-	1	-	-	-	2	1
Female	27	-	-	-	-	-	-	-	-	-	2	1
Asian or Pacific Islander 2/, both sexes	252	-	-	-	-	-	-	1	1	1	5	3
Male	113	-	-	-	-	-	-	1	1	-	2	1
Female	139	-	-	-	-	-	-	-	-	1	3	2
Alabama												
All races, both sexes	162	-	-	-	-	-	-	-	2	3	7	8
Male	78	-	-	-	-	-	-	-	2	2	3	4
Female	84	-	-	-	-	-	-	-	-	1	4	4
White, both sexes	71	-	-	-	-	-	-	-	-	2	-	3
Male	32	-	-	-	-	-	-	-	-	2	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	91	-	-	-	-	-	-	-	2	1	7	5
Male	46	-	-	-	-	-	-	-	2	-	3	3
Female	45	-	-	-	-	-	-	-	-	1	4	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Alaska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	231	-	-	-	-	1	1	-	1	-	5	1
Male	102	-	-	-	-	1	1	-	-	-	2	-
Female	129	-	-	-	-	-	-	-	1	-	3	1
White, both sexes	203	-	-	-	-	-	-	-	1	-	2	1
Male	85	-	-	-	-	-	-	-	-	-	1	-
Female	118	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	8	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	1	-	-	-	3	-
Male	6	-	-	-	-	-	1	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Arkansas												
All races, both sexes	84	-	-	-	-	-	-	-	1	-	2	2
Male	33	-	-	-	-	-	-	-	-	-	2	1
Female	51	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	36	-	-	-	-	-	-	-	1	-	2	-
Male	18	-	-	-	-	-	-	-	-	-	2	-
Female	18	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,252	-	1	-	-	1	-	-	3	5	16	26
Male	580	-	-	-	-	-	-	-	1	4	9	17
Female	672	-	1	-	-	1	-	-	2	1	7	9
White, both sexes	906	-	1	-	-	1	-	-	2	2	9	12
Male	439	-	-	-	-	-	-	-	-	2	5	7
Female	467	-	1	-	-	1	-	-	2	-	4	5
Black, both sexes	204	-	-	-	-	-	-	-	-	2	4	14
Male	85	-	-	-	-	-	-	-	-	2	2	10
Female	119	-	-	-	-	-	-	-	-	-	2	4
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	136	-	-	-	-	-	-	-	1	1	2	-
Male	52	-	-	-	-	-	-	-	1	-	1	-
Female	84	-	-	-	-	-	-	-	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Colorado												
All races, both sexes	78	-	-	-	-	-	-	-	-	-	2	-
Male	32	-	-	-	-	-	-	-	-	-	1	-
Female	46	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	67	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	84	-	-	-	-	-	-	-	-	1	-	2
Male	36	-	-	-	-	-	-	-	-	1	-	2
Female	48	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Delaware												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	31	-	-	-	-	-	-	-	-	1	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	26	-	-	-	-	-	-	-	-	1	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Florida												
All races, both sexes	849	-	-	-	-	1	2	3	4	3	10	24
Male	428	-	-	-	-	-	1	2	2	3	4	12
Female	421	-	-	-	-	1	1	1	2	-	6	12
White, both sexes	625	-	-	-	-	-	1	2	1	2	2	10
Male	320	-	-	-	-	-	-	1	-	2	1	6
Female	305	-	-	-	-	-	1	1	1	-	1	4
Black, both sexes	222	-	-	-	-	1	1	1	3	1	8	14
Male	106	-	-	-	-	-	1	1	2	1	3	6
Female	116	-	-	-	-	1	-	-	1	-	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	313	-	-	-	-	-	1	3	3	6	9	11
Male	123	-	-	-	-	-	1	2	1	3	5	6
Female	190	-	-	-	-	-	-	1	2	3	4	5
White, both sexes	107	-	-	-	-	-	-	-	-	2	1	2
Male	44	-	-	-	-	-	-	-	-	1	1	1
Female	63	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	205	-	-	-	-	-	1	3	3	4	8	9
Male	79	-	-	-	-	-	1	2	1	2	4	5
Female	126	-	-	-	-	-	-	1	2	2	4	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Hawaii												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Illinois												
All races, both sexes	346	-	-	-	-	-	-	-	2	3	5	8
Male	144	-	-	-	-	-	-	-	1	-	3	5
Female	202	-	-	-	-	-	-	-	1	3	2	3
White, both sexes	243	-	-	-	-	-	-	-	-	2	1	3
Male	98	-	-	-	-	-	-	-	-	-	1	2
Female	145	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	99	-	-	-	-	-	-	-	2	1	4	4
Male	44	-	-	-	-	-	-	-	1	-	2	3
Female	55	-	-	-	-	-	-	-	1	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Indiana												
All races, both sexes	175	-	-	-	-	-	-	-	1	1	2	3
Male	85	-	-	-	-	-	-	-	-	-	2	3
Female	90	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	137	-	-	-	-	-	-	-	-	-	-	1
Male	66	-	-	-	-	-	-	-	-	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	37	-	-	-	-	-	-	-	1	1	2	2
Male	19	-	-	-	-	-	-	-	-	-	2	2
Female	18	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Iowa												
All races, both sexes	141	-	-	-	-	-	-	-	-	-	-	2
Male	61	-	-	-	-	-	-	-	-	-	-	-
Female	80	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	137	-	-	-	-	-	-	-	-	-	-	2
Male	59	-	-	-	-	-	-	-	-	-	-	-
Female	78	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	80	-	-	-	-	-	-	-	-	-	-	4
Male	35	-	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	64	-	-	-	-	-	-	-	-	-	-	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	37	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Kentucky												
All races, both sexes	132	-	-	-	-	-	-	1	-	2	2	1
Male	64	-	-	-	-	-	-	-	-	1	1	1
Female	68	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	117	-	-	-	-	-	-	1	-	1	1	1
Male	57	-	-	-	-	-	-	-	-	1	-	1
Female	60	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	15	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	158	-	-	-	-	-	2	-	1	3	5	2
Male	64	-	-	-	-	-	1	-	-	3	3	2
Female	94	-	-	-	-	-	1	-	1	-	2	-
White, both sexes	72	-	-	-	-	-	1	-	-	-	2	-
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	48	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	85	-	-	-	-	-	1	-	1	3	3	2
Male	39	-	-	-	-	-	1	-	-	3	2	2
Female	46	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Maine												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	192	-	-	-	-	-	1	1	-	-	5	7
Male	92	-	-	-	-	-	1	1	-	-	3	4
Female	100	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	96	-	-	-	-	-	-	-	-	-	-	1
Male	45	-	-	-	-	-	-	-	-	-	-	1
Female	51	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	94	-	-	-	-	-	1	1	-	-	5	6
Male	46	-	-	-	-	-	1	1	-	-	3	3
Female	48	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Massachusetts												
All races, both sexes	117	-	-	-	-	-	-	-	1	-	-	1
Male	45	-	-	-	-	-	-	-	1	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	101	-	-	-	-	-	-	-	-	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	66	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	336	-	-	-	1	-	-	1	1	2	3	4
Male	144	-	-	-	-	-	-	1	1	2	1	2
Female	192	-	-	-	1	-	-	-	-	-	2	2
White, both sexes	235	-	-	-	1	-	-	-	-	-	1	1
Male	93	-	-	-	-	-	-	-	-	-	1	-
Female	142	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	95	-	-	-	-	-	-	-	1	2	2	3
Male	49	-	-	-	-	-	-	-	1	2	-	2
Female	46	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Minnesota												
All races, both sexes	149	-	-	-	-	-	1	-	-	2	-	-
Male	52	-	-	-	-	-	-	-	-	2	-	-
Female	97	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	140	-	-	-	-	-	1	-	-	2	-	-
Male	49	-	-	-	-	-	-	-	-	2	-	-
Female	91	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	138	-	-	-	-	-	-	1	5	-	2	4
Male	53	-	-	-	-	-	-	1	2	-	2	3
Female	85	-	-	-	-	-	-	-	3	-	-	1
White, both sexes	44	-	-	-	-	-	-	-	1	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	93	-	-	-	-	-	-	1	4	-	2	3
Male	35	-	-	-	-	-	-	1	2	-	2	2
Female	58	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Missouri												
All races, both sexes	197	-	-	-	-	-	-	-	2	1	2	5
Male	78	-	-	-	-	-	-	-	-	-	2	5
Female	119	-	-	-	-	-	-	-	2	1	-	-
White, both sexes	137	-	-	-	-	-	-	-	1	-	-	3
Male	53	-	-	-	-	-	-	-	-	-	-	3
Female	84	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	57	-	-	-	-	-	-	-	1	1	2	2
Male	24	-	-	-	-	-	-	-	-	-	2	2
Female	33	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Nebraska												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	64	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	2	2
Male	22	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Hampshire												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	186	-	-	-	-	-	-	-	-	-	4	4
Male	89	-	-	-	-	-	-	-	-	-	2	1
Female	97	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	120	-	-	-	-	-	-	-	-	-	1	2
Male	58	-	-	-	-	-	-	-	-	-	1	-
Female	62	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	61	-	-	-	-	-	-	-	-	-	3	2
Male	28	-	-	-	-	-	-	-	-	-	1	1
Female	33	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Mexico												
All races, both sexes	48	-	-	-	-	-	-	-	1	1	1	-
Male	21	-	-	-	-	-	-	-	1	1	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	1	1	-	-
Male	15	-	-	-	-	-	-	-	1	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	666	-	-	-	-	-	-	-	2	2	19	20
Male	301	-	-	-	-	-	-	-	-	1	12	12
Female	365	-	-	-	-	-	-	-	2	1	7	8
White, both sexes	446	-	-	-	-	-	-	-	-	1	6	7
Male	190	-	-	-	-	-	-	-	-	-	4	7
Female	256	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	205	-	-	-	-	-	-	-	2	1	12	11
Male	101	-	-	-	-	-	-	-	-	1	8	4
Female	104	-	-	-	-	-	-	-	2	-	4	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
North Carolina												
All races, both sexes	302	-	-	-	-	-	-	2	-	1	5	17
Male	122	-	-	-	-	-	-	1	-	1	4	8
Female	180	-	-	-	-	-	-	1	-	-	1	9
White, both sexes	153	-	-	-	-	-	-	-	-	-	-	5
Male	58	-	-	-	-	-	-	-	-	-	-	3
Female	95	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	143	-	-	-	-	-	-	2	-	1	5	10
Male	60	-	-	-	-	-	-	1	-	1	4	4
Female	83	-	-	-	-	-	-	1	-	-	1	6
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Ohio												
All races, both sexes	501	-	-	-	-	-	2	-	1	5	4	7
Male	194	-	-	-	-	-	1	-	1	5	2	4
Female	307	-	-	-	-	-	1	-	-	-	2	3
White, both sexes	381	-	-	-	-	-	2	-	1	2	2	2
Male	143	-	-	-	-	-	1	-	1	2	1	1
Female	238	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	118	-	-	-	-	-	-	-	-	3	2	5
Male	50	-	-	-	-	-	-	-	-	3	1	3
Female	68	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	105	-	-	-	-	-	-	1	1	1	1	1
Male	47	-	-	-	-	-	-	1	1	1	1	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	85	-	-	-	-	-	-	1	1	-	-	-
Male	35	-	-	-	-	-	-	1	1	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	1	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Oregon												
All races, both sexes	115	-	-	-	-	-	-	1	-	-	2	-
Male	43	-	-	-	-	-	-	1	-	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	110	-	-	-	-	-	-	1	-	-	2	-
Male	40	-	-	-	-	-	-	1	-	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	352	-	-	-	-	-	-	-	2	-	3	5
Male	143	-	-	-	-	-	-	-	-	-	3	4
Female	209	-	-	-	-	-	-	-	2	-	-	1
White, both sexes	290	-	-	-	-	-	-	-	1	-	1	1
Male	118	-	-	-	-	-	-	-	-	-	1	1
Female	172	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	59	-	-	-	-	-	-	-	1	-	2	4
Male	24	-	-	-	-	-	-	-	-	-	2	3
Female	35	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Rhode Island												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	180	-	-	-	-	-	-	-	2	-	3	4
Male	80	-	-	-	-	-	-	-	1	-	2	1
Female	100	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	85	-	-	-	-	-	-	-	-	-	-	2
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	94	-	-	-	-	-	-	-	2	-	3	2
Male	37	-	-	-	-	-	-	-	1	-	2	-
Female	57	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
South Dakota												
All races, both sexes	43	-	-	-	-	-	-	-	-	1	-	-
Male	22	-	-	-	-	-	-	-	-	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	1	-	-
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	275	-	-	-	-	-	1	-	3	8	4	6
Male	121	-	-	-	-	-	-	-	2	6	2	4
Female	154	-	-	-	-	-	1	-	1	2	2	2
White, both sexes	177	-	-	-	-	-	-	-	1	2	1	1
Male	74	-	-	-	-	-	-	-	1	2	-	1
Female	103	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	98	-	-	-	-	-	1	-	2	6	3	5
Male	47	-	-	-	-	-	-	-	1	4	2	3
Female	51	-	-	-	-	-	1	-	1	2	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Texas												
All races, both sexes	719	-	-	-	1	-	1	-	4	5	12	15
Male	313	-	-	-	1	-	1	-	3	5	6	9
Female	406	-	-	-	-	-	-	-	1	-	6	6
White, both sexes	522	-	-	-	1	-	1	-	1	2	6	8
Male	219	-	-	-	1	-	1	-	1	2	3	5
Female	303	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	193	-	-	-	-	-	-	-	3	3	6	7
Male	93	-	-	-	-	-	-	-	2	3	3	4
Female	100	-	-	-	-	-	-	-	1	-	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Vermont												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	199	-	-	-	-	-	2	1	3	3	4	4
Male	95	-	-	-	-	-	1	-	2	3	2	3
Female	104	-	-	-	-	-	1	1	1	-	2	1
White, both sexes	110	-	-	-	-	-	1	-	1	1	-	1
Male	49	-	-	-	-	-	1	-	1	1	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	87	-	-	-	-	-	1	1	2	2	4	3
Male	45	-	-	-	-	-	-	-	1	2	2	3
Female	42	-	-	-	-	-	1	1	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Washington												
All races, both sexes	198	-	-	-	-	-	-	-	2	1	-	1
Male	87	-	-	-	-	-	-	-	1	1	-	1
Female	111	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	169	-	-	-	-	-	-	-	2	1	-	-
Male	73	-	-	-	-	-	-	-	1	1	-	-
Female	96	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	68	-	-	-	-	-	-	-	2	-	-	-
Male	27	-	-	-	-	-	-	-	2	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	2	-	-	-
Male	24	-	-	-	-	-	-	-	2	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Wisconsin												
All races, both sexes	183	-	-	-	-	-	-	-	-	1	-	6
Male	69	-	-	-	-	-	-	-	-	-	-	2
Female	114	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	167	-	-	-	-	-	-	-	-	-	-	3
Male	64	-	-	-	-	-	-	-	-	-	-	1
Female	103	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	3
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
United States												
All races, both sexes	325	427	475	583	795	1,239	1,775	1,887	1,272	454	97	-
Male	194	234	255	288	367	549	802	776	429	111	17	-
Female	131	193	220	295	428	690	973	1,111	843	343	80	-
White, both sexes	125	215	249	326	505	881	1,403	1,560	1,072	366	79	-
Male	74	111	144	164	226	410	645	650	375	93	13	-
Female	51	104	105	162	279	471	758	910	697	273	66	-
Black, both sexes	189	196	201	234	261	317	323	286	158	80	13	-
Male	114	113	98	109	124	123	133	112	41	15	2	-
Female	75	83	103	125	137	194	190	174	117	65	11	-
American Indian 1/, both sexes	2	3	6	7	5	6	6	7	2	3	2	-
Male	2	3	3	4	3	5	1	1	1	1	1	-
Female	-	-	3	3	2	1	5	6	1	2	1	-
Asian or Pacific Islander 2/, both sexes	9	13	19	16	24	35	43	34	40	5	3	-
Male	4	7	10	11	14	11	23	13	12	2	1	-
Female	5	6	9	5	10	24	20	21	28	3	2	-
Alabama												
All races, both sexes	7	12	13	11	14	18	30	16	13	7	1	-
Male	6	6	6	4	5	10	13	9	7	1	-	-
Female	1	6	7	7	9	8	17	7	6	6	1	-
White, both sexes	2	4	4	6	7	6	17	9	8	3	-	-
Male	2	2	2	3	2	2	7	6	3	-	-	-
Female	-	2	2	3	5	4	10	3	5	3	-	-
Black, both sexes	5	8	9	5	7	12	13	7	5	4	1	-
Male	4	4	4	1	3	8	6	3	4	1	-	-
Female	1	4	5	4	4	4	7	4	1	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Alaska												
All races, both sexes	1	-	1	-	-	2	3	2	1	-	1	-
Male	1	-	1	-	-	2	1	-	1	-	-	-
Female	-	-	-	-	-	-	2	2	-	-	1	-
White, both sexes	-	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	1	-	-	-	1	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	3	13	8	9	24	30	50	46	27	8	4	-
Male	2	6	5	1	13	9	23	25	11	2	1	-
Female	1	7	3	8	11	21	27	21	16	6	3	-
White, both sexes	2	9	6	8	20	28	47	44	26	6	3	-
Male	2	3	4	1	10	9	20	24	10	1	-	-
Female	-	6	2	7	10	19	27	20	16	5	3	-
Black, both sexes	1	2	-	-	1	1	2	-	-	-	-	-
Male	-	2	-	-	1	-	2	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	1	1	-	-	1	-	1	1	-
Male	-	1	1	-	1	-	-	-	-	-	1	-
Female	-	-	1	1	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	1	1	1	1	1	-	-
Male	-	-	-	-	1	-	1	1	1	1	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Arkansas												
All races, both sexes	4	6	2	8	8	4	9	19	9	7	3	-
Male	-	2	1	4	4	2	5	5	3	2	2	-
Female	4	4	1	4	4	2	4	14	6	5	1	-
White, both sexes	1	2	1	6	5	3	6	13	4	4	1	-
Male	-	-	-	3	2	2	3	1	1	1	1	-
Female	1	2	1	3	3	1	3	12	3	3	-	-
Black, both sexes	3	4	1	2	3	1	3	6	5	3	2	-
Male	-	2	1	1	2	-	2	4	2	1	1	-
Female	3	2	-	1	1	1	1	2	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	42	59	74	85	91	144	221	233	177	61	13	-
Male	22	33	39	45	46	64	109	93	76	20	2	-
Female	20	26	35	40	45	80	112	140	101	41	11	-
White, both sexes	23	42	49	51	63	96	174	185	136	50	10	-
Male	13	23	28	30	33	46	88	79	66	17	2	-
Female	10	19	21	21	30	50	86	106	70	33	8	-
Black, both sexes	15	13	11	23	17	30	20	29	16	8	2	-
Male	8	9	5	7	7	11	9	9	4	2	-	-
Female	7	4	6	16	10	19	11	20	12	6	2	-
American Indian 1/, both sexes	-	-	-	3	-	-	1	1	-	-	-	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	4	14	8	11	18	26	18	25	3	1	-
Male	1	1	6	6	6	7	11	5	6	1	-	-
Female	3	3	8	2	5	11	15	13	19	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Colorado												
All races, both sexes	1	4	-	5	6	15	14	13	13	2	3	-
Male	-	3	-	2	3	6	6	6	4	1	-	-
Female	1	1	-	3	3	9	8	7	9	1	3	-
White, both sexes	1	4	-	4	3	14	12	11	13	2	3	-
Male	-	3	-	2	1	6	5	5	4	1	-	-
Female	1	1	-	2	2	8	7	6	9	1	3	-
Black, both sexes	-	-	-	1	2	-	2	2	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	1	2	2	2	7	14	24	18	7	1	-
Male	1	-	-	1	1	1	11	10	7	1	-	-
Female	2	1	2	1	1	6	3	14	11	6	1	-
White, both sexes	2	1	1	1	1	4	14	23	18	6	1	-
Male	1	-	-	1	-	1	11	10	7	1	-	-
Female	1	1	1	-	1	3	3	13	11	5	1	-
Black, both sexes	1	-	1	1	1	3	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	1	1	-	3	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Delaware												
All races, both sexes	-	2	1	2	2	3	2	4	4	-	-	-
Male	-	1	-	2	1	2	1	1	2	-	-	-
Female	-	1	1	-	1	1	1	3	2	-	-	-
White, both sexes	-	1	1	1	1	3	1	4	3	-	-	-
Male	-	-	-	1	1	2	1	1	2	-	-	-
Female	-	1	1	-	-	1	-	3	1	-	-	-
Black, both sexes	-	1	-	1	1	-	1	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	4	3	2	5	2	5	4	-	1	-	-
Male	2	4	2	-	2	1	3	1	-	1	-	-
Female	1	-	1	2	3	1	2	3	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	2	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	4	3	2	5	2	3	2	-	-	-	-
Male	2	4	2	-	2	1	1	-	-	-	-	-
Female	1	-	1	2	3	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Florida												
All races, both sexes	18	30	44	54	84	117	140	170	105	33	7	-
Male	12	21	21	26	43	62	79	78	47	13	2	-
Female	6	9	23	28	41	55	61	92	58	20	5	-
White, both sexes	8	16	22	34	59	81	116	150	91	25	5	-
Male	5	12	10	16	32	46	66	68	43	11	1	-
Female	3	4	12	18	27	35	50	82	48	14	4	-
Black, both sexes	10	14	21	20	24	36	24	20	14	8	2	-
Male	7	9	10	10	10	16	13	10	4	2	1	-
Female	3	5	11	10	14	20	11	10	10	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	15	21	18	26	33	38	49	44	23	12	1	-
Male	12	9	6	15	10	11	17	17	7	1	-	-
Female	3	12	12	11	23	27	32	27	16	11	1	-
White, both sexes	1	4	1	8	11	15	23	22	13	3	1	-
Male	1	2	1	5	5	4	11	8	4	-	-	-
Female	-	2	-	3	6	11	12	14	9	3	1	-
Black, both sexes	14	17	17	18	21	23	26	22	10	9	-	-
Male	11	7	5	10	5	7	6	9	3	1	-	-
Female	3	10	12	8	16	16	20	13	7	8	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Hawaii												
All races, both sexes	1	-	3	2	3	4	6	7	13	1	1	-
Male	1	-	2	1	2	2	4	3	7	-	-	-
Female	-	-	1	1	1	2	2	4	6	1	1	-
White, both sexes	-	-	-	1	1	-	2	3	4	-	-	-
Male	-	-	-	-	-	-	1	2	3	-	-	-
Female	-	-	-	1	1	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	1	2	4	4	4	9	1	1	-
Male	1	-	2	1	2	2	3	1	4	-	-	-
Female	-	-	1	-	-	2	1	3	5	1	1	-
Idaho												
All races, both sexes	-	1	3	4	6	12	11	12	3	2	1	-
Male	-	-	3	3	2	6	7	4	-	1	-	-
Female	-	1	-	1	4	6	4	8	3	1	1	-
White, both sexes	-	1	3	4	6	12	11	11	3	2	1	-
Male	-	-	3	3	2	6	7	3	-	1	-	-
Female	-	1	-	1	4	6	4	8	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Illinois												
All races, both sexes	16	11	19	21	35	37	63	66	41	16	3	-
Male	9	7	12	10	18	15	21	27	14	2	-	-
Female	7	4	7	11	17	22	42	39	27	14	3	-
White, both sexes	3	5	9	13	19	27	49	60	37	13	2	-
Male	3	3	7	7	9	11	17	23	13	2	-	-
Female	-	2	2	6	10	16	32	37	24	11	2	-
Black, both sexes	12	6	10	7	16	10	13	6	4	3	1	-
Male	6	4	5	2	9	4	3	4	1	-	-	-
Female	6	2	5	5	7	6	10	2	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	4	10	3	7	14	22	41	41	17	8	1	-
Male	4	7	2	4	5	14	17	23	3	1	-	-
Female	-	3	1	3	9	8	24	18	14	7	1	-
White, both sexes	3	4	2	5	10	18	31	38	16	8	1	-
Male	3	2	1	2	5	13	12	23	3	1	-	-
Female	-	2	1	3	5	5	19	15	13	7	1	-
Black, both sexes	1	6	1	2	4	3	10	3	1	-	-	-
Male	1	5	1	2	-	1	5	-	-	-	-	-
Female	-	1	-	-	4	2	5	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Iowa												
All races, both sexes	1	3	2	4	5	15	31	33	30	12	3	-
Male	-	1	-	-	3	8	15	20	11	3	-	-
Female	1	2	2	4	2	7	16	13	19	9	3	-
White, both sexes	1	2	2	3	5	15	30	32	30	12	3	-
Male	-	-	-	-	3	8	15	19	11	3	-	-
Female	1	2	2	3	2	7	15	13	19	9	3	-
Black, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	2	5	5	6	8	10	25	6	6	2	-
Male	1	-	2	3	2	5	3	11	3	3	-	-
Female	-	2	3	2	4	3	7	14	3	3	2	-
White, both sexes	1	1	4	3	4	6	9	21	5	5	2	-
Male	1	-	1	1	2	3	3	9	3	2	-	-
Female	-	1	3	2	2	3	6	12	2	3	2	-
Black, both sexes	-	1	1	2	2	1	1	4	1	1	-	-
Male	-	-	1	2	-	1	-	2	-	1	-	-
Female	-	1	-	-	2	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Kentucky												
All races, both sexes	4	3	4	9	11	22	31	19	16	6	1	-
Male	2	3	2	6	8	11	13	8	7	1	-	-
Female	2	-	2	3	3	11	18	11	9	5	1	-
White, both sexes	4	2	3	6	11	21	30	17	14	4	1	-
Male	2	2	1	4	8	10	13	7	7	1	-	-
Female	2	-	2	2	3	11	17	10	7	3	1	-
Black, both sexes	-	1	1	3	-	1	1	2	2	2	-	-
Male	-	1	1	2	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	1	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	10	10	7	11	30	28	23	18	1	1	-
Male	3	5	6	3	5	11	10	7	5	-	-	-
Female	3	5	4	4	6	19	18	16	13	1	1	-
White, both sexes	3	2	1	3	5	14	16	14	9	1	1	-
Male	2	1	-	3	1	5	4	5	2	-	-	-
Female	1	1	1	-	4	9	12	9	7	1	1	-
Black, both sexes	3	8	9	4	6	15	12	9	9	-	-	-
Male	1	4	6	-	4	5	6	2	3	-	-	-
Female	2	4	3	4	2	10	6	7	6	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Maine												
All races, both sexes	1	1	4	2	4	4	9	10	4	3	-	-
Male	1	1	3	2	1	4	3	4	2	-	-	-
Female	-	-	1	-	3	-	6	6	2	3	-	-
White, both sexes	1	1	4	2	4	4	9	10	4	3	-	-
Male	1	1	3	2	1	4	3	4	2	-	-	-
Female	-	-	1	-	3	-	6	6	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	8	5	6	18	20	25	26	46	16	7	1	-
Male	6	4	2	8	12	11	9	25	4	2	-	-
Female	2	1	4	10	8	14	17	21	12	5	1	-
White, both sexes	1	1	6	7	5	12	13	34	10	5	1	-
Male	1	1	2	2	2	5	6	19	4	2	-	-
Female	-	-	4	5	3	7	7	15	6	3	1	-
Black, both sexes	7	4	-	11	15	12	13	11	6	2	-	-
Male	5	3	-	6	10	6	3	5	-	-	-	-
Female	2	1	-	5	5	6	10	6	6	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Massachusetts												
All races, both sexes	2	1	4	5	3	14	24	29	23	9	1	-
Male	2	-	2	3	1	6	12	10	7	1	-	-
Female	-	1	2	2	2	8	12	19	16	8	1	-
White, both sexes	1	1	1	2	2	11	22	29	21	9	1	-
Male	1	-	1	-	-	5	12	10	5	1	-	-
Female	-	1	-	2	2	6	10	19	16	8	1	-
Black, both sexes	1	-	3	3	-	2	1	-	1	-	-	-
Male	1	-	1	3	-	1	-	-	1	-	-	-
Female	-	-	2	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	8	11	16	14	24	52	75	66	36	19	3	-
Male	5	6	9	8	7	24	35	32	9	2	-	-
Female	3	5	7	6	17	28	40	34	27	17	3	-
White, both sexes	2	3	8	5	14	39	57	54	29	18	3	-
Male	2	1	3	3	3	18	27	27	6	2	-	-
Female	-	2	5	2	11	21	30	27	23	16	3	-
Black, both sexes	6	8	8	7	10	11	18	11	7	1	-	-
Male	3	5	6	4	4	6	8	5	3	-	-	-
Female	3	3	2	3	6	5	10	6	4	1	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Minnesota												
All races, both sexes	1	3	5	5	9	18	26	40	26	11	2	-
Male	1	2	3	1	3	7	10	12	9	2	-	-
Female	-	1	2	4	6	11	16	28	17	9	2	-
White, both sexes	1	3	3	4	9	18	25	37	25	10	2	-
Male	1	2	2	1	3	7	10	11	9	1	-	-
Female	-	1	1	3	6	11	15	26	16	9	2	-
Black, both sexes	-	-	1	1	-	-	1	2	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	1	1	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	13	10	9	15	8	22	22	17	5	5	-	-
Male	3	5	4	6	4	9	8	2	2	2	-	-
Female	10	5	5	9	4	13	14	15	3	3	-	-
White, both sexes	3	1	1	3	3	11	10	5	2	3	-	-
Male	1	1	-	3	2	3	3	2	1	1	-	-
Female	2	-	1	-	1	8	7	3	1	2	-	-
Black, both sexes	10	9	8	11	5	11	12	12	3	2	-	-
Male	2	4	4	3	2	6	5	-	1	1	-	-
Female	8	5	4	8	3	5	7	12	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Missouri												
All races, both sexes	7	8	10	9	15	24	32	46	25	10	1	-
Male	4	4	6	3	7	9	12	13	7	5	1	-
Female	3	4	4	6	8	15	20	33	18	5	-	-
White, both sexes	2	2	6	6	8	14	24	38	24	8	1	-
Male	1	1	5	2	4	7	9	9	6	5	1	-
Female	1	1	1	4	4	7	15	29	18	3	-	-
Black, both sexes	5	5	4	3	6	9	8	8	1	2	-	-
Male	3	2	1	1	3	2	3	4	1	-	-	-
Female	2	3	3	2	3	7	5	4	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	1	1	-	2	8	2	2	-	-
Male	-	-	1	-	1	-	1	4	1	-	-	-
Female	-	-	-	1	-	-	1	4	1	2	-	-
White, both sexes	-	-	1	1	1	-	2	8	2	2	-	-
Male	-	-	1	-	1	-	1	4	1	-	-	-
Female	-	-	-	1	-	-	1	4	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Nebraska												
All races, both sexes	-	1	1	1	4	9	9	20	15	5	3	-
Male	-	-	1	1	2	2	7	7	7	-	1	-
Female	-	1	-	-	2	7	2	13	8	5	2	-
White, both sexes	-	1	-	1	3	7	9	20	15	5	3	-
Male	-	-	-	1	1	2	7	7	7	-	1	-
Female	-	1	-	-	2	5	2	13	8	5	2	-
Black, both sexes	-	-	1	-	1	2	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	1	2	3	7	5	6	4	7	2	-	-
Male	2	1	1	2	1	4	3	2	3	-	-	-
Female	1	-	1	1	6	1	3	2	4	2	-	-
White, both sexes	3	-	2	2	6	4	6	3	7	1	-	-
Male	2	-	1	1	1	4	3	2	3	-	-	-
Female	1	-	1	1	5	-	3	1	4	1	-	-
Black, both sexes	-	1	-	1	1	1	-	-	-	1	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Hampshire												
All races, both sexes	2	-	2	4	2	2	4	7	8	2	-	-
Male	2	-	1	1	2	1	2	2	2	1	-	-
Female	-	-	1	3	-	1	2	5	6	1	-	-
White, both sexes	2	-	2	3	2	2	4	7	8	2	-	-
Male	2	-	1	1	2	1	2	2	2	1	-	-
Female	-	-	1	2	-	1	2	5	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	12	10	9	13	20	43	36	20	7	1	-
Male	4	9	6	2	6	8	27	15	7	1	1	-
Female	3	3	4	7	7	12	16	21	13	6	-	-
White, both sexes	2	2	3	6	7	14	31	27	19	5	1	-
Male	1	2	3	-	2	6	21	13	7	1	1	-
Female	1	-	-	6	5	8	10	14	12	4	-	-
Black, both sexes	5	9	7	2	6	6	10	8	1	2	-	-
Male	3	6	3	1	4	2	5	2	-	-	-	-
Female	2	3	4	1	2	4	5	6	1	2	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Mexico												
All races, both sexes	2	1	3	4	8	6	8	3	7	3	-	-
Male	1	-	1	2	3	3	3	1	3	1	-	-
Female	1	1	2	2	5	3	5	2	4	2	-	-
White, both sexes	2	1	2	2	6	5	7	2	6	2	-	-
Male	1	-	1	-	2	3	3	-	2	1	-	-
Female	1	1	1	2	4	2	4	2	4	1	-	-
Black, both sexes	-	-	-	1	1	1	1	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	-	-	1	1	1	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	1	-	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	32	44	44	44	50	79	93	128	71	33	5	-
Male	20	19	26	26	23	31	49	59	16	7	-	-
Female	12	25	18	18	27	48	44	69	55	26	5	-
White, both sexes	8	25	24	22	28	54	72	102	64	29	4	-
Male	6	10	15	14	11	22	35	44	15	7	-	-
Female	2	15	9	8	17	32	37	58	49	22	4	-
Black, both sexes	22	17	20	21	22	24	19	23	6	4	1	-
Male	12	7	11	11	12	9	12	13	1	-	-	-
Female	10	10	9	10	10	15	7	10	5	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	1	-	1	2	3	1	-	-	-
Male	2	2	-	1	-	-	2	2	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
North Carolina												
All races, both sexes	12	15	16	24	27	50	56	41	25	9	2	-
Male	9	6	9	11	9	20	25	12	7	-	-	-
Female	3	9	7	13	18	30	31	29	18	9	2	-
White, both sexes	1	5	6	6	13	28	36	26	19	6	2	-
Male	1	3	5	3	4	11	14	7	7	-	-	-
Female	-	2	1	3	9	17	22	19	12	6	2	-
Black, both sexes	10	10	10	18	13	22	18	15	6	3	-	-
Male	7	3	4	8	4	9	10	5	-	-	-	-
Female	3	7	6	10	9	13	8	10	6	3	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	2	2	-	-	2	5	7	10	4	1	-
Male	-	2	1	-	-	-	2	3	1	1	-	-
Female	-	-	1	-	-	2	3	4	9	3	1	-
White, both sexes	-	2	1	-	-	2	5	7	9	4	1	-
Male	-	2	1	-	-	-	2	3	1	1	-	-
Female	-	-	-	-	-	2	3	4	8	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Ohio												
All races, both sexes	8	16	13	22	42	70	96	103	91	19	2	-
Male	6	9	9	12	13	29	37	37	21	7	1	-
Female	2	7	4	10	29	41	59	66	70	12	1	-
White, both sexes	2	11	8	12	28	49	78	86	82	14	2	-
Male	2	5	5	9	8	20	30	32	20	5	1	-
Female	-	6	3	3	20	29	48	54	62	9	1	-
Black, both sexes	6	5	5	10	14	19	18	17	9	5	-	-
Male	4	4	4	3	5	8	7	5	1	2	-	-
Female	2	1	1	7	9	11	11	12	8	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	10	7	7	8	10	17	19	13	3	2	-
Male	1	4	2	3	4	6	9	5	5	1	2	-
Female	3	6	5	4	4	4	8	14	8	2	-	-
White, both sexes	3	8	5	4	6	8	15	18	11	3	2	-
Male	-	2	1	1	3	5	8	5	5	1	2	-
Female	3	6	4	3	3	3	7	13	6	2	-	-
Black, both sexes	1	1	1	3	2	1	2	-	1	-	-	-
Male	1	1	-	2	1	-	1	-	-	-	-	-
Female	-	-	1	1	1	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	1	-	1	1	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Oregon												
All races, both sexes	1	3	2	3	12	12	19	28	25	4	3	-
Male	1	3	-	1	5	6	5	10	7	2	2	-
Female	-	-	2	2	7	6	14	18	18	2	1	-
White, both sexes	1	2	2	2	11	12	17	28	25	4	3	-
Male	1	2	-	1	4	6	4	10	7	2	2	-
Female	-	-	2	1	7	6	13	18	18	2	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	11	6	10	10	16	48	74	91	52	24	-	-
Male	8	2	4	5	6	18	30	42	18	3	-	-
Female	3	4	6	5	10	30	44	49	34	21	-	-
White, both sexes	4	2	7	7	14	40	72	78	43	20	-	-
Male	3	-	3	3	5	16	29	38	16	3	-	-
Female	1	2	4	4	9	24	43	40	27	17	-	-
Black, both sexes	7	4	3	2	1	8	2	12	9	4	-	-
Male	5	2	1	1	1	2	1	4	2	-	-	-
Female	2	2	2	1	-	6	1	8	7	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Rhode Island												
All races, both sexes	-	-	-	2	1	4	7	6	8	-	1	-
Male	-	-	-	-	1	1	5	3	2	-	-	-
Female	-	-	-	2	-	3	2	3	6	-	1	-
White, both sexes	-	-	-	2	1	4	6	6	8	-	-	-
Male	-	-	-	-	1	1	5	3	2	-	-	-
Female	-	-	-	2	-	3	1	3	6	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	7	12	12	17	10	18	36	38	17	3	1	-
Male	4	10	9	8	5	7	12	17	3	1	-	-
Female	3	2	3	9	5	11	24	21	14	2	1	-
White, both sexes	1	3	2	7	6	7	21	24	12	-	-	-
Male	-	2	2	5	3	5	9	13	3	-	-	-
Female	1	1	-	2	3	2	12	11	9	-	-	-
Black, both sexes	5	9	10	10	4	11	15	14	5	3	1	-
Male	4	8	7	3	2	2	3	4	-	1	-	-
Female	1	1	3	7	2	9	12	10	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
South Dakota												
All races, both sexes	1	-	1	1	1	6	9	6	9	5	3	-
Male	1	-	1	1	1	5	3	1	3	3	2	-
Female	-	-	-	-	-	1	6	5	6	2	1	-
White, both sexes	1	-	1	1	1	4	9	6	9	5	2	-
Male	1	-	1	1	1	3	3	1	3	3	1	-
Female	-	-	-	-	-	1	6	5	6	2	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	9	14	12	19	21	35	43	48	34	12	6	-
Male	4	9	8	9	12	12	22	17	12	2	-	-
Female	5	5	4	10	9	23	21	31	22	10	6	-
White, both sexes	4	8	4	11	11	28	29	38	27	8	4	-
Male	2	6	3	4	6	11	16	11	10	1	-	-
Female	2	2	1	7	5	17	13	27	17	7	4	-
Black, both sexes	5	6	8	8	10	7	14	10	7	4	2	-
Male	2	3	5	5	6	1	6	6	2	1	-	-
Female	3	3	3	3	4	6	8	4	5	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Texas												
All races, both sexes	38	35	45	49	72	96	126	107	88	23	2	-
Male	22	20	19	28	35	45	49	41	21	8	-	-
Female	16	15	26	21	37	51	77	66	67	15	2	-
White, both sexes	19	20	28	28	48	74	105	89	71	19	2	-
Male	8	11	15	14	20	37	42	35	17	7	-	-
Female	11	9	13	14	28	37	63	54	54	12	2	-
Black, both sexes	19	14	17	21	22	22	20	18	17	4	-	-
Male	14	9	4	14	14	8	7	6	4	1	-	-
Female	5	5	13	7	8	14	13	12	13	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	1	2	1	4	5	6	4	3	-	-
Male	-	-	1	2	-	3	1	2	1	-	-	-
Female	-	1	-	-	1	1	4	4	3	3	-	-
White, both sexes	-	-	1	2	1	4	5	6	4	3	-	-
Male	-	-	1	2	-	3	1	2	1	-	-	-
Female	-	-	-	-	1	1	4	4	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Vermont												
All races, both sexes	1	1	-	1	-	2	3	2	2	-	-	-
Male	-	1	-	1	-	2	1	1	-	-	-	-
Female	1	-	-	-	-	-	2	1	2	-	-	-
White, both sexes	1	1	-	1	-	2	2	2	2	-	-	-
Male	-	1	-	1	-	2	1	1	-	-	-	-
Female	1	-	-	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	6	8	10	23	18	39	37	20	9	1	-
Male	7	1	3	6	11	7	19	16	12	2	-	-
Female	4	5	5	4	12	11	20	21	8	7	1	-
White, both sexes	-	2	2	4	9	12	25	32	13	6	1	-
Male	-	-	1	2	4	4	11	14	8	2	-	-
Female	-	2	1	2	5	8	14	18	5	4	1	-
Black, both sexes	11	4	6	6	14	5	13	5	7	3	-	-
Male	7	1	2	4	7	3	7	2	4	-	-	-
Female	4	3	4	2	7	2	6	3	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Washington												
All races, both sexes	2	8	6	9	12	27	43	36	36	10	5	-
Male	-	5	6	-	8	13	23	15	12	2	-	-
Female	2	3	-	9	4	14	20	21	24	8	5	-
White, both sexes	2	3	4	7	11	23	38	31	32	10	5	-
Male	-	2	4	-	7	11	22	11	12	2	-	-
Female	2	1	-	7	4	12	16	20	20	8	5	-
Black, both sexes	-	2	2	2	1	2	2	4	1	-	-	-
Male	-	-	2	-	1	1	-	3	-	-	-	-
Female	-	2	-	2	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	-	-	2	2	1	3	-	-	-
Male	-	3	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	3	-	-	-
West Virginia												
All races, both sexes	1	5	4	4	6	9	12	7	13	3	2	-
Male	-	1	4	3	2	5	2	4	4	-	-	-
Female	1	4	-	1	4	4	10	3	9	3	2	-
White, both sexes	1	4	3	4	5	8	11	6	12	3	2	-
Male	-	1	3	3	2	4	2	3	4	-	-	-
Female	1	3	-	1	3	4	9	3	8	3	2	-
Black, both sexes	-	1	1	-	1	1	1	1	1	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Wisconsin												
All races, both sexes	3	3	3	6	15	17	46	41	25	15	2	-
Male	2	2	2	3	9	9	18	14	6	2	-	-
Female	1	1	1	3	6	8	28	27	19	13	2	-
White, both sexes	2	3	2	5	11	16	45	38	25	15	2	-
Male	1	2	2	3	7	9	18	13	6	2	-	-
Female	1	1	-	2	4	7	27	25	19	13	2	-
Black, both sexes	1	-	1	1	2	1	1	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	1	2	3	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	3	1	-	-	-
White, both sexes	-	-	1	-	-	1	2	3	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
United States												
All races, both sexes	2,918	-	-	-	-	1	5	8	23	35	46	75
Male	1,290	-	-	-	-	-	3	5	19	23	34	38
Female	1,628	-	-	-	-	1	2	3	4	12	12	37
White, both sexes	1,985	-	-	-	-	-	-	3	10	13	14	27
Male	836	-	-	-	-	-	-	2	9	10	12	13
Female	1,149	-	-	-	-	-	-	1	1	3	2	14
Black, both sexes	824	-	-	-	-	1	3	4	12	21	30	47
Male	404	-	-	-	-	-	2	2	9	12	21	25
Female	420	-	-	-	-	1	1	2	3	9	9	22
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	92	-	-	-	-	-	2	1	-	1	1	-
Male	42	-	-	-	-	-	1	1	-	1	1	-
Female	50	-	-	-	-	-	1	-	-	-	-	-
Alabama												
All races, both sexes	48	-	-	-	-	-	-	-	1	1	-	2
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	34	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	29	-	-	-	-	-	-	-	1	1	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Arkansas												
All races, both sexes	28	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	417	-	-	-	-	-	2	2	2	6	7	6
Male	178	-	-	-	-	-	1	2	1	4	6	2
Female	239	-	-	-	-	-	1	-	1	2	1	4
White, both sexes	286	-	-	-	-	-	-	1	2	2	4	4
Male	121	-	-	-	-	-	-	1	1	1	3	1
Female	165	-	-	-	-	-	-	-	1	1	1	3
Black, both sexes	75	-	-	-	-	-	1	-	-	4	3	2
Male	35	-	-	-	-	-	1	-	-	3	3	1
Female	40	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	1	1	-	-	-	-
Male	20	-	-	-	-	-	-	1	-	-	-	-
Female	34	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Colorado												
All races, both sexes	24	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	22	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	32	-	-	-	-	-	-	-	1	-	-	2
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	26	-	-	-	-	-	-	-	1	-	-	2
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Florida												
All races, both sexes	187	-	-	-	-	-	-	-	3	2	2	4
Male	91	-	-	-	-	-	-	-	3	2	1	2
Female	96	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	141	-	-	-	-	-	-	-	1	1	1	2
Male	66	-	-	-	-	-	-	-	1	1	1	1
Female	75	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	45	-	-	-	-	-	-	-	1	1	1	2
Male	24	-	-	-	-	-	-	-	1	1	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	90	-	-	-	-	-	-	-	-	1	5	4
Male	45	-	-	-	-	-	-	-	-	1	4	4
Female	45	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	64	-	-	-	-	-	-	-	-	1	4	3
Male	31	-	-	-	-	-	-	-	-	1	3	3
Female	33	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Hawaii												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Illinois												
All races, both sexes	98	-	-	-	-	-	-	-	1	-	2	3
Male	46	-	-	-	-	-	-	-	-	-	2	1
Female	52	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	38	-	-	-	-	-	-	-	1	-	2	3
Male	22	-	-	-	-	-	-	-	-	-	2	1
Female	16	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	53	-	-	-	-	-	-	-	1	-	-	1
Male	21	-	-	-	-	-	-	-	1	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Iowa												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Kentucky												
All races, both sexes	38	-	-	-	-	-	-	-	1	-	-	-
Male	18	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	1	-	-	-
Male	18	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	60	-	-	-	-	-	-	-	-	1	-	4
Male	24	-	-	-	-	-	-	-	-	1	-	1
Female	36	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	29	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	4
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Maine												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	67	-	-	-	-	-	-	-	-	-	2	6
Male	30	-	-	-	-	-	-	-	-	-	1	3
Female	37	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	29	-	-	-	-	-	-	-	-	-	2	5
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	16	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Massachusetts												
All races, both sexes	38	-	-	-	-	-	-	-	-	1	1	-
Male	19	-	-	-	-	-	-	-	-	1	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	1	1	-
Male	16	-	-	-	-	-	-	-	-	1	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	127	-	-	-	-	-	-	2	1	3	4	3
Male	65	-	-	-	-	-	-	1	1	2	3	2
Female	62	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	88	-	-	-	-	-	-	-	1	1	2	3
Male	45	-	-	-	-	-	-	-	1	1	1	2
Female	43	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	38	-	-	-	-	-	-	2	-	2	2	-
Male	20	-	-	-	-	-	-	1	-	1	2	-
Female	18	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Minnesota												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	59	-	-	-	-	-	1	-	1	-	2	3
Male	27	-	-	-	-	-	-	-	1	-	2	3
Female	32	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	30	-	-	-	-	-	1	-	1	-	2	2
Male	16	-	-	-	-	-	-	-	1	-	2	2
Female	14	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Missouri												
All races, both sexes	54	-	-	-	-	-	-	-	-	2	-	4
Male	22	-	-	-	-	-	-	-	-	2	-	2
Female	32	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	2	-	4
Male	7	-	-	-	-	-	-	-	-	2	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Nebraska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
New Hampshire												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	95	-	-	-	-	-	-	1	2	-	2	2
Male	44	-	-	-	-	-	-	-	2	-	1	2
Female	51	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	69	-	-	-	-	-	-	1	1	-	1	-
Male	29	-	-	-	-	-	-	-	1	-	1	-
Female	40	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	1	-	1	2
Male	14	-	-	-	-	-	-	-	1	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
New Mexico												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	220	-	-	-	-	-	1	3	2	3	3	7
Male	102	-	-	-	-	-	1	2	2	2	2	3
Female	118	-	-	-	-	-	-	1	-	1	1	4
White, both sexes	129	-	-	-	-	-	-	1	1	3	-	5
Male	55	-	-	-	-	-	-	1	1	2	-	2
Female	74	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	81	-	-	-	-	-	-	2	1	-	3	2
Male	40	-	-	-	-	-	-	1	1	-	2	1
Female	41	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
North Carolina												
All races, both sexes	60	-	-	-	-	-	1	-	-	3	2	3
Male	25	-	-	-	-	-	1	-	-	1	1	2
Female	35	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	29	-	-	-	-	-	-	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	31	-	-	-	-	-	1	-	-	2	2	1
Male	16	-	-	-	-	-	1	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Ohio												
All races, both sexes	165	-	-	-	-	-	-	-	2	1	2	6
Male	77	-	-	-	-	-	-	-	2	-	1	5
Female	88	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	121	-	-	-	-	-	-	-	1	-	1	3
Male	51	-	-	-	-	-	-	-	1	-	1	2
Female	70	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	43	-	-	-	-	-	-	-	1	1	1	3
Male	26	-	-	-	-	-	-	-	1	-	-	3
Female	17	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	20	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Oregon												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	102	-	-	-	-	1	-	-	-	1	-	2
Male	50	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	1	-	-	-	1	-	1
White, both sexes	84	-	-	-	-	-	-	-	-	-	-	-
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	1	-	-	-	1	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	1	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Rhode Island												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	1	1
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	33	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
South Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	100	-	-	-	-	-	-	-	1	1	2	2
Male	44	-	-	-	-	-	-	-	-	-	2	2
Female	56	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	32	-	-	-	-	-	-	-	1	1	2	2
Male	14	-	-	-	-	-	-	-	-	-	2	2
Female	18	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Texas												
All races, both sexes	192	-	-	-	-	-	-	-	3	4	5	3
Male	83	-	-	-	-	-	-	-	2	4	4	1
Female	109	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	130	-	-	-	-	-	-	-	1	2	1	1
Male	55	-	-	-	-	-	-	-	1	2	1	1
Female	75	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	55	-	-	-	-	-	-	-	2	2	3	2
Male	24	-	-	-	-	-	-	-	1	2	2	-
Female	31	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Vermont												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Washington												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	71	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Wisconsin												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
United States												
All races, both sexes	104	123	136	150	209	317	489	554	439	178	26	-
Male	63	81	81	76	101	148	217	205	138	51	7	-
Female	41	42	55	74	108	169	272	349	301	127	19	-
White, both sexes	36	54	57	77	113	200	387	455	368	151	20	-
Male	22	35	35	35	53	91	177	170	121	45	6	-
Female	14	19	22	42	60	109	210	285	247	106	14	-
Black, both sexes	63	66	74	65	88	100	89	83	48	24	6	-
Male	38	44	43	35	44	51	34	28	10	5	1	-
Female	25	22	31	30	44	49	55	55	38	19	5	-
American Indian 1/, both sexes	1	-	2	1	2	1	2	2	3	-	-	-
Male	-	-	1	1	2	-	1	1	1	-	-	-
Female	1	-	1	-	-	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	3	7	6	16	11	14	20	3	-	-
Male	3	2	2	5	2	6	5	6	6	1	-	-
Female	1	1	1	2	4	10	6	8	14	2	-	-
Alabama												
All races, both sexes	-	2	2	3	3	11	11	4	3	3	2	-
Male	-	2	1	-	-	3	3	1	1	1	1	-
Female	-	-	1	3	3	8	8	3	2	2	1	-
White, both sexes	-	-	1	1	1	5	4	1	2	2	2	-
Male	-	-	1	-	-	1	-	-	1	1	1	-
Female	-	-	-	1	1	4	4	1	1	1	1	-
Black, both sexes	-	2	1	2	2	6	7	3	1	1	-	-
Male	-	2	-	-	-	2	3	1	-	-	-	-
Female	-	-	1	2	2	4	4	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	2	-	1	-	5	9	6	1	1	-	-
Male	-	2	-	1	-	2	5	2	1	1	-	-
Female	-	-	-	-	-	3	4	4	-	-	-	-
White, both sexes	-	2	-	1	-	5	9	6	1	1	-	-
Male	-	2	-	1	-	2	5	2	1	1	-	-
Female	-	-	-	-	-	3	4	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Arkansas												
All races, both sexes	3	1	1	1	6	3	5	2	1	4	-	-
Male	2	-	1	1	2	2	4	1	-	-	-	-
Female	1	1	-	-	4	1	1	1	1	4	-	-
White, both sexes	1	1	1	1	2	2	4	2	1	3	-	-
Male	-	-	1	1	-	1	4	1	-	-	-	-
Female	1	1	-	-	2	1	-	1	1	3	-	-
Black, both sexes	2	-	-	-	4	1	1	-	-	1	-	-
Male	2	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	22	21	30	26	41	81	72	70	19	2	-
Male	4	13	10	13	11	17	40	23	25	5	1	-
Female	4	9	11	17	15	24	41	49	45	14	1	-
White, both sexes	3	14	12	21	16	26	64	55	45	15	2	-
Male	1	7	6	7	7	13	33	16	18	5	1	-
Female	2	7	6	14	9	13	31	39	27	10	1	-
Black, both sexes	2	7	8	5	5	7	10	9	9	3	-	-
Male	1	6	4	3	3	2	3	3	2	-	-	-
Female	1	1	4	2	2	5	7	6	7	3	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	3	5	8	7	7	16	1	-	-
Male	2	-	-	2	1	2	4	3	5	-	-	-
Female	1	1	1	1	4	6	3	4	11	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Colorado												
All races, both sexes	1	1	1	1	2	1	2	7	4	2	-	-
Male	-	1	-	-	1	-	-	1	1	-	-	-
Female	1	-	1	1	1	1	2	6	3	2	-	-
White, both sexes	-	1	1	1	1	1	1	6	4	2	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	1	1	1	1	5	3	2	-	-
Black, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	2	-	3	4	6	4	1	-	-
Male	1	-	-	1	-	2	2	1	-	1	-	-
Female	-	-	-	1	-	1	2	5	4	-	-	-
White, both sexes	-	-	-	2	-	2	4	6	4	1	-	-
Male	-	-	-	1	-	1	2	1	-	1	-	-
Female	-	-	-	1	-	1	2	5	4	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Delaware												
All races, both sexes	1	-	2	1	1	-	-	2	1	-	-	-
Male	1	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	2	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	1	-	1	1	1	-	-	-	1	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	3	1	1	5	4	4	4	2	1	-	-
Male	1	-	-	1	2	3	2	1	2	-	-	-
Female	3	3	1	-	3	1	2	3	-	1	-	-
White, both sexes	-	1	-	-	1	-	1	1	2	-	-	-
Male	-	-	-	-	-	-	1	-	2	-	-	-
Female	-	1	-	-	1	-	-	1	-	-	-	-
Black, both sexes	4	2	1	1	4	4	3	3	-	1	-	-
Male	1	-	-	1	2	3	1	1	-	-	-	-
Female	3	2	1	-	2	1	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Florida												
All races, both sexes	6	5	10	6	14	25	42	25	32	11	-	-
Male	4	5	7	2	7	13	17	11	12	5	-	-
Female	2	-	3	4	7	12	25	14	20	6	-	-
White, both sexes	2	4	6	3	9	19	32	22	29	10	-	-
Male	2	4	3	-	5	8	15	9	12	4	-	-
Female	-	-	3	3	4	11	17	13	17	6	-	-
Black, both sexes	4	1	4	3	5	6	10	3	3	1	-	-
Male	2	1	4	2	2	5	2	2	-	1	-	-
Female	2	-	-	1	3	1	8	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	7	4	8	7	13	7	10	13	7	4	-	-
Male	4	2	6	3	8	4	4	2	3	-	-	-
Female	3	2	2	4	5	3	6	11	4	4	-	-
White, both sexes	-	1	1	1	3	3	5	3	4	2	-	-
Male	-	-	1	-	3	2	2	1	2	-	-	-
Female	-	1	-	1	-	1	3	2	2	2	-	-
Black, both sexes	7	2	7	6	10	4	5	10	3	2	-	-
Male	4	1	5	3	5	2	2	1	1	-	-	-
Female	3	1	2	3	5	2	3	9	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Hawaii												
All races, both sexes	-	-	-	-	-	1	2	3	1	1	-	-
Male	-	-	-	-	-	-	1	2	-	1	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	3	1	1	-	-
Male	-	-	-	-	-	-	1	2	-	1	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Idaho												
All races, both sexes	-	-	-	-	2	-	2	-	3	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	2	-	-	-	2	-	-	-
White, both sexes	-	-	-	-	2	-	2	-	3	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	2	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Illinois												
All races, both sexes	4	2	8	9	9	14	15	12	13	5	1	-
Male	2	2	6	5	5	7	7	5	3	1	-	-
Female	2	-	2	4	4	7	8	7	10	4	1	-
White, both sexes	1	1	2	2	6	10	14	7	11	5	1	-
Male	-	1	1	1	3	5	7	2	3	1	-	-
Female	1	-	1	1	3	5	7	5	8	4	1	-
Black, both sexes	3	1	6	7	3	4	1	5	2	-	-	-
Male	2	1	5	4	2	2	-	3	-	-	-	-
Female	1	-	1	3	1	2	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	1	2	1	11	12	8	10	5	-	-
Male	1	-	1	1	-	4	6	4	3	-	-	-
Female	-	-	-	1	1	7	6	4	7	5	-	-
White, both sexes	-	-	-	2	1	5	12	7	7	4	-	-
Male	-	-	-	1	-	2	6	3	2	-	-	-
Female	-	-	-	1	1	3	6	4	5	4	-	-
Black, both sexes	1	-	1	-	-	6	-	1	3	1	-	-
Male	1	-	1	-	-	2	-	1	1	-	-	-
Female	-	-	-	-	-	4	-	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Iowa												
All races, both sexes	-	1	-	2	3	5	8	10	12	5	-	-
Male	-	-	-	1	1	1	3	7	1	1	-	-
Female	-	1	-	1	2	4	5	3	11	4	-	-
White, both sexes	-	1	-	2	2	5	7	10	12	4	-	-
Male	-	-	-	1	1	1	3	7	1	1	-	-
Female	-	1	-	1	1	4	4	3	11	3	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	2	-	6	6	2	2	2	-
Male	-	-	-	-	1	-	3	4	1	1	-	-
Female	-	-	-	-	1	-	3	2	1	1	2	-
White, both sexes	-	-	-	-	1	-	4	5	2	2	2	-
Male	-	-	-	-	-	-	2	3	1	1	-	-
Female	-	-	-	-	1	-	2	2	1	1	2	-
Black, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Kentucky												
All races, both sexes	1	2	1	2	3	3	9	6	5	4	1	-
Male	-	2	-	1	2	2	6	1	2	1	-	-
Female	1	-	1	1	1	1	3	5	3	3	1	-
White, both sexes	1	2	1	2	3	2	8	5	5	4	1	-
Male	-	2	-	1	2	2	6	1	2	1	-	-
Female	1	-	1	1	1	-	2	4	3	3	1	-
Black, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	1	3	4	4	10	4	13	9	2	-	-
Male	3	1	2	2	2	5	3	2	1	1	-	-
Female	2	-	1	2	2	5	1	11	8	1	-	-
White, both sexes	2	-	2	-	2	5	3	7	6	1	-	-
Male	1	-	2	-	1	3	2	1	-	-	-	-
Female	1	-	-	-	1	2	1	6	6	1	-	-
Black, both sexes	3	1	1	4	2	5	1	6	3	1	-	-
Male	2	1	-	2	1	2	1	1	1	1	-	-
Female	1	-	1	2	1	3	-	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Maine												
All races, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	4	5	3	2	5	11	14	7	2	2	-
Male	3	3	2	3	-	-	6	6	3	-	-	-
Female	1	1	3	-	2	5	5	8	4	2	2	-
White, both sexes	-	1	3	-	1	3	9	10	5	2	1	-
Male	-	1	1	-	-	-	5	5	2	-	-	-
Female	-	-	2	-	1	3	4	5	3	2	1	-
Black, both sexes	4	2	2	3	1	2	2	4	1	-	1	-
Male	3	1	1	3	-	-	1	1	-	-	-	-
Female	1	1	1	-	1	2	1	3	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Massachusetts												
All races, both sexes	-	1	2	-	2	3	8	8	7	4	1	-
Male	-	1	1	-	2	2	4	5	1	1	-	-
Female	-	-	1	-	-	1	4	3	6	3	1	-
White, both sexes	-	1	2	-	1	1	6	8	7	4	1	-
Male	-	1	1	-	1	1	3	5	1	1	-	-
Female	-	-	1	-	-	-	3	3	6	3	1	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	5	9	9	10	10	12	15	24	9	8	3	-
Male	4	5	4	3	6	5	7	12	4	4	2	-
Female	1	4	5	7	4	7	8	12	5	4	1	-
White, both sexes	3	4	-	6	9	10	12	21	7	8	1	-
Male	2	2	-	1	5	4	6	11	4	4	1	-
Female	1	2	-	5	4	6	6	10	3	4	-	-
Black, both sexes	2	5	8	4	1	2	3	3	2	-	2	-
Male	2	3	4	2	1	1	1	1	-	-	1	-
Female	-	2	4	2	-	1	2	2	2	-	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Minnesota												
All races, both sexes	-	1	2	1	1	3	3	8	6	4	-	-
Male	-	-	2	1	1	1	2	4	3	-	-	-
Female	-	1	-	-	-	2	1	4	3	4	-	-
White, both sexes	-	1	-	1	1	2	3	8	6	3	-	-
Male	-	-	-	1	1	1	2	4	3	-	-	-
Female	-	1	-	-	-	1	1	4	3	3	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Mississippi												
All races, both sexes	1	1	3	2	7	12	4	13	6	2	1	-
Male	1	1	1	-	2	7	2	6	1	-	-	-
Female	-	-	2	2	5	5	2	7	5	2	1	-
White, both sexes	-	-	3	1	3	6	3	6	4	2	-	-
Male	-	-	1	-	1	3	2	2	1	-	-	-
Female	-	-	2	1	2	3	1	4	3	2	-	-
Black, both sexes	1	1	-	1	4	6	1	7	2	-	1	-
Male	1	1	-	-	1	4	-	4	-	-	-	-
Female	-	-	-	1	3	2	1	3	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Missouri												
All races, both sexes	2	-	-	1	4	2	8	18	12	1	-	-
Male	-	-	-	-	1	1	4	6	5	1	-	-
Female	2	-	-	1	3	1	4	12	7	-	-	-
White, both sexes	2	-	-	1	1	2	7	16	9	1	-	-
Male	-	-	-	-	-	1	3	6	3	1	-	-
Female	2	-	-	1	1	1	4	10	6	-	-	-
Black, both sexes	-	-	-	-	3	-	1	2	2	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	2	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	2	1	2	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	2	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Nebraska												
All races, both sexes	-	-	1	-	1	3	2	1	-	2	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	1	-	-	2	2	1	-	1	-	-
White, both sexes	-	-	1	-	1	2	2	1	-	2	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	1	-	-	1	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	-	-	-	1	2	2	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	1	-	-	-
Female	2	-	-	-	1	2	2	-	-	-	-	-
White, both sexes	3	-	-	-	1	2	2	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	1	-	-	-
Female	2	-	-	-	1	2	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
New Hampshire												
All races, both sexes	-	-	-	1	-	-	2	3	3	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	3	-	-	-
White, both sexes	-	-	-	1	-	-	2	3	3	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	5	3	5	4	7	17	20	15	8	1	-
Male	2	4	2	3	2	4	4	11	6	1	-	-
Female	1	1	1	2	2	3	13	9	9	7	1	-
White, both sexes	1	2	-	1	2	4	16	18	15	7	-	-
Male	-	1	-	1	1	3	4	10	6	1	-	-
Female	1	1	-	-	1	1	12	8	9	6	-	-
Black, both sexes	2	3	3	3	2	3	1	2	-	1	1	-
Male	2	3	2	1	1	1	-	1	-	-	-	-
Female	-	-	1	2	1	2	1	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
New Mexico												
All races, both sexes	-	-	-	1	1	-	1	-	3	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	3	1	-	-
White, both sexes	-	-	-	1	1	-	1	-	3	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	10	15	17	9	25	30	26	33	21	12	3	-
Male	7	8	13	4	17	15	9	7	4	4	2	-
Female	3	7	4	5	8	15	17	26	17	8	1	-
White, both sexes	1	4	6	2	10	17	19	29	17	11	3	-
Male	1	2	6	1	7	8	7	7	4	4	2	-
Female	-	2	-	1	3	9	12	22	13	7	1	-
Black, both sexes	9	11	9	6	14	10	6	3	4	1	-	-
Male	6	6	5	2	9	5	2	-	-	-	-	-
Female	3	5	4	4	5	5	4	3	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	1	3	1	1	-	-	-	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
North Carolina												
All races, both sexes	3	2	2	4	7	6	6	9	7	5	-	-
Male	3	1	-	4	4	2	2	3	1	-	-	-
Female	-	1	2	-	3	4	4	6	6	5	-	-
White, both sexes	-	-	-	1	4	3	4	5	6	3	-	-
Male	-	-	-	1	3	1	1	1	-	-	-	-
Female	-	-	-	-	1	2	3	4	6	3	-	-
Black, both sexes	3	2	2	3	3	3	2	4	1	2	-	-
Male	3	1	-	3	1	1	1	2	1	-	-	-
Female	-	1	2	-	2	2	1	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	1	4	1	-	-
Male	-	-	-	-	-	-	-	1	3	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	4	1	-	-
Male	-	-	-	-	-	-	-	1	3	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Ohio												
All races, both sexes	3	6	5	8	7	13	37	42	21	10	2	-
Male	2	4	5	6	4	10	11	16	8	3	-	-
Female	1	2	-	2	3	3	26	26	13	7	2	-
White, both sexes	1	1	3	4	3	9	28	39	18	8	2	-
Male	1	1	3	3	2	6	7	15	7	2	-	-
Female	-	-	-	1	1	3	21	24	11	6	2	-
Black, both sexes	2	5	2	4	4	4	9	2	3	2	-	-
Male	1	3	2	3	2	4	4	1	1	1	-	-
Female	1	2	-	1	2	-	5	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	-	-	1	3	2	3	3	5	-	-
Male	-	-	-	-	1	2	2	1	-	1	-	-
Female	-	2	-	-	-	1	-	2	3	4	-	-
White, both sexes	-	-	-	-	1	3	-	3	1	5	-	-
Male	-	-	-	-	1	2	-	1	-	1	-	-
Female	-	-	-	-	-	1	-	2	1	4	-	-
Black, both sexes	-	2	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Oregon												
All races, both sexes	-	-	-	-	1	5	7	12	14	4	-	-
Male	-	-	-	-	-	-	6	5	2	1	-	-
Female	-	-	-	-	1	5	1	7	12	3	-	-
White, both sexes	-	-	-	-	1	5	7	11	13	4	-	-
Male	-	-	-	-	-	-	6	5	2	1	-	-
Female	-	-	-	-	1	5	1	6	11	3	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	-	6	3	1	5	8	20	25	24	6	-	-
Male	-	6	3	1	2	5	8	11	9	4	-	-
Female	-	-	-	-	3	3	12	14	15	2	-	-
White, both sexes	-	3	2	1	4	6	16	24	23	5	-	-
Male	-	3	2	1	1	3	6	11	8	4	-	-
Female	-	-	-	-	3	3	10	13	15	1	-	-
Black, both sexes	-	3	1	-	1	2	4	1	1	1	-	-
Male	-	3	1	-	1	2	2	-	1	-	-	-
Female	-	-	-	-	-	-	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Rhode Island												
All races, both sexes	-	-	-	-	-	3	1	2	1	1	-	-
Male	-	-	-	-	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	2	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	3	1	2	1	1	-	-
Male	-	-	-	-	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	5	3	2	4	6	10	10	5	3	-	-
Male	-	4	1	1	3	3	7	2	2	1	-	-
Female	-	1	2	1	1	3	3	8	3	2	-	-
White, both sexes	-	-	-	-	-	1	5	6	4	1	-	-
Male	-	-	-	-	-	-	3	2	2	-	-	-
Female	-	-	-	-	-	1	2	4	2	1	-	-
Black, both sexes	-	5	3	2	4	5	5	4	1	2	-	-
Male	-	4	1	1	3	3	4	-	-	1	-	-
Female	-	1	2	1	1	2	1	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	2	6	-	-	-	-
Male	-	-	-	-	-	-	-	4	-	-	-	-
Female	-	-	-	-	-	-	2	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	5	-	-	-	-
Male	-	-	-	-	-	-	-	3	-	-	-	-
Female	-	-	-	-	-	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	3	7	8	6	12	19	16	14	4	-	-
Male	2	1	5	4	2	5	8	6	5	2	-	-
Female	3	2	2	4	4	7	11	10	9	2	-	-
White, both sexes	3	2	3	3	2	6	16	16	13	4	-	-
Male	2	-	2	2	1	2	8	6	5	2	-	-
Female	1	2	1	1	1	4	8	10	8	2	-	-
Black, both sexes	2	1	4	5	4	6	3	-	1	-	-	-
Male	-	1	3	2	1	3	-	-	-	-	-	-
Female	2	-	1	3	3	3	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Texas												
All races, both sexes	15	11	8	14	17	19	26	28	30	8	1	-
Male	8	8	3	7	4	10	10	8	11	2	1	-
Female	7	3	5	7	13	9	16	20	19	6	-	-
White, both sexes	6	5	2	8	11	12	22	26	26	6	1	-
Male	5	5	1	4	2	7	8	6	10	1	1	-
Female	1	-	1	4	9	5	14	20	16	5	-	-
Black, both sexes	8	6	6	4	6	5	3	2	4	2	-	-
Male	3	3	2	2	2	1	2	2	1	1	-	-
Female	5	3	4	2	4	4	1	-	3	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	2	1	-	-	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	1	3	2	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	3	2	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	3	5	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	3	3	-	-	-
White, both sexes	-	-	-	-	-	-	-	3	5	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	3	2	1	5	5	12	14	7	3	-	-
Male	2	2	1	-	3	1	7	3	2	1	-	-
Female	1	1	1	1	2	4	5	11	5	2	-	-
White, both sexes	1	-	-	1	2	2	8	10	6	3	-	-
Male	1	-	-	-	1	1	6	2	2	1	-	-
Female	-	-	-	1	1	1	2	8	4	2	-	-
Black, both sexes	1	3	2	-	3	3	4	4	1	-	-	-
Male	-	2	1	-	2	-	1	1	-	-	-	-
Female	1	1	1	-	1	3	3	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Washington												
All races, both sexes	2	2	1	5	-	8	11	20	17	9	1	-
Male	2	2	-	4	-	4	3	7	4	2	-	-
Female	-	-	1	1	-	4	8	13	13	7	1	-
White, both sexes	2	1	1	4	-	7	11	19	16	9	1	-
Male	2	1	-	3	-	3	3	6	4	2	-	-
Female	-	-	1	1	-	4	8	13	12	7	1	-
Black, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	1	1	1	-	1	3	1	3	4	2	1	-
Male	1	1	-	-	1	1	1	-	-	1	-	-
Female	-	-	1	-	-	2	-	3	4	1	1	-
White, both sexes	1	1	1	-	1	3	1	3	4	2	-	-
Male	1	1	-	-	1	1	1	-	-	1	-	-
Female	-	-	1	-	-	2	-	3	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Wisconsin												
All races, both sexes	2	-	3	2	3	3	5	11	8	2	2	-
Male	2	-	2	2	1	3	1	6	2	1	-	-
Female	-	-	1	-	2	-	4	5	6	1	2	-
White, both sexes	2	-	2	2	3	1	5	9	8	2	2	-
Male	2	-	2	2	1	1	1	6	2	1	-	-
Female	-	-	-	-	2	-	4	3	6	1	2	-
Black, both sexes	-	-	1	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
United States												
All races, both sexes	425,425	22	7	3	10	38	100	286	768	1,872	4,827	9,627
Male	224,510	15	5	2	5	30	76	224	605	1,410	3,665	7,249
Female	200,915	7	2	1	5	8	24	62	163	462	1,162	2,378
White, both sexes	371,445	15	5	3	7	29	77	207	563	1,399	3,810	7,501
Male	196,396	12	3	2	5	26	60	168	452	1,083	2,982	5,809
Female	175,049	3	2	1	2	3	17	39	111	316	828	1,692
Black, both sexes	44,530	6	2	-	2	9	23	68	166	411	867	1,831
Male	22,713	2	2	-	-	4	16	47	119	272	561	1,204
Female	21,817	4	-	-	2	5	7	21	47	139	306	627
American Indian 1/, both sexes	1,880	-	-	-	-	-	-	3	11	12	55	88
Male	1,082	-	-	-	-	-	-	2	9	11	38	68
Female	798	-	-	-	-	-	-	1	2	1	17	20
Asian or Pacific Islander 2/, both sexes	7,570	1	-	-	1	-	-	8	28	50	95	207
Male	4,319	1	-	-	-	-	-	7	25	44	84	168
Female	3,251	-	-	-	1	-	-	1	3	6	11	39
Alabama												
All races, both sexes	6,038	1	-	-	-	1	6	11	23	48	110	207
Male	3,206	1	-	-	-	1	5	8	19	32	74	146
Female	2,832	-	-	-	-	-	1	3	4	16	36	61
White, both sexes	4,815	1	-	-	-	1	4	8	12	34	82	143
Male	2,583	1	-	-	-	1	3	7	9	23	60	110
Female	2,232	-	-	-	-	-	1	1	3	11	22	33
Black, both sexes	1,204	-	-	-	-	-	2	3	11	14	28	61
Male	613	-	-	-	-	-	2	1	10	9	14	34
Female	591	-	-	-	-	-	-	2	1	5	14	27
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Alaska												
All races, both sexes	351	-	-	-	-	1	-	-	-	3	8	13
Male	234	-	-	-	-	1	-	-	-	3	7	11
Female	117	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	285	-	-	-	-	1	-	-	-	2	5	10
Male	191	-	-	-	-	1	-	-	-	2	5	9
Female	94	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	52	-	-	-	-	-	-	-	-	1	1	3
Male	35	-	-	-	-	-	-	-	-	1	1	2
Female	17	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	7,806	-	-	1	-	-	3	4	14	29	73	143
Male	4,331	-	-	-	-	-	3	3	12	23	60	111
Female	3,475	-	-	1	-	-	-	1	2	6	13	32
White, both sexes	7,380	-	-	1	-	-	3	2	11	25	62	128
Male	4,069	-	-	-	-	-	3	2	9	19	51	97
Female	3,311	-	-	1	-	-	-	-	2	6	11	31
Black, both sexes	211	-	-	-	-	-	-	1	2	2	2	9
Male	137	-	-	-	-	-	-	-	2	2	2	9
Female	74	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	158	-	-	-	-	-	-	1	1	2	8	3
Male	96	-	-	-	-	-	-	1	1	2	6	2
Female	62	-	-	-	-	-	-	-	-	-	2	1
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	-	1	3
Male	29	-	-	-	-	-	-	-	-	-	1	3
Female	28	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Arkansas												
All races, both sexes	5,100	1	-	1	-	3	2	12	23	44	87	171
Male	2,709	1	-	1	-	2	2	9	15	29	63	116
Female	2,391	-	-	-	-	1	-	3	8	15	24	55
White, both sexes	4,411	1	-	1	-	2	2	8	20	33	69	125
Male	2,364	1	-	1	-	2	2	6	14	24	50	87
Female	2,047	-	-	-	-	-	-	2	6	9	19	38
Black, both sexes	664	-	-	-	-	1	-	4	3	11	18	44
Male	329	-	-	-	-	-	-	3	1	5	13	27
Female	335	-	-	-	-	1	-	1	2	6	5	17
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	46,584	-	1	1	2	3	4	19	52	150	392	892
Male	24,619	-	1	1	1	2	3	16	40	118	308	690
Female	21,965	-	-	-	1	1	1	3	12	32	84	202
White, both sexes	39,222	-	1	1	1	3	3	15	36	106	299	693
Male	20,703	-	1	1	1	2	3	14	26	83	236	547
Female	18,519	-	-	-	-	1	-	1	10	23	63	146
Black, both sexes	3,593	-	-	-	-	-	1	3	8	25	55	120
Male	1,846	-	-	-	-	-	-	1	7	17	36	79
Female	1,747	-	-	-	-	-	1	2	1	8	19	41
American Indian 1/, both sexes	210	-	-	-	-	-	-	-	-	2	4	5
Male	113	-	-	-	-	-	-	-	-	2	4	3
Female	97	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	3,559	-	-	-	1	-	-	1	8	17	34	74
Male	1,957	-	-	-	-	-	-	1	7	16	32	61
Female	1,602	-	-	-	1	-	-	-	1	1	2	13

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Colorado												
All races, both sexes	3,922	1	1	-	-	1	-	5	10	19	53	107
Male	2,220	1	1	-	-	1	-	4	10	16	41	89
Female	1,702	-	-	-	-	-	-	1	-	3	12	18
White, both sexes	3,786	1	1	-	-	-	-	4	9	19	52	96
Male	2,138	1	1	-	-	-	-	3	9	16	40	81
Female	1,648	-	-	-	-	-	-	1	-	3	12	15
Black, both sexes	93	-	-	-	-	1	-	1	1	-	-	10
Male	62	-	-	-	-	1	-	1	1	-	-	7
Female	31	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4,630	-	-	-	-	-	2	3	5	20	37	73
Male	2,293	-	-	-	-	-	2	3	4	17	29	60
Female	2,337	-	-	-	-	-	-	-	1	3	8	13
White, both sexes	4,341	-	-	-	-	-	2	3	4	16	31	53
Male	2,142	-	-	-	-	-	2	3	4	13	25	43
Female	2,199	-	-	-	-	-	-	-	-	3	6	10
Black, both sexes	254	-	-	-	-	-	-	-	1	4	6	19
Male	132	-	-	-	-	-	-	-	-	4	4	16
Female	122	-	-	-	-	-	-	-	1	-	2	3
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Delaware												
All races, both sexes	1,305	-	-	-	-	-	-	-	2	3	14	44
Male	682	-	-	-	-	-	-	-	1	2	11	33
Female	623	-	-	-	-	-	-	-	1	1	3	11
White, both sexes	1,112	-	-	-	-	-	-	-	-	3	12	30
Male	577	-	-	-	-	-	-	-	-	2	9	23
Female	535	-	-	-	-	-	-	-	-	1	3	7
Black, both sexes	185	-	-	-	-	-	-	-	2	-	2	14
Male	102	-	-	-	-	-	-	-	1	-	2	10
Female	83	-	-	-	-	-	-	-	1	-	-	4
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1,144	-	-	-	-	-	1	1	1	3	9	27
Male	569	-	-	-	-	-	1	1	1	2	6	19
Female	575	-	-	-	-	-	-	-	-	1	3	8
White, both sexes	278	-	-	-	-	-	-	-	-	-	3	6
Male	158	-	-	-	-	-	-	-	-	-	2	6
Female	120	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	856	-	-	-	-	-	1	1	1	3	6	21
Male	404	-	-	-	-	-	1	1	1	2	4	13
Female	452	-	-	-	-	-	-	-	-	1	2	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Florida												
All races, both sexes	32,868	1	-	-	1	2	4	11	33	102	306	563
Male	17,673	1	-	-	1	1	2	6	28	74	238	433
Female	15,195	-	-	-	-	1	2	5	5	28	68	130
White, both sexes	29,949	1	-	-	1	1	3	8	22	79	239	452
Male	16,203	1	-	-	1	1	2	5	19	58	191	365
Female	13,746	-	-	-	-	-	1	3	3	21	48	87
Black, both sexes	2,747	-	-	-	-	1	1	3	9	23	63	102
Male	1,371	-	-	-	-	-	-	1	7	16	43	61
Female	1,376	-	-	-	-	1	1	2	2	7	20	41
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	1	-	-	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	147	-	-	-	-	-	-	-	1	-	4	9
Male	88	-	-	-	-	-	-	-	1	-	4	7
Female	59	-	-	-	-	-	-	-	-	-	-	2
Georgia												
All races, both sexes	8,371	-	-	-	1	1	4	11	25	59	153	303
Male	4,646	-	-	-	-	1	4	7	16	44	120	235
Female	3,725	-	-	-	1	-	-	4	9	15	33	68
White, both sexes	6,327	-	-	-	-	1	3	7	15	27	104	184
Male	3,597	-	-	-	-	1	3	4	12	20	85	147
Female	2,730	-	-	-	-	-	-	3	3	7	19	37
Black, both sexes	1,964	-	-	-	1	-	1	4	10	30	46	115
Male	1,005	-	-	-	-	-	1	3	4	22	32	86
Female	959	-	-	-	1	-	-	1	6	8	14	29
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	66	-	-	-	-	-	-	-	-	2	3	3
Male	33	-	-	-	-	-	-	-	-	2	3	1
Female	33	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Hawaii												
All races, both sexes	1,298	-	-	-	-	-	1	1	5	8	12	29
Male	827	-	-	-	-	-	1	1	5	5	7	23
Female	471	-	-	-	-	-	-	-	-	3	5	6
White, both sexes	371	-	-	-	-	-	1	-	-	1	2	6
Male	236	-	-	-	-	-	1	-	-	1	1	5
Female	135	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	9	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	1	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	917	-	-	-	-	-	-	1	4	6	10	22
Male	584	-	-	-	-	-	-	1	4	3	6	17
Female	333	-	-	-	-	-	-	-	-	3	4	5
Idaho												
All races, both sexes	1,565	-	-	-	-	1	-	1	2	7	21	41
Male	929	-	-	-	-	1	-	1	2	5	16	29
Female	636	-	-	-	-	-	-	-	-	2	5	12
White, both sexes	1,540	-	-	-	-	1	-	1	2	7	20	41
Male	915	-	-	-	-	1	-	1	2	5	16	29
Female	625	-	-	-	-	-	-	-	-	2	4	12
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Illinois												
All races, both sexes	17,747	3	-	-	-	2	3	12	32	66	197	412
Male	9,206	1	-	-	-	1	-	8	29	49	142	304
Female	8,541	2	-	-	-	1	3	4	3	17	55	108
White, both sexes	15,092	-	-	-	-	2	2	9	22	46	152	306
Male	7,769	-	-	-	-	1	-	5	19	37	118	238
Female	7,323	-	-	-	-	1	2	4	3	9	34	68
Black, both sexes	2,430	3	-	-	-	-	1	3	10	20	42	99
Male	1,312	1	-	-	-	-	-	3	10	12	22	60
Female	1,118	2	-	-	-	-	1	-	-	8	20	39
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	212	-	-	-	-	-	-	-	-	-	3	7
Male	118	-	-	-	-	-	-	-	-	-	2	6
Female	94	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	9,210	-	-	-	-	1	2	8	29	52	147	250
Male	4,960	-	-	-	-	-	1	8	19	35	117	189
Female	4,250	-	-	-	-	1	1	-	10	17	30	61
White, both sexes	8,514	-	-	-	-	-	2	8	22	46	130	224
Male	4,572	-	-	-	-	-	1	8	16	32	104	174
Female	3,942	-	-	-	-	-	1	-	6	14	26	50
Black, both sexes	668	-	-	-	-	1	-	-	7	6	16	24
Male	368	-	-	-	-	-	-	-	3	3	12	13
Female	300	-	-	-	-	1	-	-	4	3	4	11
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Iowa												
All races, both sexes	5,469	-	1	-	-	-	1	6	8	13	48	100
Male	2,811	-	1	-	-	-	1	6	6	11	34	79
Female	2,658	-	-	-	-	-	-	-	2	2	14	21
White, both sexes	5,384	-	-	-	-	-	1	5	8	9	46	94
Male	2,760	-	-	-	-	-	1	5	6	9	32	76
Female	2,624	-	-	-	-	-	-	-	2	-	14	18
Black, both sexes	69	-	1	-	-	-	-	-	-	3	2	4
Male	40	-	1	-	-	-	-	-	-	1	2	2
Female	29	-	-	-	-	-	-	-	-	2	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	1	-	1	-	2
Male	11	-	-	-	-	-	-	1	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	3,565	-	-	-	-	1	1	1	8	6	36	83
Male	1,939	-	-	-	-	1	-	1	7	4	25	59
Female	1,626	-	-	-	-	-	1	-	1	2	11	24
White, both sexes	3,371	-	-	-	-	-	1	1	7	6	33	71
Male	1,830	-	-	-	-	-	-	1	7	4	24	50
Female	1,541	-	-	-	-	-	1	-	-	2	9	21
Black, both sexes	144	-	-	-	-	1	-	-	1	-	2	7
Male	78	-	-	-	-	1	-	-	-	-	-	5
Female	66	-	-	-	-	-	-	-	1	-	2	2
American Indian 1/, both sexes	34	-	-	-	-	-	-	-	-	-	1	4
Male	24	-	-	-	-	-	-	-	-	-	1	3
Female	10	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Kentucky												
All races, both sexes	6,530	1	-	-	-	2	3	10	23	61	129	210
Male	3,526	-	-	-	-	1	1	6	20	44	103	169
Female	3,004	1	-	-	-	1	2	4	3	17	26	41
White, both sexes	6,120	1	-	-	-	2	3	8	18	56	108	195
Male	3,297	-	-	-	-	1	1	5	15	41	89	158
Female	2,823	1	-	-	-	1	2	3	3	15	19	37
Black, both sexes	393	-	-	-	-	-	-	2	4	5	19	14
Male	220	-	-	-	-	-	-	1	4	3	12	10
Female	173	-	-	-	-	-	-	1	-	2	7	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	2	1
Male	9	-	-	-	-	-	-	-	1	-	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5,919	-	-	-	-	-	2	7	25	40	130	219
Male	3,318	-	-	-	-	-	2	7	14	30	92	159
Female	2,601	-	-	-	-	-	-	-	11	10	38	60
White, both sexes	4,241	-	-	-	-	-	1	4	15	20	81	130
Male	2,425	-	-	-	-	-	1	4	8	16	60	99
Female	1,816	-	-	-	-	-	-	-	7	4	21	31
Black, both sexes	1,643	-	-	-	-	-	1	3	10	20	48	88
Male	871	-	-	-	-	-	1	3	6	14	31	59
Female	772	-	-	-	-	-	-	-	4	6	17	29
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Maine												
All races, both sexes	1,816	-	-	-	-	-	-	2	6	6	18	44
Male	998	-	-	-	-	-	-	2	5	4	15	36
Female	818	-	-	-	-	-	-	-	1	2	3	8
White, both sexes	1,805	-	-	-	-	-	-	2	5	6	16	43
Male	989	-	-	-	-	-	-	2	4	4	14	35
Female	816	-	-	-	-	-	-	-	1	2	2	8
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	2	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7,744	-	-	-	-	-	-	6	10	34	73	172
Male	3,876	-	-	-	-	-	-	4	6	30	54	124
Female	3,868	-	-	-	-	-	-	2	4	4	19	48
White, both sexes	5,678	-	-	-	-	-	-	4	4	17	44	104
Male	2,852	-	-	-	-	-	-	3	3	15	31	78
Female	2,826	-	-	-	-	-	-	1	1	2	13	26
Black, both sexes	1,939	-	-	-	-	-	-	-	4	16	27	64
Male	950	-	-	-	-	-	-	-	2	14	21	44
Female	989	-	-	-	-	-	-	-	2	2	6	20
American Indian 1/, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	122	-	-	-	-	-	-	1	2	1	2	4
Male	72	-	-	-	-	-	-	1	1	1	2	2
Female	50	-	-	-	-	-	-	-	1	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Massachusetts												
All races, both sexes	8,015	-	-	-	-	-	1	5	9	20	72	126
Male	4,023	-	-	-	-	-	1	4	9	16	57	99
Female	3,992	-	-	-	-	-	-	1	-	4	15	27
White, both sexes	7,698	-	-	-	-	-	1	5	9	16	65	115
Male	3,842	-	-	-	-	-	1	4	9	13	52	89
Female	3,856	-	-	-	-	-	-	1	-	3	13	26
Black, both sexes	231	-	-	-	-	-	-	-	-	3	6	7
Male	131	-	-	-	-	-	-	-	-	2	4	6
Female	100	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	80	-	-	-	-	-	-	-	-	1	1	3
Male	45	-	-	-	-	-	-	-	-	1	1	3
Female	35	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	16,782	-	-	-	1	1	2	5	30	74	191	405
Male	8,706	-	-	-	-	1	1	4	20	54	155	295
Female	8,076	-	-	-	1	-	1	1	10	20	36	110
White, both sexes	14,142	-	-	-	-	1	2	3	23	55	148	310
Male	7,348	-	-	-	-	1	1	2	17	38	128	234
Female	6,794	-	-	-	-	-	1	1	6	17	20	76
Black, both sexes	2,479	-	-	-	1	-	-	2	5	18	41	90
Male	1,285	-	-	-	-	-	-	2	3	15	26	59
Female	1,194	-	-	-	1	-	-	-	2	3	15	31
American Indian 1/, both sexes	91	-	-	-	-	-	-	-	2	1	2	4
Male	36	-	-	-	-	-	-	-	-	1	1	1
Female	55	-	-	-	-	-	-	-	2	-	1	3
Asian or Pacific Islander 2/, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	37	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Minnesota												
All races, both sexes	4,430	-	-	-	-	-	2	3	6	24	62	109
Male	2,561	-	-	-	-	-	2	3	3	20	48	96
Female	1,869	-	-	-	-	-	-	-	3	4	14	13
White, both sexes	4,270	-	-	-	-	-	1	3	5	22	55	95
Male	2,455	-	-	-	-	-	1	3	3	18	43	84
Female	1,815	-	-	-	-	-	-	-	2	4	12	11
Black, both sexes	70	-	-	-	-	-	1	-	-	1	2	6
Male	43	-	-	-	-	-	1	-	-	1	2	4
Female	27	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	57	-	-	-	-	-	-	-	-	-	3	7
Male	41	-	-	-	-	-	-	-	-	-	2	7
Female	16	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	1	1	2	1
Male	22	-	-	-	-	-	-	-	-	1	1	1
Female	11	-	-	-	-	-	-	-	1	-	1	-
Mississippi												
All races, both sexes	4,354	-	-	-	-	1	1	7	25	44	97	168
Male	2,291	-	-	-	-	-	1	5	20	30	68	114
Female	2,063	-	-	-	-	1	-	2	5	14	29	54
White, both sexes	3,099	-	-	-	-	-	-	3	16	30	58	97
Male	1,613	-	-	-	-	-	-	2	14	21	40	69
Female	1,486	-	-	-	-	-	-	1	2	9	18	28
Black, both sexes	1,239	-	-	-	-	1	1	4	9	13	39	68
Male	671	-	-	-	-	-	1	3	6	8	28	45
Female	568	-	-	-	-	1	-	1	3	5	11	23
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	2
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Missouri												
All races, both sexes	10,206	-	-	-	1	2	4	10	26	43	121	262
Male	5,251	-	-	-	1	2	3	8	19	31	91	200
Female	4,955	-	-	-	-	-	1	2	7	12	30	62
White, both sexes	9,261	-	-	-	1	1	3	8	21	38	102	229
Male	4,785	-	-	-	1	1	2	6	15	27	76	175
Female	4,476	-	-	-	-	-	1	2	6	11	26	54
Black, both sexes	890	-	-	-	-	1	1	1	5	5	18	30
Male	436	-	-	-	-	1	1	1	4	4	14	22
Female	454	-	-	-	-	-	-	-	1	1	4	8
American Indian 1/, both sexes	23	-	-	-	-	-	-	1	-	-	-	1
Male	14	-	-	-	-	-	-	1	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	-	1	2
Male	16	-	-	-	-	-	-	-	-	-	1	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1,093	-	-	-	-	-	-	-	3	6	20	37
Male	647	-	-	-	-	-	-	-	3	4	17	28
Female	446	-	-	-	-	-	-	-	-	2	3	9
White, both sexes	1,051	-	-	-	-	-	-	-	2	5	18	36
Male	622	-	-	-	-	-	-	-	2	4	16	27
Female	429	-	-	-	-	-	-	-	-	1	2	9
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	39	-	-	-	-	-	-	-	1	-	2	1
Male	24	-	-	-	-	-	-	-	1	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Nebraska												
All races, both sexes	1,861	-	-	-	-	-	1	-	3	4	21	41
Male	982	-	-	-	-	-	1	-	2	4	20	31
Female	879	-	-	-	-	-	-	-	1	-	1	10
White, both sexes	1,803	-	-	-	-	-	1	-	3	2	20	37
Male	950	-	-	-	-	-	1	-	2	2	19	28
Female	853	-	-	-	-	-	-	-	1	-	1	9
Black, both sexes	41	-	-	-	-	-	-	-	-	2	-	4
Male	23	-	-	-	-	-	-	-	-	2	-	3
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2,649	-	-	-	-	-	2	1	8	14	41	68
Male	1,654	-	-	-	-	-	2	1	7	12	34	52
Female	995	-	-	-	-	-	-	-	1	2	7	16
White, both sexes	2,377	-	-	-	-	-	2	-	8	10	35	59
Male	1,490	-	-	-	-	-	2	-	7	8	29	47
Female	887	-	-	-	-	-	-	-	1	2	6	12
Black, both sexes	158	-	-	-	-	-	-	1	-	3	5	4
Male	93	-	-	-	-	-	-	1	-	3	4	2
Female	65	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	1	-	2
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	87	-	-	-	-	-	-	-	-	-	1	3
Male	58	-	-	-	-	-	-	-	-	-	1	2
Female	29	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
New Hampshire												
All races, both sexes	1,629	-	-	-	-	-	-	-	3	4	19	43
Male	868	-	-	-	-	-	-	-	3	3	14	34
Female	761	-	-	-	-	-	-	-	-	1	5	9
White, both sexes	1,617	-	-	-	-	-	-	-	3	4	19	41
Male	863	-	-	-	-	-	-	-	3	3	14	33
Female	754	-	-	-	-	-	-	-	-	1	5	8
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
New Jersey												
All races, both sexes	13,684	1	-	-	-	1	2	7	11	48	114	205
Male	6,686	1	-	-	-	1	1	5	10	39	95	147
Female	6,998	-	-	-	-	-	1	2	1	9	19	58
White, both sexes	12,103	1	-	-	-	1	1	5	9	34	84	154
Male	5,892	1	-	-	-	1	1	4	8	27	77	118
Female	6,211	-	-	-	-	-	-	1	1	7	7	36
Black, both sexes	1,344	-	-	-	-	-	1	2	2	11	25	46
Male	653	-	-	-	-	-	-	1	2	9	13	25
Female	691	-	-	-	-	-	1	1	-	2	12	21
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	223	-	-	-	-	-	-	-	-	3	4	5
Male	134	-	-	-	-	-	-	-	-	3	4	4
Female	89	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
New Mexico												
All races, both sexes	2,277	1	-	-	-	1	-	1	6	8	21	36
Male	1,273	1	-	-	-	1	-	1	5	7	16	21
Female	1,004	-	-	-	-	-	-	-	1	1	5	15
White, both sexes	2,149	1	-	-	-	1	-	1	6	8	19	32
Male	1,201	1	-	-	-	1	-	1	5	7	16	17
Female	948	-	-	-	-	-	-	-	1	1	3	15
Black, both sexes	43	-	-	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	76	-	-	-	-	-	-	-	-	-	1	2
Male	50	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	39,385	1	1	-	1	2	11	23	32	116	267	552
Male	18,321	1	1	-	-	1	10	13	29	91	195	426
Female	21,064	-	-	-	1	1	1	10	3	25	72	126
White, both sexes	33,335	1	-	-	1	1	8	11	19	78	205	390
Male	15,610	1	-	-	-	1	7	7	17	65	156	317
Female	17,725	-	-	-	1	-	1	4	2	13	49	73
Black, both sexes	5,112	-	1	-	-	1	3	10	8	32	53	137
Male	2,202	-	1	-	-	-	3	5	7	20	31	88
Female	2,910	-	-	-	-	1	-	5	1	12	22	49
American Indian 1/, both sexes	43	-	-	-	-	-	-	-	1	-	-	1
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	895	-	-	-	-	-	-	2	4	6	9	24
Male	493	-	-	-	-	-	-	1	4	6	8	20
Female	402	-	-	-	-	-	-	1	-	-	1	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
North Carolina												
All races, both sexes	11,173	1	-	-	-	1	3	7	20	70	188	301
Male	6,228	-	-	-	-	1	1	7	17	53	140	236
Female	4,945	1	-	-	-	-	2	-	3	17	48	65
White, both sexes	8,877	-	-	-	-	1	1	3	14	49	133	197
Male	5,010	-	-	-	-	1	-	3	13	38	104	162
Female	3,867	-	-	-	-	-	1	-	1	11	29	35
Black, both sexes	2,136	1	-	-	-	-	2	4	6	21	51	95
Male	1,124	-	-	-	-	-	1	4	4	15	33	67
Female	1,012	1	-	-	-	-	1	-	2	6	18	28
American Indian 1/, both sexes	124	-	-	-	-	-	-	-	-	-	2	9
Male	75	-	-	-	-	-	-	-	-	-	1	7
Female	49	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	-	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	2	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1,115	-	-	-	-	-	-	2	-	5	10	18
Male	608	-	-	-	-	-	-	2	-	4	7	13
Female	507	-	-	-	-	-	-	-	-	1	3	5
White, both sexes	1,079	-	-	-	-	-	-	2	-	4	10	17
Male	584	-	-	-	-	-	-	2	-	3	7	12
Female	495	-	-	-	-	-	-	-	-	1	3	5
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	35	-	-	-	-	-	-	-	-	1	-	1
Male	23	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Ohio												
All races, both sexes	19,820	2	-	-	-	1	8	17	26	91	196	432
Male	10,321	2	-	-	-	1	6	13	21	65	152	330
Female	9,499	-	-	-	-	-	2	4	5	26	44	102
White, both sexes	17,883	2	-	-	-	1	8	15	24	75	165	357
Male	9,302	2	-	-	-	1	6	11	19	56	132	282
Female	8,581	-	-	-	-	-	2	4	5	19	33	75
Black, both sexes	1,867	-	-	-	-	-	-	2	2	15	30	74
Male	977	-	-	-	-	-	-	2	2	8	19	47
Female	890	-	-	-	-	-	-	-	-	7	11	27
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	55	-	-	-	-	-	-	-	-	1	-	1
Male	33	-	-	-	-	-	-	-	-	1	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	6,930	-	-	-	-	-	1	6	21	32	88	149
Male	3,742	-	-	-	-	-	1	4	18	25	55	110
Female	3,188	-	-	-	-	-	-	2	3	7	33	39
White, both sexes	6,130	-	-	-	-	-	1	4	16	24	67	127
Male	3,287	-	-	-	-	-	1	3	13	21	41	95
Female	2,843	-	-	-	-	-	-	1	3	3	26	32
Black, both sexes	379	-	-	-	-	-	-	2	2	7	10	6
Male	204	-	-	-	-	-	-	1	2	3	6	4
Female	175	-	-	-	-	-	-	1	-	4	4	2
American Indian 1/, both sexes	396	-	-	-	-	-	-	-	3	1	11	16
Male	237	-	-	-	-	-	-	-	3	1	8	11
Female	159	-	-	-	-	-	-	-	-	-	3	5
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Oregon												
All races, both sexes	4,070	1	-	-	-	2	-	-	4	17	35	87
Male	2,443	1	-	-	-	2	-	-	3	13	27	68
Female	1,627	-	-	-	-	-	-	-	1	4	8	19
White, both sexes	3,955	1	-	-	-	2	-	-	4	16	33	80
Male	2,375	1	-	-	-	2	-	-	3	12	27	62
Female	1,580	-	-	-	-	-	-	-	1	4	6	18
Black, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	56	-	-	-	-	-	-	-	-	1	1	5
Male	31	-	-	-	-	-	-	-	-	1	-	4
Female	25	-	-	-	-	-	-	-	-	-	1	1
Pennsylvania												
All races, both sexes	22,030	1	-	-	1	-	4	8	24	83	190	409
Male	11,228	-	-	-	1	-	3	6	20	69	142	315
Female	10,802	1	-	-	-	-	1	2	4	14	48	94
White, both sexes	20,210	1	-	-	1	-	2	5	15	67	152	345
Male	10,258	-	-	-	1	-	1	4	14	59	115	269
Female	9,952	1	-	-	-	-	1	1	1	8	37	76
Black, both sexes	1,732	-	-	-	-	-	2	3	7	14	35	60
Male	914	-	-	-	-	-	2	2	4	8	24	42
Female	818	-	-	-	-	-	-	1	3	6	11	18
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	80	-	-	-	-	-	-	-	2	2	3	4
Male	50	-	-	-	-	-	-	-	2	2	3	4
Female	30	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Rhode Island												
All races, both sexes	2,187	-	-	-	-	-	-	1	-	2	17	29
Male	1,027	-	-	-	-	-	-	1	-	2	14	24
Female	1,160	-	-	-	-	-	-	-	-	-	3	5
White, both sexes	2,128	-	-	-	-	-	-	1	-	1	16	28
Male	1,002	-	-	-	-	-	-	1	-	1	14	23
Female	1,126	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5,398	-	-	-	1	-	2	4	18	37	92	190
Male	3,156	-	-	-	1	-	1	4	14	26	73	140
Female	2,242	-	-	-	-	-	1	-	4	11	19	50
White, both sexes	4,005	-	-	-	1	-	1	4	10	26	61	130
Male	2,411	-	-	-	1	-	1	4	8	20	54	103
Female	1,594	-	-	-	-	-	-	-	2	6	7	27
Black, both sexes	1,376	-	-	-	-	-	1	-	7	11	31	59
Male	734	-	-	-	-	-	-	-	5	6	19	36
Female	642	-	-	-	-	-	1	-	2	5	12	23
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
South Dakota												
All races, both sexes	1,397	-	-	-	-	-	1	2	1	5	14	31
Male	744	-	-	-	-	-	1	2	1	3	9	26
Female	653	-	-	-	-	-	-	-	-	2	5	5
White, both sexes	1,325	-	-	-	-	-	1	1	1	4	8	27
Male	705	-	-	-	-	-	1	1	1	3	5	23
Female	620	-	-	-	-	-	-	-	-	1	3	4
Black, both sexes	8	-	-	-	-	-	-	1	-	-	1	1
Male	7	-	-	-	-	-	-	1	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	63	-	-	-	-	-	-	-	-	1	5	3
Male	31	-	-	-	-	-	-	-	-	-	3	2
Female	32	-	-	-	-	-	-	-	-	1	2	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10,602	3	2	-	-	-	1	6	30	67	192	353
Male	5,768	2	-	-	-	-	1	6	26	55	135	270
Female	4,834	1	2	-	-	-	-	-	4	12	57	83
White, both sexes	9,213	1	2	-	-	-	1	4	25	47	158	277
Male	5,061	1	-	-	-	-	1	4	22	42	114	218
Female	4,152	-	2	-	-	-	-	-	3	5	44	59
Black, both sexes	1,355	2	-	-	-	-	-	2	5	20	33	71
Male	684	1	-	-	-	-	-	2	4	13	20	47
Female	671	1	-	-	-	-	-	-	1	7	13	24
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	3
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Texas												
All races, both sexes	25,933	1	1	-	1	3	7	17	59	137	358	780
Male	14,120	1	1	-	-	3	5	13	44	106	270	539
Female	11,813	-	-	-	1	-	2	4	15	31	88	241
White, both sexes	22,818	1	1	-	1	3	6	13	44	106	289	605
Male	12,436	1	1	-	-	3	4	10	31	84	222	429
Female	10,382	-	-	-	1	-	2	3	13	22	67	176
Black, both sexes	2,867	-	-	-	-	-	1	4	13	29	64	162
Male	1,542	-	-	-	-	-	1	3	11	20	44	98
Female	1,325	-	-	-	-	-	-	1	2	9	20	64
American Indian 1/, both sexes	47	-	-	-	-	-	-	-	1	-	3	5
Male	23	-	-	-	-	-	-	-	1	-	2	5
Female	24	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	201	-	-	-	-	-	-	-	1	2	2	8
Male	119	-	-	-	-	-	-	-	1	2	2	7
Female	82	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	1,462	-	-	-	-	-	-	-	1	9	20	34
Male	925	-	-	-	-	-	-	-	1	7	14	27
Female	537	-	-	-	-	-	-	-	-	2	6	7
White, both sexes	1,433	-	-	-	-	-	-	-	1	7	19	34
Male	905	-	-	-	-	-	-	-	1	5	13	27
Female	528	-	-	-	-	-	-	-	-	2	6	7
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	1	1	-
Male	12	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Vermont												
All races, both sexes	880	-	-	-	-	-	-	-	1	1	11	22
Male	475	-	-	-	-	-	-	-	1	-	10	20
Female	405	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	879	-	-	-	-	-	-	-	1	1	11	22
Male	474	-	-	-	-	-	-	-	1	-	10	20
Female	405	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8,486	1	-	-	-	2	2	8	19	55	129	258
Male	4,675	-	-	-	-	2	2	8	14	34	106	191
Female	3,811	1	-	-	-	-	-	-	5	21	23	67
White, both sexes	6,807	1	-	-	-	1	1	5	15	33	97	182
Male	3,782	-	-	-	-	1	1	5	12	25	86	134
Female	3,025	1	-	-	-	-	-	-	3	8	11	48
Black, both sexes	1,573	-	-	-	-	1	1	1	3	22	31	70
Male	832	-	-	-	-	1	1	1	1	9	19	52
Female	741	-	-	-	-	-	-	-	2	13	12	18
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	101	-	-	-	-	-	-	2	1	-	1	6
Male	59	-	-	-	-	-	-	2	1	-	1	5
Female	42	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Washington												
All races, both sexes	7,303	-	-	-	-	-	2	4	19	34	68	143
Male	4,027	-	-	-	-	-	2	4	16	25	53	111
Female	3,276	-	-	-	-	-	-	-	3	9	15	32
White, both sexes	6,873	-	-	-	-	-	2	4	16	31	59	125
Male	3,770	-	-	-	-	-	2	4	13	23	45	94
Female	3,103	-	-	-	-	-	-	-	3	8	14	31
Black, both sexes	142	-	-	-	-	-	-	-	1	1	2	8
Male	86	-	-	-	-	-	-	-	1	1	2	8
Female	56	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	77	-	-	-	-	-	-	-	-	-	3	3
Male	40	-	-	-	-	-	-	-	-	-	2	3
Female	37	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	211	-	-	-	-	-	-	-	2	2	4	7
Male	131	-	-	-	-	-	-	-	2	1	4	6
Female	80	-	-	-	-	-	-	-	-	1	-	1
West Virginia												
All races, both sexes	3,548	-	-	-	-	-	2	4	10	21	53	89
Male	1,920	-	-	-	-	-	2	3	6	14	35	65
Female	1,628	-	-	-	-	-	-	1	4	7	18	24
White, both sexes	3,444	-	-	-	-	-	2	4	9	21	53	86
Male	1,867	-	-	-	-	-	2	3	5	14	35	63
Female	1,577	-	-	-	-	-	-	1	4	7	18	23
Black, both sexes	99	-	-	-	-	-	-	-	1	-	-	2
Male	49	-	-	-	-	-	-	-	1	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Wisconsin												
All races, both sexes	7,183	1	-	-	-	2	2	6	16	23	61	134
Male	3,946	1	-	-	-	2	1	5	13	18	48	101
Female	3,237	-	-	-	-	-	1	1	3	5	13	33
White, both sexes	6,906	-	-	-	-	2	1	5	11	23	56	120
Male	3,810	-	-	-	-	2	1	5	11	18	46	92
Female	3,096	-	-	-	-	-	-	-	-	5	10	28
Black, both sexes	220	-	-	-	-	-	1	1	5	-	4	11
Male	108	-	-	-	-	-	-	-	2	-	1	7
Female	112	-	-	-	-	-	1	1	3	-	3	4
American Indian 1/, both sexes	44	-	-	-	-	-	-	-	-	-	1	2
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	22	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	13	1	-	-	-	-	-	-	-	-	-	1
Male	6	1	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	561	-	-	-	-	-	-	1	1	5	6	13
Male	322	-	-	-	-	-	-	1	1	3	3	6
Female	239	-	-	-	-	-	-	-	-	2	3	7
White, both sexes	553	-	-	-	-	-	-	1	1	4	5	13
Male	316	-	-	-	-	-	-	1	1	2	3	6
Female	237	-	-	-	-	-	-	-	-	2	2	7
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
United States												
All races, both sexes	15,363	21,524	25,182	28,956	37,729	53,618	71,607	73,652	53,357	21,915	4,931	31
Male	11,723	15,921	17,672	19,361	23,487	30,824	36,451	31,702	17,998	5,256	807	22
Female	3,640	5,603	7,510	9,595	14,242	22,794	35,156	41,950	35,359	16,659	4,124	9
White, both sexes	12,109	17,229	20,680	23,865	31,856	46,869	64,121	67,463	49,176	20,096	4,339	26
Male	9,485	13,016	14,756	16,257	20,177	27,359	33,093	29,352	16,749	4,830	701	19
Female	2,624	4,213	5,924	7,608	11,679	19,510	31,028	38,111	32,427	15,266	3,638	7
Black, both sexes	2,873	3,703	3,823	4,240	4,919	5,506	5,992	4,833	3,275	1,483	493	5
Male	1,936	2,450	2,440	2,535	2,765	2,776	2,608	1,728	881	292	72	3
Female	937	1,253	1,383	1,705	2,154	2,730	3,384	3,105	2,394	1,191	421	2
American Indian 1/, both sexes	123	166	191	204	216	242	248	173	109	33	6	-
Male	96	117	124	129	127	138	110	71	32	8	2	-
Female	27	49	67	75	89	104	138	102	77	25	4	-
Asian or Pacific Islander 2/, both sexes	258	426	488	647	738	1,001	1,246	1,183	797	303	93	-
Male	206	338	352	440	418	551	640	551	336	126	32	-
Female	52	88	136	207	320	450	606	632	461	177	61	-
Alabama												
All races, both sexes	281	371	422	463	637	845	913	876	546	225	52	-
Male	204	263	303	313	402	452	421	346	166	45	5	-
Female	77	108	119	150	235	393	492	530	380	180	47	-
White, both sexes	201	259	320	365	514	668	763	753	465	190	32	-
Male	149	182	236	250	337	366	355	302	148	42	2	-
Female	52	77	84	115	177	302	408	451	317	148	30	-
Black, both sexes	77	110	101	98	120	175	148	122	79	35	20	-
Male	53	80	66	63	64	85	65	44	17	3	3	-
Female	24	30	35	35	56	90	83	78	62	32	17	-
American Indian 1/, both sexes	1	1	-	-	1	-	1	1	1	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	-	2	2	1	-	1	-	-	-
Male	1	-	1	-	-	1	1	-	1	-	-	-
Female	1	1	-	-	2	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Alaska												
All races, both sexes	31	31	25	36	40	54	57	29	12	9	2	-
Male	26	22	19	26	28	40	26	16	6	3	-	-
Female	5	9	6	10	12	14	31	13	6	6	2	-
White, both sexes	26	25	19	29	29	45	48	27	10	7	2	-
Male	21	18	14	21	20	34	24	16	4	2	-	-
Female	5	7	5	8	9	11	24	11	6	5	2	-
Black, both sexes	1	-	-	-	-	1	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	3	4	6	6	9	7	8	1	2	1	-	-
Male	3	2	5	5	6	6	2	-	2	-	-	-
Female	-	2	1	1	3	1	6	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	1	2	1	1	1	-	-	-	-
Male	1	2	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
Arizona												
All races, both sexes	282	392	521	571	730	1,047	1,318	1,370	881	353	66	8
Male	226	293	391	379	462	596	701	631	328	87	19	6
Female	56	99	130	192	268	451	617	739	553	266	47	2
White, both sexes	257	365	479	527	691	993	1,256	1,317	855	340	62	6
Male	203	272	359	354	437	569	668	604	318	81	18	5
Female	54	93	120	173	254	424	588	713	537	259	44	1
Black, both sexes	17	13	23	19	27	21	25	27	10	8	3	2
Male	16	10	18	13	19	8	16	13	4	4	-	1
Female	1	3	5	6	8	13	9	14	6	4	3	1
American Indian 1/, both sexes	7	13	17	19	6	24	26	18	11	2	-	-
Male	6	10	13	9	3	17	11	10	4	1	-	-
Female	1	3	4	10	3	7	15	8	7	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	6	6	9	11	8	5	3	1	-
Male	1	1	1	3	3	2	6	4	2	1	1	-
Female	-	-	1	3	3	7	5	4	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Arkansas												
All races, both sexes	218	308	381	389	495	633	773	735	554	226	43	1
Male	166	235	268	253	290	326	394	293	184	56	6	-
Female	52	73	113	136	205	307	379	442	370	170	37	1
White, both sexes	167	249	319	337	419	557	691	681	492	200	37	1
Male	131	194	224	229	254	294	360	273	165	49	4	-
Female	36	55	95	108	165	263	331	408	327	151	33	1
Black, both sexes	49	56	57	49	73	74	80	53	61	25	6	-
Male	33	39	40	24	34	31	33	19	19	6	2	-
Female	16	17	17	25	39	43	47	34	42	19	4	-
American Indian 1/, both sexes	2	2	2	2	2	-	2	-	1	-	-	-
Male	2	1	2	-	1	-	1	-	-	-	-	-
Female	-	1	-	2	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	1	1	2	-	1	-	1	-	-
Male	-	1	2	-	1	1	-	1	-	1	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
California												
All races, both sexes	1,452	2,214	2,483	2,969	3,928	5,649	8,168	8,564	6,349	2,660	627	5
Male	1,155	1,630	1,756	1,935	2,415	3,343	4,208	3,895	2,298	689	112	3
Female	297	584	727	1,034	1,513	2,306	3,960	4,669	4,051	1,971	515	2
White, both sexes	1,155	1,710	1,998	2,347	3,193	4,670	6,992	7,515	5,602	2,351	526	5
Male	932	1,282	1,435	1,557	1,996	2,783	3,620	3,467	2,040	589	85	3
Female	223	428	563	790	1,197	1,887	3,372	4,048	3,562	1,762	441	2
Black, both sexes	194	310	282	335	393	456	505	422	303	134	47	-
Male	141	202	184	189	221	268	239	145	80	31	6	-
Female	53	108	98	146	172	188	266	277	223	103	41	-
American Indian 1/, both sexes	10	19	17	29	22	32	32	25	9	4	-	-
Male	9	14	7	19	13	18	14	7	2	1	-	-
Female	1	5	10	10	9	14	18	18	7	3	-	-
Asian or Pacific Islander 2/, both sexes	93	175	186	258	320	491	639	602	435	171	54	-
Male	73	132	130	170	185	274	335	276	176	68	21	-
Female	20	43	56	88	135	217	304	326	259	103	33	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Colorado												
All races, both sexes	179	224	258	275	358	481	599	653	470	192	36	-
Male	147	184	187	185	233	283	308	312	170	45	3	-
Female	32	40	71	90	125	198	291	341	300	147	33	-
White, both sexes	172	212	246	263	340	460	581	645	463	187	35	-
Male	140	175	181	178	221	272	302	306	166	43	3	-
Female	32	37	65	85	119	188	279	339	297	144	32	-
Black, both sexes	5	10	9	12	13	10	9	4	3	4	1	-
Male	5	7	5	7	11	7	3	4	2	1	-	-
Female	-	3	4	5	2	3	6	-	1	3	1	-
American Indian 1/, both sexes	1	-	1	-	1	5	3	1	1	-	-	-
Male	1	-	-	-	1	3	1	1	1	-	-	-
Female	-	-	1	-	-	2	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	-	4	6	6	3	3	1	-	-
Male	1	2	1	-	-	1	2	1	1	1	-	-
Female	-	-	1	-	4	5	4	2	2	-	-	-
Connecticut												
All races, both sexes	114	177	222	233	335	523	809	967	699	331	79	1
Male	89	132	155	162	213	288	410	422	225	71	10	1
Female	25	45	67	71	122	235	399	545	474	260	69	-
White, both sexes	98	153	191	204	303	480	775	940	686	325	76	1
Male	78	116	138	144	194	273	394	410	224	70	10	1
Female	20	37	53	60	109	207	381	530	462	255	66	-
Black, both sexes	15	24	27	25	30	34	29	22	10	5	3	-
Male	10	16	13	15	17	12	14	9	1	1	-	-
Female	5	8	14	10	13	22	15	13	9	4	3	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	1	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	4	2	8	4	4	1	1	-	-
Male	1	-	3	3	2	3	2	3	-	-	-	-
Female	-	-	-	1	-	5	2	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Delaware												
All races, both sexes	54	50	83	87	124	155	224	241	159	47	17	1
Male	39	40	55	57	81	93	103	111	39	15	1	1
Female	15	10	28	30	43	62	121	130	120	32	16	-
White, both sexes	43	42	68	67	98	139	195	221	141	39	13	1
Male	31	34	45	46	62	82	90	102	35	14	1	1
Female	12	8	23	21	36	57	105	119	106	25	12	-
Black, both sexes	10	8	14	18	24	15	29	20	17	8	4	-
Male	8	6	9	11	17	11	13	9	4	1	-	-
Female	2	2	5	7	7	4	16	11	13	7	4	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	1	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	52	60	79	115	100	145	184	171	127	51	18	-
Male	34	39	49	70	58	80	83	66	44	14	2	-
Female	18	21	30	45	42	65	101	105	83	37	16	-
White, both sexes	5	10	15	17	14	34	55	50	44	20	5	-
Male	4	9	11	11	9	25	26	28	18	9	-	-
Female	1	1	4	6	5	9	29	22	26	11	5	-
Black, both sexes	47	49	64	97	85	109	128	119	81	31	13	-
Male	30	29	38	58	49	54	56	36	25	5	2	-
Female	17	20	26	39	36	55	72	83	56	26	11	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	1	1	-	-	-	-
Male	-	1	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Florida												
All races, both sexes	874	1,325	1,637	2,023	2,727	4,249	5,976	6,224	4,527	1,870	412	1
Male	679	978	1,169	1,407	1,753	2,499	3,184	2,856	1,757	508	99	-
Female	195	347	468	616	974	1,750	2,792	3,368	2,770	1,362	313	1
White, both sexes	700	1,113	1,413	1,738	2,378	3,898	5,565	5,910	4,287	1,765	375	1
Male	557	855	1,028	1,229	1,549	2,326	2,998	2,750	1,694	480	94	-
Female	143	258	385	509	829	1,572	2,567	3,160	2,593	1,285	281	1
Black, both sexes	160	205	214	267	335	325	383	293	231	97	35	-
Male	110	119	135	164	195	158	173	99	61	25	4	-
Female	50	86	79	103	140	167	210	194	170	72	31	-
American Indian 1/, both sexes	3	2	3	3	2	2	4	3	1	1	-	-
Male	3	-	2	2	1	-	2	-	-	-	-	-
Female	-	2	1	1	1	2	2	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	11	5	7	15	12	24	24	18	8	7	2	-
Male	9	4	4	12	8	15	11	7	2	3	1	-
Female	2	1	3	3	4	9	13	11	6	4	1	-
Georgia												
All races, both sexes	433	568	675	723	840	992	1,299	1,140	817	281	45	1
Male	308	437	490	481	545	527	669	469	222	63	7	1
Female	125	131	185	242	295	465	630	671	595	218	38	-
White, both sexes	274	386	482	536	616	763	1,069	935	657	236	31	1
Male	214	312	367	375	416	426	567	406	181	58	2	1
Female	60	74	115	161	200	337	502	529	476	178	29	-
Black, both sexes	158	178	185	181	215	212	219	196	155	45	13	-
Male	93	121	120	101	125	92	95	61	40	5	4	-
Female	65	57	65	80	90	120	124	135	115	40	9	-
American Indian 1/, both sexes	-	-	3	1	1	4	2	1	1	-	-	-
Male	-	-	1	1	1	4	2	1	-	-	-	-
Female	-	-	2	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	5	5	8	13	9	8	4	-	1	-
Male	1	4	2	4	3	5	5	1	1	-	1	-
Female	-	-	3	1	5	8	4	7	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Hawaii												
All races, both sexes	57	79	96	107	99	170	216	208	149	45	16	-
Male	46	61	70	80	57	108	145	107	76	28	7	-
Female	11	18	26	27	42	62	71	101	73	17	9	-
White, both sexes	22	25	29	27	25	58	58	59	40	14	4	-
Male	18	18	23	20	14	38	42	31	17	6	1	-
Female	4	7	6	7	11	20	16	28	23	8	3	-
Black, both sexes	-	-	-	-	3	1	1	1	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	53	67	80	71	111	157	148	109	31	12	-
Male	28	42	47	60	41	69	103	76	59	22	6	-
Female	7	11	20	20	30	42	54	72	50	9	6	-
Idaho												
All races, both sexes	51	91	95	110	150	198	265	251	203	72	6	-
Male	38	72	70	78	98	136	151	130	75	24	3	-
Female	13	19	25	32	52	62	114	121	128	48	3	-
White, both sexes	47	89	93	108	148	195	262	247	202	71	6	-
Male	36	70	69	77	97	135	149	127	75	23	3	-
Female	11	19	24	31	51	60	113	120	127	48	3	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	2	1	1	2	1	1	1	-	-	-
Male	1	1	1	1	-	1	1	1	-	-	-	-
Female	1	-	1	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	1	1	3	-	1	-	-
Male	-	1	-	-	1	-	-	2	-	1	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Illinois												
All races, both sexes	678	931	1,030	1,166	1,508	2,187	2,981	3,024	2,326	976	212	1
Male	531	696	730	786	953	1,235	1,470	1,284	732	227	28	-
Female	147	235	300	380	555	952	1,511	1,740	1,594	749	184	1
White, both sexes	514	688	794	898	1,218	1,853	2,607	2,751	2,155	888	187	-
Male	406	530	571	612	789	1,064	1,298	1,169	679	207	26	-
Female	108	158	223	286	429	789	1,309	1,582	1,476	681	161	-
Black, both sexes	152	229	220	238	262	308	353	232	156	78	23	1
Male	117	155	148	154	148	158	158	99	48	17	2	-
Female	35	74	72	84	114	150	195	133	108	61	21	1
American Indian 1/, both sexes	2	1	2	2	-	4	1	1	-	-	-	-
Male	1	1	1	2	-	1	-	1	-	-	-	-
Female	1	-	1	-	-	3	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	13	14	28	28	22	20	40	15	10	2	-
Male	7	10	10	18	16	12	14	15	5	3	-	-
Female	3	3	4	10	12	10	6	25	10	7	2	-
Indiana												
All races, both sexes	355	484	572	682	893	1,204	1,508	1,501	1,015	409	98	-
Male	265	336	397	463	555	712	776	631	344	97	15	-
Female	90	148	175	219	338	492	732	870	671	312	83	-
White, both sexes	318	413	512	607	816	1,124	1,414	1,430	963	392	93	-
Male	235	294	352	414	504	668	727	606	330	93	14	-
Female	83	119	160	193	312	456	687	824	633	299	79	-
Black, both sexes	37	69	57	69	71	77	92	68	52	17	5	-
Male	30	41	42	44	47	43	47	24	14	4	1	-
Female	7	28	15	25	24	34	45	44	38	13	4	-
American Indian 1/, both sexes	-	-	-	1	2	1	-	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	3	5	4	2	2	3	-	-	-	-
Male	-	1	3	5	2	-	2	1	-	-	-	-
Female	-	1	-	-	2	2	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Iowa												
All races, both sexes	173	221	257	344	388	624	934	1,035	826	392	98	-
Male	129	162	172	243	252	356	487	482	281	98	11	-
Female	44	59	85	101	136	268	447	553	545	294	87	-
White, both sexes	159	214	252	334	380	614	927	1,028	823	392	98	-
Male	119	157	170	237	249	347	484	478	281	98	11	-
Female	40	57	82	97	131	267	443	550	542	294	87	-
Black, both sexes	13	7	4	8	5	8	4	7	3	-	-	-
Male	9	5	1	4	2	7	2	4	-	-	-	-
Female	4	2	3	4	3	1	2	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	3	2	1	-	-	-	-	-
Male	1	-	1	2	1	2	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
Kansas												
All races, both sexes	144	182	205	247	305	444	595	629	445	188	45	-
Male	120	145	143	166	189	259	335	281	161	38	5	-
Female	24	37	62	81	116	185	260	348	284	150	40	-
White, both sexes	133	167	185	230	280	409	575	609	437	184	43	-
Male	110	136	129	158	173	244	323	271	159	36	5	-
Female	23	31	56	72	107	165	252	338	278	148	38	-
Black, both sexes	9	14	14	10	17	26	13	17	7	4	2	-
Male	8	8	10	4	11	11	7	9	2	2	-	-
Female	1	6	4	6	6	15	6	8	5	2	2	-
American Indian 1/, both sexes	1	1	4	6	4	6	5	1	1	-	-	-
Male	1	1	3	3	3	4	4	1	-	-	-	-
Female	-	-	1	3	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	4	3	2	2	-	-	-	-
Male	1	-	1	1	2	-	1	-	-	-	-	-
Female	-	-	1	-	2	3	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Kentucky												
All races, both sexes	355	375	457	529	701	832	1,019	941	622	212	48	-
Male	264	261	310	349	420	489	475	377	176	54	7	-
Female	91	114	147	180	281	343	544	564	446	158	41	-
White, both sexes	318	348	426	484	648	787	981	891	602	198	46	-
Male	236	247	289	322	384	467	458	358	168	52	6	-
Female	82	101	137	162	264	320	523	533	434	146	40	-
Black, both sexes	36	26	29	44	50	44	36	49	19	14	2	-
Male	28	14	20	27	35	21	16	19	7	2	1	-
Female	8	12	9	17	15	23	20	30	12	12	1	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	2	1	2	1	1	-	-	-
Male	-	-	1	-	1	1	1	-	1	-	-	-
Female	-	1	1	1	1	-	1	1	-	-	-	-
Louisiana												
All races, both sexes	344	438	524	519	608	711	879	789	461	176	47	-
Male	246	302	360	350	387	415	460	312	138	35	9	-
Female	98	136	164	169	221	296	419	477	323	141	38	-
White, both sexes	198	266	345	341	447	537	692	629	366	139	30	-
Male	153	187	251	245	297	321	381	254	110	31	7	-
Female	45	79	94	96	150	216	311	375	256	108	23	-
Black, both sexes	143	168	175	173	155	173	183	156	93	37	17	-
Male	90	111	106	103	87	93	77	56	28	4	2	-
Female	53	57	69	70	68	80	106	100	65	33	15	-
American Indian 1/, both sexes	1	-	2	3	3	-	2	-	-	-	-	-
Male	1	-	2	1	2	-	2	-	-	-	-	-
Female	-	-	-	2	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	2	2	3	1	2	4	2	-	-	-
Male	2	4	1	1	1	1	-	2	-	-	-	-
Female	-	-	1	1	2	-	2	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Maine												
All races, both sexes	56	93	117	120	155	215	333	309	231	86	25	-
Male	47	73	88	83	100	129	183	119	88	24	2	-
Female	9	20	29	37	55	86	150	190	143	62	23	-
White, both sexes	53	91	117	120	155	215	332	309	231	85	25	-
Male	44	71	88	83	100	129	183	119	88	23	2	-
Female	9	20	29	37	55	86	149	190	143	62	23	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	1	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	319	413	498	544	706	982	1,307	1,284	931	374	90	1
Male	217	281	335	334	426	543	609	530	276	94	12	1
Female	102	132	163	210	280	439	698	754	655	280	78	-
White, both sexes	190	245	310	337	464	733	1,012	1,069	763	311	70	1
Male	138	170	217	216	286	407	497	462	239	80	9	1
Female	52	75	93	121	178	326	515	607	524	231	61	-
Black, both sexes	126	162	173	196	226	234	276	199	159	58	19	-
Male	77	106	109	107	129	126	104	60	34	14	3	-
Female	49	56	64	89	97	108	172	139	125	44	16	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	6	14	11	16	14	18	16	8	5	1	-
Male	2	5	8	11	11	9	8	8	3	-	-	-
Female	1	1	6	-	5	5	10	8	5	5	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Massachusetts												
All races, both sexes	231	312	393	435	625	948	1,416	1,594	1,161	520	147	-
Male	197	248	272	307	391	542	744	630	385	104	17	-
Female	34	64	121	128	234	406	672	964	776	416	130	-
White, both sexes	214	292	367	404	583	917	1,370	1,550	1,137	510	143	-
Male	183	230	252	284	374	525	719	611	379	101	16	-
Female	31	62	115	120	209	392	651	939	758	409	127	-
Black, both sexes	13	16	20	23	30	20	37	32	15	8	1	-
Male	10	14	16	16	11	9	22	13	5	2	1	-
Female	3	2	4	7	19	11	15	19	10	6	-	-
American Indian 1/, both sexes	2	-	1	-	-	-	1	-	-	1	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	4	5	8	12	11	8	12	9	1	3	-
Male	2	4	3	7	6	8	2	6	1	1	-	-
Female	-	-	2	1	6	3	6	6	8	-	3	-
Michigan												
All races, both sexes	685	959	1,001	1,074	1,486	2,129	2,879	2,913	2,002	766	179	-
Male	495	694	687	680	888	1,193	1,423	1,251	657	175	33	-
Female	190	265	314	394	598	936	1,456	1,662	1,345	591	146	-
White, both sexes	496	701	770	850	1,197	1,812	2,492	2,617	1,826	687	152	-
Male	378	520	539	551	731	1,043	1,257	1,122	601	157	28	-
Female	118	181	231	299	466	769	1,235	1,495	1,225	530	124	-
Black, both sexes	178	246	219	209	265	300	365	274	161	78	27	-
Male	109	166	143	124	142	144	160	117	52	18	5	-
Female	69	80	76	85	123	156	205	157	109	60	22	-
American Indian 1/, both sexes	6	8	6	12	7	13	11	10	8	1	-	-
Male	3	5	3	4	4	4	3	4	3	-	-	-
Female	3	3	3	8	3	9	8	6	5	1	-	-
Asian or Pacific Islander 2/, both sexes	5	4	6	3	17	4	11	12	7	-	-	-
Male	5	3	2	1	11	2	3	8	1	-	-	-
Female	-	1	4	2	6	2	8	4	6	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Minnesota												
All races, both sexes	191	206	226	245	358	488	759	806	629	250	66	-
Male	160	162	160	175	251	304	456	402	244	65	10	-
Female	31	44	66	70	107	184	303	404	385	185	56	-
White, both sexes	171	180	206	229	339	474	748	799	627	250	66	-
Male	146	142	147	165	238	298	449	399	244	65	10	-
Female	25	38	59	64	101	176	299	400	383	185	56	-
Black, both sexes	10	14	10	3	8	6	4	5	-	-	-	-
Male	7	10	5	-	5	3	3	2	-	-	-	-
Female	3	4	5	3	3	3	1	3	-	-	-	-
American Indian 1/, both sexes	7	6	7	11	4	5	4	2	1	-	-	-
Male	4	4	5	8	3	3	4	1	-	-	-	-
Female	3	2	2	3	1	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	6	3	2	7	3	3	-	1	-	-	-
Male	3	6	3	2	5	-	-	-	-	-	-	-
Female	-	-	-	-	2	3	3	-	1	-	-	-
Mississippi												
All races, both sexes	263	301	332	376	440	553	581	569	397	167	32	-
Male	193	199	224	256	264	306	273	204	100	32	2	-
Female	70	102	108	120	176	247	308	365	297	135	30	-
White, both sexes	138	196	223	265	298	421	447	449	314	125	19	-
Male	102	130	150	189	182	240	213	160	77	23	1	-
Female	36	66	73	76	116	181	234	289	237	102	18	-
Black, both sexes	125	101	108	108	141	131	132	120	83	42	13	-
Male	91	67	73	65	82	66	59	44	23	9	1	-
Female	34	34	35	43	59	65	73	76	60	33	12	-
American Indian 1/, both sexes	-	1	1	1	1	1	1	-	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	2	-	-	1	-	-	-	-	-
Male	-	2	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Missouri												
All races, both sexes	411	538	607	732	881	1,201	1,657	1,769	1,282	546	113	-
Male	302	392	405	492	543	683	824	708	411	119	17	-
Female	109	146	202	240	338	518	833	1,061	871	427	96	-
White, both sexes	350	464	531	638	761	1,091	1,528	1,663	1,214	512	106	-
Male	260	341	356	439	479	627	778	677	394	115	16	-
Female	90	123	175	199	282	464	750	986	820	397	90	-
Black, both sexes	58	70	71	90	111	101	124	100	65	33	6	-
Male	39	49	46	52	58	50	45	29	16	4	1	-
Female	19	21	25	38	53	51	79	71	49	29	5	-
American Indian 1/, both sexes	2	1	4	-	4	3	4	3	-	-	-	-
Male	2	-	2	-	4	3	1	-	-	-	-	-
Female	-	1	2	-	-	-	3	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	4	5	6	1	3	3	1	1	-
Male	1	2	1	1	2	3	-	2	1	-	-	-
Female	-	1	-	3	3	3	1	1	2	1	1	-
Montana												
All races, both sexes	40	64	69	81	99	151	163	175	132	48	5	-
Male	31	49	52	57	64	90	90	87	59	14	2	-
Female	9	15	17	24	35	61	73	88	73	34	3	-
White, both sexes	37	60	65	75	94	146	160	172	128	48	5	-
Male	28	46	49	52	61	89	87	87	58	14	2	-
Female	9	14	16	23	33	57	73	85	70	34	3	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	3	4	6	5	5	3	3	3	-	-	-
Male	3	2	3	5	3	1	3	-	1	-	-	-
Female	-	1	1	1	2	4	-	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Nebraska												
All races, both sexes	61	71	90	102	143	203	341	323	281	147	29	-
Male	49	58	62	72	90	120	196	138	96	38	5	-
Female	12	13	28	30	53	83	145	185	185	109	24	-
White, both sexes	56	65	82	100	140	197	334	317	277	143	29	-
Male	44	54	56	72	89	117	193	135	95	38	5	-
Female	12	11	26	28	51	80	141	182	182	105	24	-
Black, both sexes	4	6	2	1	3	6	3	3	4	3	-	-
Male	4	4	2	-	1	3	2	1	1	-	-	-
Female	-	2	-	1	2	3	1	2	3	3	-	-
American Indian 1/, both sexes	1	-	5	1	-	-	1	-	-	1	-	-
Male	1	-	3	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	3	3	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Nevada												
All races, both sexes	147	211	230	265	335	359	382	326	187	60	13	-
Male	121	159	166	196	226	213	222	155	69	18	1	-
Female	26	52	64	69	109	146	160	171	118	42	12	-
White, both sexes	130	181	201	229	296	321	353	312	176	56	8	-
Male	109	140	145	173	202	194	204	146	67	16	1	-
Female	21	41	56	56	94	127	149	166	109	40	7	-
Black, both sexes	13	17	14	21	23	23	14	6	8	2	4	-
Male	8	11	9	14	16	10	10	3	1	1	-	-
Female	5	6	5	7	7	13	4	3	7	1	4	-
American Indian 1/, both sexes	1	4	5	3	3	2	3	1	-	1	1	-
Male	1	-	3	2	1	1	2	1	-	-	-	-
Female	-	4	2	1	2	1	1	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	3	9	10	12	13	13	12	7	3	1	-	-
Male	3	8	9	7	7	8	6	5	1	1	-	-
Female	-	1	1	5	6	5	6	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
New Hampshire												
All races, both sexes	53	90	75	91	139	212	277	293	215	96	19	-
Male	42	73	57	75	86	117	134	128	73	25	4	-
Female	11	17	18	16	53	95	143	165	142	71	15	-
White, both sexes	53	88	73	91	139	210	275	291	215	96	19	-
Male	42	72	56	75	86	116	134	127	73	25	4	-
Female	11	16	17	16	53	94	141	164	142	71	15	-
Black, both sexes	-	2	1	-	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
New Jersey												
All races, both sexes	371	493	634	767	1,083	1,747	2,461	2,681	1,992	871	193	2
Male	286	356	437	515	680	949	1,195	1,099	646	202	20	2
Female	85	137	197	252	403	798	1,266	1,582	1,346	669	173	-
White, both sexes	293	385	497	607	876	1,549	2,236	2,505	1,863	824	177	2
Male	234	282	361	414	560	851	1,090	1,035	613	194	19	2
Female	59	103	136	193	316	698	1,146	1,470	1,250	630	158	-
Black, both sexes	69	94	112	130	183	168	191	145	109	43	13	-
Male	44	62	57	80	107	83	88	50	25	7	-	-
Female	25	32	55	50	76	85	103	95	84	36	13	-
American Indian 1/, both sexes	-	2	1	2	1	1	3	1	2	-	-	-
Male	-	2	-	-	-	1	1	1	1	-	-	-
Female	-	-	1	2	1	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	9	12	24	28	23	29	31	30	18	4	3	-
Male	8	10	19	21	13	14	16	13	7	1	1	-
Female	1	2	5	7	10	15	15	17	11	3	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
New Mexico												
All races, both sexes	83	129	132	161	242	303	353	381	281	115	23	-
Male	70	97	94	113	157	180	195	180	110	22	3	-
Female	13	32	38	48	85	123	158	201	171	93	20	-
White, both sexes	75	119	123	149	216	292	338	370	271	107	21	-
Male	63	91	86	104	145	174	186	176	106	20	2	-
Female	12	28	37	45	71	118	152	194	165	87	19	-
Black, both sexes	2	3	-	8	8	3	5	4	3	5	-	-
Male	2	1	-	6	3	1	1	1	-	-	-	-
Female	-	2	-	2	5	2	4	3	3	5	-	-
American Indian 1/, both sexes	6	6	9	3	16	7	7	7	7	3	2	-
Male	5	4	8	2	8	5	6	3	4	2	1	-
Female	1	2	1	1	8	2	1	4	3	1	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	1	3	-	-	-	-	-
Male	-	1	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
New York												
All races, both sexes	885	1,407	1,780	2,227	3,075	4,809	6,825	7,723	6,167	2,780	700	1
Male	641	1,037	1,206	1,404	1,774	2,592	3,189	3,079	1,913	606	112	1
Female	244	370	574	823	1,301	2,217	3,636	4,644	4,254	2,174	588	-
White, both sexes	653	1,082	1,350	1,663	2,414	4,008	5,867	6,864	5,612	2,489	618	1
Male	509	831	931	1,079	1,450	2,222	2,832	2,776	1,755	551	102	1
Female	144	251	419	584	964	1,786	3,035	4,088	3,857	1,938	516	-
Black, both sexes	200	280	375	500	572	684	787	704	446	244	75	-
Male	107	169	230	286	272	306	287	232	107	42	9	-
Female	93	111	145	214	300	378	500	472	339	202	66	-
American Indian 1/, both sexes	-	1	4	3	7	8	5	7	2	4	-	-
Male	-	1	3	2	2	2	2	2	-	-	-	-
Female	-	-	1	1	5	6	3	5	2	4	-	-
Asian or Pacific Islander 2/, both sexes	32	44	51	61	82	109	166	148	107	43	7	-
Male	25	36	42	37	50	62	68	69	51	13	1	-
Female	7	8	9	24	32	47	98	79	56	30	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
North Carolina												
All races, both sexes	512	711	848	920	1,115	1,436	1,735	1,686	1,129	394	96	-
Male	401	540	615	627	737	820	918	696	323	82	14	-
Female	111	171	233	293	378	616	817	990	806	312	82	-
White, both sexes	358	517	647	689	873	1,165	1,424	1,432	957	340	77	-
Male	289	401	481	472	596	687	788	608	284	74	9	-
Female	69	116	166	217	277	478	636	824	673	266	68	-
Black, both sexes	143	174	190	215	224	257	284	238	161	51	19	-
Male	101	120	126	143	132	128	118	85	34	8	5	-
Female	42	54	64	72	92	129	166	153	127	43	14	-
American Indian 1/, both sexes	8	16	7	13	14	10	22	12	9	2	-	-
Male	8	15	6	11	9	4	9	1	4	-	-	-
Female	-	1	1	2	5	6	13	11	5	2	-	-
Asian or Pacific Islander 2/, both sexes	3	4	4	3	4	4	5	4	2	1	-	-
Male	3	4	2	1	-	1	3	2	1	-	-	-
Female	-	-	2	2	4	3	2	2	1	1	-	-
North Dakota												
All races, both sexes	42	40	50	70	91	112	184	214	188	70	19	-
Male	34	35	41	49	55	76	100	94	68	25	5	-
Female	8	5	9	21	36	36	84	120	120	45	14	-
White, both sexes	38	37	45	67	83	107	183	211	186	70	19	-
Male	31	32	38	46	51	73	99	92	68	25	5	-
Female	7	5	7	21	32	34	84	119	118	45	14	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	3	4	3	8	5	1	3	2	-	-	-
Male	3	3	2	3	4	3	1	2	-	-	-	-
Female	1	-	2	-	4	2	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Ohio												
All races, both sexes	740	1,041	1,106	1,312	1,744	2,627	3,474	3,477	2,431	910	185	-
Male	578	769	773	852	1,051	1,492	1,745	1,457	784	200	30	-
Female	162	272	333	460	693	1,135	1,729	2,020	1,647	710	155	-
White, both sexes	615	869	960	1,135	1,535	2,361	3,190	3,271	2,290	848	162	-
Male	491	651	684	744	927	1,359	1,604	1,380	743	185	25	-
Female	124	218	276	391	608	1,002	1,586	1,891	1,547	663	137	-
Black, both sexes	121	164	140	169	204	258	270	201	135	59	23	-
Male	85	111	84	102	122	129	134	75	39	13	5	-
Female	36	53	56	67	82	129	136	126	96	46	18	-
American Indian 1/, both sexes	2	2	3	1	-	2	3	1	-	-	-	-
Male	1	1	2	1	-	1	1	1	-	-	-	-
Female	1	1	1	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	6	3	7	5	6	11	4	6	3	-	-
Male	1	6	3	5	2	3	6	1	2	2	-	-
Female	1	-	-	2	3	3	5	3	4	1	-	-
Oklahoma												
All races, both sexes	308	407	501	529	681	862	1,114	1,078	760	335	57	1
Male	217	300	345	350	428	512	580	457	259	71	9	1
Female	91	107	156	179	253	350	534	621	501	264	48	-
White, both sexes	247	338	430	444	585	765	1,013	999	706	310	53	1
Male	178	253	297	291	368	458	530	420	243	66	8	1
Female	69	85	133	153	217	307	483	579	463	244	45	-
Black, both sexes	37	35	31	37	36	44	50	34	26	20	2	-
Male	23	22	20	25	23	22	23	15	10	4	1	-
Female	14	13	11	12	13	22	27	19	16	16	1	-
American Indian 1/, both sexes	23	31	38	43	56	50	47	44	27	5	1	-
Male	15	23	27	29	35	31	25	22	6	1	-	-
Female	8	8	11	14	21	19	22	22	21	4	1	-
Asian or Pacific Islander 2/, both sexes	1	3	2	5	4	3	4	1	1	-	1	-
Male	1	2	1	5	2	1	2	-	-	-	-	-
Female	-	1	1	-	2	2	2	1	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Oregon												
All races, both sexes	150	212	234	317	376	493	681	739	504	188	30	-
Male	127	167	173	227	249	317	403	387	209	64	6	-
Female	23	45	61	90	127	176	278	352	295	124	24	-
White, both sexes	144	209	220	305	364	479	669	720	496	184	29	-
Male	122	165	165	216	241	309	399	376	207	62	6	-
Female	22	44	55	89	123	170	270	344	289	122	23	-
Black, both sexes	4	3	6	-	4	3	3	6	2	2	-	-
Male	3	2	2	-	3	2	2	3	1	1	-	-
Female	1	1	4	-	1	1	1	3	1	1	-	-
American Indian 1/, both sexes	2	-	4	4	3	3	-	6	1	-	-	-
Male	2	-	2	3	3	2	-	4	-	-	-	-
Female	-	-	2	1	-	1	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	4	8	5	8	9	7	5	2	1	-
Male	-	-	4	8	2	4	2	4	1	1	-	-
Female	-	-	-	-	3	4	7	3	4	1	1	-
Pennsylvania												
All races, both sexes	619	943	1,167	1,345	1,872	2,899	3,943	4,190	2,957	1,155	218	2
Male	467	701	831	884	1,167	1,661	1,993	1,729	933	277	28	1
Female	152	242	336	461	705	1,238	1,950	2,461	2,024	878	190	1
White, both sexes	500	790	1,007	1,166	1,667	2,658	3,698	3,998	2,836	1,101	199	2
Male	391	595	724	776	1,047	1,529	1,878	1,660	901	267	26	1
Female	109	195	283	390	620	1,129	1,820	2,338	1,935	834	173	1
Black, both sexes	116	148	152	169	194	230	236	180	116	52	18	-
Male	74	102	101	101	113	124	111	63	32	9	2	-
Female	42	46	51	68	81	106	125	117	84	43	16	-
American Indian 1/, both sexes	-	1	1	-	2	1	1	-	-	1	1	-
Male	-	1	1	-	1	1	1	-	-	1	-	-
Female	-	-	-	-	1	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	4	7	10	9	10	8	12	5	1	-	-
Male	2	3	5	7	6	7	3	6	-	-	-	-
Female	1	1	2	3	3	3	5	6	5	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Rhode Island												
All races, both sexes	58	65	91	111	131	273	368	469	369	164	39	-
Male	44	50	57	72	73	153	197	191	113	27	9	-
Female	14	15	34	39	58	120	171	278	256	137	30	-
White, both sexes	54	60	87	104	124	260	361	466	365	162	39	-
Male	42	47	55	69	70	148	194	190	113	26	9	-
Female	12	13	32	35	54	112	167	276	252	136	30	-
Black, both sexes	2	4	2	7	6	8	5	3	3	1	-	-
Male	1	2	1	3	3	4	2	1	-	-	-	-
Female	1	2	1	4	3	4	3	2	3	1	-	-
American Indian 1/, both sexes	2	-	-	-	1	1	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	-	4	1	-	-	1	-	-
Male	-	1	1	-	-	1	1	-	-	1	-	-
Female	-	-	1	-	-	3	-	-	-	-	-	-
South Carolina												
All races, both sexes	293	421	467	468	579	711	809	740	390	139	33	4
Male	217	304	317	322	371	444	418	327	143	27	3	4
Female	76	117	150	146	208	267	391	413	247	112	30	-
White, both sexes	201	287	317	343	419	529	634	591	323	110	16	2
Male	151	210	219	247	281	345	345	273	122	22	3	2
Female	50	77	98	96	138	184	289	318	201	88	13	-
Black, both sexes	91	130	148	123	158	180	175	147	67	29	17	2
Male	66	90	97	74	89	98	73	53	21	5	-	2
Female	25	40	51	49	69	82	102	94	46	24	17	-
American Indian 1/, both sexes	-	3	2	1	1	2	-	-	-	-	-	-
Male	-	3	1	-	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	-	-	2	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
South Dakota												
All races, both sexes	41	65	75	77	93	175	227	245	212	109	24	-
Male	31	50	56	52	61	110	128	112	74	23	5	-
Female	10	15	19	25	32	65	99	133	138	86	19	-
White, both sexes	35	56	69	70	87	167	217	241	210	107	24	-
Male	25	45	54	49	59	104	124	109	74	23	5	-
Female	10	11	15	21	28	63	93	132	136	84	19	-
Black, both sexes	-	-	-	1	-	2	2	-	-	-	-	-
Male	-	-	-	1	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	6	9	5	6	6	6	8	4	2	2	-	-
Male	6	5	1	2	2	4	3	3	-	-	-	-
Female	-	4	4	4	4	2	5	1	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	471	685	803	828	1,094	1,342	1,629	1,536	1,076	390	94	-
Male	343	518	554	550	693	759	790	629	335	86	16	-
Female	128	167	249	278	401	583	839	907	741	304	78	-
White, both sexes	363	563	689	712	948	1,169	1,458	1,412	956	353	75	-
Male	268	437	483	477	613	670	727	588	303	78	15	-
Female	95	126	206	235	335	499	731	824	653	275	60	-
Black, both sexes	106	120	111	111	142	171	168	121	116	37	19	-
Male	73	80	68	69	78	88	61	40	31	8	1	-
Female	33	40	43	42	64	83	107	81	85	29	18	-
American Indian 1/, both sexes	-	1	1	2	-	2	1	-	1	-	-	-
Male	-	1	1	2	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	3	4	-	2	3	3	-	-	-
Male	2	-	2	2	2	-	1	1	1	-	-	-
Female	-	1	-	1	2	-	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Texas												
All races, both sexes	1,121	1,643	1,844	2,063	2,511	3,436	4,134	3,824	2,694	1,064	234	1
Male	852	1,190	1,268	1,346	1,548	1,978	2,075	1,607	951	279	44	-
Female	269	453	576	717	963	1,458	2,059	2,217	1,743	785	190	1
White, both sexes	903	1,341	1,551	1,753	2,186	3,057	3,753	3,539	2,508	954	203	1
Male	697	980	1,089	1,166	1,361	1,777	1,892	1,508	890	255	36	-
Female	206	361	462	587	825	1,280	1,861	2,031	1,618	699	167	1
Black, both sexes	206	269	273	283	299	349	349	257	172	107	30	-
Male	145	185	165	164	177	183	171	88	55	24	8	-
Female	61	84	108	119	122	166	178	169	117	83	22	-
American Indian 1/, both sexes	1	7	2	6	4	3	7	5	2	1	-	-
Male	-	4	1	4	2	2	2	-	-	-	-	-
Female	1	3	1	2	2	1	5	5	2	1	-	-
Asian or Pacific Islander 2/, both sexes	11	26	18	21	22	27	25	23	12	2	1	-
Male	10	21	13	12	8	16	10	11	6	-	-	-
Female	1	5	5	9	14	11	15	12	6	2	1	-
Utah												
All races, both sexes	56	79	93	117	117	192	245	268	161	61	9	-
Male	48	66	70	95	76	128	152	145	72	23	1	-
Female	8	13	23	22	41	64	93	123	89	38	8	-
White, both sexes	52	77	91	112	113	188	243	267	159	61	9	-
Male	45	65	68	92	73	125	151	145	71	23	1	-
Female	7	12	23	20	40	63	92	122	88	38	8	-
Black, both sexes	2	1	-	-	-	1	1	-	1	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	1	1	1	1	1	2	-	-	-	-	-	-
Male	1	-	1	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	4	3	1	1	1	1	-	-	-
Male	1	-	1	2	3	1	1	-	1	-	-	-
Female	-	-	-	2	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Vermont												
All races, both sexes	25	44	43	58	74	126	137	161	114	49	14	-
Male	20	36	32	45	44	70	76	72	37	9	3	-
Female	5	8	11	13	30	56	61	89	77	40	11	-
White, both sexes	25	44	43	58	73	126	137	161	114	49	14	-
Male	20	36	32	45	43	70	76	72	37	9	3	-
Female	5	8	11	13	30	56	61	89	77	40	11	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	349	517	608	635	815	1,110	1,367	1,362	845	339	65	-
Male	264	382	442	444	495	675	671	587	277	74	7	-
Female	85	135	166	191	320	435	696	775	568	265	58	-
White, both sexes	258	384	466	475	647	898	1,135	1,140	731	283	55	-
Male	200	290	340	343	404	562	574	493	244	64	4	-
Female	58	94	126	132	243	336	561	647	487	219	51	-
Black, both sexes	86	122	135	149	158	196	216	206	111	55	10	-
Male	60	82	97	94	85	104	91	90	32	10	3	-
Female	26	40	38	55	73	92	125	116	79	45	7	-
American Indian 1/, both sexes	-	1	-	-	-	1	3	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	10	7	11	10	15	13	16	3	1	-	-
Male	4	9	5	7	6	8	6	4	1	-	-	-
Female	1	1	2	4	4	7	7	12	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Washington												
All races, both sexes	232	306	436	495	644	911	1,207	1,280	1,022	419	81	-
Male	186	233	312	340	432	538	653	599	379	124	20	-
Female	46	73	124	155	212	373	554	681	643	295	61	-
White, both sexes	209	278	385	444	605	849	1,158	1,233	991	406	78	-
Male	170	213	279	304	408	508	631	576	362	119	19	-
Female	39	65	106	140	197	341	527	657	629	287	59	-
Black, both sexes	10	13	23	19	9	21	11	11	10	3	-	-
Male	8	10	14	14	6	9	6	3	4	-	-	-
Female	2	3	9	5	3	12	5	8	6	3	-	-
American Indian 1/, both sexes	3	5	9	5	12	13	13	6	4	-	1	-
Male	2	3	4	3	7	7	3	2	3	-	1	-
Female	1	2	5	2	5	6	10	4	1	-	-	-
Asian or Pacific Islander 2/, both sexes	10	10	19	27	18	28	25	30	17	10	2	-
Male	6	7	15	19	11	14	13	18	10	5	-	-
Female	4	3	4	8	7	14	12	12	7	5	2	-
West Virginia												
All races, both sexes	159	187	200	309	357	493	602	529	374	136	23	-
Male	128	143	140	208	212	288	298	231	117	28	2	-
Female	31	44	60	101	145	205	304	298	257	108	21	-
White, both sexes	153	178	191	306	345	482	579	520	363	132	20	-
Male	123	136	135	205	204	281	288	229	114	28	2	-
Female	30	42	56	101	141	201	291	291	249	104	18	-
Black, both sexes	5	8	9	3	12	11	21	9	11	4	3	-
Male	4	6	5	3	8	7	9	2	3	-	-	-
Female	1	2	4	-	4	4	12	7	8	4	3	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Wisconsin												
All races, both sexes	242	320	371	440	638	901	1,220	1,281	997	423	105	-
Male	186	250	279	319	424	554	665	613	354	97	16	-
Female	56	70	92	121	214	347	555	668	643	326	89	-
White, both sexes	216	289	344	412	610	863	1,191	1,259	983	417	104	-
Male	172	232	262	303	407	537	655	605	351	95	16	-
Female	44	57	82	109	203	326	536	654	632	322	88	-
Black, both sexes	20	24	22	22	23	30	23	19	11	4	-	-
Male	10	13	14	11	14	14	10	8	3	1	-	-
Female	10	11	8	11	9	16	13	11	8	3	-	-
American Indian 1/, both sexes	6	6	4	3	4	6	6	2	2	2	-	-
Male	4	4	2	3	3	3	-	-	-	1	-	-
Female	2	2	2	-	1	3	6	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	1	2	-	1	1	-	1	-
Male	-	1	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	2	-	1	1	-	1	-
Wyoming												
All races, both sexes	22	30	39	54	64	76	80	79	58	29	4	-
Male	17	21	28	37	40	51	50	38	21	5	-	-
Female	5	9	11	17	24	25	30	41	37	24	4	-
White, both sexes	22	29	37	54	63	76	80	78	58	28	4	-
Male	17	20	26	37	39	51	50	38	21	4	-	-
Female	5	9	11	17	24	25	30	40	37	24	4	-
Black, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	1	-	-
Male	-	1	1	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
United States												
All races, both sexes	141,462	11	5	1	5	20	40	124	298	757	1,919	3,761
Male	76,089	8	4	1	2	15	27	90	225	558	1,453	2,787
Female	65,373	3	1	-	3	5	13	34	73	199	466	974
White, both sexes	123,549	10	3	1	3	16	33	92	224	595	1,545	3,014
Male	66,946	8	2	1	2	14	24	67	173	453	1,198	2,299
Female	56,603	2	1	-	1	2	9	25	51	142	347	715
Black, both sexes	14,776	1	2	-	2	4	7	29	63	146	325	652
Male	7,392	-	2	-	-	1	3	21	43	91	217	414
Female	7,384	1	-	-	2	3	4	8	20	55	108	238
American Indian 1/, both sexes	665	-	-	-	-	-	-	3	4	3	23	31
Male	371	-	-	-	-	-	-	2	3	2	15	24
Female	294	-	-	-	-	-	-	1	1	1	8	7
Asian or Pacific Islander 2/, both sexes	2,472	-	-	-	-	-	-	-	7	13	26	64
Male	1,380	-	-	-	-	-	-	-	6	12	23	50
Female	1,092	-	-	-	-	-	-	-	1	1	3	14
Alabama												
All races, both sexes	2,645	-	-	-	-	-	3	8	15	31	75	116
Male	1,436	-	-	-	-	-	2	6	12	19	50	85
Female	1,209	-	-	-	-	-	1	2	3	12	25	31
White, both sexes	2,089	-	-	-	-	-	2	7	8	23	54	86
Male	1,140	-	-	-	-	-	1	6	6	14	39	64
Female	949	-	-	-	-	-	1	1	2	9	15	22
Black, both sexes	546	-	-	-	-	-	1	1	7	8	21	28
Male	292	-	-	-	-	-	1	-	6	5	11	20
Female	254	-	-	-	-	-	-	1	1	3	10	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Alaska												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	1	3
Male	52	-	-	-	-	-	-	-	-	-	1	2
Female	31	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	62	-	-	-	-	-	-	-	-	-	1	2
Male	40	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,864	-	-	-	-	-	-	2	2	2	17	28
Male	1,017	-	-	-	-	-	-	2	2	2	13	24
Female	847	-	-	-	-	-	-	-	-	-	4	4
White, both sexes	1,756	-	-	-	-	-	-	1	1	1	13	24
Male	948	-	-	-	-	-	-	1	1	1	10	21
Female	808	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	43	-	-	-	-	-	-	-	-	1	-	3
Male	27	-	-	-	-	-	-	-	-	1	-	3
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	59	-	-	-	-	-	-	1	1	-	4	1
Male	38	-	-	-	-	-	-	1	1	-	3	-
Female	21	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Arkansas												
All races, both sexes	3,068	1	-	1	-	2	2	6	17	32	70	119
Male	1,683	1	-	1	-	1	2	4	12	20	52	77
Female	1,385	-	-	-	-	1	-	2	5	12	18	42
White, both sexes	2,653	1	-	1	-	1	2	5	15	24	57	89
Male	1,464	1	-	1	-	1	2	3	11	18	43	60
Female	1,189	-	-	-	-	-	-	2	4	6	14	29
Black, both sexes	399	-	-	-	-	1	-	1	2	8	13	29
Male	210	-	-	-	-	-	-	1	1	2	9	16
Female	189	-	-	-	-	1	-	-	1	6	4	13
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13,147	-	1	-	-	-	2	6	22	39	103	251
Male	6,921	-	1	-	-	-	1	4	15	28	78	190
Female	6,226	-	-	-	-	-	1	2	7	11	25	61
White, both sexes	10,881	-	1	-	-	-	1	5	16	31	79	193
Male	5,770	-	1	-	-	-	1	4	11	22	59	150
Female	5,111	-	-	-	-	-	-	1	5	9	20	43
Black, both sexes	1,046	-	-	-	-	-	1	1	3	6	15	32
Male	500	-	-	-	-	-	-	-	2	4	11	20
Female	546	-	-	-	-	-	1	1	1	2	4	12
American Indian 1/, both sexes	63	-	-	-	-	-	-	-	-	-	1	1
Male	35	-	-	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1,157	-	-	-	-	-	-	-	3	2	8	25
Male	616	-	-	-	-	-	-	-	2	2	7	19
Female	541	-	-	-	-	-	-	-	1	-	1	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Colorado												
All races, both sexes	1,235	-	-	-	-	-	-	1	2	5	23	40
Male	691	-	-	-	-	-	-	1	2	4	17	34
Female	544	-	-	-	-	-	-	-	-	1	6	6
White, both sexes	1,203	-	-	-	-	-	-	-	2	5	23	36
Male	671	-	-	-	-	-	-	-	2	4	17	31
Female	532	-	-	-	-	-	-	-	-	1	6	5
Black, both sexes	23	-	-	-	-	-	-	1	-	-	-	3
Male	15	-	-	-	-	-	-	1	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,283	-	-	-	-	-	1	2	1	4	11	32
Male	627	-	-	-	-	-	1	2	-	4	10	25
Female	656	-	-	-	-	-	-	-	1	-	1	7
White, both sexes	1,203	-	-	-	-	-	1	2	-	3	8	25
Male	584	-	-	-	-	-	1	2	-	3	7	19
Female	619	-	-	-	-	-	-	-	-	-	1	6
Black, both sexes	68	-	-	-	-	-	-	-	1	1	3	7
Male	37	-	-	-	-	-	-	-	-	1	3	6
Female	31	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Delaware												
All races, both sexes	362	-	-	-	-	-	-	-	-	-	3	12
Male	204	-	-	-	-	-	-	-	-	-	3	10
Female	158	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	306	-	-	-	-	-	-	-	-	-	2	8
Male	176	-	-	-	-	-	-	-	-	-	2	7
Female	130	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	55	-	-	-	-	-	-	-	-	-	1	4
Male	27	-	-	-	-	-	-	-	-	-	1	3
Female	28	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	236	-	-	-	-	-	-	-	-	1	-	6
Male	109	-	-	-	-	-	-	-	-	1	-	5
Female	127	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	177	-	-	-	-	-	-	-	-	1	-	5
Male	71	-	-	-	-	-	-	-	-	1	-	4
Female	106	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Florida												
All races, both sexes	8,740	1	-	-	-	1	-	4	9	41	115	172
Male	4,684	1	-	-	-	1	-	2	7	26	91	131
Female	4,056	-	-	-	-	-	-	2	2	15	24	41
White, both sexes	7,928	1	-	-	-	1	-	2	6	35	97	140
Male	4,294	1	-	-	-	1	-	1	4	23	79	111
Female	3,634	-	-	-	-	-	-	1	2	12	18	29
Black, both sexes	753	-	-	-	-	-	-	2	2	6	18	30
Male	360	-	-	-	-	-	-	1	2	3	12	19
Female	393	-	-	-	-	-	-	1	-	3	6	11
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	-	-	-	-	-	-	1	-	-	2
Male	26	-	-	-	-	-	-	-	1	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	3,170	-	-	-	1	1	2	3	9	23	62	123
Male	1,773	-	-	-	-	1	2	1	7	17	46	90
Female	1,397	-	-	-	1	-	-	2	2	6	16	33
White, both sexes	2,407	-	-	-	-	1	1	2	7	14	44	82
Male	1,380	-	-	-	-	1	1	1	6	10	34	61
Female	1,027	-	-	-	-	-	-	1	1	4	10	21
Black, both sexes	726	-	-	-	1	-	1	1	2	7	18	38
Male	373	-	-	-	-	-	1	-	1	5	12	27
Female	353	-	-	-	1	-	-	1	1	2	6	11
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	2	-	2
Male	15	-	-	-	-	-	-	-	-	2	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Hawaii												
All races, both sexes	435	-	-	-	-	-	1	-	-	2	2	9
Male	255	-	-	-	-	-	1	-	-	2	1	7
Female	180	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	112	-	-	-	-	-	1	-	-	1	-	2
Male	65	-	-	-	-	-	1	-	-	1	-	2
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	320	-	-	-	-	-	-	-	-	1	2	7
Male	188	-	-	-	-	-	-	-	-	1	1	5
Female	132	-	-	-	-	-	-	-	-	-	1	2
Idaho												
All races, both sexes	807	-	-	-	-	1	-	-	2	4	16	34
Male	498	-	-	-	-	1	-	-	2	2	12	25
Female	309	-	-	-	-	-	-	-	-	2	4	9
White, both sexes	787	-	-	-	-	1	-	-	2	4	15	34
Male	487	-	-	-	-	1	-	-	2	2	12	25
Female	300	-	-	-	-	-	-	-	-	2	3	9
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Illinois												
All races, both sexes	6,855	-	-	-	-	2	3	5	18	31	80	183
Male	3,605	-	-	-	-	1	-	2	15	22	62	134
Female	3,250	-	-	-	-	1	3	3	3	9	18	49
White, both sexes	5,881	-	-	-	-	2	2	5	14	20	66	148
Male	3,107	-	-	-	-	1	-	2	11	15	52	115
Female	2,774	-	-	-	-	1	2	3	3	5	14	33
Black, both sexes	873	-	-	-	-	-	1	-	4	11	12	33
Male	446	-	-	-	-	-	-	-	4	7	8	17
Female	427	-	-	-	-	-	1	-	-	4	4	16
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	93	-	-	-	-	-	-	-	-	-	2	2
Male	48	-	-	-	-	-	-	-	-	-	2	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3,532	-	-	-	-	-	-	2	11	23	54	110
Male	1,972	-	-	-	-	-	-	2	8	16	46	83
Female	1,560	-	-	-	-	-	-	-	3	7	8	27
White, both sexes	3,321	-	-	-	-	-	-	2	9	19	50	102
Male	1,855	-	-	-	-	-	-	2	7	13	42	80
Female	1,466	-	-	-	-	-	-	-	2	6	8	22
Black, both sexes	205	-	-	-	-	-	-	-	2	4	4	8
Male	113	-	-	-	-	-	-	-	1	3	4	3
Female	92	-	-	-	-	-	-	-	1	1	-	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Iowa												
All races, both sexes	1,766	-	1	-	-	-	-	4	2	5	15	36
Male	945	-	1	-	-	-	-	4	2	4	10	31
Female	821	-	-	-	-	-	-	-	-	1	5	5
White, both sexes	1,739	-	-	-	-	-	-	4	2	3	15	35
Male	930	-	-	-	-	-	-	4	2	3	10	30
Female	809	-	-	-	-	-	-	-	-	-	5	5
Black, both sexes	23	-	1	-	-	-	-	-	-	1	-	1
Male	13	-	1	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,241	-	-	-	-	-	1	-	3	3	16	24
Male	662	-	-	-	-	-	-	-	3	1	13	18
Female	579	-	-	-	-	-	1	-	-	2	3	6
White, both sexes	1,178	-	-	-	-	-	1	-	3	3	15	19
Male	627	-	-	-	-	-	-	-	3	1	12	14
Female	551	-	-	-	-	-	1	-	-	2	3	5
Black, both sexes	38	-	-	-	-	-	-	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Kentucky												
All races, both sexes	3,419	1	-	-	-	2	2	8	15	39	80	138
Male	1,951	-	-	-	-	1	1	4	12	30	63	109
Female	1,468	1	-	-	-	1	1	4	3	9	17	29
White, both sexes	3,190	1	-	-	-	2	2	6	13	37	67	126
Male	1,811	-	-	-	-	1	1	3	10	29	53	101
Female	1,379	1	-	-	-	1	1	3	3	8	14	25
Black, both sexes	220	-	-	-	-	-	-	2	2	2	11	11
Male	135	-	-	-	-	-	-	1	2	1	8	7
Female	85	-	-	-	-	-	-	1	-	1	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	2	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2,727	-	-	-	-	-	2	5	14	21	75	123
Male	1,597	-	-	-	-	-	2	5	7	15	56	88
Female	1,130	-	-	-	-	-	-	-	7	6	19	35
White, both sexes	1,879	-	-	-	-	-	1	3	9	13	44	75
Male	1,136	-	-	-	-	-	1	3	4	10	36	57
Female	743	-	-	-	-	-	-	-	5	3	8	18
Black, both sexes	827	-	-	-	-	-	1	2	5	8	30	47
Male	447	-	-	-	-	-	1	2	3	5	19	30
Female	380	-	-	-	-	-	-	-	2	3	11	17
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Maine												
All races, both sexes	654	-	-	-	-	-	-	-	2	2	6	20
Male	357	-	-	-	-	-	-	-	2	2	5	17
Female	297	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	652	-	-	-	-	-	-	-	2	2	6	20
Male	356	-	-	-	-	-	-	-	2	2	5	17
Female	296	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,462	-	-	-	-	-	-	3	5	13	23	56
Male	1,236	-	-	-	-	-	-	1	3	12	16	40
Female	1,226	-	-	-	-	-	-	2	2	1	7	16
White, both sexes	1,812	-	-	-	-	-	-	2	1	5	13	34
Male	905	-	-	-	-	-	-	1	1	5	8	25
Female	907	-	-	-	-	-	-	1	-	-	5	9
Black, both sexes	602	-	-	-	-	-	-	-	4	8	10	20
Male	305	-	-	-	-	-	-	-	2	7	8	14
Female	297	-	-	-	-	-	-	-	2	1	2	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	46	-	-	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Massachusetts												
All races, both sexes	2,535	-	-	-	-	-	-	-	-	7	15	47
Male	1,287	-	-	-	-	-	-	-	-	7	12	35
Female	1,248	-	-	-	-	-	-	-	-	-	3	12
White, both sexes	2,421	-	-	-	-	-	-	-	-	6	13	43
Male	1,220	-	-	-	-	-	-	-	-	6	10	32
Female	1,201	-	-	-	-	-	-	-	-	-	3	11
Black, both sexes	80	-	-	-	-	-	-	-	-	1	1	3
Male	43	-	-	-	-	-	-	-	-	1	1	2
Female	37	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4,790	-	-	-	1	1	-	-	10	24	52	128
Male	2,466	-	-	-	-	1	-	-	3	18	47	94
Female	2,324	-	-	-	1	-	-	-	7	6	5	34
White, both sexes	4,244	-	-	-	-	1	-	-	6	18	47	105
Male	2,215	-	-	-	-	1	-	-	2	12	43	78
Female	2,029	-	-	-	-	-	-	-	4	6	4	27
Black, both sexes	491	-	-	-	1	-	-	-	3	6	5	22
Male	226	-	-	-	-	-	-	-	1	6	4	16
Female	265	-	-	-	1	-	-	-	2	-	1	6
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	1	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Minnesota												
All races, both sexes	1,308	-	-	-	-	-	-	-	-	8	21	33
Male	740	-	-	-	-	-	-	-	-	7	15	28
Female	568	-	-	-	-	-	-	-	-	1	6	5
White, both sexes	1,256	-	-	-	-	-	-	-	-	6	18	27
Male	707	-	-	-	-	-	-	-	-	5	14	24
Female	549	-	-	-	-	-	-	-	-	1	4	3
Black, both sexes	13	-	-	-	-	-	-	-	-	1	-	3
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	2	3
Male	19	-	-	-	-	-	-	-	-	-	1	3
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	1	1	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Mississippi												
All races, both sexes	1,933	-	-	-	-	1	-	6	10	22	48	91
Male	1,103	-	-	-	-	-	-	4	8	13	34	59
Female	830	-	-	-	-	1	-	2	2	9	14	32
White, both sexes	1,294	-	-	-	-	-	-	3	5	15	25	51
Male	743	-	-	-	-	-	-	2	5	10	18	37
Female	551	-	-	-	-	-	-	1	-	5	7	14
Black, both sexes	633	-	-	-	-	1	-	3	5	7	23	40
Male	357	-	-	-	-	-	-	2	3	3	16	22
Female	276	-	-	-	-	1	-	1	2	4	7	18
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Missouri												
All races, both sexes	4,537	-	-	-	1	1	1	7	12	29	64	141
Male	2,468	-	-	-	1	1	1	5	10	22	46	111
Female	2,069	-	-	-	-	-	-	2	2	7	18	30
White, both sexes	4,129	-	-	-	1	-	1	5	11	26	54	134
Male	2,269	-	-	-	1	-	1	3	9	20	38	106
Female	1,860	-	-	-	-	-	-	2	2	6	16	28
Black, both sexes	383	-	-	-	-	1	-	1	1	3	9	6
Male	185	-	-	-	-	1	-	1	1	2	7	4
Female	198	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	10	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	359	-	-	-	-	-	-	-	-	3	3	16
Male	208	-	-	-	-	-	-	-	-	3	3	12
Female	151	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	344	-	-	-	-	-	-	-	-	3	3	15
Male	198	-	-	-	-	-	-	-	-	3	3	11
Female	146	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Nebraska												
All races, both sexes	543	-	-	-	-	-	-	-	2	1	12	13
Male	302	-	-	-	-	-	-	-	1	1	12	11
Female	241	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	530	-	-	-	-	-	-	-	2	1	11	13
Male	294	-	-	-	-	-	-	-	1	1	11	11
Female	236	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	578	-	-	-	-	-	-	-	3	5	10	14
Male	362	-	-	-	-	-	-	-	3	4	9	9
Female	216	-	-	-	-	-	-	-	-	1	1	5
White, both sexes	529	-	-	-	-	-	-	-	3	5	9	13
Male	336	-	-	-	-	-	-	-	3	4	8	9
Female	193	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
New Hampshire												
All races, both sexes	449	-	-	-	-	-	-	-	1	1	7	15
Male	248	-	-	-	-	-	-	-	1	1	5	13
Female	201	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	446	-	-	-	-	-	-	-	1	1	7	15
Male	248	-	-	-	-	-	-	-	1	1	5	13
Female	198	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,163	1	-	-	-	1	2	2	2	15	37	75
Male	2,099	1	-	-	-	1	1	1	2	10	32	50
Female	2,064	-	-	-	-	-	1	1	-	5	5	25
White, both sexes	3,554	1	-	-	-	1	1	1	1	12	21	52
Male	1,798	1	-	-	-	1	1	-	1	8	20	40
Female	1,756	-	-	-	-	-	-	1	-	4	1	12
Black, both sexes	520	-	-	-	-	-	1	1	1	2	12	22
Male	239	-	-	-	-	-	-	1	1	1	8	10
Female	281	-	-	-	-	-	1	-	-	1	4	12
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	84	-	-	-	-	-	-	-	-	1	3	1
Male	59	-	-	-	-	-	-	-	-	1	3	-
Female	25	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
New Mexico												
All races, both sexes	511	1	-	-	-	-	-	1	-	1	6	7
Male	308	1	-	-	-	-	-	1	-	1	3	2
Female	203	-	-	-	-	-	-	-	-	-	3	5
White, both sexes	482	1	-	-	-	-	-	1	-	1	6	7
Male	293	1	-	-	-	-	-	1	-	1	3	2
Female	189	-	-	-	-	-	-	-	-	-	3	5
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	9,427	1	1	-	-	2	2	7	10	38	94	157
Male	4,551	1	1	-	-	1	1	5	9	28	66	119
Female	4,876	-	-	-	-	1	1	2	1	10	28	38
White, both sexes	8,100	1	-	-	-	1	2	3	6	25	74	118
Male	3,960	1	-	-	-	1	1	2	6	18	54	92
Female	4,140	-	-	-	-	-	1	1	-	7	20	26
Black, both sexes	1,133	-	1	-	-	1	-	4	2	10	20	32
Male	482	-	1	-	-	-	-	3	1	7	12	21
Female	651	-	-	-	-	1	-	1	1	3	8	11
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	175	-	-	-	-	-	-	-	1	3	-	6
Male	100	-	-	-	-	-	-	-	1	3	-	5
Female	75	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
North Carolina												
All races, both sexes	3,784	1	-	-	-	-	1	1	3	22	61	104
Male	2,069	-	-	-	-	-	-	1	3	18	45	76
Female	1,715	1	-	-	-	-	1	-	-	4	16	28
White, both sexes	3,020	-	-	-	-	-	1	1	2	16	50	66
Male	1,693	-	-	-	-	-	-	1	2	15	38	53
Female	1,327	-	-	-	-	-	1	-	-	1	12	13
Black, both sexes	711	1	-	-	-	-	-	-	1	6	10	34
Male	344	-	-	-	-	-	-	-	1	3	6	19
Female	367	1	-	-	-	-	-	-	-	3	4	15
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-	-	-	4
Male	25	-	-	-	-	-	-	-	-	-	-	4
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	465	-	-	-	-	-	-	1	-	2	4	7
Male	270	-	-	-	-	-	-	1	-	2	2	5
Female	195	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	451	-	-	-	-	-	-	1	-	2	4	7
Male	260	-	-	-	-	-	-	1	-	2	2	5
Female	191	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Ohio												
All races, both sexes	6,384	1	-	-	-	1	4	9	11	36	82	178
Male	3,433	1	-	-	-	1	4	6	8	24	65	138
Female	2,951	-	-	-	-	-	-	3	3	12	17	40
White, both sexes	5,836	1	-	-	-	1	4	8	10	30	72	150
Male	3,151	1	-	-	-	1	4	5	7	22	57	119
Female	2,685	-	-	-	-	-	-	3	3	8	15	31
Black, both sexes	522	-	-	-	-	-	-	1	1	6	9	27
Male	263	-	-	-	-	-	-	1	1	2	7	18
Female	259	-	-	-	-	-	-	-	-	4	2	9
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,809	-	-	-	-	-	1	2	5	8	29	39
Male	993	-	-	-	-	-	1	1	4	7	19	30
Female	816	-	-	-	-	-	-	1	1	1	10	9
White, both sexes	1,593	-	-	-	-	-	1	2	5	7	23	34
Male	882	-	-	-	-	-	1	1	4	6	17	26
Female	711	-	-	-	-	-	-	1	1	1	6	8
Black, both sexes	102	-	-	-	-	-	-	-	-	-	3	2
Male	53	-	-	-	-	-	-	-	-	-	-	1
Female	49	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	107	-	-	-	-	-	-	-	-	1	3	3
Male	56	-	-	-	-	-	-	-	-	1	2	3
Female	51	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Oregon												
All races, both sexes	1,311	1	-	-	-	-	-	-	2	3	7	24
Male	771	1	-	-	-	-	-	-	2	2	4	16
Female	540	-	-	-	-	-	-	-	-	1	3	8
White, both sexes	1,267	1	-	-	-	-	-	-	2	3	6	21
Male	742	1	-	-	-	-	-	-	2	2	4	13
Female	525	-	-	-	-	-	-	-	-	1	2	8
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	10	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	7,715	-	-	-	-	-	1	4	9	36	87	174
Male	4,124	-	-	-	-	-	-	3	8	31	65	138
Female	3,591	-	-	-	-	-	1	1	1	5	22	36
White, both sexes	7,204	-	-	-	-	-	1	3	6	33	75	160
Male	3,857	-	-	-	-	-	-	2	6	29	57	126
Female	3,347	-	-	-	-	-	1	1	-	4	18	34
Black, both sexes	481	-	-	-	-	-	-	1	2	3	11	14
Male	247	-	-	-	-	-	-	1	1	2	7	12
Female	234	-	-	-	-	-	-	-	1	1	4	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	1	-	1	-
Male	17	-	-	-	-	-	-	-	1	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Rhode Island												
All races, both sexes	758	-	-	-	-	-	-	1	-	1	4	13
Male	387	-	-	-	-	-	-	1	-	1	2	9
Female	371	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	731	-	-	-	-	-	-	1	-	1	3	13
Male	376	-	-	-	-	-	-	1	-	1	2	9
Female	355	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,350	-	-	-	1	-	1	1	12	14	43	92
Male	1,362	-	-	-	1	-	1	1	10	11	35	76
Female	988	-	-	-	-	-	-	-	2	3	8	16
White, both sexes	1,686	-	-	-	1	-	1	1	8	11	28	58
Male	1,012	-	-	-	1	-	1	1	7	9	25	52
Female	674	-	-	-	-	-	-	-	1	2	3	6
Black, both sexes	655	-	-	-	-	-	-	-	3	3	15	34
Male	344	-	-	-	-	-	-	-	2	2	10	24
Female	311	-	-	-	-	-	-	-	1	1	5	10
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
South Dakota												
All races, both sexes	634	-	-	-	-	-	1	2	-	5	9	16
Male	359	-	-	-	-	-	1	2	-	3	5	13
Female	275	-	-	-	-	-	-	-	-	2	4	3
White, both sexes	593	-	-	-	-	-	1	1	-	4	4	14
Male	337	-	-	-	-	-	1	1	-	3	2	12
Female	256	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	5	-	-	-	-	-	-	1	-	-	1	-
Male	4	-	-	-	-	-	-	1	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	35	-	-	-	-	-	-	-	-	1	4	2
Male	17	-	-	-	-	-	-	-	-	-	2	1
Female	18	-	-	-	-	-	-	-	-	1	2	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4,381	1	1	-	-	-	-	4	15	34	87	152
Male	2,428	1	-	-	-	-	-	4	12	30	66	111
Female	1,953	-	1	-	-	-	-	-	3	4	21	41
White, both sexes	3,825	1	1	-	-	-	-	2	12	27	75	124
Male	2,160	1	-	-	-	-	-	2	10	25	58	94
Female	1,665	-	1	-	-	-	-	-	2	2	17	30
Black, both sexes	545	-	-	-	-	-	-	2	3	7	12	26
Male	259	-	-	-	-	-	-	2	2	5	8	15
Female	286	-	-	-	-	-	-	-	1	2	4	11
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Texas												
All races, both sexes	10,678	-	1	-	1	1	4	7	27	67	153	345
Male	5,716	-	1	-	-	1	3	6	19	49	113	226
Female	4,962	-	-	-	1	-	1	1	8	18	40	119
White, both sexes	9,458	-	1	-	1	1	4	5	21	55	125	285
Male	5,089	-	1	-	-	1	3	4	14	40	93	186
Female	4,369	-	-	-	1	-	1	1	7	15	32	99
Black, both sexes	1,121	-	-	-	-	-	-	2	5	11	25	56
Male	570	-	-	-	-	-	-	2	4	8	17	36
Female	551	-	-	-	-	-	-	-	1	3	8	20
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	81	-	-	-	-	-	-	-	-	1	2	3
Male	50	-	-	-	-	-	-	-	-	1	2	3
Female	31	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	529	-	-	-	-	-	-	-	1	2	10	12
Male	325	-	-	-	-	-	-	-	1	1	6	9
Female	204	-	-	-	-	-	-	-	-	1	4	3
White, both sexes	518	-	-	-	-	-	-	-	1	2	10	12
Male	319	-	-	-	-	-	-	-	1	1	6	9
Female	199	-	-	-	-	-	-	-	-	1	4	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Vermont												
All races, both sexes	251	-	-	-	-	-	-	-	-	-	2	5
Male	138	-	-	-	-	-	-	-	-	-	2	5
Female	113	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	251	-	-	-	-	-	-	-	-	-	2	5
Male	138	-	-	-	-	-	-	-	-	-	2	5
Female	113	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,040	1	-	-	-	1	-	2	5	18	48	79
Male	1,665	-	-	-	-	1	-	2	5	12	39	56
Female	1,375	1	-	-	-	-	-	-	-	6	9	23
White, both sexes	2,443	1	-	-	-	1	-	1	5	10	39	55
Male	1,368	-	-	-	-	1	-	1	5	9	34	41
Female	1,075	1	-	-	-	-	-	-	-	1	5	14
Black, both sexes	556	-	-	-	-	-	-	1	-	8	9	23
Male	277	-	-	-	-	-	-	1	-	3	5	14
Female	279	-	-	-	-	-	-	-	-	5	4	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Washington												
All races, both sexes	2,155	-	-	-	-	-	-	3	2	12	21	48
Male	1,204	-	-	-	-	-	-	3	-	8	16	37
Female	951	-	-	-	-	-	-	-	2	4	5	11
White, both sexes	2,017	-	-	-	-	-	-	3	2	11	18	41
Male	1,127	-	-	-	-	-	-	3	-	8	14	31
Female	890	-	-	-	-	-	-	-	2	3	4	10
Black, both sexes	39	-	-	-	-	-	-	-	-	-	1	3
Male	26	-	-	-	-	-	-	-	-	-	1	3
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	72	-	-	-	-	-	-	-	-	1	1	3
Male	38	-	-	-	-	-	-	-	-	-	1	2
Female	34	-	-	-	-	-	-	-	-	1	-	1
West Virginia												
All races, both sexes	1,173	-	-	-	-	-	1	3	2	6	19	29
Male	638	-	-	-	-	-	1	2	1	4	12	22
Female	535	-	-	-	-	-	-	1	1	2	7	7
White, both sexes	1,136	-	-	-	-	-	1	3	2	6	19	27
Male	618	-	-	-	-	-	1	2	1	4	12	20
Female	518	-	-	-	-	-	-	1	1	2	7	7
Black, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Wisconsin												
All races, both sexes	2,933	-	-	-	-	2	2	2	4	14	34	64
Male	1,652	-	-	-	-	2	1	1	3	11	26	47
Female	1,281	-	-	-	-	-	1	1	1	3	8	17
White, both sexes	2,854	-	-	-	-	2	1	1	2	14	32	60
Male	1,621	-	-	-	-	2	1	1	2	11	25	45
Female	1,233	-	-	-	-	-	-	-	-	3	7	15
Black, both sexes	48	-	-	-	-	-	1	1	2	-	1	3
Male	18	-	-	-	-	-	-	-	1	-	-	2
Female	30	-	-	-	-	-	1	1	1	-	1	1
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	248	-	-	-	-	-	-	-	1	2	3	8
Male	138	-	-	-	-	-	-	-	1	2	2	5
Female	110	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	243	-	-	-	-	-	-	-	1	1	3	8
Male	134	-	-	-	-	-	-	-	1	1	2	5
Female	109	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
United States												
All races, both sexes	5,985	8,503	9,870	11,097	13,698	18,600	23,024	22,202	14,766	5,635	1,136	5
Male	4,491	6,189	6,842	7,343	8,424	10,333	11,486	9,327	4,874	1,413	193	4
Female	1,494	2,314	3,028	3,754	5,274	8,267	11,538	12,875	9,892	4,222	943	1
White, both sexes	4,865	7,008	8,333	9,351	11,714	16,308	20,642	20,184	13,506	5,118	980	4
Male	3,750	5,214	5,864	6,331	7,358	9,231	10,429	8,574	4,497	1,293	161	3
Female	1,115	1,794	2,469	3,020	4,356	7,077	10,213	11,610	9,009	3,825	819	1
Black, both sexes	987	1,292	1,304	1,456	1,653	1,859	1,894	1,562	991	421	125	1
Male	641	828	820	834	884	874	807	533	273	83	22	1
Female	346	464	484	622	769	985	1,087	1,029	718	338	103	-
American Indian 1/, both sexes	50	64	67	72	70	91	84	55	36	11	1	-
Male	41	39	47	43	33	50	37	21	8	5	1	-
Female	9	25	20	29	37	41	47	34	28	6	-	-
Asian or Pacific Islander 2/, both sexes	83	139	166	218	261	342	404	401	233	85	30	-
Male	59	108	111	135	149	178	213	199	96	32	9	-
Female	24	31	55	83	112	164	191	202	137	53	21	-
Alabama												
All races, both sexes	162	203	237	230	259	370	342	325	171	82	16	-
Male	112	139	165	158	157	193	155	130	41	12	-	-
Female	50	64	72	72	102	177	187	195	130	70	16	-
White, both sexes	121	140	187	173	205	289	297	276	141	69	11	-
Male	88	96	132	122	130	152	132	112	35	11	-	-
Female	33	44	55	51	75	137	165	164	106	58	11	-
Black, both sexes	40	62	49	57	53	79	44	49	29	13	5	-
Male	24	43	32	36	27	40	22	18	6	1	-	-
Female	16	19	17	21	26	39	22	31	23	12	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	1	2	1	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	1	1	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Alaska												
All races, both sexes	6	10	6	5	9	15	19	5	2	2	-	-
Male	6	7	5	1	6	8	10	4	1	1	-	-
Female	-	3	1	4	3	7	9	1	1	1	-	-
White, both sexes	4	8	4	2	4	11	17	5	2	2	-	-
Male	4	5	3	-	3	6	10	4	1	1	-	-
Female	-	3	1	2	1	5	7	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	2	2	4	3	2	-	-	-	-	-
Male	1	1	2	1	2	2	-	-	-	-	-	-
Female	-	-	-	1	2	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	1	-	-	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Arizona												
All races, both sexes	54	83	124	142	209	272	333	327	197	64	8	-
Male	40	62	92	95	129	145	172	143	70	23	3	-
Female	14	21	32	47	80	127	161	184	127	41	5	-
White, both sexes	46	77	115	134	198	254	314	316	191	63	8	-
Male	33	58	85	91	122	134	160	139	67	22	3	-
Female	13	19	30	43	76	120	154	177	124	41	5	-
Black, both sexes	4	2	3	1	8	6	8	5	1	1	-	-
Male	4	1	2	-	5	2	6	2	-	1	-	-
Female	-	1	1	1	3	4	2	3	1	-	-	-
American Indian 1/, both sexes	4	4	6	6	3	12	8	4	5	-	-	-
Male	3	3	5	3	2	9	4	1	3	-	-	-
Female	1	1	1	3	1	3	4	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	3	2	-	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Arkansas												
All races, both sexes	173	232	271	255	287	380	431	379	283	106	20	1
Male	134	176	192	172	164	188	220	146	92	26	3	-
Female	39	56	79	83	123	192	211	233	191	80	17	1
White, both sexes	140	189	230	227	247	330	382	350	256	91	15	1
Male	110	145	161	159	143	165	201	134	82	23	1	-
Female	30	44	69	68	104	165	181	216	174	68	14	1
Black, both sexes	32	42	36	25	39	49	48	28	26	15	5	-
Male	23	31	27	13	20	22	19	11	10	3	2	-
Female	9	11	9	12	19	27	29	17	16	12	3	-
American Indian 1/, both sexes	1	1	2	2	1	-	1	-	1	-	-	-
Male	1	-	2	-	1	-	-	-	-	-	-	-
Female	-	1	-	2	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	1	-	1	-	1	-	-	-	-
Male	-	-	2	-	-	1	-	1	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
California												
All races, both sexes	417	661	764	915	1,211	1,742	2,341	2,320	1,610	621	121	-
Male	307	472	511	576	724	1,007	1,192	1,056	561	178	20	-
Female	110	189	253	339	487	735	1,149	1,264	1,049	443	101	-
White, both sexes	334	518	611	727	972	1,418	1,960	1,980	1,403	536	96	-
Male	254	376	415	479	597	826	1,008	911	492	152	12	-
Female	80	142	196	248	375	592	952	1,069	911	384	84	-
Black, both sexes	51	85	79	96	110	143	165	134	81	34	10	-
Male	31	54	50	45	53	81	69	47	21	10	2	-
Female	20	31	29	51	57	62	96	87	60	24	8	-
American Indian 1/, both sexes	4	6	7	7	7	11	8	8	2	1	-	-
Male	4	4	3	4	2	6	4	4	1	1	-	-
Female	-	2	4	3	5	5	4	4	1	-	-	-
Asian or Pacific Islander 2/, both sexes	28	52	67	85	122	170	208	198	124	50	15	-
Male	18	38	43	48	72	94	111	94	47	15	6	-
Female	10	14	24	37	50	76	97	104	77	35	9	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Colorado												
All races, both sexes	59	74	84	114	123	153	178	212	127	31	9	-
Male	52	56	59	83	79	77	88	94	40	5	-	-
Female	7	18	25	31	44	76	90	118	87	26	9	-
White, both sexes	58	73	82	111	119	148	173	209	125	30	9	-
Male	51	56	57	82	75	75	86	91	39	5	-	-
Female	7	17	25	29	44	73	87	118	86	25	9	-
Black, both sexes	1	1	2	3	4	1	3	2	1	1	-	-
Male	1	-	2	1	4	1	1	2	-	-	-	-
Female	-	1	-	2	-	-	2	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Connecticut												
All races, both sexes	41	49	75	72	98	134	226	279	171	73	13	1
Male	31	34	55	49	61	69	112	111	46	15	1	1
Female	10	15	20	23	37	65	114	168	125	58	12	-
White, both sexes	37	48	65	66	81	126	214	273	169	71	13	1
Male	29	33	48	44	53	67	108	108	46	14	1	1
Female	8	15	17	22	28	59	106	165	123	57	12	-
Black, both sexes	4	1	8	5	16	5	10	3	2	2	-	-
Male	2	1	5	4	7	1	4	2	-	1	-	-
Female	2	-	3	1	9	4	6	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	1	3	2	2	-	-	-	-
Male	-	-	2	1	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Delaware												
All races, both sexes	17	15	22	24	43	45	64	67	37	11	2	-
Male	11	14	13	18	32	28	33	28	12	2	-	-
Female	6	1	9	6	11	17	31	39	25	9	2	-
White, both sexes	15	13	20	16	36	40	52	62	33	8	1	-
Male	9	12	12	13	28	25	28	27	11	2	-	-
Female	6	1	8	3	8	15	24	35	22	6	1	-
Black, both sexes	2	2	2	8	6	5	12	5	4	3	1	-
Male	2	2	1	5	3	3	5	1	1	-	-	-
Female	-	-	1	3	3	2	7	4	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	12	14	12	25	18	31	38	29	32	15	3	-
Male	9	10	9	10	7	15	17	13	8	5	-	-
Female	3	4	3	15	11	16	21	16	24	10	3	-
White, both sexes	2	3	1	2	1	6	14	10	5	8	2	-
Male	2	3	1	2	1	5	9	7	1	3	-	-
Female	-	-	-	-	-	1	5	3	4	5	2	-
Black, both sexes	10	10	11	22	16	25	24	18	27	7	1	-
Male	7	6	8	7	6	10	8	5	7	2	-	-
Female	3	4	3	15	10	15	16	13	20	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Florida												
All races, both sexes	241	438	521	638	826	1,208	1,528	1,522	1,000	397	78	-
Male	176	313	355	430	530	643	789	698	364	104	23	-
Female	65	125	166	208	296	565	739	824	636	293	55	-
White, both sexes	200	378	454	545	713	1,104	1,438	1,424	946	371	73	-
Male	149	278	314	382	466	595	750	671	351	95	23	-
Female	51	100	140	163	247	509	688	753	595	276	50	-
Black, both sexes	37	58	64	86	106	93	82	89	53	23	4	-
Male	24	33	39	42	60	44	37	24	13	7	-	-
Female	13	25	25	44	46	49	45	65	40	16	4	-
American Indian 1/, both sexes	2	-	2	-	2	2	-	2	-	-	-	-
Male	2	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	1	2	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	7	5	9	8	7	1	3	1	-
Male	1	2	1	6	3	4	2	3	-	2	-	-
Female	1	-	-	1	2	5	6	4	1	1	1	-
Georgia												
All races, both sexes	193	247	282	311	348	345	464	386	284	74	12	-
Male	142	188	204	198	216	173	243	150	78	15	2	-
Female	51	59	78	113	132	172	221	236	206	59	10	-
White, both sexes	131	173	211	229	249	271	389	314	218	63	8	-
Male	104	139	158	153	159	144	208	129	57	14	1	-
Female	27	34	53	76	90	127	181	185	161	49	7	-
Black, both sexes	62	72	68	80	95	68	69	66	63	11	4	-
Male	38	47	45	43	56	27	30	19	20	1	1	-
Female	24	25	23	37	39	41	39	47	43	10	3	-
American Indian 1/, both sexes	-	-	-	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	3	1	4	5	5	5	3	-	-	-
Male	-	2	1	1	1	1	4	1	1	-	-	-
Female	-	-	2	-	3	4	1	4	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Hawaii												
All races, both sexes	18	26	28	31	21	60	80	79	60	12	6	-
Male	14	20	18	20	11	32	51	37	29	9	3	-
Female	4	6	10	11	10	28	29	42	31	3	3	-
White, both sexes	4	5	9	6	6	19	23	19	14	2	1	-
Male	3	3	5	6	4	9	16	10	5	-	-	-
Female	1	2	4	-	2	10	7	9	9	2	1	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	20	19	25	14	41	57	59	46	10	5	-
Male	11	16	13	14	6	23	35	27	24	9	3	-
Female	3	4	6	11	8	18	22	32	22	1	2	-
Idaho												
All races, both sexes	38	73	66	69	92	99	117	92	75	28	1	-
Male	31	56	45	50	59	68	66	48	23	9	1	-
Female	7	17	21	19	33	31	51	44	52	19	-	-
White, both sexes	34	71	64	67	90	97	115	90	74	28	1	-
Male	29	54	44	49	58	67	64	47	23	9	1	-
Female	5	17	20	18	32	30	51	43	51	19	-	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	2	1	1	2	1	-	1	-	-	-
Male	1	1	1	1	-	1	1	-	-	-	-	-
Female	1	-	1	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	-	2	-	-	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Illinois												
All races, both sexes	271	420	445	478	648	865	1,151	1,107	748	332	68	-
Male	198	306	320	321	420	463	568	439	240	83	11	-
Female	73	114	125	157	228	402	583	668	508	249	57	-
White, both sexes	222	332	365	380	532	739	1,000	1,003	691	300	60	-
Male	162	253	267	262	351	409	498	399	221	78	11	-
Female	60	79	98	118	181	330	502	604	470	222	49	-
Black, both sexes	39	83	76	86	102	113	144	83	51	28	7	-
Male	29	48	51	51	61	48	66	34	17	5	-	-
Female	10	35	25	35	41	65	78	49	34	23	7	-
American Indian 1/, both sexes	2	1	-	1	-	3	1	-	-	-	-	-
Male	1	1	-	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	4	4	11	14	10	6	21	6	4	1	-
Male	6	4	2	7	8	5	4	6	2	-	-	-
Female	2	-	2	4	6	5	2	15	4	4	1	-
Indiana												
All races, both sexes	173	240	276	301	384	463	520	518	316	122	19	-
Male	132	165	189	203	237	274	254	212	119	27	5	-
Female	41	75	87	98	147	189	266	306	197	95	14	-
White, both sexes	162	217	255	286	349	435	501	496	301	119	18	-
Male	123	152	175	193	215	258	244	207	114	26	4	-
Female	39	65	80	93	134	177	257	289	187	93	14	-
Black, both sexes	11	23	20	14	32	27	19	22	15	3	1	-
Male	9	13	13	9	21	15	10	5	5	1	1	-
Female	2	10	7	5	11	12	9	17	10	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Iowa												
All races, both sexes	69	90	108	130	126	210	323	302	227	101	17	-
Male	53	73	75	91	79	116	154	144	84	22	2	-
Female	16	17	33	39	47	94	169	158	143	79	15	-
White, both sexes	66	87	107	125	123	208	322	299	225	101	17	-
Male	51	70	74	89	78	114	154	143	84	22	2	-
Female	15	17	33	36	45	94	168	156	141	79	15	-
Black, both sexes	3	3	1	4	2	2	-	3	2	-	-	-
Male	2	3	1	1	1	2	-	1	-	-	-	-
Female	1	-	-	3	1	-	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	57	67	73	88	112	183	215	193	131	65	10	-
Male	46	52	46	55	73	100	124	77	43	10	1	-
Female	11	15	27	33	39	83	91	116	88	55	9	-
White, both sexes	52	65	66	82	103	170	209	189	127	65	9	-
Male	42	50	40	52	67	95	121	77	42	10	1	-
Female	10	15	26	30	36	75	88	112	85	55	8	-
Black, both sexes	4	2	4	4	4	9	2	3	3	-	1	-
Male	3	2	3	2	3	4	1	-	1	-	-	-
Female	1	-	1	2	1	5	1	3	2	-	1	-
American Indian 1/, both sexes	-	-	3	2	2	3	2	-	1	-	-	-
Male	-	-	3	1	1	1	1	-	-	-	-	-
Female	-	-	-	1	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	3	1	2	1	-	-	-	-
Male	1	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Kentucky												
All races, both sexes	228	249	277	315	382	426	489	422	254	76	16	-
Male	169	177	193	210	235	248	224	173	71	27	4	-
Female	59	72	84	105	147	178	265	249	183	49	12	-
White, both sexes	205	234	255	287	354	402	472	395	245	73	14	-
Male	153	169	177	190	214	236	219	160	66	26	3	-
Female	52	65	78	97	140	166	253	235	179	47	11	-
Black, both sexes	23	14	21	28	27	24	15	26	9	3	2	-
Male	16	8	15	20	21	12	4	13	5	1	1	-
Female	7	6	6	8	6	12	11	13	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	1	1	-	-	-	-
Louisiana												
All races, both sexes	204	249	307	267	282	326	341	297	155	44	15	-
Male	142	170	212	184	181	192	175	111	46	7	4	-
Female	62	79	95	83	101	134	166	186	109	37	11	-
White, both sexes	116	145	195	180	205	241	268	225	116	33	10	-
Male	87	98	143	130	139	146	151	87	35	5	4	-
Female	29	47	52	50	66	95	117	138	81	28	6	-
Black, both sexes	86	101	109	83	75	84	72	70	38	11	5	-
Male	53	69	67	52	41	45	24	23	11	2	-	-
Female	33	32	42	31	34	39	48	47	27	9	5	-
American Indian 1/, both sexes	1	-	2	2	1	-	-	-	-	-	-	-
Male	1	-	2	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	2	1	1	1	2	1	-	-	-
Male	1	3	-	1	-	1	-	1	-	-	-	-
Female	-	-	1	1	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Maine												
All races, both sexes	31	43	53	38	69	78	111	98	65	32	6	-
Male	27	34	39	26	42	46	50	31	26	9	1	-
Female	4	9	14	12	27	32	61	67	39	23	5	-
White, both sexes	31	42	53	38	69	78	110	98	65	32	6	-
Male	27	33	39	26	42	46	50	31	26	9	1	-
Female	4	9	14	12	27	32	60	67	39	23	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	93	139	174	194	238	342	442	365	258	92	25	-
Male	60	91	112	122	141	183	204	154	76	17	4	-
Female	33	48	62	72	97	159	238	211	182	75	21	-
White, both sexes	58	80	108	125	164	260	349	312	204	75	22	-
Male	36	53	76	82	100	143	162	132	65	13	3	-
Female	22	27	32	43	64	117	187	180	139	62	19	-
Black, both sexes	35	57	58	68	63	75	86	47	52	16	3	-
Male	24	37	32	39	34	35	39	18	11	4	1	-
Female	11	20	26	29	29	40	47	29	41	12	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	8	1	11	6	7	6	2	1	-	-
Male	-	1	4	1	7	4	3	4	-	-	-	-
Female	-	1	4	-	4	2	4	2	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Massachusetts												
All races, both sexes	83	116	134	180	188	319	455	493	334	126	38	-
Male	75	90	91	121	119	176	237	180	114	26	4	-
Female	8	26	43	59	69	143	218	313	220	100	34	-
White, both sexes	78	107	127	165	175	309	441	476	322	123	36	-
Male	71	83	86	109	112	170	228	173	111	26	3	-
Female	7	24	41	56	63	139	213	303	211	97	33	-
Black, both sexes	3	8	5	9	7	8	11	13	7	3	1	-
Male	2	6	4	6	3	4	7	4	2	-	1	-
Female	1	2	1	3	4	4	4	9	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	6	6	2	3	4	5	-	1	-
Male	2	1	1	6	4	2	2	3	1	-	-	-
Female	-	-	1	-	2	-	1	1	4	-	1	-
Michigan												
All races, both sexes	199	268	321	313	465	621	827	842	509	166	43	-
Male	140	192	225	193	268	344	395	353	146	39	8	-
Female	59	76	96	120	197	277	432	489	363	127	35	-
White, both sexes	161	224	278	279	397	547	750	775	473	147	36	-
Male	119	165	196	176	233	310	370	329	138	35	8	-
Female	42	59	82	103	164	237	380	446	335	112	28	-
Black, both sexes	33	40	36	30	62	69	71	57	31	18	7	-
Male	16	25	26	16	31	33	23	18	7	4	-	-
Female	17	15	10	14	31	36	48	39	24	14	7	-
American Indian 1/, both sexes	3	3	3	3	-	4	3	3	3	1	-	-
Male	3	2	2	1	-	1	1	1	1	-	-	-
Female	-	1	1	2	-	3	2	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	2	1	4	1	6	1	3	7	2	-	-	-
Male	2	-	1	-	4	-	1	5	-	-	-	-
Female	-	1	3	1	2	1	2	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Minnesota												
All races, both sexes	68	60	74	90	118	156	216	240	151	55	18	-
Male	54	49	54	63	83	83	117	108	54	20	5	-
Female	14	11	20	27	35	73	99	132	97	35	13	-
White, both sexes	62	53	72	83	110	151	214	237	150	55	18	-
Male	50	43	52	59	78	80	116	107	54	20	5	-
Female	12	10	20	24	32	71	98	130	96	35	13	-
Black, both sexes	2	2	-	-	2	1	-	2	-	-	-	-
Male	1	2	-	-	1	1	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	4	3	2	6	2	2	1	1	1	-	-	-
Male	3	2	2	3	2	2	1	-	-	-	-	-
Female	1	1	-	3	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	4	2	1	-	-	-	-	-
Male	-	2	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	2	2	1	-	-	-	-	-
Mississippi												
All races, both sexes	141	174	184	193	198	241	247	215	103	53	6	-
Male	96	119	136	133	116	139	125	80	31	10	-	-
Female	45	55	48	60	82	102	122	135	72	43	6	-
White, both sexes	69	113	120	126	133	178	177	163	82	31	3	-
Male	49	76	89	93	81	104	88	61	23	7	-	-
Female	20	37	31	33	52	74	89	102	59	24	3	-
Black, both sexes	72	60	63	65	65	63	68	52	21	22	3	-
Male	47	43	46	39	35	35	36	19	8	3	-	-
Female	25	17	17	26	30	28	32	33	13	19	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Missouri												
All races, both sexes	237	282	329	382	422	576	710	668	451	193	31	-
Male	174	203	222	261	258	338	348	260	152	51	4	-
Female	63	79	107	121	164	238	362	408	299	142	27	-
White, both sexes	209	252	297	340	363	522	652	622	429	182	29	-
Male	155	182	203	239	231	310	325	247	145	50	4	-
Female	54	70	94	101	132	212	327	375	284	132	25	-
Black, both sexes	28	27	28	39	55	52	56	44	21	11	1	-
Male	19	20	16	21	24	27	23	12	6	1	-	-
Female	9	7	12	18	31	25	33	32	15	10	1	-
American Indian 1/, both sexes	-	1	3	-	2	1	1	1	-	-	-	-
Male	-	-	2	-	2	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	3	2	1	1	1	1	-	1	-
Male	-	1	1	1	1	-	-	1	1	-	-	-
Female	-	1	-	2	1	1	1	-	-	-	1	-
Montana												
All races, both sexes	18	21	24	22	32	59	58	44	47	10	2	-
Male	13	15	20	17	21	29	34	18	17	5	1	-
Female	5	6	4	5	11	30	24	26	30	5	1	-
White, both sexes	17	19	21	21	31	57	55	44	46	10	2	-
Male	12	14	18	16	21	29	31	18	16	5	1	-
Female	5	5	3	5	10	28	24	26	30	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	3	1	1	2	3	-	1	-	-	-
Male	1	1	2	1	-	-	3	-	1	-	-	-
Female	-	1	1	-	1	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Nebraska												
All races, both sexes	30	30	35	42	60	70	85	75	57	28	3	-
Male	26	26	23	23	38	42	41	32	21	5	-	-
Female	4	4	12	19	22	28	44	43	36	23	3	-
White, both sexes	28	28	34	41	59	68	82	75	57	28	3	-
Male	24	25	22	23	38	40	40	32	21	5	-	-
Female	4	3	12	18	21	28	42	43	36	23	3	-
Black, both sexes	1	2	-	1	1	2	1	-	-	-	-	-
Male	1	1	-	-	-	2	1	-	-	-	-	-
Female	-	1	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Nevada												
All races, both sexes	37	44	51	58	82	76	79	74	31	12	2	-
Male	29	32	34	44	51	45	52	34	10	6	-	-
Female	8	12	17	14	31	31	27	40	21	6	2	-
White, both sexes	36	38	43	53	76	68	73	71	29	11	1	-
Male	29	28	28	41	49	42	48	32	9	6	-	-
Female	7	10	15	12	27	26	25	39	20	5	1	-
Black, both sexes	1	3	2	2	3	4	4	-	2	1	1	-
Male	-	2	1	1	1	1	3	-	1	-	-	-
Female	1	1	1	1	2	3	1	-	1	1	1	-
American Indian 1/, both sexes	-	1	2	1	1	-	1	-	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	4	2	2	4	1	3	-	-	-	-
Male	-	2	4	1	1	2	-	2	-	-	-	-
Female	-	-	-	1	1	2	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
New Hampshire												
All races, both sexes	15	32	22	27	37	64	76	81	42	24	5	-
Male	11	24	14	21	24	35	39	39	17	4	-	-
Female	4	8	8	6	13	29	37	42	25	20	5	-
White, both sexes	15	31	21	27	37	64	75	81	42	24	5	-
Male	11	24	14	21	24	35	39	39	17	4	-	-
Female	4	7	7	6	13	29	36	42	25	20	5	-
Black, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	133	178	228	280	377	574	741	729	521	226	41	-
Male	99	121	161	188	241	298	344	297	188	60	4	-
Female	34	57	67	92	136	276	397	432	333	166	37	-
White, both sexes	100	141	171	209	297	495	659	666	477	213	36	-
Male	79	99	126	146	196	267	302	270	179	58	4	-
Female	21	42	45	63	101	228	357	396	298	155	32	-
Black, both sexes	31	32	47	58	70	67	71	52	35	13	5	-
Male	18	17	28	33	38	26	32	20	4	2	-	-
Female	13	15	19	25	32	41	39	32	31	11	5	-
American Indian 1/, both sexes	-	1	1	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	9	13	9	12	11	10	9	-	-	-
Male	2	4	7	9	7	5	10	6	5	-	-	-
Female	-	-	2	4	2	7	1	4	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
New Mexico												
All races, both sexes	14	32	41	48	63	81	82	72	43	17	2	-
Male	13	24	28	32	42	55	47	36	20	2	1	-
Female	1	8	13	16	21	26	35	36	23	15	1	-
White, both sexes	13	31	39	45	56	78	80	69	40	14	1	-
Male	12	23	26	31	39	53	46	34	20	1	-	-
Female	1	8	13	14	17	25	34	35	20	13	1	-
Black, both sexes	-	-	-	2	3	-	1	-	1	1	-	-
Male	-	-	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	1	1	-	-
American Indian 1/, both sexes	1	1	2	1	3	3	1	3	2	2	1	-
Male	1	1	2	-	-	2	1	2	-	1	1	-
Female	-	-	-	1	3	1	-	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	248	418	516	591	772	1,235	1,607	1,777	1,292	546	112	1
Male	188	302	339	388	441	638	763	708	403	134	15	1
Female	60	116	177	203	331	597	844	1,069	889	412	97	-
White, both sexes	187	338	423	474	621	1,058	1,418	1,584	1,180	490	96	1
Male	150	253	282	313	377	566	693	643	370	124	13	1
Female	37	85	141	161	244	492	725	941	810	366	83	-
Black, both sexes	54	71	82	103	135	149	153	164	94	44	14	-
Male	32	42	48	66	55	58	57	46	25	6	2	-
Female	22	29	34	37	80	91	96	118	69	38	12	-
American Indian 1/, both sexes	-	1	1	2	2	2	3	2	2	2	-	-
Male	-	1	1	2	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	2	1	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	7	8	10	12	14	26	33	27	16	10	2	-
Male	6	6	8	7	8	14	11	19	8	4	-	-
Female	1	2	2	5	6	12	22	8	8	6	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
North Carolina												
All races, both sexes	182	260	288	337	385	503	618	538	341	111	28	-
Male	138	182	195	227	232	273	331	222	95	25	6	-
Female	44	78	93	110	153	230	287	316	246	86	22	-
White, both sexes	132	194	226	246	303	415	506	456	288	94	24	-
Male	105	140	162	170	188	232	280	199	81	23	4	-
Female	27	54	64	76	115	183	226	257	207	71	20	-
Black, both sexes	46	60	61	85	74	84	104	75	49	17	4	-
Male	29	36	32	53	41	40	45	22	13	2	2	-
Female	17	24	29	32	33	44	59	53	36	15	2	-
American Indian 1/, both sexes	2	4	1	5	6	3	6	7	3	-	-	-
Male	2	4	1	4	3	1	4	1	1	-	-	-
Female	-	-	-	1	3	2	2	6	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	1	2	1	2	-	1	-	-	-
Male	2	2	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	2	1	-	-	1	-	-	-
North Dakota												
All races, both sexes	19	19	31	43	40	53	73	83	60	22	8	-
Male	15	18	26	33	25	37	39	39	19	7	2	-
Female	4	1	5	10	15	16	34	44	41	15	6	-
White, both sexes	17	19	28	42	35	52	73	82	59	22	8	-
Male	13	18	24	32	22	36	39	38	19	7	2	-
Female	4	1	4	10	13	16	34	44	40	15	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	2	1	5	1	-	1	1	-	-	-
Male	2	-	1	1	3	1	-	1	-	-	-	-
Female	-	-	1	-	2	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Ohio												
All races, both sexes	264	406	433	494	614	915	999	1,023	645	220	49	-
Male	214	298	309	321	383	521	473	416	202	38	11	-
Female	50	108	124	173	231	394	526	607	443	182	38	-
White, both sexes	245	349	398	443	566	833	913	962	605	204	42	-
Male	199	261	287	290	354	486	431	393	189	37	8	-
Female	46	88	111	153	212	347	482	569	416	167	34	-
Black, both sexes	19	51	32	49	47	82	77	59	39	16	7	-
Male	15	32	19	30	29	35	36	21	13	1	3	-
Female	4	19	13	19	18	47	41	38	26	15	4	-
American Indian 1/, both sexes	-	1	1	-	-	-	3	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	5	2	2	1	-	6	1	1	-	-	-
Male	-	5	2	1	-	-	5	1	-	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
Oklahoma												
All races, both sexes	78	123	140	149	184	230	327	245	184	60	5	-
Male	57	87	97	91	114	124	162	117	62	19	1	-
Female	21	36	43	58	70	106	165	128	122	41	4	-
White, both sexes	62	98	125	124	163	203	288	228	173	53	4	-
Male	47	73	86	76	106	110	143	109	59	17	1	-
Female	15	25	39	48	57	93	145	119	114	36	3	-
Black, both sexes	10	10	7	8	10	12	21	7	6	6	-	-
Male	6	7	5	5	6	4	11	4	3	1	-	-
Female	4	3	2	3	4	8	10	3	3	5	-	-
American Indian 1/, both sexes	6	13	8	17	10	14	16	10	5	1	-	-
Male	4	6	6	10	2	10	7	4	-	1	-	-
Female	2	7	2	7	8	4	9	6	5	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	1	2	-	-	-	1	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Oregon												
All races, both sexes	40	74	81	115	153	170	222	222	136	48	13	-
Male	35	56	61	76	102	102	133	102	55	22	2	-
Female	5	18	20	39	51	68	89	120	81	26	11	-
White, both sexes	38	74	79	111	147	160	218	216	132	47	12	-
Male	34	56	60	72	97	97	131	97	53	21	2	-
Female	4	18	19	39	50	63	87	119	79	26	10	-
Black, both sexes	1	-	1	-	2	2	1	3	1	-	-	-
Male	-	-	-	-	2	1	1	3	1	-	-	-
Female	1	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	3	-	-	-	-	-	-
Male	1	-	-	1	2	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	2	5	3	3	3	1	1	-
Male	-	-	1	3	1	2	1	2	1	1	-	-
Female	-	-	-	-	1	3	2	1	2	-	1	-
Pennsylvania												
All races, both sexes	288	396	520	597	746	1,087	1,352	1,288	804	279	47	-
Male	220	304	372	385	460	619	664	508	272	67	8	-
Female	68	92	148	212	286	468	688	780	532	212	39	-
White, both sexes	253	364	473	533	689	1,019	1,281	1,232	775	265	42	-
Male	200	283	340	349	424	586	634	486	262	65	8	-
Female	53	81	133	184	265	433	647	746	513	200	34	-
Black, both sexes	34	31	45	60	51	65	67	52	28	12	5	-
Male	20	20	30	34	31	31	27	20	10	1	-	-
Female	14	11	15	26	20	34	40	32	18	11	5	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	4	5	3	3	4	1	1	-	-
Male	-	1	1	2	5	2	2	2	-	-	-	-
Female	1	-	-	2	-	1	1	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Rhode Island												
All races, both sexes	25	26	41	50	54	114	135	141	107	39	7	-
Male	20	20	27	33	28	62	75	59	41	7	2	-
Female	5	6	14	17	26	52	60	82	66	32	5	-
White, both sexes	22	23	39	47	50	107	133	141	105	39	7	-
Male	18	18	26	31	27	59	75	59	41	7	2	-
Female	4	5	13	16	23	48	58	82	64	32	5	-
Black, both sexes	1	2	1	3	4	3	1	-	1	-	-	-
Male	1	1	1	2	1	2	-	-	-	-	-	-
Female	-	1	-	1	3	1	1	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	4	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	3	-	-	-	-	-	-
South Carolina												
All races, both sexes	146	212	249	223	270	272	305	310	136	49	12	2
Male	106	152	172	155	160	150	151	130	41	8	-	2
Female	40	60	77	68	110	122	154	180	95	41	12	-
White, both sexes	105	144	158	167	197	184	233	241	108	34	6	1
Male	75	107	112	120	124	110	123	106	32	6	-	1
Female	30	37	46	47	73	74	110	135	76	28	6	-
Black, both sexes	41	67	89	55	72	87	72	67	28	15	6	1
Male	31	44	59	34	35	40	28	23	9	2	-	1
Female	10	23	30	21	37	47	44	44	19	13	6	-
American Indian 1/, both sexes	-	1	2	-	1	1	-	-	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
South Dakota												
All races, both sexes	21	35	42	47	44	88	103	95	78	41	7	-
Male	15	27	35	31	31	52	60	44	28	11	1	-
Female	6	8	7	16	13	36	43	51	50	30	6	-
White, both sexes	17	30	39	44	42	83	98	93	76	40	7	-
Male	11	24	34	30	31	49	57	42	28	11	1	-
Female	6	6	5	14	11	34	41	51	48	29	6	-
Black, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	4	5	2	2	2	5	3	2	2	1	-	-
Male	4	3	-	-	-	3	2	2	-	-	-	-
Female	-	2	2	2	2	2	1	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	208	306	359	407	477	565	635	563	396	137	34	-
Male	150	230	248	277	298	294	308	234	126	35	4	-
Female	58	76	111	130	179	271	327	329	270	102	30	-
White, both sexes	167	257	322	362	425	491	559	504	347	121	28	-
Male	122	200	226	248	273	263	277	211	114	32	4	-
Female	45	57	96	114	152	228	282	293	233	89	24	-
Black, both sexes	41	49	36	43	51	73	75	58	47	16	6	-
Male	28	30	21	27	24	31	30	22	11	3	-	-
Female	13	19	15	16	27	42	45	36	36	13	6	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	1	1	1	-	-	-
Male	-	-	1	1	1	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Texas												
All races, both sexes	490	744	819	904	1,095	1,401	1,640	1,479	1,016	402	82	-
Male	367	523	546	570	671	786	793	588	334	103	17	-
Female	123	221	273	334	424	615	847	891	682	299	65	-
White, both sexes	404	621	718	787	981	1,241	1,485	1,352	939	362	70	-
Male	310	441	491	505	606	704	725	548	314	90	13	-
Female	94	180	227	282	375	537	760	804	625	272	57	-
Black, both sexes	82	109	93	108	104	145	144	114	74	38	11	-
Male	54	73	49	59	61	72	65	34	19	13	4	-
Female	28	36	44	49	43	73	79	80	55	25	7	-
American Indian 1/, both sexes	1	5	-	3	-	1	2	2	-	1	-	-
Male	-	2	-	2	-	-	-	-	-	-	-	-
Female	1	3	-	1	-	1	2	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	9	8	6	10	14	9	11	3	1	1	-
Male	3	7	6	4	4	10	3	6	1	-	-	-
Female	-	2	2	2	6	4	6	5	2	1	1	-
Utah												
All races, both sexes	21	37	36	45	46	79	91	90	45	13	1	-
Male	17	31	26	33	30	49	57	43	18	4	-	-
Female	4	6	10	12	16	30	34	47	27	9	1	-
White, both sexes	20	35	35	44	43	79	90	89	44	13	1	-
Male	17	30	25	32	28	49	56	43	18	4	-	-
Female	3	5	10	12	15	30	34	46	26	9	1	-
Black, both sexes	1	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	1	1	-	-	-	-
Male	-	-	-	1	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Vermont												
All races, both sexes	6	8	15	19	26	39	45	42	29	13	2	-
Male	6	6	9	18	15	22	25	21	6	3	-	-
Female	-	2	6	1	11	17	20	21	23	10	2	-
White, both sexes	6	8	15	19	26	39	45	42	29	13	2	-
Male	6	6	9	18	15	22	25	21	6	3	-	-
Female	-	2	6	1	11	17	20	21	23	10	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	147	197	231	249	312	414	475	462	278	98	23	-
Male	112	143	157	171	179	236	221	214	89	26	2	-
Female	35	54	74	78	133	178	254	248	189	72	21	-
White, both sexes	111	160	179	190	249	337	389	378	237	82	19	-
Male	86	120	123	138	147	199	189	176	78	21	-	-
Female	25	40	56	52	102	138	200	202	159	61	19	-
Black, both sexes	33	35	49	54	59	72	77	77	40	15	4	-
Male	24	21	32	30	30	35	29	35	11	5	2	-
Female	9	14	17	24	29	37	48	42	29	10	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	3	5	4	5	8	7	1	1	-	-
Male	2	2	2	3	2	2	3	3	-	-	-	-
Female	1	-	1	2	2	3	5	4	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Washington												
All races, both sexes	82	112	139	175	224	287	356	332	238	107	17	-
Male	61	86	100	114	150	168	188	158	85	27	3	-
Female	21	26	39	61	74	119	168	174	153	80	14	-
White, both sexes	74	105	122	157	211	269	338	316	230	104	16	-
Male	57	82	90	102	142	160	180	150	79	26	3	-
Female	17	23	32	55	69	109	158	166	151	78	13	-
Black, both sexes	2	-	9	5	4	6	3	3	3	-	-	-
Male	2	-	6	3	2	3	3	1	2	-	-	-
Female	-	-	3	2	2	3	-	2	1	-	-	-
American Indian 1/, both sexes	1	2	2	2	4	5	6	2	1	-	-	-
Male	1	1	1	1	3	3	-	1	1	-	-	-
Female	-	1	1	1	1	2	6	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	5	6	11	5	7	9	11	4	3	1	-
Male	1	3	3	8	3	2	5	6	3	1	-	-
Female	4	2	3	3	2	5	4	5	1	2	1	-
West Virginia												
All races, both sexes	51	76	71	126	128	172	197	159	99	29	5	-
Male	45	59	45	78	73	100	92	66	29	8	1	-
Female	6	17	26	48	55	72	105	93	70	21	4	-
White, both sexes	49	69	70	125	122	170	190	156	95	28	4	-
Male	43	54	44	77	69	100	88	65	29	8	1	-
Female	6	15	26	48	53	70	102	91	66	20	3	-
Black, both sexes	1	6	1	1	6	2	6	3	4	1	1	-
Male	1	4	1	1	4	-	3	1	-	-	-	-
Female	-	2	-	-	2	2	3	2	4	1	1	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Wisconsin												
All races, both sexes	124	177	187	220	277	367	479	464	368	123	25	-
Male	95	140	145	154	174	234	254	206	122	34	3	-
Female	29	37	42	66	103	133	225	258	246	89	22	-
White, both sexes	117	168	180	210	267	360	472	457	364	122	25	-
Male	92	136	141	148	168	232	254	204	122	34	3	-
Female	25	32	39	62	99	128	218	253	242	88	22	-
Black, both sexes	4	7	5	5	6	3	3	5	1	1	-	-
Male	1	3	3	2	3	1	-	2	-	-	-	-
Female	3	4	2	3	3	2	3	3	1	1	-	-
American Indian 1/, both sexes	3	2	2	2	4	3	4	2	2	-	-	-
Male	2	1	1	2	3	1	-	-	-	-	-	-
Female	1	1	1	-	1	2	4	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	-	1	-	-	1	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Wyoming												
All races, both sexes	13	18	24	29	24	25	36	32	22	8	3	-
Male	8	15	16	20	14	14	17	16	7	1	-	-
Female	5	3	8	9	10	11	19	16	15	7	3	-
White, both sexes	13	17	24	29	23	25	36	31	22	7	3	-
Male	8	14	16	20	13	14	17	16	7	-	-	-
Female	5	3	8	9	10	11	19	15	15	7	3	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
United States												
All races, both sexes	3,932	2	1	-	-	4	3	6	15	31	61	129
Male	2,016	1	-	-	-	4	2	5	12	14	40	90
Female	1,916	1	1	-	-	-	1	1	3	17	21	39
White, both sexes	3,289	2	1	-	-	3	2	6	9	16	44	88
Male	1,664	1	-	-	-	3	1	5	8	6	30	65
Female	1,625	1	1	-	-	-	1	1	1	10	14	23
Black, both sexes	534	-	-	-	-	1	1	-	6	14	14	32
Male	291	-	-	-	-	1	1	-	4	7	8	19
Female	243	-	-	-	-	-	-	-	2	7	6	13
American Indian 1/, both sexes	52	-	-	-	-	-	-	-	-	-	3	5
Male	29	-	-	-	-	-	-	-	-	-	2	3
Female	23	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	1	-	4
Male	32	-	-	-	-	-	-	-	-	1	-	3
Female	25	-	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	29	-	-	-	-	-	-	-	1	-	-	-
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Arkansas												
All races, both sexes	20	-	-	-	-	-	-	1	-	-	1	-
Male	12	-	-	-	-	-	-	1	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	1	-	-	1	-
Male	12	-	-	-	-	-	-	1	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	184	-	-	-	-	-	-	-	-	1	4	4
Male	90	-	-	-	-	-	-	-	-	1	2	3
Female	94	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	141	-	-	-	-	-	-	-	-	-	4	2
Male	68	-	-	-	-	-	-	-	-	-	2	2
Female	73	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	1	-	1
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Colorado												
All races, both sexes	29	-	-	-	-	-	-	-	2	-	-	2
Male	24	-	-	-	-	-	-	-	2	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	2	-	-	2
Male	23	-	-	-	-	-	-	-	2	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Delaware												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Florida												
All races, both sexes	121	-	-	-	-	-	-	-	-	-	1	4
Male	68	-	-	-	-	-	-	-	-	-	-	4
Female	53	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	110	-	-	-	-	-	-	-	-	-	-	2
Male	63	-	-	-	-	-	-	-	-	-	-	2
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	60	-	-	-	-	-	-	1	-	2	-	4
Male	32	-	-	-	-	-	-	1	-	2	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	42	-	-	-	-	-	-	1	-	-	-	2
Male	20	-	-	-	-	-	-	1	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	18	-	-	-	-	-	-	-	-	2	-	2
Male	12	-	-	-	-	-	-	-	-	2	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Hawaii												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Illinois												
All races, both sexes	133	-	-	-	-	-	-	-	-	-	-	4
Male	68	-	-	-	-	-	-	-	-	-	-	4
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	112	-	-	-	-	-	-	-	-	-	-	3
Male	56	-	-	-	-	-	-	-	-	-	-	3
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	53	-	-	-	-	-	-	-	1	2	2	2
Male	28	-	-	-	-	-	-	-	-	2	2	2
Female	25	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	2	2	2
Male	23	-	-	-	-	-	-	-	-	2	2	2
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Iowa												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	21	-	-	-	-	1	-	-	-	-	-	-
Male	9	-	-	-	-	1	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Kentucky												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	65	-	-	-	-	-	-	-	-	1	2	2
Male	28	-	-	-	-	-	-	-	-	1	1	2
Female	37	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Maine												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	41	-	-	-	-	-	-	1	-	-	-	1
Male	17	-	-	-	-	-	-	1	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Massachusetts												
All races, both sexes	46	-	-	-	-	-	-	-	-	1	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	1	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Minnesota												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	16	-	-	-	-	-	1	-	-	-	-	3
Male	8	-	-	-	-	-	1	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	1	-	-	-	-	2
Male	4	-	-	-	-	-	1	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Missouri												
All races, both sexes	80	-	-	-	-	-	-	-	-	-	-	2
Male	34	-	-	-	-	-	-	-	-	-	-	2
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	74	-	-	-	-	-	-	-	-	-	-	2
Male	31	-	-	-	-	-	-	-	-	-	-	2
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Nebraska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
New Hampshire												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
New Jersey												
All races, both sexes	110	-	-	-	-	-	-	1	-	-	-	3
Male	39	-	-	-	-	-	-	1	-	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	100	-	-	-	-	-	-	1	-	-	-	3
Male	35	-	-	-	-	-	-	1	-	-	-	1
Female	65	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
New Mexico												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	188	-	-	-	-	-	-	-	1	-	2	1
Male	88	-	-	-	-	-	-	-	1	-	1	-
Female	100	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	173	-	-	-	-	-	-	-	1	-	2	-
Male	81	-	-	-	-	-	-	-	1	-	1	-
Female	92	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
North Carolina												
All races, both sexes	93	-	-	-	-	1	-	-	-	1	4	1
Male	52	-	-	-	-	1	-	-	-	-	4	1
Female	41	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	70	-	-	-	-	1	-	-	-	1	3	-
Male	39	-	-	-	-	1	-	-	-	-	3	-
Female	31	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Ohio												
All races, both sexes	117	1	-	-	-	-	1	-	-	-	3	5
Male	67	1	-	-	-	-	-	-	-	-	3	4
Female	50	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	110	1	-	-	-	-	1	-	-	-	3	5
Male	64	1	-	-	-	-	-	-	-	-	3	4
Female	46	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	953	-	-	-	-	-	-	-	6	2	13	22
Male	491	-	-	-	-	-	-	-	5	1	7	17
Female	462	-	-	-	-	-	-	-	1	1	6	5
White, both sexes	864	-	-	-	-	-	-	-	5	-	8	17
Male	438	-	-	-	-	-	-	-	4	-	3	14
Female	426	-	-	-	-	-	-	-	1	-	5	3
Black, both sexes	46	-	-	-	-	-	-	-	1	2	2	2
Male	25	-	-	-	-	-	-	-	1	1	2	1
Female	21	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	39	-	-	-	-	-	-	-	-	-	3	3
Male	25	-	-	-	-	-	-	-	-	-	2	2
Female	14	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Oregon												
All races, both sexes	22	-	-	-	-	1	-	-	-	-	1	-
Male	13	-	-	-	-	1	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	1	-	-	-	-	1	-
Male	12	-	-	-	-	1	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	158	1	-	-	-	-	-	-	-	2	2	6
Male	85	-	-	-	-	-	-	-	-	-	2	5
Female	73	1	-	-	-	-	-	-	-	2	-	1
White, both sexes	142	1	-	-	-	-	-	-	-	2	2	6
Male	77	-	-	-	-	-	-	-	-	-	2	5
Female	65	1	-	-	-	-	-	-	-	2	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Rhode Island												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	171	-	-	-	-	-	-	1	2	4	6	11
Male	102	-	-	-	-	-	-	1	2	2	4	5
Female	69	-	-	-	-	-	-	-	-	2	2	6
White, both sexes	78	-	-	-	-	-	-	1	-	2	3	5
Male	49	-	-	-	-	-	-	1	-	2	2	3
Female	29	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	91	-	-	-	-	-	-	-	2	2	3	6
Male	53	-	-	-	-	-	-	-	2	-	2	2
Female	38	-	-	-	-	-	-	-	-	2	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	47	-	1	-	-	-	-	-	-	1	1	3
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	1	-	-	-	-	-	-	1	1	2
White, both sexes	40	-	1	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Texas												
All races, both sexes	188	-	-	-	-	1	-	1	-	1	5	3
Male	77	-	-	-	-	1	-	-	-	-	1	1
Female	111	-	-	-	-	-	-	1	-	1	4	2
White, both sexes	166	-	-	-	-	1	-	1	-	1	3	1
Male	65	-	-	-	-	1	-	-	-	-	-	1
Female	101	-	-	-	-	-	-	1	-	1	3	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	2	2
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	543	-	-	-	-	-	1	-	2	12	13	35
Male	305	-	-	-	-	-	1	-	1	5	11	25
Female	238	-	-	-	-	-	-	-	1	7	2	10
White, both sexes	393	-	-	-	-	-	1	-	-	6	10	23
Male	216	-	-	-	-	-	1	-	-	2	9	16
Female	177	-	-	-	-	-	-	-	-	4	1	7
Black, both sexes	146	-	-	-	-	-	-	-	2	6	3	10
Male	86	-	-	-	-	-	-	-	1	3	2	7
Female	60	-	-	-	-	-	-	-	1	3	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Washington												
All races, both sexes	30	-	-	-	-	-	-	-	-	1	-	3
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	1	-	3
Male	13	-	-	-	-	-	-	-	-	-	-	2
Female	15	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Wisconsin												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
United States												
All races, both sexes	225	266	330	296	380	472	569	539	378	183	39	3
Male	164	172	228	176	226	248	263	218	108	39	3	3
Female	61	94	102	120	154	224	306	321	270	144	36	-
White, both sexes	162	209	265	241	307	405	497	482	348	167	33	2
Male	123	135	185	147	184	211	227	191	98	39	3	2
Female	39	74	80	94	123	194	270	291	250	128	30	-
Black, both sexes	54	51	58	47	61	54	60	41	21	12	6	1
Male	35	34	39	24	36	28	28	19	7	-	-	1
Female	19	17	19	23	25	26	32	22	14	12	6	-
American Indian 1/, both sexes	6	4	5	3	5	7	5	4	4	1	-	-
Male	3	2	3	1	2	6	3	2	2	-	-	-
Female	3	2	2	2	3	1	2	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	3	2	2	5	7	6	7	12	5	3	-	-
Male	3	1	1	4	4	3	5	6	1	-	-	-
Female	-	1	1	1	3	3	2	6	4	3	-	-
Alabama												
All races, both sexes	1	2	5	4	4	5	4	1	2	-	-	-
Male	1	1	4	3	2	2	2	-	-	-	-	-
Female	-	1	1	1	2	3	2	1	2	-	-	-
White, both sexes	1	1	3	3	3	4	3	1	1	-	-	-
Male	1	-	2	2	2	1	2	-	-	-	-	-
Female	-	1	1	1	1	3	1	1	1	-	-	-
Black, both sexes	-	1	2	1	1	1	1	-	1	-	-	-
Male	-	1	2	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Alaska												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	1	1	4	3	3	5	4	3	1	-	-
Male	1	1	1	3	2	3	4	-	-	-	-	-
Female	-	-	-	1	1	-	1	4	3	1	-	-
White, both sexes	1	1	1	4	3	2	5	4	2	-	-	-
Male	1	1	1	3	2	2	4	-	-	-	-	-
Female	-	-	-	1	1	-	1	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Arkansas												
All races, both sexes	-	1	-	3	2	3	5	2	1	1	-	-
Male	-	1	-	2	1	-	3	2	-	1	-	-
Female	-	-	-	1	1	3	2	-	1	-	-	-
White, both sexes	-	1	-	3	2	2	4	2	1	1	-	-
Male	-	1	-	2	1	-	3	2	-	1	-	-
Female	-	-	-	1	1	2	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	13	11	13	19	20	28	39	15	10	3	-
Male	2	6	7	6	10	10	20	15	6	2	-	-
Female	2	7	4	7	9	10	8	24	9	8	3	-
White, both sexes	3	10	9	9	13	17	24	28	12	7	3	-
Male	2	4	6	3	6	8	17	10	6	2	-	-
Female	1	6	3	6	7	9	7	18	6	5	3	-
Black, both sexes	-	2	2	2	3	-	1	6	1	-	-	-
Male	-	2	1	1	2	-	1	2	-	-	-	-
Female	-	-	1	1	1	-	-	4	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	3	3	3	5	2	3	-	-
Male	-	-	-	2	2	2	2	3	-	-	-	-
Female	-	1	-	-	1	1	1	2	2	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Colorado												
All races, both sexes	3	2	3	2	2	3	4	4	2	-	-	-
Male	3	2	3	1	1	2	4	3	1	-	-	-
Female	-	-	-	1	1	1	-	1	1	-	-	-
White, both sexes	2	2	3	2	2	2	4	4	2	-	-	-
Male	2	2	3	1	1	2	4	3	1	-	-	-
Female	-	-	-	1	1	-	-	1	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	-	3	3	4	6	1	1	-	-
Male	-	-	-	-	2	2	1	4	1	1	-	-
Female	-	1	-	-	1	1	3	2	-	-	-	-
White, both sexes	-	-	-	-	3	3	4	6	1	1	-	-
Male	-	-	-	-	2	2	1	4	1	1	-	-
Female	-	-	-	-	1	1	3	2	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Delaware												
All races, both sexes	-	-	1	-	-	-	-	1	2	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
White, both sexes	-	-	1	-	-	-	-	1	2	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Florida												
All races, both sexes	4	5	5	8	15	17	16	29	11	4	2	-
Male	3	3	4	6	9	10	10	10	5	3	1	-
Female	1	2	1	2	6	7	6	19	6	1	1	-
White, both sexes	4	5	5	7	13	17	13	29	10	4	1	-
Male	3	3	4	6	8	10	9	10	4	3	1	-
Female	1	2	1	1	5	7	4	19	6	1	-	-
Black, both sexes	-	-	-	1	2	-	3	-	1	-	1	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	1	5	5	10	7	3	8	6	1	2	-
Male	4	1	2	4	9	5	1	1	1	-	-	-
Female	1	-	3	1	1	2	2	7	5	1	2	-
White, both sexes	2	-	3	5	8	6	1	7	5	1	1	-
Male	2	-	1	4	7	4	-	-	1	-	-	-
Female	-	-	2	1	1	2	1	7	4	1	1	-
Black, both sexes	3	1	2	-	2	1	2	1	1	-	1	-
Male	2	1	1	-	2	1	1	1	-	-	-	-
Female	1	-	1	-	-	-	1	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Hawaii												
All races, both sexes	1	-	-	-	1	-	2	2	-	-	-	-
Male	1	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	2	2	-	-	-	-
Male	1	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Idaho												
All races, both sexes	-	-	1	2	-	-	3	2	1	-	-	-
Male	-	-	1	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	3	1	1	-	-	-
White, both sexes	-	-	1	2	-	-	3	2	1	-	-	-
Male	-	-	1	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Illinois												
All races, both sexes	5	3	11	11	14	17	19	14	23	10	2	-
Male	4	3	4	6	9	7	11	10	8	2	-	-
Female	1	-	7	5	5	10	8	4	15	8	2	-
White, both sexes	3	2	10	11	11	14	16	11	20	10	1	-
Male	2	2	3	6	8	7	9	7	7	2	-	-
Female	1	-	7	5	3	7	7	4	13	8	1	-
Black, both sexes	2	1	1	-	2	3	3	2	2	-	1	-
Male	2	1	1	-	1	-	2	2	1	-	-	-
Female	-	-	-	-	1	3	1	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Indiana												
All races, both sexes	6	4	3	3	2	6	7	7	6	2	-	-
Male	3	4	3	2	1	1	1	3	4	-	-	-
Female	3	-	-	1	1	5	6	4	2	2	-	-
White, both sexes	5	3	2	1	2	4	7	7	4	1	-	-
Male	3	3	2	1	1	-	1	3	3	-	-	-
Female	2	-	-	-	1	4	6	4	1	1	-	-
Black, both sexes	1	1	1	2	-	2	-	-	2	1	-	-
Male	-	1	1	1	-	1	-	-	1	-	-	-
Female	1	-	-	1	-	1	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Iowa												
All races, both sexes	1	-	1	4	3	1	2	2	2	1	-	-
Male	1	-	-	2	2	1	1	1	-	-	-	-
Female	-	-	1	2	1	-	1	1	2	1	-	-
White, both sexes	-	-	1	4	3	1	2	2	2	1	-	-
Male	-	-	-	2	2	1	1	1	-	-	-	-
Female	-	-	1	2	1	-	1	1	2	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	2	1	1	2	2	3	3	2	2	1	-
Male	1	1	1	-	1	-	2	-	1	1	-	-
Female	-	1	-	1	1	2	1	3	1	1	1	-
White, both sexes	1	2	1	1	2	2	3	3	2	2	1	-
Male	1	1	1	-	1	-	2	-	1	1	-	-
Female	-	1	-	1	1	2	1	3	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Kentucky												
All races, both sexes	1	-	4	1	1	5	4	1	4	-	-	-
Male	1	-	4	1	-	2	-	1	-	-	-	-
Female	-	-	-	-	1	3	4	-	4	-	-	-
White, both sexes	1	-	4	1	1	4	4	-	4	-	-	-
Male	1	-	4	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	2	4	-	4	-	-	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	7	4	4	5	5	12	11	6	1	-	-
Male	4	2	3	3	3	1	5	3	-	-	-	-
Female	1	5	1	1	2	4	7	8	6	1	-	-
White, both sexes	1	4	2	-	5	5	9	8	6	-	-	-
Male	1	1	1	-	3	1	4	2	-	-	-	-
Female	-	3	1	-	2	4	5	6	6	-	-	-
Black, both sexes	4	3	2	4	-	-	3	3	-	1	-	-
Male	3	1	2	3	-	-	1	1	-	-	-	-
Female	1	2	-	1	-	-	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Maine												
All races, both sexes	1	2	3	2	-	3	3	1	3	4	-	-
Male	1	2	3	2	-	3	1	1	-	2	-	-
Female	-	-	-	-	-	-	2	-	3	2	-	-
White, both sexes	1	2	3	2	-	3	3	1	3	4	-	-
Male	1	2	3	2	-	3	1	1	-	2	-	-
Female	-	-	-	-	-	-	2	-	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	6	3	4	3	5	4	4	4	2	2	-
Male	-	5	2	1	1	2	-	2	1	1	-	-
Female	2	1	1	3	2	3	4	2	3	1	2	-
White, both sexes	1	2	-	4	2	5	3	2	4	2	1	-
Male	-	2	-	1	1	2	-	2	1	1	-	-
Female	1	-	-	3	1	3	3	-	3	1	1	-
Black, both sexes	1	3	3	-	1	-	1	1	-	-	1	-
Male	-	2	2	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	1	-	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Massachusetts												
All races, both sexes	1	3	2	2	1	9	10	7	7	-	2	-
Male	1	2	2	1	1	5	3	4	2	-	-	-
Female	-	1	-	1	-	4	7	3	5	-	2	-
White, both sexes	1	3	2	2	1	8	10	7	6	-	2	-
Male	1	2	2	1	1	4	3	4	2	-	-	-
Female	-	1	-	1	-	4	7	3	4	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Michigan												
All races, both sexes	4	4	6	3	13	9	11	10	5	4	4	-
Male	3	2	4	3	5	2	6	3	1	-	-	-
Female	1	2	2	-	8	7	5	7	4	4	4	-
White, both sexes	2	3	3	3	10	5	8	9	4	4	4	-
Male	2	2	3	3	4	2	4	3	1	-	-	-
Female	-	1	-	-	6	3	4	6	3	4	4	-
Black, both sexes	1	-	3	-	3	4	3	1	1	-	-	-
Male	1	-	1	-	1	-	2	-	-	-	-	-
Female	-	-	2	-	2	4	1	1	1	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Minnesota												
All races, both sexes	-	5	5	2	4	2	7	4	11	3	-	-
Male	-	4	3	1	3	2	4	3	4	1	-	-
Female	-	1	2	1	1	-	3	1	7	2	-	-
White, both sexes	-	5	4	2	4	2	7	4	11	3	-	-
Male	-	4	3	1	3	2	4	3	4	1	-	-
Female	-	1	1	1	1	-	3	1	7	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	3	1	1	-	-	2	-	2	-	-	-
Male	3	1	1	-	-	-	1	-	-	-	-	-
Female	-	2	-	1	-	-	1	-	2	-	-	-
White, both sexes	2	2	1	1	-	-	2	-	2	-	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	2	-	1	-	-	1	-	2	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Missouri												
All races, both sexes	2	3	4	3	6	6	18	11	17	7	1	-
Male	1	3	3	1	4	4	5	5	5	-	1	-
Female	1	-	1	2	2	2	13	6	12	7	-	-
White, both sexes	2	2	3	3	4	6	16	11	17	7	1	-
Male	1	2	2	1	3	4	5	5	5	-	1	-
Female	1	-	1	2	1	2	11	6	12	7	-	-
Black, both sexes	-	1	1	-	2	-	2	-	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	1	3	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	3	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Nebraska												
All races, both sexes	-	-	-	1	-	1	1	1	1	2	1	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	-	-	1	-	-	1	1	1	-
White, both sexes	-	-	-	1	-	1	1	1	1	2	1	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	-	-	1	-	-	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	3	2	4	1	5	2	-	-	-	-
Male	-	-	1	2	3	1	3	1	-	-	-	-
Female	-	1	2	-	1	-	2	1	-	-	-	-
White, both sexes	-	1	3	2	4	1	4	2	-	-	-	-
Male	-	-	1	2	3	1	3	1	-	-	-	-
Female	-	1	2	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
New Hampshire												
All races, both sexes	-	-	1	-	1	1	1	2	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	1	-	1	1	1	2	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	4	8	5	12	15	29	19	10	1	-
Male	-	1	2	4	4	5	5	12	2	1	1	-
Female	1	1	2	4	1	7	10	17	17	9	-	-
White, both sexes	1	2	3	7	3	11	13	27	19	9	1	-
Male	-	1	2	4	3	4	4	11	2	1	1	-
Female	1	1	1	3	-	7	9	16	17	8	-	-
Black, both sexes	-	-	1	1	2	1	2	-	-	1	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
New Mexico												
All races, both sexes	-	1	-	1	1	2	2	5	2	1	-	-
Male	-	1	-	1	-	2	1	3	1	-	-	-
Female	-	-	-	-	1	-	1	2	1	1	-	-
White, both sexes	-	1	-	1	-	2	2	4	2	1	-	-
Male	-	1	-	1	-	2	1	3	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5	12	12	10	13	24	35	29	28	14	2	-
Male	5	9	9	5	7	12	15	14	8	2	-	-
Female	-	3	3	5	6	12	20	15	20	12	2	-
White, both sexes	5	10	11	10	12	20	33	27	27	13	2	-
Male	5	8	9	5	7	10	14	12	7	2	-	-
Female	-	2	2	5	5	10	19	15	20	11	2	-
Black, both sexes	-	2	1	-	1	3	1	2	-	1	-	-
Male	-	1	-	-	-	2	-	2	-	-	-	-
Female	-	1	1	-	1	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
North Carolina												
All races, both sexes	6	7	5	14	8	12	11	12	7	2	2	-
Male	5	6	5	10	5	7	3	3	2	-	-	-
Female	1	1	-	4	3	5	8	9	5	2	2	-
White, both sexes	1	7	5	8	6	10	8	12	6	1	1	-
Male	1	6	5	7	3	6	2	3	2	-	-	-
Female	-	1	-	1	3	4	6	9	4	1	1	-
Black, both sexes	5	-	-	6	2	2	2	-	1	1	1	-
Male	4	-	-	3	2	1	1	-	-	-	-	-
Female	1	-	-	3	-	1	1	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	1	-	1	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
White, both sexes	-	-	-	-	1	-	1	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Ohio												
All races, both sexes	7	7	10	7	13	18	17	16	8	3	1	-
Male	7	6	8	2	8	11	8	7	-	2	-	-
Female	-	1	2	5	5	7	9	9	8	1	1	-
White, both sexes	7	7	9	5	11	18	15	16	8	3	1	-
Male	7	6	8	2	7	11	6	7	-	2	-	-
Female	-	1	1	3	4	7	9	9	8	1	1	-
Black, both sexes	-	-	1	2	2	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	1	2	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	72	75	88	70	92	112	124	123	89	57	7	1
Male	46	45	59	37	57	64	59	49	31	13	-	1
Female	26	30	29	33	35	48	65	74	58	44	7	-
White, both sexes	61	67	81	64	84	102	114	116	83	54	7	1
Male	40	41	56	34	50	55	53	46	28	13	-	1
Female	21	26	25	30	34	47	61	70	55	41	7	-
Black, both sexes	6	6	1	3	4	4	6	4	2	3	-	-
Male	2	3	-	1	4	4	3	2	1	-	-	-
Female	4	3	1	2	-	-	3	2	1	3	-	-
American Indian 1/, both sexes	4	2	5	2	3	6	4	3	4	-	-	-
Male	3	1	3	1	2	5	3	1	2	-	-	-
Female	1	1	2	1	1	1	1	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	-	-	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Oregon												
All races, both sexes	1	-	2	3	2	2	3	4	2	1	-	-
Male	1	-	1	3	1	1	1	1	1	1	-	-
Female	-	-	1	-	1	1	2	3	1	-	-	-
White, both sexes	1	-	2	3	2	2	3	3	2	1	-	-
Male	1	-	1	3	1	1	1	-	1	1	-	-
Female	-	-	1	-	1	1	2	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	8	8	11	26	23	25	24	9	7	1	-
Male	2	5	4	8	17	14	17	8	2	1	-	-
Female	3	3	4	3	9	9	8	16	7	6	1	-
White, both sexes	4	7	6	10	22	22	22	22	8	7	1	-
Male	2	4	4	7	15	14	15	6	2	1	-	-
Female	2	3	2	3	7	8	7	16	6	6	1	-
Black, both sexes	1	1	2	1	4	1	3	2	1	-	-	-
Male	-	1	-	1	2	-	2	2	-	-	-	-
Female	1	-	2	-	2	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Rhode Island												
All races, both sexes	-	-	1	1	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
White, both sexes	-	-	1	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	16	22	17	17	26	17	8	5	2	1	2
Male	8	10	16	9	9	17	8	5	4	-	-	2
Female	6	6	6	8	8	9	9	3	1	2	1	-
White, both sexes	6	6	9	7	6	13	10	5	4	-	-	1
Male	5	4	5	4	3	8	5	3	3	-	-	1
Female	1	2	4	3	3	5	5	2	1	-	-	-
Black, both sexes	8	10	13	9	10	13	7	3	1	2	1	1
Male	3	6	11	5	6	9	3	2	1	-	-	1
Female	5	4	2	4	4	4	4	1	-	2	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
South Dakota												
All races, both sexes	-	-	2	1	-	2	1	1	2	1	-	-
Male	-	-	1	1	-	1	1	1	1	-	-	-
Female	-	-	1	-	-	1	-	-	1	1	-	-
White, both sexes	-	-	2	1	-	2	1	1	2	1	-	-
Male	-	-	1	1	-	1	1	1	1	-	-	-
Female	-	-	1	-	-	1	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	7	5	1	6	5	6	6	-	1	-	-
Male	3	4	3	-	4	2	3	2	-	-	-	-
Female	1	3	2	1	2	3	3	4	-	1	-	-
White, both sexes	4	6	3	1	5	5	6	6	-	1	-	-
Male	3	4	2	-	3	2	3	2	-	-	-	-
Female	1	2	1	1	2	3	3	4	-	1	-	-
Black, both sexes	-	1	2	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Texas												
All races, both sexes	9	7	15	16	19	29	31	24	20	7	-	-
Male	7	5	10	8	11	11	14	6	1	1	-	-
Female	2	2	5	8	8	18	17	18	19	6	-	-
White, both sexes	5	5	12	14	16	28	30	23	20	6	-	-
Male	4	3	7	7	9	11	14	6	1	1	-	-
Female	1	2	5	7	7	17	16	17	19	5	-	-
Black, both sexes	3	2	3	2	2	1	1	1	-	1	-	-
Male	2	2	3	1	2	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	2	2	1	4	2	3	1	2	1	-
Male	1	-	2	2	-	2	-	2	-	-	-	-
Female	-	-	-	-	1	2	2	1	1	2	1	-
White, both sexes	1	-	1	2	1	4	2	3	1	2	1	-
Male	1	-	1	2	-	2	-	2	-	-	-	-
Female	-	-	-	-	1	2	2	1	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Vermont												
All races, both sexes	-	-	-	1	1	1	-	2	1	1	-	-
Male	-	-	-	1	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	1	-	-
White, both sexes	-	-	-	1	1	1	-	2	1	1	-	-
Male	-	-	-	1	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	46	50	66	41	49	52	69	58	35	11	3	-
Male	39	32	46	29	28	27	26	22	11	2	-	-
Female	7	18	20	12	21	25	43	36	24	9	3	-
White, both sexes	30	35	50	29	32	36	54	45	29	10	3	-
Male	26	21	34	22	19	19	18	18	9	2	-	-
Female	4	14	16	7	13	17	36	27	20	8	3	-
Black, both sexes	16	14	16	12	17	16	15	12	6	1	-	-
Male	13	10	12	7	9	8	8	4	2	-	-	-
Female	3	4	4	5	8	8	7	8	4	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Washington												
All races, both sexes	1	-	2	3	-	5	6	5	2	2	-	-
Male	1	-	2	1	-	4	2	2	-	-	-	-
Female	-	-	-	2	-	1	4	3	2	2	-	-
White, both sexes	1	-	2	1	-	5	6	5	2	2	-	-
Male	1	-	2	-	-	4	2	2	-	-	-	-
Female	-	-	-	1	-	1	4	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	2	2	3	3	3	1	2	-	-	-	-
Male	-	2	2	2	1	1	1	-	-	-	-	-
Female	1	-	-	1	2	2	-	2	-	-	-	-
White, both sexes	1	2	2	3	3	3	1	2	-	-	-	-
Male	-	2	2	2	1	1	1	-	-	-	-	-
Female	1	-	-	1	2	2	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Wisconsin												
All races, both sexes	1	3	-	-	2	3	10	9	9	3	-	-
Male	1	2	-	-	1	1	5	2	4	1	-	-
Female	-	1	-	-	1	2	5	7	5	2	-	-
White, both sexes	1	3	-	-	2	3	10	9	9	3	-	-
Male	1	2	-	-	1	1	5	2	4	1	-	-
Female	-	1	-	-	1	2	5	7	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	2	1	4	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	1	3	-	-	1	-	-
White, both sexes	-	-	-	-	2	1	4	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	1	3	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
United States												
All races, both sexes	280,031	9	1	2	5	14	57	156	455	1,084	2,847	5,737
Male	146,405	6	1	1	3	11	47	129	368	838	2,172	4,372
Female	133,626	3	-	1	2	3	10	27	87	246	675	1,365
White, both sexes	244,607	3	1	2	4	10	42	109	330	788	2,221	4,399
Male	127,786	3	1	1	3	9	35	96	271	624	1,754	3,445
Female	116,821	-	-	1	1	1	7	13	59	164	467	954
Black, both sexes	29,220	5	-	-	-	4	15	39	97	251	528	1,147
Male	15,030	2	-	-	-	2	12	26	72	174	336	771
Female	14,190	3	-	-	-	2	3	13	25	77	192	376
American Indian 1/, both sexes	1,163	-	-	-	-	-	-	-	7	9	29	52
Male	682	-	-	-	-	-	-	-	6	9	21	41
Female	481	-	-	-	-	-	-	-	1	-	8	11
Asian or Pacific Islander 2/, both sexes	5,041	1	-	-	1	-	-	8	21	36	69	139
Male	2,907	1	-	-	-	-	-	7	19	31	61	115
Female	2,134	-	-	-	1	-	-	1	2	5	8	24
Alabama												
All races, both sexes	3,364	1	-	-	-	1	3	3	7	17	35	91
Male	1,754	1	-	-	-	1	3	2	6	13	24	61
Female	1,610	-	-	-	-	-	-	1	1	4	11	30
White, both sexes	2,705	1	-	-	-	1	2	1	3	11	28	57
Male	1,432	1	-	-	-	1	2	1	2	9	21	46
Female	1,273	-	-	-	-	-	-	-	1	2	7	11
Black, both sexes	650	-	-	-	-	-	1	2	4	6	7	33
Male	316	-	-	-	-	-	1	1	4	4	3	14
Female	334	-	-	-	-	-	-	1	-	2	4	19
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Alaska												
All races, both sexes	267	-	-	-	-	1	-	-	-	3	7	10
Male	182	-	-	-	-	1	-	-	-	3	6	9
Female	85	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	222	-	-	-	-	1	-	-	-	2	4	8
Male	151	-	-	-	-	1	-	-	-	2	4	7
Female	71	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	36	-	-	-	-	-	-	-	-	1	1	2
Male	26	-	-	-	-	-	-	-	-	1	1	2
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5,916	-	-	1	-	-	3	2	12	27	56	115
Male	3,299	-	-	-	-	-	3	1	10	21	47	87
Female	2,617	-	-	1	-	-	-	1	2	6	9	28
White, both sexes	5,601	-	-	1	-	-	3	1	10	24	49	104
Male	3,107	-	-	-	-	-	3	1	8	18	41	76
Female	2,494	-	-	1	-	-	-	-	2	6	8	28
Black, both sexes	167	-	-	-	-	-	-	1	2	1	2	6
Male	110	-	-	-	-	-	-	-	2	1	2	6
Female	57	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	97	-	-	-	-	-	-	-	-	2	4	2
Male	57	-	-	-	-	-	-	-	-	2	3	2
Female	40	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	-	-	-	-	1	3
Male	25	-	-	-	-	-	-	-	-	-	1	3
Female	26	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Arkansas												
All races, both sexes	2,012	-	-	-	-	1	-	5	6	12	16	52
Male	1,014	-	-	-	-	1	-	4	3	9	10	39
Female	998	-	-	-	-	-	-	1	3	3	6	13
White, both sexes	1,740	-	-	-	-	1	-	2	5	9	11	36
Male	888	-	-	-	-	1	-	2	3	6	6	27
Female	852	-	-	-	-	-	-	-	2	3	5	9
Black, both sexes	263	-	-	-	-	-	-	3	1	3	5	15
Male	119	-	-	-	-	-	-	2	-	3	4	11
Female	144	-	-	-	-	-	-	1	1	-	1	4
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	33,253	-	-	1	2	3	2	13	30	110	285	637
Male	17,608	-	-	1	1	2	2	12	25	89	228	497
Female	15,645	-	-	-	1	1	-	1	5	21	57	140
White, both sexes	28,200	-	-	1	1	3	2	10	20	75	216	498
Male	14,865	-	-	1	1	2	2	10	15	61	175	395
Female	13,335	-	-	-	-	1	-	-	5	14	41	103
Black, both sexes	2,530	-	-	-	-	-	-	2	5	19	40	88
Male	1,337	-	-	-	-	-	-	1	5	13	25	59
Female	1,193	-	-	-	-	-	-	1	-	6	15	29
American Indian 1/, both sexes	145	-	-	-	-	-	-	-	-	2	3	3
Male	78	-	-	-	-	-	-	-	-	2	3	2
Female	67	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2,378	-	-	-	1	-	-	1	5	14	26	48
Male	1,328	-	-	-	-	-	-	1	5	13	25	41
Female	1,050	-	-	-	1	-	-	-	-	1	1	7

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Colorado												
All races, both sexes	2,658	1	1	-	-	1	-	4	6	14	30	65
Male	1,505	1	1	-	-	1	-	3	6	12	24	53
Female	1,153	-	-	-	-	-	-	1	-	2	6	12
White, both sexes	2,556	1	1	-	-	-	-	4	5	14	29	58
Male	1,444	1	1	-	-	-	-	3	5	12	23	48
Female	1,112	-	-	-	-	-	-	1	-	2	6	10
Black, both sexes	69	-	-	-	-	1	-	-	1	-	-	7
Male	46	-	-	-	-	1	-	-	1	-	-	5
Female	23	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3,328	-	-	-	-	-	1	1	4	16	26	41
Male	1,655	-	-	-	-	-	1	1	4	13	19	35
Female	1,673	-	-	-	-	-	-	-	-	3	7	6
White, both sexes	3,120	-	-	-	-	-	1	1	4	13	23	28
Male	1,547	-	-	-	-	-	1	1	4	10	18	24
Female	1,573	-	-	-	-	-	-	-	-	3	5	4
Black, both sexes	185	-	-	-	-	-	-	-	-	3	3	12
Male	95	-	-	-	-	-	-	-	-	3	1	10
Female	90	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Delaware												
All races, both sexes	939	-	-	-	-	-	-	-	2	3	11	32
Male	476	-	-	-	-	-	-	-	1	2	8	23
Female	463	-	-	-	-	-	-	-	1	1	3	9
White, both sexes	802	-	-	-	-	-	-	-	-	3	10	22
Male	399	-	-	-	-	-	-	-	-	2	7	16
Female	403	-	-	-	-	-	-	-	-	1	3	6
Black, both sexes	130	-	-	-	-	-	-	-	2	-	1	10
Male	75	-	-	-	-	-	-	-	1	-	1	7
Female	55	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	906	-	-	-	-	-	1	1	1	2	9	21
Male	460	-	-	-	-	-	1	1	1	1	6	14
Female	446	-	-	-	-	-	-	-	-	1	3	7
White, both sexes	223	-	-	-	-	-	-	-	-	-	3	5
Male	123	-	-	-	-	-	-	-	-	-	2	5
Female	100	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	677	-	-	-	-	-	1	1	1	2	6	16
Male	333	-	-	-	-	-	1	1	1	1	4	9
Female	344	-	-	-	-	-	-	-	-	1	2	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Florida												
All races, both sexes	24,007	-	-	-	1	1	4	7	24	61	190	387
Male	12,921	-	-	-	1	-	2	4	21	48	147	298
Female	11,086	-	-	-	-	1	2	3	3	13	43	89
White, both sexes	21,911	-	-	-	1	-	3	6	16	44	142	310
Male	11,846	-	-	-	1	-	2	4	15	35	112	252
Female	10,065	-	-	-	-	-	1	2	1	9	30	58
Black, both sexes	1,983	-	-	-	-	1	1	1	7	17	44	70
Male	1,006	-	-	-	-	-	-	-	5	13	31	40
Female	977	-	-	-	-	1	1	1	2	4	13	30
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	98	-	-	-	-	-	-	-	-	-	4	7
Male	62	-	-	-	-	-	-	-	-	-	4	6
Female	36	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	5,141	-	-	-	-	-	2	7	16	34	91	176
Male	2,841	-	-	-	-	-	2	5	9	25	74	144
Female	2,300	-	-	-	-	-	-	2	7	9	17	32
White, both sexes	3,878	-	-	-	-	-	2	4	8	13	60	100
Male	2,197	-	-	-	-	-	2	2	6	10	51	86
Female	1,681	-	-	-	-	-	-	2	2	3	9	14
Black, both sexes	1,220	-	-	-	-	-	-	3	8	21	28	75
Male	620	-	-	-	-	-	-	3	3	15	20	58
Female	600	-	-	-	-	-	-	-	5	6	8	17
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	-	-	3	1
Male	18	-	-	-	-	-	-	-	-	-	3	-
Female	16	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Hawaii												
All races, both sexes	857	-	-	-	-	-	-	1	5	6	10	20
Male	567	-	-	-	-	-	-	1	5	3	6	16
Female	290	-	-	-	-	-	-	-	-	3	4	4
White, both sexes	259	-	-	-	-	-	-	-	-	-	2	4
Male	171	-	-	-	-	-	-	-	-	-	1	3
Female	88	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	1	1	-	1
Male	5	-	-	-	-	-	-	-	1	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	591	-	-	-	-	-	-	1	4	5	8	15
Male	391	-	-	-	-	-	-	1	4	2	5	12
Female	200	-	-	-	-	-	-	-	-	3	3	3
Idaho												
All races, both sexes	749	-	-	-	-	-	-	1	-	3	5	7
Male	427	-	-	-	-	-	-	1	-	3	4	4
Female	322	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	744	-	-	-	-	-	-	1	-	3	5	7
Male	424	-	-	-	-	-	-	1	-	3	4	4
Female	320	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Illinois												
All races, both sexes	10,759	3	-	-	-	-	-	7	14	35	117	225
Male	5,533	1	-	-	-	-	-	6	14	27	80	166
Female	5,226	2	-	-	-	-	-	1	-	8	37	59
White, both sexes	9,099	-	-	-	-	-	-	4	8	26	86	155
Male	4,606	-	-	-	-	-	-	3	8	22	66	120
Female	4,493	-	-	-	-	-	-	1	-	4	20	35
Black, both sexes	1,539	3	-	-	-	-	-	3	6	9	30	65
Male	855	1	-	-	-	-	-	3	6	5	14	42
Female	684	2	-	-	-	-	-	-	-	4	16	23
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	116	-	-	-	-	-	-	-	-	-	1	5
Male	69	-	-	-	-	-	-	-	-	-	-	4
Female	47	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	5,625	-	-	-	-	1	2	6	17	27	91	138
Male	2,960	-	-	-	-	-	1	6	11	17	69	104
Female	2,665	-	-	-	-	1	1	-	6	10	22	34
White, both sexes	5,151	-	-	-	-	-	2	6	13	25	78	120
Male	2,694	-	-	-	-	-	1	6	9	17	60	92
Female	2,457	-	-	-	-	-	1	-	4	8	18	28
Black, both sexes	452	-	-	-	-	1	-	-	4	2	12	16
Male	250	-	-	-	-	-	-	-	2	-	8	10
Female	202	-	-	-	-	1	-	-	2	2	4	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Iowa												
All races, both sexes	3,686	-	-	-	-	-	1	2	6	8	33	64
Male	1,858	-	-	-	-	-	1	2	4	7	24	48
Female	1,828	-	-	-	-	-	-	-	2	1	9	16
White, both sexes	3,629	-	-	-	-	-	1	1	6	6	31	59
Male	1,823	-	-	-	-	-	1	1	4	6	22	46
Female	1,806	-	-	-	-	-	-	-	2	-	9	13
Black, both sexes	45	-	-	-	-	-	-	-	-	2	2	3
Male	26	-	-	-	-	-	-	-	-	1	2	1
Female	19	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	1	-	-	-	2
Male	9	-	-	-	-	-	-	1	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	2,303	-	-	-	-	-	-	1	5	3	20	59
Male	1,268	-	-	-	-	-	-	1	4	3	12	41
Female	1,035	-	-	-	-	-	-	-	1	-	8	18
White, both sexes	2,173	-	-	-	-	-	-	1	4	3	18	52
Male	1,195	-	-	-	-	-	-	1	4	3	12	36
Female	978	-	-	-	-	-	-	-	-	-	6	16
Black, both sexes	105	-	-	-	-	-	-	-	1	-	2	5
Male	57	-	-	-	-	-	-	-	-	-	-	4
Female	48	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Kentucky												
All races, both sexes	3,090	-	-	-	-	-	1	2	8	22	49	72
Male	1,566	-	-	-	-	-	-	2	8	14	40	60
Female	1,524	-	-	-	-	-	1	-	-	8	9	12
White, both sexes	2,911	-	-	-	-	-	1	2	5	19	41	69
Male	1,478	-	-	-	-	-	-	2	5	12	36	57
Female	1,433	-	-	-	-	-	1	-	-	7	5	12
Black, both sexes	171	-	-	-	-	-	-	-	2	3	8	3
Male	84	-	-	-	-	-	-	-	2	2	4	3
Female	87	-	-	-	-	-	-	-	-	1	4	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,127	-	-	-	-	-	-	2	11	18	53	94
Male	1,693	-	-	-	-	-	-	2	7	14	35	69
Female	1,434	-	-	-	-	-	-	-	4	4	18	25
White, both sexes	2,320	-	-	-	-	-	-	1	6	7	36	54
Male	1,274	-	-	-	-	-	-	1	4	6	23	41
Female	1,046	-	-	-	-	-	-	-	2	1	13	13
Black, both sexes	793	-	-	-	-	-	-	1	5	11	17	40
Male	411	-	-	-	-	-	-	1	3	8	12	28
Female	382	-	-	-	-	-	-	-	2	3	5	12
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Maine												
All races, both sexes	1,140	-	-	-	-	-	-	2	4	4	12	24
Male	626	-	-	-	-	-	-	2	3	2	10	19
Female	514	-	-	-	-	-	-	-	1	2	2	5
White, both sexes	1,131	-	-	-	-	-	-	2	3	4	10	23
Male	618	-	-	-	-	-	-	2	2	2	9	18
Female	513	-	-	-	-	-	-	-	1	2	1	5
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	2	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5,241	-	-	-	-	-	-	2	5	21	50	115
Male	2,623	-	-	-	-	-	-	2	3	18	38	83
Female	2,618	-	-	-	-	-	-	-	2	3	12	32
White, both sexes	3,838	-	-	-	-	-	-	1	3	12	31	69
Male	1,935	-	-	-	-	-	-	1	2	10	23	52
Female	1,903	-	-	-	-	-	-	-	1	2	8	17
Black, both sexes	1,326	-	-	-	-	-	-	-	-	8	17	44
Male	641	-	-	-	-	-	-	-	-	7	13	30
Female	685	-	-	-	-	-	-	-	-	1	4	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	74	-	-	-	-	-	-	1	2	1	2	2
Male	46	-	-	-	-	-	-	1	1	1	2	1
Female	28	-	-	-	-	-	-	-	1	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Massachusetts												
All races, both sexes	5,434	-	-	-	-	-	1	5	9	12	57	78
Male	2,714	-	-	-	-	-	1	4	9	9	45	63
Female	2,720	-	-	-	-	-	-	1	-	3	12	15
White, both sexes	5,233	-	-	-	-	-	1	5	9	9	52	71
Male	2,601	-	-	-	-	-	1	4	9	7	42	56
Female	2,632	-	-	-	-	-	-	1	-	2	10	15
Black, both sexes	151	-	-	-	-	-	-	-	-	2	5	4
Male	88	-	-	-	-	-	-	-	-	1	3	4
Female	63	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	44	-	-	-	-	-	-	-	-	1	-	2
Male	20	-	-	-	-	-	-	-	-	1	-	2
Female	24	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	11,918	-	-	-	-	-	2	5	20	50	139	276
Male	6,210	-	-	-	-	-	1	4	17	36	108	200
Female	5,708	-	-	-	-	-	1	1	3	14	31	76
White, both sexes	9,843	-	-	-	-	-	2	3	17	37	101	205
Male	5,109	-	-	-	-	-	1	2	15	26	85	156
Female	4,734	-	-	-	-	-	1	1	2	11	16	49
Black, both sexes	1,971	-	-	-	-	-	-	2	2	12	36	67
Male	1,053	-	-	-	-	-	-	2	2	9	22	42
Female	918	-	-	-	-	-	-	-	-	3	14	25
American Indian 1/, both sexes	61	-	-	-	-	-	-	-	1	1	2	3
Male	24	-	-	-	-	-	-	-	-	1	1	1
Female	37	-	-	-	-	-	-	-	1	-	1	2
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Minnesota												
All races, both sexes	3,078	-	-	-	-	-	2	3	6	16	41	75
Male	1,795	-	-	-	-	-	2	3	3	13	33	67
Female	1,283	-	-	-	-	-	-	-	3	3	8	8
White, both sexes	2,972	-	-	-	-	-	1	3	5	16	37	68
Male	1,723	-	-	-	-	-	1	3	3	13	29	60
Female	1,249	-	-	-	-	-	-	-	2	3	8	8
Black, both sexes	56	-	-	-	-	-	1	-	-	-	2	3
Male	35	-	-	-	-	-	1	-	-	-	2	3
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	-	1	3
Male	21	-	-	-	-	-	-	-	-	-	1	3
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	1	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	1	-	-	-
Mississippi												
All races, both sexes	2,405	-	-	-	-	-	-	1	15	22	49	74
Male	1,180	-	-	-	-	-	-	1	12	17	34	54
Female	1,225	-	-	-	-	-	-	-	3	5	15	20
White, both sexes	1,794	-	-	-	-	-	-	-	11	15	33	45
Male	866	-	-	-	-	-	-	-	9	11	22	32
Female	928	-	-	-	-	-	-	-	2	4	11	13
Black, both sexes	601	-	-	-	-	-	-	1	4	6	16	26
Male	310	-	-	-	-	-	-	1	3	5	12	22
Female	291	-	-	-	-	-	-	-	1	1	4	4
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	2
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Missouri												
All races, both sexes	5,589	-	-	-	-	1	3	3	14	14	57	119
Male	2,749	-	-	-	-	1	2	3	9	9	45	87
Female	2,840	-	-	-	-	-	1	-	5	5	12	32
White, both sexes	5,058	-	-	-	-	1	2	3	10	12	48	93
Male	2,485	-	-	-	-	1	1	3	6	7	38	67
Female	2,573	-	-	-	-	-	1	-	4	5	10	26
Black, both sexes	501	-	-	-	-	-	1	-	4	2	9	24
Male	248	-	-	-	-	-	1	-	3	2	7	18
Female	253	-	-	-	-	-	-	-	1	-	2	6
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	729	-	-	-	-	-	-	-	3	3	17	21
Male	437	-	-	-	-	-	-	-	3	1	14	16
Female	292	-	-	-	-	-	-	-	-	2	3	5
White, both sexes	702	-	-	-	-	-	-	-	2	2	15	21
Male	422	-	-	-	-	-	-	-	2	1	13	16
Female	280	-	-	-	-	-	-	-	-	1	2	5
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	1	-	2	-
Male	14	-	-	-	-	-	-	-	1	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Nebraska												
All races, both sexes	1,308	-	-	-	-	-	1	-	1	3	8	27
Male	675	-	-	-	-	-	1	-	1	3	8	19
Female	633	-	-	-	-	-	-	-	-	-	-	8
White, both sexes	1,263	-	-	-	-	-	1	-	1	1	8	23
Male	651	-	-	-	-	-	1	-	1	1	8	16
Female	612	-	-	-	-	-	-	-	-	-	-	7
Black, both sexes	33	-	-	-	-	-	-	-	-	2	-	4
Male	18	-	-	-	-	-	-	-	-	2	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2,052	-	-	-	-	-	2	1	5	9	31	53
Male	1,280	-	-	-	-	-	2	1	4	8	25	42
Female	772	-	-	-	-	-	-	-	1	1	6	11
White, both sexes	1,830	-	-	-	-	-	2	-	5	5	26	45
Male	1,142	-	-	-	-	-	2	-	4	4	21	37
Female	688	-	-	-	-	-	-	-	1	1	5	8
Black, both sexes	134	-	-	-	-	-	-	1	-	3	4	4
Male	82	-	-	-	-	-	-	1	-	3	3	2
Female	52	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	68	-	-	-	-	-	-	-	-	-	1	3
Male	46	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
New Hampshire												
All races, both sexes	1,172	-	-	-	-	-	-	-	2	3	12	26
Male	617	-	-	-	-	-	-	-	2	2	9	21
Female	555	-	-	-	-	-	-	-	-	1	3	5
White, both sexes	1,164	-	-	-	-	-	-	-	2	3	12	25
Male	612	-	-	-	-	-	-	-	2	2	9	20
Female	552	-	-	-	-	-	-	-	-	1	3	5
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9,411	-	-	-	-	-	-	4	9	33	77	127
Male	4,548	-	-	-	-	-	-	3	8	29	63	96
Female	4,863	-	-	-	-	-	-	1	1	4	14	31
White, both sexes	8,449	-	-	-	-	-	-	3	8	22	63	99
Male	4,059	-	-	-	-	-	-	3	7	19	57	77
Female	4,390	-	-	-	-	-	-	-	1	3	6	22
Black, both sexes	816	-	-	-	-	-	-	1	1	9	13	24
Male	411	-	-	-	-	-	-	-	1	8	5	15
Female	405	-	-	-	-	-	-	1	-	1	8	9
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	137	-	-	-	-	-	-	-	-	2	1	4
Male	74	-	-	-	-	-	-	-	-	2	1	4
Female	63	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
New Mexico												
All races, both sexes	1,751	-	-	-	-	1	-	-	6	7	15	29
Male	956	-	-	-	-	1	-	-	5	6	13	19
Female	795	-	-	-	-	-	-	-	1	1	2	10
White, both sexes	1,654	-	-	-	-	1	-	-	6	7	13	25
Male	899	-	-	-	-	1	-	-	5	6	13	15
Female	755	-	-	-	-	-	-	-	1	1	-	10
Black, both sexes	34	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	55	-	-	-	-	-	-	-	-	-	1	2
Male	39	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	29,770	-	-	-	1	-	9	16	21	78	171	394
Male	13,682	-	-	-	-	-	9	8	19	63	128	307
Female	16,088	-	-	-	1	-	-	8	2	15	43	87
White, both sexes	25,062	-	-	-	1	-	6	8	12	53	129	272
Male	11,569	-	-	-	-	-	6	5	10	47	101	225
Female	13,493	-	-	-	1	-	-	3	2	6	28	47
Black, both sexes	3,967	-	-	-	-	-	3	6	6	22	33	104
Male	1,715	-	-	-	-	-	3	2	6	13	19	67
Female	2,252	-	-	-	-	-	-	4	-	9	14	37
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	717	-	-	-	-	-	-	2	3	3	9	18
Male	391	-	-	-	-	-	-	1	3	3	8	15
Female	326	-	-	-	-	-	-	1	-	-	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
North Carolina												
All races, both sexes	7,296	-	-	-	-	-	2	6	17	47	123	196
Male	4,107	-	-	-	-	-	1	6	14	35	91	159
Female	3,189	-	-	-	-	-	1	-	3	12	32	37
White, both sexes	5,787	-	-	-	-	-	-	2	12	32	80	131
Male	3,278	-	-	-	-	-	-	2	11	23	63	109
Female	2,509	-	-	-	-	-	-	-	1	9	17	22
Black, both sexes	1,403	-	-	-	-	-	2	4	5	15	40	60
Male	767	-	-	-	-	-	1	4	3	12	26	47
Female	636	-	-	-	-	-	1	-	2	3	14	13
American Indian 1/, both sexes	82	-	-	-	-	-	-	-	-	-	2	5
Male	50	-	-	-	-	-	-	-	-	-	1	3
Female	32	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	646	-	-	-	-	-	-	1	-	3	6	11
Male	337	-	-	-	-	-	-	1	-	2	5	8
Female	309	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	624	-	-	-	-	-	-	1	-	2	6	10
Male	323	-	-	-	-	-	-	1	-	1	5	7
Female	301	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Ohio												
All races, both sexes	13,319	-	-	-	-	-	3	8	15	55	111	249
Male	6,821	-	-	-	-	-	2	7	13	41	84	188
Female	6,498	-	-	-	-	-	1	1	2	14	27	61
White, both sexes	11,937	-	-	-	-	-	3	7	14	45	90	202
Male	6,087	-	-	-	-	-	2	6	12	34	72	159
Female	5,850	-	-	-	-	-	1	1	2	11	18	43
Black, both sexes	1,338	-	-	-	-	-	-	1	1	9	21	47
Male	711	-	-	-	-	-	-	1	1	6	12	29
Female	627	-	-	-	-	-	-	-	-	3	9	18
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	-	1	-	-
Male	18	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4,168	-	-	-	-	-	-	4	10	22	46	88
Male	2,258	-	-	-	-	-	-	3	9	17	29	63
Female	1,910	-	-	-	-	-	-	1	1	5	17	25
White, both sexes	3,673	-	-	-	-	-	-	2	6	17	36	76
Male	1,967	-	-	-	-	-	-	2	5	15	21	55
Female	1,706	-	-	-	-	-	-	-	1	2	15	21
Black, both sexes	231	-	-	-	-	-	-	2	1	5	5	2
Male	126	-	-	-	-	-	-	1	1	2	4	2
Female	105	-	-	-	-	-	-	1	-	3	1	-
American Indian 1/, both sexes	250	-	-	-	-	-	-	-	3	-	5	10
Male	156	-	-	-	-	-	-	-	3	-	4	6
Female	94	-	-	-	-	-	-	-	-	-	1	4
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Oregon												
All races, both sexes	2,737	-	-	-	-	1	-	-	2	14	27	63
Male	1,659	-	-	-	-	1	-	-	1	11	22	52
Female	1,078	-	-	-	-	-	-	-	1	3	5	11
White, both sexes	2,667	-	-	-	-	1	-	-	2	13	26	59
Male	1,621	-	-	-	-	1	-	-	1	10	22	49
Female	1,046	-	-	-	-	-	-	-	1	3	4	10
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	1	1	2
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	15	-	-	-	-	-	-	-	-	-	1	1
Pennsylvania												
All races, both sexes	14,157	-	-	-	1	-	3	4	15	45	101	229
Male	7,019	-	-	-	1	-	3	3	12	38	75	172
Female	7,138	-	-	-	-	-	-	1	3	7	26	57
White, both sexes	12,864	-	-	-	1	-	1	2	9	32	75	179
Male	6,324	-	-	-	1	-	1	2	8	30	56	138
Female	6,540	-	-	-	-	-	-	-	1	2	19	41
Black, both sexes	1,235	-	-	-	-	-	2	2	5	11	24	46
Male	659	-	-	-	-	-	2	1	3	6	17	30
Female	576	-	-	-	-	-	-	1	2	5	7	16
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	-	-	1	2	2	4
Male	33	-	-	-	-	-	-	-	1	2	2	4
Female	21	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Rhode Island												
All races, both sexes	1,423	-	-	-	-	-	-	-	-	1	13	16
Male	638	-	-	-	-	-	-	-	-	1	12	15
Female	785	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	1,392	-	-	-	-	-	-	-	-	-	13	15
Male	624	-	-	-	-	-	-	-	-	-	12	14
Female	768	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,877	-	-	-	-	-	1	2	4	19	43	87
Male	1,692	-	-	-	-	-	-	2	2	13	34	59
Female	1,185	-	-	-	-	-	1	-	2	6	9	28
White, both sexes	2,241	-	-	-	-	-	-	2	2	13	30	67
Male	1,350	-	-	-	-	-	-	2	1	9	27	48
Female	891	-	-	-	-	-	-	-	1	4	3	19
Black, both sexes	630	-	-	-	-	-	1	-	2	6	13	19
Male	337	-	-	-	-	-	-	-	1	4	7	10
Female	293	-	-	-	-	-	1	-	1	2	6	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
South Dakota												
All races, both sexes	753	-	-	-	-	-	-	-	1	-	5	15
Male	379	-	-	-	-	-	-	-	1	-	4	13
Female	374	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	722	-	-	-	-	-	-	-	1	-	4	13
Male	362	-	-	-	-	-	-	-	1	-	3	11
Female	360	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6,174	2	-	-	-	-	1	2	15	32	104	198
Male	3,318	1	-	-	-	-	1	2	14	25	69	158
Female	2,856	1	-	-	-	-	-	-	1	7	35	40
White, both sexes	5,348	-	-	-	-	-	1	2	13	20	83	151
Male	2,882	-	-	-	-	-	1	2	12	17	56	124
Female	2,466	-	-	-	-	-	-	-	1	3	27	27
Black, both sexes	803	2	-	-	-	-	-	-	2	12	20	44
Male	422	1	-	-	-	-	-	-	2	8	12	31
Female	381	1	-	-	-	-	-	-	-	4	8	13
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Texas												
All races, both sexes	15,067	1	-	-	-	1	3	9	32	69	200	432
Male	8,327	1	-	-	-	1	2	7	25	57	156	312
Female	6,740	-	-	-	-	-	1	2	7	12	44	120
White, both sexes	13,194	1	-	-	-	1	2	7	23	50	161	319
Male	7,282	1	-	-	-	1	1	6	17	44	129	242
Female	5,912	-	-	-	-	-	1	1	6	6	32	77
Black, both sexes	1,726	-	-	-	-	-	1	2	8	18	37	104
Male	961	-	-	-	-	-	1	1	7	12	26	62
Female	765	-	-	-	-	-	-	1	1	6	11	42
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	2	4
Male	16	-	-	-	-	-	-	-	-	-	1	4
Female	12	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	119	-	-	-	-	-	-	-	1	1	-	5
Male	68	-	-	-	-	-	-	-	1	1	-	4
Female	51	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	914	-	-	-	-	-	-	-	-	7	10	22
Male	591	-	-	-	-	-	-	-	-	6	8	18
Female	323	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	897	-	-	-	-	-	-	-	-	5	9	22
Male	578	-	-	-	-	-	-	-	-	4	7	18
Female	319	-	-	-	-	-	-	-	-	1	2	4
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Vermont												
All races, both sexes	622	-	-	-	-	-	-	-	1	1	9	17
Male	334	-	-	-	-	-	-	-	1	-	8	15
Female	288	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	621	-	-	-	-	-	-	-	1	1	9	17
Male	333	-	-	-	-	-	-	-	1	-	8	15
Female	288	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4,903	-	-	-	-	1	1	6	12	25	68	144
Male	2,705	-	-	-	-	1	1	6	8	17	56	110
Female	2,198	-	-	-	-	-	-	-	4	8	12	34
White, both sexes	3,971	-	-	-	-	-	-	4	10	17	48	104
Male	2,198	-	-	-	-	-	-	4	7	14	43	77
Female	1,773	-	-	-	-	-	-	-	3	3	5	27
Black, both sexes	871	-	-	-	-	1	1	-	1	8	19	37
Male	469	-	-	-	-	1	1	-	-	3	12	31
Female	402	-	-	-	-	-	-	-	1	5	7	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	58	-	-	-	-	-	-	2	1	-	1	3
Male	37	-	-	-	-	-	-	2	1	-	1	2
Female	21	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Washington												
All races, both sexes	5,118	-	-	-	-	-	2	1	17	21	47	92
Male	2,809	-	-	-	-	-	2	1	16	17	37	72
Female	2,309	-	-	-	-	-	-	-	1	4	10	20
White, both sexes	4,828	-	-	-	-	-	2	1	14	19	41	81
Male	2,630	-	-	-	-	-	2	1	13	15	31	61
Female	2,198	-	-	-	-	-	-	-	1	4	10	20
Black, both sexes	103	-	-	-	-	-	-	-	1	1	1	5
Male	60	-	-	-	-	-	-	-	1	1	1	5
Female	43	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	50	-	-	-	-	-	-	-	-	-	2	2
Male	27	-	-	-	-	-	-	-	-	-	2	2
Female	23	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	137	-	-	-	-	-	-	-	2	1	3	4
Male	92	-	-	-	-	-	-	-	2	1	3	4
Female	45	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2,358	-	-	-	-	-	1	1	8	15	34	60
Male	1,273	-	-	-	-	-	1	1	5	10	23	43
Female	1,085	-	-	-	-	-	-	-	3	5	11	17
White, both sexes	2,291	-	-	-	-	-	1	1	7	15	34	59
Male	1,240	-	-	-	-	-	1	1	4	10	23	43
Female	1,051	-	-	-	-	-	-	-	3	5	11	16
Black, both sexes	66	-	-	-	-	-	-	-	1	-	-	1
Male	33	-	-	-	-	-	-	-	1	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Wisconsin												
All races, both sexes	4,209	1	-	-	-	-	-	4	12	9	27	69
Male	2,276	1	-	-	-	-	-	4	10	7	22	53
Female	1,933	-	-	-	-	-	-	-	2	2	5	16
White, both sexes	4,011	-	-	-	-	-	-	4	9	9	24	59
Male	2,171	-	-	-	-	-	-	4	9	7	21	46
Female	1,840	-	-	-	-	-	-	-	-	2	3	13
Black, both sexes	172	-	-	-	-	-	-	-	3	-	3	8
Male	90	-	-	-	-	-	-	-	1	-	1	5
Female	82	-	-	-	-	-	-	-	2	-	2	3
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	1
Male	4	1	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	305	-	-	-	-	-	-	1	-	3	3	5
Male	183	-	-	-	-	-	-	1	-	1	1	1
Female	122	-	-	-	-	-	-	-	-	2	2	4
White, both sexes	302	-	-	-	-	-	-	1	-	3	2	5
Male	181	-	-	-	-	-	-	1	-	1	1	1
Female	121	-	-	-	-	-	-	-	-	2	1	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
United States												
All races, both sexes	9,153	12,755	14,982	17,563	23,651	34,546	48,014	50,911	38,213	16,097	3,756	23
Male	7,068	9,560	10,602	11,842	14,837	20,243	24,702	22,157	13,016	3,804	611	15
Female	2,085	3,195	4,380	5,721	8,814	14,303	23,312	28,754	25,197	12,293	3,145	8
White, both sexes	7,082	10,012	12,082	14,273	19,835	30,156	42,982	46,797	35,322	14,811	3,326	20
Male	5,612	7,667	8,707	9,779	12,635	17,917	22,437	20,587	12,154	3,498	537	14
Female	1,470	2,345	3,375	4,494	7,200	12,239	20,545	26,210	23,168	11,313	2,789	6
Black, both sexes	1,832	2,360	2,461	2,737	3,205	3,593	4,038	3,230	2,263	1,050	362	3
Male	1,260	1,588	1,581	1,677	1,845	1,874	1,773	1,176	601	209	50	1
Female	572	772	880	1,060	1,360	1,719	2,265	2,054	1,662	841	312	2
American Indian 1/, both sexes	67	98	119	129	141	144	159	114	69	21	5	-
Male	52	76	74	85	92	82	70	48	22	3	1	-
Female	15	22	45	44	49	62	89	66	47	18	4	-
Asian or Pacific Islander 2/, both sexes	172	285	320	424	470	653	835	770	559	215	63	-
Male	144	229	240	301	265	370	422	346	239	94	23	-
Female	28	56	80	123	205	283	413	424	320	121	40	-
Alabama												
All races, both sexes	118	166	180	229	374	470	567	550	373	143	36	-
Male	91	123	134	152	243	257	264	216	125	33	5	-
Female	27	43	46	77	131	213	303	334	248	110	31	-
White, both sexes	79	118	130	189	306	375	463	476	323	121	21	-
Male	60	86	102	126	205	213	221	190	113	31	2	-
Female	19	32	28	63	101	162	242	286	210	90	19	-
Black, both sexes	37	47	50	40	66	95	103	73	49	22	15	-
Male	29	36	32	26	37	44	43	26	11	2	3	-
Female	8	11	18	14	29	51	60	47	38	20	12	-
American Indian 1/, both sexes	1	1	-	-	1	-	1	1	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Alaska												
All races, both sexes	25	21	19	30	31	39	38	24	10	7	2	-
Male	20	15	14	25	22	32	16	12	5	2	-	-
Female	5	6	5	5	9	7	22	12	5	5	2	-
White, both sexes	22	17	15	26	25	34	31	22	8	5	2	-
Male	17	13	11	21	17	28	14	12	3	1	-	-
Female	5	4	4	5	8	6	17	10	5	4	2	-
Black, both sexes	1	-	-	-	-	1	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	4	4	5	4	6	1	2	1	-	-
Male	2	1	3	4	4	4	2	-	2	-	-	-
Female	-	2	1	-	1	-	4	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Arizona												
All races, both sexes	227	308	396	425	518	772	980	1,039	681	288	58	8
Male	185	230	298	281	331	448	525	488	258	64	16	6
Female	42	78	98	144	187	324	455	551	423	224	42	2
White, both sexes	210	287	363	389	490	737	937	997	662	277	54	6
Male	169	213	273	260	313	433	504	465	251	59	15	5
Female	41	74	90	129	177	304	433	532	411	218	39	1
Black, both sexes	13	11	20	18	19	15	17	22	8	7	3	2
Male	12	9	16	13	14	6	10	11	4	3	-	1
Female	1	2	4	5	5	9	7	11	4	4	3	1
American Indian 1/, both sexes	3	9	11	13	3	11	18	14	6	1	-	-
Male	3	7	8	6	1	7	7	9	1	1	-	-
Female	-	2	3	7	2	4	11	5	5	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	5	6	9	8	6	5	3	1	-
Male	1	1	1	2	3	2	4	3	2	1	1	-
Female	-	-	1	3	3	7	4	3	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Arkansas												
All races, both sexes	45	75	110	131	206	250	337	354	270	119	23	-
Male	32	58	76	79	125	138	171	145	92	29	3	-
Female	13	17	34	52	81	112	166	209	178	90	20	-
White, both sexes	27	59	89	107	170	225	305	329	235	108	22	-
Male	21	48	63	68	110	129	156	137	83	25	3	-
Female	6	11	26	39	60	96	149	192	152	83	19	-
Black, both sexes	17	14	21	24	34	24	31	25	35	10	1	-
Male	10	8	13	11	14	9	14	8	9	3	-	-
Female	7	6	8	13	20	15	17	17	26	7	1	-
American Indian 1/, both sexes	1	1	-	-	1	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	-	1	-	-
Male	-	1	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
California												
All races, both sexes	1,031	1,540	1,708	2,041	2,698	3,887	5,799	6,205	4,724	2,029	503	5
Male	846	1,152	1,238	1,353	1,681	2,326	2,996	2,824	1,731	509	92	3
Female	185	388	470	688	1,017	1,561	2,803	3,381	2,993	1,520	411	2
White, both sexes	818	1,182	1,378	1,611	2,208	3,235	5,008	5,507	4,187	1,808	427	5
Male	676	902	1,014	1,075	1,393	1,949	2,595	2,546	1,542	435	73	3
Female	142	280	364	536	815	1,286	2,413	2,961	2,645	1,373	354	2
Black, both sexes	143	223	201	237	280	313	339	282	221	100	37	-
Male	110	146	133	143	166	187	169	96	59	21	4	-
Female	33	77	68	94	114	126	170	186	162	79	33	-
American Indian 1/, both sexes	5	13	10	22	15	21	24	17	7	3	-	-
Male	5	10	4	15	11	12	10	3	1	-	-	-
Female	-	3	6	7	4	9	14	14	6	3	-	-
Asian or Pacific Islander 2/, both sexes	65	122	119	171	195	318	428	399	309	118	39	-
Male	55	94	87	120	111	178	222	179	129	53	15	-
Female	10	28	32	51	84	140	206	220	180	65	24	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Colorado												
All races, both sexes	117	148	171	159	233	325	417	437	341	161	27	-
Male	92	126	125	101	153	204	216	215	129	40	3	-
Female	25	22	46	58	80	121	201	222	212	121	24	-
White, both sexes	112	137	161	150	219	310	404	432	336	157	26	-
Male	87	117	121	95	145	195	212	212	126	38	3	-
Female	25	20	40	55	74	115	192	220	210	119	23	-
Black, both sexes	3	9	7	9	9	9	6	2	2	3	1	-
Male	3	7	3	6	7	6	2	2	2	1	-	-
Female	-	2	4	3	2	3	4	-	-	2	1	-
American Indian 1/, both sexes	1	-	1	-	1	4	2	1	1	-	-	-
Male	1	-	-	-	1	3	-	1	1	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	-	4	2	5	2	2	1	-	-
Male	1	2	1	-	-	-	2	-	-	1	-	-
Female	-	-	1	-	4	2	3	2	2	-	-	-
Connecticut												
All races, both sexes	73	127	147	161	234	386	579	682	527	257	66	-
Male	58	98	100	113	150	217	297	307	178	55	9	-
Female	15	29	47	48	84	169	282	375	349	202	57	-
White, both sexes	61	105	126	138	219	351	557	661	516	253	63	-
Male	49	83	90	100	139	204	285	298	177	55	9	-
Female	12	22	36	38	80	147	272	363	339	198	54	-
Black, both sexes	11	22	19	20	14	29	19	19	8	3	3	-
Male	8	15	8	11	10	11	10	7	1	-	-	-
Female	3	7	11	9	4	18	9	12	7	3	3	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	-	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	1	5	2	2	1	1	-	-
Male	1	-	1	2	1	2	2	2	-	-	-	-
Female	-	-	-	1	-	3	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Delaware												
All races, both sexes	37	35	60	63	81	110	160	173	120	36	15	1
Male	28	26	41	39	49	65	70	82	27	13	1	1
Female	9	9	19	24	32	45	90	91	93	23	14	-
White, both sexes	28	29	47	51	62	99	143	158	106	31	12	1
Male	22	22	32	33	34	57	62	74	24	12	1	1
Female	6	7	15	18	28	42	81	84	82	19	11	-
Black, both sexes	8	6	12	10	18	10	17	15	13	5	3	-
Male	6	4	8	6	14	8	8	8	3	1	-	-
Female	2	2	4	4	4	2	9	7	10	4	3	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	1	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	40	46	67	90	82	114	145	141	95	36	15	-
Male	25	29	40	60	51	65	66	53	36	9	2	-
Female	15	17	27	30	31	49	79	88	59	27	13	-
White, both sexes	3	7	14	15	13	28	41	40	39	12	3	-
Male	2	6	10	9	8	20	17	21	17	6	-	-
Female	1	1	4	6	5	8	24	19	22	6	3	-
Black, both sexes	37	39	53	75	69	84	103	100	54	24	12	-
Male	23	23	30	51	43	44	48	31	18	3	2	-
Female	14	16	23	24	26	40	55	69	36	21	10	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Florida												
All races, both sexes	629	882	1,111	1,377	1,886	3,024	4,432	4,673	3,516	1,469	332	1
Male	500	662	810	971	1,214	1,846	2,385	2,148	1,388	401	75	-
Female	129	220	301	406	672	1,178	2,047	2,525	2,128	1,068	257	1
White, both sexes	496	730	954	1,186	1,652	2,777	4,114	4,457	3,331	1,390	301	1
Male	405	574	710	841	1,075	1,721	2,239	2,069	1,339	382	70	-
Female	91	156	244	345	577	1,056	1,875	2,388	1,992	1,008	231	1
Black, both sexes	123	147	150	180	227	232	298	204	177	74	30	-
Male	86	86	96	122	134	114	135	75	47	18	4	-
Female	37	61	54	58	93	118	163	129	130	56	26	-
American Indian 1/, both sexes	1	2	1	3	-	-	4	1	1	1	-	-
Male	1	-	1	2	-	-	2	-	-	-	-	-
Female	-	2	-	1	-	-	2	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	9	3	6	8	7	15	16	11	7	4	1	-
Male	8	2	3	6	5	11	9	4	2	1	1	-
Female	1	1	3	2	2	4	7	7	5	3	-	-
Georgia												
All races, both sexes	235	320	388	407	482	640	832	746	527	206	31	1
Male	162	248	284	279	320	349	425	318	143	48	5	1
Female	73	72	104	128	162	291	407	428	384	158	26	-
White, both sexes	141	213	268	302	359	486	679	614	434	172	22	1
Male	108	173	208	218	250	278	359	277	123	44	1	1
Female	33	40	60	84	109	208	320	337	311	128	21	-
Black, both sexes	93	105	115	101	118	143	148	129	91	34	8	-
Male	53	73	74	58	67	64	64	41	20	4	3	-
Female	40	32	41	43	51	79	84	88	71	30	5	-
American Indian 1/, both sexes	-	-	3	-	1	3	1	-	1	-	-	-
Male	-	-	1	-	1	3	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	4	4	8	4	3	1	-	1	-
Male	1	2	1	3	2	4	1	-	-	-	1	-
Female	-	-	1	1	2	4	3	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Hawaii												
All races, both sexes	38	53	68	76	77	110	134	127	89	33	10	-
Male	31	41	52	60	45	76	92	69	47	19	4	-
Female	7	12	16	16	32	34	42	58	42	14	6	-
White, both sexes	18	20	20	21	19	39	35	40	26	12	3	-
Male	15	15	18	14	10	29	26	21	12	6	1	-
Female	3	5	2	7	9	10	9	19	14	6	2	-
Black, both sexes	-	-	-	-	2	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	33	48	55	56	70	98	87	63	21	7	-
Male	16	26	34	46	34	46	66	48	35	13	3	-
Female	4	7	14	9	22	24	32	39	28	8	4	-
Idaho												
All races, both sexes	13	18	28	39	58	99	145	157	127	44	5	-
Male	7	16	24	26	39	68	85	81	52	15	2	-
Female	6	2	4	13	19	31	60	76	75	29	3	-
White, both sexes	13	18	28	39	58	98	144	155	127	43	5	-
Male	7	16	24	26	39	68	85	79	52	14	2	-
Female	6	2	4	13	19	30	59	76	75	29	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Illinois												
All races, both sexes	402	508	574	677	846	1,305	1,811	1,903	1,555	634	142	1
Male	329	387	406	459	524	765	891	835	484	142	17	-
Female	73	121	168	218	322	540	920	1,068	1,071	492	125	1
White, both sexes	289	354	419	507	675	1,100	1,591	1,737	1,444	578	126	-
Male	242	275	301	344	430	648	791	763	451	127	15	-
Female	47	79	118	163	245	452	800	974	993	451	111	-
Black, both sexes	111	145	143	152	158	192	206	147	103	50	15	1
Male	86	106	96	103	86	110	90	63	30	12	2	-
Female	25	39	47	49	72	82	116	84	73	38	13	1
American Indian 1/, both sexes	-	-	2	1	-	1	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	9	10	17	13	12	14	18	8	6	1	-
Male	1	6	8	11	8	7	10	8	3	3	-	-
Female	1	3	2	6	5	5	4	10	5	3	1	-
Indiana												
All races, both sexes	176	240	293	378	507	735	981	976	693	285	79	-
Male	130	167	205	258	317	437	521	416	221	70	10	-
Female	46	73	88	120	190	298	460	560	472	215	69	-
White, both sexes	151	193	255	320	465	685	906	927	658	272	75	-
Male	109	139	175	220	288	410	482	396	213	67	10	-
Female	42	54	80	100	177	275	424	531	445	205	65	-
Black, both sexes	25	45	36	53	39	48	73	46	35	13	4	-
Male	21	27	28	34	26	27	37	19	8	3	-	-
Female	4	18	8	19	13	21	36	27	27	10	4	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	4	2	2	2	3	-	-	-	-
Male	-	1	2	4	2	-	2	1	-	-	-	-
Female	-	1	-	-	-	2	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Iowa												
All races, both sexes	103	131	148	210	259	413	609	731	597	290	81	-
Male	75	89	97	150	171	239	332	337	197	76	9	-
Female	28	42	51	60	88	174	277	394	400	214	72	-
White, both sexes	93	127	144	205	254	405	603	727	596	290	81	-
Male	68	87	96	146	169	232	329	334	197	76	9	-
Female	25	40	48	59	85	173	274	393	399	214	72	-
Black, both sexes	9	4	3	4	3	6	4	4	1	-	-	-
Male	6	2	-	3	1	5	2	3	-	-	-	-
Female	3	2	3	1	2	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	2	2	1	-	-	-	-	-
Male	1	-	1	1	1	2	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	86	113	131	158	191	259	377	433	312	121	34	-
Male	73	92	96	111	115	159	209	204	117	27	4	-
Female	13	21	35	47	76	100	168	229	195	94	30	-
White, both sexes	80	100	118	147	175	237	363	417	308	117	33	-
Male	67	85	88	106	105	149	200	194	116	25	4	-
Female	13	15	30	41	70	88	163	223	192	92	29	-
Black, both sexes	5	12	10	6	13	17	11	14	4	4	1	-
Male	5	6	7	2	8	7	6	9	1	2	-	-
Female	-	6	3	4	5	10	5	5	3	2	1	-
American Indian 1/, both sexes	1	1	1	4	2	3	3	1	-	-	-	-
Male	1	1	-	2	2	3	3	1	-	-	-	-
Female	-	-	1	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	1	2	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Kentucky												
All races, both sexes	126	126	176	213	318	401	526	518	364	136	32	-
Male	94	84	113	138	185	239	251	203	105	27	3	-
Female	32	42	63	75	133	162	275	315	259	109	29	-
White, both sexes	112	114	167	196	293	381	505	496	353	125	32	-
Male	82	78	108	131	170	229	239	198	102	26	3	-
Female	30	36	59	65	123	152	266	298	251	99	29	-
Black, both sexes	13	12	8	16	23	19	21	22	10	11	-	-
Male	12	6	5	7	14	9	12	5	2	1	-	-
Female	1	6	3	9	9	10	9	17	8	10	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	135	182	213	248	321	380	526	481	300	131	32	-
Male	100	130	145	163	203	222	280	198	92	28	5	-
Female	35	52	68	85	118	158	246	283	208	103	27	-
White, both sexes	81	117	148	161	237	291	415	396	244	106	20	-
Male	65	88	107	115	155	174	226	165	75	26	3	-
Female	16	29	41	46	82	117	189	231	169	80	17	-
Black, both sexes	53	64	64	86	80	89	108	83	55	25	12	-
Male	34	41	37	48	46	48	52	32	17	2	2	-
Female	19	23	27	38	34	41	56	51	38	23	10	-
American Indian 1/, both sexes	-	-	-	1	2	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	2	-	1	2	1	-	-	-
Male	1	1	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Maine												
All races, both sexes	24	48	61	80	86	134	219	210	163	50	19	-
Male	19	37	46	55	58	80	132	87	62	13	1	-
Female	5	11	15	25	28	54	87	123	101	37	18	-
White, both sexes	21	47	61	80	86	134	219	210	163	49	19	-
Male	16	36	46	55	58	80	132	87	62	12	1	-
Female	5	11	15	25	28	54	87	123	101	37	18	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	224	268	321	346	465	635	861	915	669	280	63	1
Male	157	185	221	211	284	358	405	374	199	76	8	1
Female	67	83	100	135	181	277	456	541	470	204	55	-
White, both sexes	131	163	202	208	298	468	660	755	555	234	47	1
Male	102	115	141	133	185	262	335	328	173	66	6	1
Female	29	48	61	75	113	206	325	427	382	168	41	-
Black, both sexes	90	102	112	128	162	159	189	151	107	42	15	-
Male	53	67	75	68	95	91	65	42	23	10	2	-
Female	37	35	37	60	67	68	124	109	84	32	13	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	6	10	5	8	11	9	6	4	1	-
Male	2	3	4	10	4	5	5	4	3	-	-	-
Female	1	-	2	-	1	3	6	5	3	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Massachusetts												
All races, both sexes	147	193	257	253	436	620	951	1,094	820	394	107	-
Male	121	156	179	185	271	361	504	446	269	78	13	-
Female	26	37	78	68	165	259	447	648	551	316	94	-
White, both sexes	135	182	238	237	407	600	919	1,067	809	387	105	-
Male	111	145	164	174	261	351	488	434	266	75	13	-
Female	24	37	74	63	146	249	431	633	543	312	92	-
Black, both sexes	10	8	15	14	23	12	26	19	8	5	-	-
Male	8	8	12	10	8	5	15	9	3	2	-	-
Female	2	-	3	4	15	7	11	10	5	3	-	-
American Indian 1/, both sexes	2	-	1	-	-	-	1	-	-	1	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	3	3	2	6	8	5	8	3	1	2	-
Male	-	3	2	1	2	5	-	3	-	1	-	-
Female	-	-	1	1	4	3	5	5	3	-	2	-
Michigan												
All races, both sexes	482	687	674	758	1,008	1,499	2,041	2,061	1,488	596	132	-
Male	352	500	458	484	615	847	1,022	895	510	136	25	-
Female	130	187	216	274	393	652	1,019	1,166	978	460	107	-
White, both sexes	333	474	489	568	790	1,260	1,734	1,833	1,349	536	112	-
Male	257	353	340	372	494	731	883	790	462	122	20	-
Female	76	121	149	196	296	529	851	1,043	887	414	92	-
Black, both sexes	144	206	180	179	200	227	291	216	129	60	20	-
Male	92	141	116	108	110	111	135	99	45	14	5	-
Female	52	65	64	71	90	116	156	117	84	46	15	-
American Indian 1/, both sexes	2	4	3	9	7	9	8	7	5	-	-	-
Male	-	3	1	3	4	3	2	3	2	-	-	-
Female	2	1	2	6	3	6	6	4	3	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	2	2	11	3	8	5	5	-	-	-
Male	3	3	1	1	7	2	2	3	1	-	-	-
Female	-	-	1	1	4	1	6	2	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Minnesota												
All races, both sexes	123	141	147	153	236	330	536	562	467	192	48	-
Male	106	109	103	111	165	219	335	291	186	44	5	-
Female	17	32	44	42	71	111	201	271	281	148	43	-
White, both sexes	109	122	130	144	225	321	527	558	466	192	48	-
Male	96	95	92	105	157	216	329	289	186	44	5	-
Female	13	27	38	39	68	105	198	269	280	148	43	-
Black, both sexes	8	12	9	3	6	5	4	3	-	-	-	-
Male	6	8	5	-	4	2	3	1	-	-	-	-
Female	2	4	4	3	2	3	1	2	-	-	-	-
American Indian 1/, both sexes	3	3	5	5	2	3	3	1	-	-	-	-
Male	1	2	3	5	1	1	3	1	-	-	-	-
Female	2	1	2	-	1	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	3	1	3	1	2	-	1	-	-	-
Male	3	4	3	1	3	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
Mississippi												
All races, both sexes	119	124	147	182	242	312	332	354	292	114	26	-
Male	94	79	87	123	148	167	147	124	69	22	2	-
Female	25	45	60	59	94	145	185	230	223	92	24	-
White, both sexes	67	81	102	138	165	243	268	286	230	94	16	-
Male	51	54	60	96	101	136	124	99	54	16	1	-
Female	16	27	42	42	64	107	144	187	176	78	15	-
Black, both sexes	52	40	45	43	76	68	64	68	62	20	10	-
Male	43	23	27	26	47	31	23	25	15	6	1	-
Female	9	17	18	17	29	37	41	43	47	14	9	-
American Indian 1/, both sexes	-	1	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Missouri												
All races, both sexes	172	253	274	347	453	619	929	1,090	814	346	81	-
Male	127	186	180	230	281	341	471	443	254	68	12	-
Female	45	67	94	117	172	278	458	647	560	278	69	-
White, both sexes	139	210	231	295	394	563	860	1,030	768	323	76	-
Male	104	157	151	199	245	313	448	425	244	65	11	-
Female	35	53	80	96	149	250	412	605	524	258	65	-
Black, both sexes	30	42	42	51	54	49	66	56	44	22	5	-
Male	20	28	29	31	33	23	22	17	10	3	1	-
Female	10	14	13	20	21	26	44	39	34	19	4	-
American Indian 1/, both sexes	2	-	1	-	2	2	3	2	-	-	-	-
Male	2	-	-	-	2	2	1	-	-	-	-	-
Female	-	-	1	-	-	-	2	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	3	5	-	2	2	1	-	-
Male	1	1	-	-	1	3	-	1	-	-	-	-
Female	-	-	-	1	2	2	-	1	2	1	-	-
Montana												
All races, both sexes	22	43	45	59	66	91	102	131	85	38	3	-
Male	18	34	32	40	42	61	55	69	42	9	1	-
Female	4	9	13	19	24	30	47	62	43	29	2	-
White, both sexes	20	41	44	54	62	88	102	128	82	38	3	-
Male	16	32	31	36	39	60	55	69	42	9	1	-
Female	4	9	13	18	23	28	47	59	40	29	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	1	5	4	3	-	3	2	-	-	-
Male	2	1	1	4	3	1	-	-	-	-	-	-
Female	-	-	-	1	1	2	-	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Nebraska												
All races, both sexes	31	41	55	59	83	132	255	247	223	117	25	-
Male	23	32	39	48	52	78	154	105	75	32	5	-
Female	8	9	16	11	31	54	101	142	148	85	20	-
White, both sexes	28	37	48	58	81	128	251	241	219	113	25	-
Male	20	29	34	48	51	77	152	102	74	32	5	-
Female	8	8	14	10	30	51	99	139	145	81	20	-
Black, both sexes	3	4	2	-	2	4	2	3	4	3	-	-
Male	3	3	2	-	1	1	1	1	1	-	-	-
Female	-	1	-	-	1	3	1	2	3	3	-	-
American Indian 1/, both sexes	-	-	4	1	-	-	1	-	-	1	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	3	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	110	166	176	205	249	282	298	250	156	48	11	-
Male	92	127	131	150	172	167	167	120	59	12	1	-
Female	18	39	45	55	77	115	131	130	97	36	10	-
White, both sexes	94	142	155	174	216	252	276	239	147	45	7	-
Male	80	112	116	130	150	151	153	113	58	10	1	-
Female	14	30	39	44	66	101	123	126	89	35	6	-
Black, both sexes	12	14	12	19	20	19	10	6	6	1	3	-
Male	8	9	8	13	15	9	7	3	-	1	-	-
Female	4	5	4	6	5	10	3	3	6	-	3	-
American Indian 1/, both sexes	1	3	3	2	2	2	2	1	-	1	1	-
Male	1	-	2	1	1	1	1	1	-	-	-	-
Female	-	3	1	1	1	1	1	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	3	7	6	10	11	9	10	4	3	1	-	-
Male	3	6	5	6	6	6	6	3	1	1	-	-
Female	-	1	1	4	5	3	4	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
New Hampshire												
All races, both sexes	38	58	52	64	101	147	200	210	173	72	14	-
Male	31	49	42	54	61	82	95	88	56	21	4	-
Female	7	9	10	10	40	65	105	122	117	51	10	-
White, both sexes	38	57	51	64	101	145	199	208	173	72	14	-
Male	31	48	41	54	61	81	95	87	56	21	4	-
Female	7	9	10	10	40	64	104	121	117	51	10	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
New Jersey												
All races, both sexes	237	313	402	479	701	1,161	1,705	1,923	1,452	635	151	2
Male	187	234	274	323	435	646	846	790	456	141	15	2
Female	50	79	128	156	266	515	859	1,133	996	494	136	-
White, both sexes	192	242	323	391	576	1,043	1,564	1,812	1,367	602	140	2
Male	155	182	233	264	361	580	784	754	432	135	14	2
Female	37	60	90	127	215	463	780	1,058	935	467	126	-
Black, both sexes	38	62	64	71	111	100	118	93	74	29	8	-
Male	26	45	29	47	68	56	55	30	21	5	-	-
Female	12	17	35	24	43	44	63	63	53	24	8	-
American Indian 1/, both sexes	-	1	-	2	-	1	3	-	2	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	2	-	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	8	15	15	14	17	20	18	9	4	3	-
Male	6	6	12	12	6	9	6	6	2	1	1	-
Female	1	2	3	3	8	8	14	12	7	3	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
New Mexico												
All races, both sexes	69	96	91	112	178	220	269	304	236	97	21	-
Male	57	72	66	80	115	123	147	141	89	20	2	-
Female	12	24	25	32	63	97	122	163	147	77	19	-
White, both sexes	62	87	84	103	160	212	256	297	229	92	20	-
Male	51	67	60	72	106	119	139	139	85	19	2	-
Female	11	20	24	31	54	93	117	158	144	73	18	-
Black, both sexes	2	3	-	6	5	3	4	3	2	4	-	-
Male	2	1	-	5	1	1	1	1	-	-	-	-
Female	-	2	-	1	4	2	3	2	2	4	-	-
American Indian 1/, both sexes	5	5	7	2	12	4	6	4	5	1	1	-
Male	4	3	6	2	8	3	5	1	4	1	-	-
Female	1	2	1	-	4	1	1	3	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	1	3	-	-	-	-	-
Male	-	1	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
New York												
All races, both sexes	632	977	1,252	1,626	2,290	3,550	5,183	5,917	4,847	2,220	586	-
Male	448	726	858	1,011	1,326	1,942	2,411	2,357	1,502	470	97	-
Female	184	251	394	615	964	1,608	2,772	3,560	3,345	1,750	489	-
White, both sexes	461	734	916	1,179	1,781	2,930	4,416	5,253	4,405	1,986	520	-
Male	354	570	640	761	1,066	1,646	2,125	2,121	1,378	425	89	-
Female	107	164	276	418	715	1,284	2,291	3,132	3,027	1,561	431	-
Black, both sexes	146	207	292	397	436	532	633	538	352	199	61	-
Male	75	126	182	220	217	246	230	184	82	36	7	-
Female	71	81	110	177	219	286	403	354	270	163	54	-
American Indian 1/, both sexes	-	-	3	1	5	6	2	5	-	2	-	-
Male	-	-	2	-	1	2	-	2	-	-	-	-
Female	-	-	1	1	4	4	2	3	-	2	-	-
Asian or Pacific Islander 2/, both sexes	25	36	41	49	68	82	132	121	90	33	5	-
Male	19	30	34	30	42	48	56	50	42	9	1	-
Female	6	6	7	19	26	34	76	71	48	24	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
North Carolina												
All races, both sexes	324	444	555	569	722	921	1,106	1,136	781	281	66	-
Male	258	352	415	390	500	540	584	471	226	57	8	-
Female	66	92	140	179	222	381	522	665	555	224	58	-
White, both sexes	225	316	416	435	564	740	910	964	663	245	52	-
Male	183	255	314	295	405	449	506	406	201	51	5	-
Female	42	61	102	140	159	291	404	558	462	194	47	-
Black, both sexes	92	114	129	124	148	171	178	163	111	33	14	-
Male	68	84	94	87	89	87	72	63	21	6	3	-
Female	24	30	35	37	59	84	106	100	90	27	11	-
American Indian 1/, both sexes	6	12	6	8	8	7	15	5	6	2	-	-
Male	6	11	5	7	6	3	5	-	3	-	-	-
Female	-	1	1	1	2	4	10	5	3	2	-	-
Asian or Pacific Islander 2/, both sexes	1	2	4	2	2	3	3	4	1	1	-	-
Male	1	2	2	1	-	1	1	2	1	-	-	-
Female	-	-	2	1	2	2	2	2	-	1	-	-
North Dakota												
All races, both sexes	23	21	19	27	50	59	110	131	127	47	11	-
Male	19	17	15	16	29	39	61	55	49	18	3	-
Female	4	4	4	11	21	20	49	76	78	29	8	-
White, both sexes	21	18	17	25	47	55	109	129	126	47	11	-
Male	18	14	14	14	28	37	60	54	49	18	3	-
Female	3	4	3	11	19	18	49	75	77	29	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	2	2	3	4	1	2	1	-	-	-
Male	1	3	1	2	1	2	1	1	-	-	-	-
Female	1	-	1	-	2	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Ohio												
All races, both sexes	469	628	663	811	1,117	1,694	2,458	2,438	1,778	687	135	-
Male	357	465	456	529	660	960	1,264	1,034	582	160	19	-
Female	112	163	207	282	457	734	1,194	1,404	1,196	527	116	-
White, both sexes	363	513	553	687	958	1,510	2,262	2,293	1,677	641	119	-
Male	285	384	389	452	566	862	1,167	980	554	146	17	-
Female	78	129	164	235	392	648	1,095	1,313	1,123	495	102	-
Black, both sexes	102	113	107	118	155	176	191	142	96	43	16	-
Male	70	79	65	72	92	94	96	54	26	12	2	-
Female	32	34	42	46	63	82	95	88	70	31	14	-
American Indian 1/, both sexes	2	1	2	1	-	2	-	-	-	-	-	-
Male	1	1	1	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	5	4	6	5	3	5	3	-	-
Male	1	1	1	4	2	3	1	-	2	2	-	-
Female	1	-	-	1	2	3	4	3	3	1	-	-
Oklahoma												
All races, both sexes	158	209	273	310	405	520	663	710	487	218	45	-
Male	114	168	189	222	257	324	359	291	166	39	8	-
Female	44	41	84	88	148	196	304	419	321	179	37	-
White, both sexes	124	173	224	256	338	460	611	655	450	203	42	-
Male	91	139	155	181	212	293	334	265	156	36	7	-
Female	33	34	69	75	126	167	277	390	294	167	35	-
Black, both sexes	21	19	23	26	22	28	23	23	18	11	2	-
Male	15	12	15	19	13	14	9	9	6	3	1	-
Female	6	7	8	7	9	14	14	14	12	8	1	-
American Indian 1/, both sexes	13	16	25	24	43	30	27	31	18	4	1	-
Male	8	16	18	18	31	16	15	17	4	-	-	-
Female	5	-	7	6	12	14	12	14	14	4	1	-
Asian or Pacific Islander 2/, both sexes	-	1	1	4	2	2	2	1	1	-	-	-
Male	-	1	1	4	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Oregon												
All races, both sexes	109	138	151	199	221	321	456	513	366	139	17	-
Male	91	111	111	148	146	214	269	284	153	41	4	-
Female	18	27	40	51	75	107	187	229	213	98	13	-
White, both sexes	105	135	139	191	215	317	448	501	362	136	17	-
Male	87	109	104	141	143	211	267	279	153	40	4	-
Female	18	26	35	50	72	106	181	222	209	96	13	-
Black, both sexes	3	3	5	-	2	1	2	3	1	2	-	-
Male	3	2	2	-	1	1	1	-	-	1	-	-
Female	-	1	3	-	1	-	1	3	1	1	-	-
American Indian 1/, both sexes	1	-	4	3	1	-	-	5	1	-	-	-
Male	1	-	2	2	1	-	-	3	-	-	-	-
Female	-	-	2	1	-	-	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	5	3	3	6	4	2	1	-	-
Male	-	-	3	5	1	2	1	2	-	-	-	-
Female	-	-	-	-	2	1	5	2	2	1	-	-
Pennsylvania												
All races, both sexes	326	539	639	737	1,100	1,789	2,566	2,878	2,144	869	170	2
Male	245	392	455	491	690	1,028	1,312	1,213	659	209	20	1
Female	81	147	184	246	410	761	1,254	1,665	1,485	660	150	1
White, both sexes	243	419	528	623	956	1,617	2,395	2,744	2,053	829	156	2
Male	189	308	380	420	608	929	1,229	1,168	637	201	18	1
Female	54	111	148	203	348	688	1,166	1,576	1,416	628	138	1
Black, both sexes	81	116	105	108	139	164	166	126	87	40	13	-
Male	54	81	71	66	80	93	82	41	22	8	2	-
Female	27	35	34	42	59	71	84	85	65	32	11	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	-	-	1	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	3	6	6	4	7	5	8	4	-	-	-
Male	2	2	4	5	1	5	1	4	-	-	-	-
Female	-	1	2	1	3	2	4	4	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Rhode Island												
All races, both sexes	33	39	49	60	77	158	231	328	261	125	32	-
Male	24	30	30	39	45	90	121	132	72	20	7	-
Female	9	9	19	21	32	68	110	196	189	105	25	-
White, both sexes	32	37	47	57	74	152	226	325	259	123	32	-
Male	24	29	29	38	43	88	118	131	72	19	7	-
Female	8	8	18	19	31	64	108	194	187	104	25	-
Black, both sexes	1	2	1	3	2	5	4	3	2	1	-	-
Male	-	1	-	1	2	2	2	1	-	-	-	-
Female	1	1	1	2	-	3	2	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	133	193	196	228	292	413	487	422	249	88	20	-
Male	103	142	129	158	202	277	259	192	98	19	3	-
Female	30	51	67	70	90	136	228	230	151	69	17	-
White, both sexes	90	137	150	169	216	332	391	345	211	76	10	-
Male	71	99	102	123	154	227	217	164	87	16	3	-
Female	19	38	48	46	62	105	174	181	124	60	7	-
Black, both sexes	42	53	46	59	76	80	96	77	38	12	10	-
Male	32	40	27	35	48	49	42	28	11	3	-	-
Female	10	13	19	24	28	31	54	49	27	9	10	-
American Indian 1/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
South Dakota												
All races, both sexes	20	30	31	29	49	85	123	149	132	67	17	-
Male	16	23	20	20	30	57	67	67	45	12	4	-
Female	4	7	11	9	19	28	56	82	87	55	13	-
White, both sexes	18	26	28	25	45	82	118	147	132	66	17	-
Male	14	21	19	18	28	54	66	66	45	12	4	-
Female	4	5	9	7	17	28	52	81	87	54	13	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	4	3	4	4	1	5	2	-	1	-	-
Male	2	2	1	2	2	1	1	1	-	-	-	-
Female	-	2	2	2	2	-	4	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	259	372	439	420	611	772	988	967	680	252	60	-
Male	190	284	303	273	391	463	479	393	209	51	12	-
Female	69	88	136	147	220	309	509	574	471	201	48	-
White, both sexes	192	300	364	349	518	673	893	902	609	231	47	-
Male	143	233	255	229	337	405	447	375	189	46	11	-
Female	49	67	109	120	181	268	446	527	420	185	36	-
Black, both sexes	65	70	73	68	90	98	93	63	69	21	13	-
Male	45	50	46	42	53	57	31	18	20	5	1	-
Female	20	20	27	26	37	41	62	45	49	16	12	-
American Indian 1/, both sexes	-	1	1	1	-	1	1	-	-	-	-	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	2	3	-	1	2	2	-	-	-
Male	2	-	1	1	1	-	-	-	-	-	-	-
Female	-	1	-	1	2	-	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Texas												
All races, both sexes	622	892	1,010	1,143	1,397	2,006	2,463	2,321	1,658	655	152	1
Male	478	662	712	768	866	1,181	1,268	1,013	616	175	27	-
Female	144	230	298	375	531	825	1,195	1,308	1,042	480	125	1
White, both sexes	494	715	821	952	1,189	1,788	2,238	2,164	1,549	586	133	1
Male	383	536	591	654	746	1,062	1,153	954	575	164	23	-
Female	111	179	230	298	443	726	1,085	1,210	974	422	110	1
Black, both sexes	121	158	177	173	193	203	204	142	98	68	19	-
Male	89	110	113	104	114	111	106	54	36	11	4	-
Female	32	48	64	69	79	92	98	88	62	57	15	-
American Indian 1/, both sexes	-	2	2	3	3	2	5	3	2	-	-	-
Male	-	2	1	2	2	2	2	-	-	-	-	-
Female	-	-	1	1	1	-	3	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	7	17	10	15	12	13	16	12	9	1	-	-
Male	6	14	7	8	4	6	7	5	5	-	-	-
Female	1	3	3	7	8	7	9	7	4	1	-	-
Utah												
All races, both sexes	34	42	55	70	70	109	152	175	115	46	7	-
Male	30	35	42	60	46	77	95	100	54	19	1	-
Female	4	7	13	10	24	32	57	75	61	27	6	-
White, both sexes	31	42	55	66	69	105	151	175	114	46	7	-
Male	27	35	42	58	45	74	95	100	53	19	1	-
Female	4	7	13	8	24	31	56	75	61	27	6	-
Black, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	2	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	3	1	1	-	-	1	-	-	-
Male	1	-	-	1	1	1	-	-	1	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Vermont												
All races, both sexes	19	36	28	38	47	86	92	117	84	35	12	-
Male	14	30	23	26	29	48	51	49	31	6	3	-
Female	5	6	5	12	18	38	41	68	53	29	9	-
White, both sexes	19	36	28	38	46	86	92	117	84	35	12	-
Male	14	30	23	26	28	48	51	49	31	6	3	-
Female	5	6	5	12	18	38	41	68	53	29	9	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	156	270	311	345	454	644	823	842	532	230	39	-
Male	113	207	239	244	288	412	424	351	177	46	5	-
Female	43	63	72	101	166	232	399	491	355	184	34	-
White, both sexes	117	189	237	256	366	525	692	717	465	191	33	-
Male	88	149	183	183	238	344	367	299	157	41	4	-
Female	29	40	54	73	128	181	325	418	308	150	29	-
Black, both sexes	37	73	70	83	82	108	124	117	65	39	6	-
Male	23	51	53	57	46	61	54	51	19	5	1	-
Female	14	22	17	26	36	47	70	66	46	34	5	-
American Indian 1/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	8	4	6	6	10	5	8	2	-	-	-
Male	2	7	3	4	4	6	3	1	1	-	-	-
Female	-	1	1	2	2	4	2	7	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Washington												
All races, both sexes	149	194	295	317	420	619	845	943	782	310	64	-
Male	124	147	210	225	282	366	463	439	294	97	17	-
Female	25	47	85	92	138	253	382	504	488	213	47	-
White, both sexes	134	173	261	286	394	575	814	912	759	300	62	-
Male	112	131	187	202	266	344	449	424	283	93	16	-
Female	22	42	74	84	128	231	365	488	476	207	46	-
Black, both sexes	8	13	14	14	5	15	8	8	7	3	-	-
Male	6	10	8	11	4	6	3	2	2	-	-	-
Female	2	3	6	3	1	9	5	6	5	3	-	-
American Indian 1/, both sexes	2	3	7	3	8	8	7	4	3	-	1	-
Male	1	2	3	2	4	4	3	1	2	-	1	-
Female	1	1	4	1	4	4	4	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	5	13	14	13	21	16	19	13	7	1	-
Male	5	4	12	10	8	12	8	12	7	4	-	-
Female	-	1	1	4	5	9	8	7	6	3	1	-
West Virginia												
All races, both sexes	107	109	127	180	226	318	404	368	275	107	18	-
Male	83	82	93	128	138	187	205	165	88	20	1	-
Female	24	27	34	52	88	131	199	203	187	87	17	-
White, both sexes	103	107	119	178	220	309	388	362	268	104	16	-
Male	80	80	89	126	134	180	199	164	85	20	1	-
Female	23	27	30	52	86	129	189	198	183	84	15	-
Black, both sexes	4	2	8	2	6	9	15	6	7	3	2	-
Male	3	2	4	2	4	7	6	1	3	-	-	-
Female	1	-	4	-	2	2	9	5	4	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Wisconsin												
All races, both sexes	117	140	184	220	359	531	731	808	620	297	80	-
Male	90	108	134	165	249	319	406	405	228	62	13	-
Female	27	32	50	55	110	212	325	403	392	235	67	-
White, both sexes	98	118	164	202	341	500	709	793	610	292	79	-
Male	79	94	121	155	238	304	396	399	225	60	13	-
Female	19	24	43	47	103	196	313	394	385	232	66	-
Black, both sexes	16	17	17	17	17	27	20	14	10	3	-	-
Male	9	10	11	9	11	13	10	6	3	1	-	-
Female	7	7	6	8	6	14	10	8	7	2	-	-
American Indian 1/, both sexes	3	4	2	1	-	3	2	-	-	2	-	-
Male	2	3	1	1	-	2	-	-	-	1	-	-
Female	1	1	1	-	-	1	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	1	-	1	-	-	1	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	1	-
Wyoming												
All races, both sexes	9	12	15	25	38	50	40	47	36	20	1	-
Male	9	6	12	17	26	37	32	22	14	4	-	-
Female	-	6	3	8	12	13	8	25	22	16	1	-
White, both sexes	9	12	13	25	38	50	40	47	36	20	1	-
Male	9	6	10	17	26	37	32	22	14	4	-	-
Female	-	6	3	8	12	13	8	25	22	16	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
United States												
All races, both sexes	61,030	-	-	-	-	1	19	47	163	408	1,033	2,149
Male	34,157	-	-	-	-	1	16	38	135	330	789	1,678
Female	26,873	-	-	-	-	-	3	9	28	78	244	471
White, both sexes	50,146	-	-	-	-	1	15	28	110	298	774	1,608
Male	27,777	-	-	-	-	1	12	27	94	243	614	1,288
Female	22,369	-	-	-	-	-	3	1	16	55	160	320
Black, both sexes	9,362	-	-	-	-	-	4	17	41	94	223	471
Male	5,410	-	-	-	-	-	4	9	31	72	144	326
Female	3,952	-	-	-	-	-	-	8	10	22	79	145
American Indian 1/, both sexes	317	-	-	-	-	-	-	-	4	3	10	22
Male	210	-	-	-	-	-	-	-	3	3	7	18
Female	107	-	-	-	-	-	-	-	1	-	3	4
Asian or Pacific Islander 2/, both sexes	1,205	-	-	-	-	-	-	2	8	13	26	48
Male	760	-	-	-	-	-	-	2	7	12	24	46
Female	445	-	-	-	-	-	-	-	1	1	2	2
Alabama												
All races, both sexes	744	-	-	-	-	-	2	1	3	10	15	41
Male	378	-	-	-	-	-	2	1	3	9	12	25
Female	366	-	-	-	-	-	-	-	-	1	3	16
White, both sexes	553	-	-	-	-	-	1	1	1	6	13	25
Male	286	-	-	-	-	-	1	1	1	5	12	19
Female	267	-	-	-	-	-	-	-	-	1	1	6
Black, both sexes	188	-	-	-	-	-	1	-	2	4	2	15
Male	89	-	-	-	-	-	1	-	2	4	-	5
Female	99	-	-	-	-	-	-	-	-	-	2	10
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Alaska												
All races, both sexes	134	-	-	-	-	-	-	-	-	2	4	8
Male	98	-	-	-	-	-	-	-	-	2	4	7
Female	36	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	111	-	-	-	-	-	-	-	-	2	2	6
Male	83	-	-	-	-	-	-	-	-	2	2	5
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,000	-	-	-	-	-	-	1	6	17	28	61
Male	1,165	-	-	-	-	-	-	-	4	14	25	46
Female	835	-	-	-	-	-	-	1	2	3	3	15
White, both sexes	1,857	-	-	-	-	-	-	-	5	15	24	53
Male	1,068	-	-	-	-	-	-	-	3	12	21	38
Female	789	-	-	-	-	-	-	-	2	3	3	15
Black, both sexes	86	-	-	-	-	-	-	1	1	1	1	4
Male	61	-	-	-	-	-	-	-	1	1	1	4
Female	25	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	37	-	-	-	-	-	-	-	-	1	2	2
Male	26	-	-	-	-	-	-	-	-	1	2	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Arkansas												
All races, both sexes	274	-	-	-	-	1	-	4	3	7	8	25
Male	153	-	-	-	-	1	-	3	2	5	6	22
Female	121	-	-	-	-	-	-	1	1	2	2	3
White, both sexes	207	-	-	-	-	1	-	2	3	5	6	16
Male	118	-	-	-	-	1	-	2	2	3	4	14
Female	89	-	-	-	-	-	-	-	1	2	2	2
Black, both sexes	61	-	-	-	-	-	-	2	-	2	2	8
Male	30	-	-	-	-	-	-	1	-	2	2	7
Female	31	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8,007	-	-	-	-	-	-	5	6	29	87	219
Male	4,614	-	-	-	-	-	-	4	5	26	72	180
Female	3,393	-	-	-	-	-	-	1	1	3	15	39
White, both sexes	6,549	-	-	-	-	-	-	4	2	20	64	175
Male	3,733	-	-	-	-	-	-	4	1	18	55	145
Female	2,816	-	-	-	-	-	-	-	1	2	9	30
Black, both sexes	878	-	-	-	-	-	-	1	3	4	12	27
Male	525	-	-	-	-	-	-	-	3	3	7	20
Female	353	-	-	-	-	-	-	1	-	1	5	7
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	1	-	2
Male	21	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	551	-	-	-	-	-	-	-	1	4	11	15
Male	335	-	-	-	-	-	-	-	1	4	10	14
Female	216	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Colorado												
All races, both sexes	590	-	-	-	-	-	-	-	2	5	8	17
Male	358	-	-	-	-	-	-	-	2	4	5	14
Female	232	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	563	-	-	-	-	-	-	-	2	5	8	15
Male	342	-	-	-	-	-	-	-	2	4	5	13
Female	221	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	730	-	-	-	-	-	-	-	1	3	9	18
Male	388	-	-	-	-	-	-	-	1	3	5	16
Female	342	-	-	-	-	-	-	-	-	-	4	2
White, both sexes	668	-	-	-	-	-	-	-	1	2	9	14
Male	349	-	-	-	-	-	-	-	1	2	5	12
Female	319	-	-	-	-	-	-	-	-	-	4	2
Black, both sexes	54	-	-	-	-	-	-	-	-	1	-	4
Male	36	-	-	-	-	-	-	-	-	1	-	4
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Delaware												
All races, both sexes	292	-	-	-	-	-	-	-	1	1	1	16
Male	159	-	-	-	-	-	-	-	-	1	1	10
Female	133	-	-	-	-	-	-	-	1	-	-	6
White, both sexes	244	-	-	-	-	-	-	-	-	1	1	12
Male	126	-	-	-	-	-	-	-	-	1	1	8
Female	118	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	47	-	-	-	-	-	-	-	1	-	-	4
Male	33	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	613	-	-	-	-	-	1	-	-	2	6	17
Male	326	-	-	-	-	-	1	-	-	1	5	11
Female	287	-	-	-	-	-	-	-	-	1	1	6
White, both sexes	146	-	-	-	-	-	-	-	-	-	2	3
Male	81	-	-	-	-	-	-	-	-	-	2	3
Female	65	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	462	-	-	-	-	-	1	-	-	2	4	14
Male	242	-	-	-	-	-	1	-	-	1	3	8
Female	220	-	-	-	-	-	-	-	-	1	1	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Florida												
All races, both sexes	4,961	-	-	-	-	-	1	1	7	21	49	118
Male	2,720	-	-	-	-	-	1	1	7	18	41	97
Female	2,241	-	-	-	-	-	-	-	-	3	8	21
White, both sexes	4,508	-	-	-	-	-	1	1	5	16	35	97
Male	2,467	-	-	-	-	-	1	1	5	14	29	83
Female	2,041	-	-	-	-	-	-	-	-	2	6	14
Black, both sexes	428	-	-	-	-	-	-	-	2	5	13	20
Male	238	-	-	-	-	-	-	-	2	4	11	13
Female	190	-	-	-	-	-	-	-	-	1	2	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,049	-	-	-	-	-	2	-	2	9	25	48
Male	623	-	-	-	-	-	2	-	1	8	19	41
Female	426	-	-	-	-	-	-	-	1	1	6	7
White, both sexes	716	-	-	-	-	-	2	-	-	4	15	25
Male	423	-	-	-	-	-	2	-	-	3	12	21
Female	293	-	-	-	-	-	-	-	-	1	3	4
Black, both sexes	326	-	-	-	-	-	-	-	2	5	9	23
Male	195	-	-	-	-	-	-	-	1	5	6	20
Female	131	-	-	-	-	-	-	-	1	-	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hawaii												
All races, both sexes	182	-	-	-	-	-	-	-	3	1	2	7
Male	129	-	-	-	-	-	-	-	3	-	1	7
Female	53	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	57	-	-	-	-	-	-	-	-	-	2	3
Male	44	-	-	-	-	-	-	-	-	-	1	3
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	123	-	-	-	-	-	-	-	2	1	-	4
Male	84	-	-	-	-	-	-	-	2	-	-	4
Female	39	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	81	-	-	-	-	-	-	-	-	2	1	1
Male	39	-	-	-	-	-	-	-	-	2	1	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	81	-	-	-	-	-	-	-	-	2	1	1
Male	39	-	-	-	-	-	-	-	-	2	1	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Illinois												
All races, both sexes	1,796	-	-	-	-	-	-	-	4	2	13	71
Male	1,096	-	-	-	-	-	-	-	4	1	10	52
Female	700	-	-	-	-	-	-	-	-	1	3	19
White, both sexes	1,293	-	-	-	-	-	-	-	4	2	10	46
Male	763	-	-	-	-	-	-	-	4	1	9	36
Female	530	-	-	-	-	-	-	-	-	1	1	10
Black, both sexes	476	-	-	-	-	-	-	-	-	-	3	22
Male	318	-	-	-	-	-	-	-	-	-	1	13
Female	158	-	-	-	-	-	-	-	-	-	2	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	-	3
Male	14	-	-	-	-	-	-	-	-	-	-	3
Female	12	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	624	-	-	-	-	-	1	2	2	6	29	37
Male	328	-	-	-	-	-	1	2	2	3	18	33
Female	296	-	-	-	-	-	-	-	-	3	11	4
White, both sexes	560	-	-	-	-	-	1	2	2	6	24	33
Male	285	-	-	-	-	-	1	2	2	3	16	29
Female	275	-	-	-	-	-	-	-	-	3	8	4
Black, both sexes	60	-	-	-	-	-	-	-	-	-	5	3
Male	40	-	-	-	-	-	-	-	-	-	2	3
Female	20	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Iowa												
All races, both sexes	1,012	-	-	-	-	-	-	-	2	4	16	23
Male	542	-	-	-	-	-	-	-	1	4	9	18
Female	470	-	-	-	-	-	-	-	1	-	7	5
White, both sexes	991	-	-	-	-	-	-	-	2	4	14	22
Male	528	-	-	-	-	-	-	-	1	4	7	17
Female	463	-	-	-	-	-	-	-	1	-	7	5
Black, both sexes	19	-	-	-	-	-	-	-	-	-	2	1
Male	13	-	-	-	-	-	-	-	-	-	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	352	-	-	-	-	-	-	-	-	1	2	17
Male	208	-	-	-	-	-	-	-	-	1	2	14
Female	144	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	323	-	-	-	-	-	-	-	-	1	2	16
Male	188	-	-	-	-	-	-	-	-	1	2	13
Female	135	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Kentucky												
All races, both sexes	439	-	-	-	-	-	-	-	3	6	9	16
Male	205	-	-	-	-	-	-	-	3	3	6	13
Female	234	-	-	-	-	-	-	-	-	3	3	3
White, both sexes	413	-	-	-	-	-	-	-	2	6	7	14
Male	192	-	-	-	-	-	-	-	2	3	6	11
Female	221	-	-	-	-	-	-	-	-	3	1	3
Black, both sexes	26	-	-	-	-	-	-	-	1	-	2	2
Male	13	-	-	-	-	-	-	-	1	-	-	2
Female	13	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	926	-	-	-	-	-	-	1	5	9	27	36
Male	487	-	-	-	-	-	-	1	2	7	21	27
Female	439	-	-	-	-	-	-	-	3	2	6	9
White, both sexes	663	-	-	-	-	-	-	-	4	3	19	21
Male	354	-	-	-	-	-	-	-	2	3	14	14
Female	309	-	-	-	-	-	-	-	2	-	5	7
Black, both sexes	257	-	-	-	-	-	-	1	1	6	8	15
Male	129	-	-	-	-	-	-	1	-	4	7	13
Female	128	-	-	-	-	-	-	-	1	2	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Maine												
All races, both sexes	174	-	-	-	-	-	-	-	1	-	4	5
Male	96	-	-	-	-	-	-	-	1	-	3	3
Female	78	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	170	-	-	-	-	-	-	-	-	-	2	5
Male	93	-	-	-	-	-	-	-	-	-	2	3
Female	77	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	2	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,367	-	-	-	-	-	-	2	4	14	42	80
Male	1,206	-	-	-	-	-	-	2	2	13	32	56
Female	1,161	-	-	-	-	-	-	-	2	1	10	24
White, both sexes	1,638	-	-	-	-	-	-	1	3	9	27	51
Male	840	-	-	-	-	-	-	1	2	8	20	38
Female	798	-	-	-	-	-	-	-	1	1	7	13
Black, both sexes	696	-	-	-	-	-	-	-	-	4	13	28
Male	344	-	-	-	-	-	-	-	-	4	10	17
Female	352	-	-	-	-	-	-	-	-	-	3	11
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	1	1	1	2	1
Male	22	-	-	-	-	-	-	1	-	1	2	1
Female	11	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Massachusetts												
All races, both sexes	1,005	-	-	-	-	-	1	1	3	3	22	28
Male	557	-	-	-	-	-	1	1	3	3	16	25
Female	448	-	-	-	-	-	-	-	-	-	6	3
White, both sexes	955	-	-	-	-	-	1	1	3	2	21	27
Male	526	-	-	-	-	-	1	1	3	2	16	24
Female	429	-	-	-	-	-	-	-	-	-	5	3
Black, both sexes	36	-	-	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3,750	-	-	-	-	-	2	3	10	24	89	162
Male	2,187	-	-	-	-	-	1	3	9	20	72	125
Female	1,563	-	-	-	-	-	1	-	1	4	17	37
White, both sexes	2,493	-	-	-	-	-	2	1	8	17	58	112
Male	1,452	-	-	-	-	-	1	1	8	14	51	89
Female	1,041	-	-	-	-	-	1	-	-	3	7	23
Black, both sexes	1,209	-	-	-	-	-	-	2	1	7	30	48
Male	709	-	-	-	-	-	-	2	1	6	20	35
Female	500	-	-	-	-	-	-	-	-	1	10	13
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	1	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Minnesota												
All races, both sexes	318	-	-	-	-	-	-	1	-	2	1	7
Male	178	-	-	-	-	-	-	1	-	2	1	7
Female	140	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	311	-	-	-	-	-	-	1	-	2	1	6
Male	174	-	-	-	-	-	-	1	-	2	1	6
Female	137	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	600	-	-	-	-	-	-	-	2	2	3	13
Male	216	-	-	-	-	-	-	-	2	2	1	10
Female	384	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	449	-	-	-	-	-	-	-	1	1	1	8
Male	152	-	-	-	-	-	-	-	1	1	1	6
Female	297	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	149	-	-	-	-	-	-	-	1	1	2	5
Male	64	-	-	-	-	-	-	-	1	1	-	4
Female	85	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Missouri												
All races, both sexes	583	-	-	-	-	-	-	1	1	2	8	11
Male	285	-	-	-	-	-	-	1	1	2	4	8
Female	298	-	-	-	-	-	-	-	-	-	4	3
White, both sexes	521	-	-	-	-	-	-	1	-	2	8	8
Male	255	-	-	-	-	-	-	1	-	2	4	7
Female	266	-	-	-	-	-	-	-	-	-	4	1
Black, both sexes	58	-	-	-	-	-	-	-	1	-	-	3
Male	28	-	-	-	-	-	-	-	1	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	167	-	-	-	-	-	-	-	-	2	5	4
Male	104	-	-	-	-	-	-	-	-	1	5	3
Female	63	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	165	-	-	-	-	-	-	-	-	2	5	4
Male	103	-	-	-	-	-	-	-	-	1	5	3
Female	62	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Nebraska												
All races, both sexes	112	-	-	-	-	-	-	-	1	-	1	2
Male	47	-	-	-	-	-	-	-	1	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	111	-	-	-	-	-	-	-	1	-	1	2
Male	46	-	-	-	-	-	-	-	1	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,091	-	-	-	-	-	-	-	3	4	14	31
Male	707	-	-	-	-	-	-	-	2	3	11	25
Female	384	-	-	-	-	-	-	-	1	1	3	6
White, both sexes	966	-	-	-	-	-	-	-	3	3	12	26
Male	622	-	-	-	-	-	-	-	2	2	10	21
Female	344	-	-	-	-	-	-	-	1	1	2	5
Black, both sexes	82	-	-	-	-	-	-	-	-	-	2	3
Male	55	-	-	-	-	-	-	-	-	-	1	2
Female	27	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	-	-	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Hampshire												
All races, both sexes	279	-	-	-	-	-	-	-	1	1	3	11
Male	169	-	-	-	-	-	-	-	1	1	2	8
Female	110	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	277	-	-	-	-	-	-	-	1	1	3	10
Male	168	-	-	-	-	-	-	-	1	1	2	7
Female	109	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,952	-	-	-	-	-	-	3	4	11	21	47
Male	1,039	-	-	-	-	-	-	2	3	11	18	38
Female	913	-	-	-	-	-	-	1	1	-	3	9
White, both sexes	1,661	-	-	-	-	-	-	2	4	7	17	33
Male	861	-	-	-	-	-	-	2	3	7	16	26
Female	800	-	-	-	-	-	-	-	1	-	1	7
Black, both sexes	245	-	-	-	-	-	-	1	-	4	4	11
Male	149	-	-	-	-	-	-	-	-	4	2	9
Female	96	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	44	-	-	-	-	-	-	-	-	-	-	3
Male	29	-	-	-	-	-	-	-	-	-	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Mexico												
All races, both sexes	687	-	-	-	-	-	-	-	3	6	10	16
Male	402	-	-	-	-	-	-	-	2	5	8	10
Female	285	-	-	-	-	-	-	-	1	1	2	6
White, both sexes	640	-	-	-	-	-	-	-	3	6	8	15
Male	375	-	-	-	-	-	-	-	2	5	8	9
Female	265	-	-	-	-	-	-	-	1	1	-	6
Black, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3,961	-	-	-	-	-	1	3	11	34	76	156
Male	2,289	-	-	-	-	-	1	2	11	24	58	128
Female	1,672	-	-	-	-	-	-	1	-	10	18	28
White, both sexes	2,925	-	-	-	-	-	1	1	4	18	56	106
Male	1,698	-	-	-	-	-	1	1	4	15	45	91
Female	1,227	-	-	-	-	-	-	-	-	3	11	15
Black, both sexes	922	-	-	-	-	-	-	2	5	14	15	45
Male	508	-	-	-	-	-	-	1	5	7	8	32
Female	414	-	-	-	-	-	-	1	-	7	7	13
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	111	-	-	-	-	-	-	-	2	2	5	5
Male	80	-	-	-	-	-	-	-	2	2	5	5
Female	31	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
North Carolina												
All races, both sexes	1,151	-	-	-	-	-	1	1	4	14	32	36
Male	593	-	-	-	-	-	1	1	2	11	26	30
Female	558	-	-	-	-	-	-	-	2	3	6	6
White, both sexes	929	-	-	-	-	-	-	-	1	11	23	22
Male	459	-	-	-	-	-	-	-	1	8	18	19
Female	470	-	-	-	-	-	-	-	-	3	5	3
Black, both sexes	210	-	-	-	-	-	1	1	3	3	9	13
Male	126	-	-	-	-	-	1	1	1	3	8	10
Female	84	-	-	-	-	-	-	-	2	-	1	3
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	132	-	-	-	-	-	-	-	-	2	4	6
Male	61	-	-	-	-	-	-	-	-	1	4	4
Female	71	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	126	-	-	-	-	-	-	-	-	2	4	6
Male	57	-	-	-	-	-	-	-	-	1	4	4
Female	69	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Ohio												
All races, both sexes	2,022	-	-	-	-	-	2	3	5	20	34	74
Male	1,081	-	-	-	-	-	1	2	4	13	21	54
Female	941	-	-	-	-	-	1	1	1	7	13	20
White, both sexes	1,743	-	-	-	-	-	2	2	5	15	26	62
Male	929	-	-	-	-	-	1	1	4	9	18	50
Female	814	-	-	-	-	-	1	1	1	6	8	12
Black, both sexes	270	-	-	-	-	-	-	1	-	4	8	12
Male	148	-	-	-	-	-	-	1	-	3	3	4
Female	122	-	-	-	-	-	-	-	-	1	5	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,052	-	-	-	-	-	-	2	9	7	17	40
Male	589	-	-	-	-	-	-	1	8	6	12	27
Female	463	-	-	-	-	-	-	1	1	1	5	13
White, both sexes	892	-	-	-	-	-	-	1	6	5	14	32
Male	488	-	-	-	-	-	-	1	5	5	9	21
Female	404	-	-	-	-	-	-	-	1	-	5	11
Black, both sexes	78	-	-	-	-	-	-	1	1	2	2	1
Male	44	-	-	-	-	-	-	-	1	1	2	1
Female	34	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	78	-	-	-	-	-	-	-	2	-	1	7
Male	55	-	-	-	-	-	-	-	2	-	1	5
Female	23	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Oregon												
All races, both sexes	280	-	-	-	-	-	-	-	-	5	4	11
Male	159	-	-	-	-	-	-	-	-	4	2	10
Female	121	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	276	-	-	-	-	-	-	-	-	4	3	11
Male	157	-	-	-	-	-	-	-	-	3	2	10
Female	119	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Pennsylvania												
All races, both sexes	3,540	-	-	-	-	-	1	3	6	25	55	107
Male	1,815	-	-	-	-	-	1	2	5	22	42	81
Female	1,725	-	-	-	-	-	-	1	1	3	13	26
White, both sexes	2,953	-	-	-	-	-	-	1	2	23	36	84
Male	1,445	-	-	-	-	-	-	1	2	21	28	64
Female	1,508	-	-	-	-	-	-	-	-	2	8	20
Black, both sexes	563	-	-	-	-	-	1	2	3	2	17	23
Male	355	-	-	-	-	-	1	1	2	1	12	17
Female	208	-	-	-	-	-	-	1	1	1	5	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	1	-	2	-
Male	14	-	-	-	-	-	-	-	1	-	2	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Rhode Island												
All races, both sexes	490	-	-	-	-	-	-	-	-	-	4	10
Male	225	-	-	-	-	-	-	-	-	-	4	9
Female	265	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	477	-	-	-	-	-	-	-	-	-	4	9
Male	219	-	-	-	-	-	-	-	-	-	4	8
Female	258	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	402	-	-	-	-	-	-	-	4	3	8	19
Male	228	-	-	-	-	-	-	-	2	2	5	14
Female	174	-	-	-	-	-	-	-	2	1	3	5
White, both sexes	293	-	-	-	-	-	-	-	2	1	4	14
Male	168	-	-	-	-	-	-	-	1	-	3	10
Female	125	-	-	-	-	-	-	-	1	1	1	4
Black, both sexes	108	-	-	-	-	-	-	-	2	2	4	5
Male	60	-	-	-	-	-	-	-	1	2	2	4
Female	48	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
South Dakota												
All races, both sexes	94	-	-	-	-	-	-	-	-	-	1	-
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	88	-	-	-	-	-	-	-	-	-	1	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,806	-	-	-	-	-	1	2	10	15	50	102
Male	1,044	-	-	-	-	-	1	2	10	12	32	83
Female	762	-	-	-	-	-	-	-	-	3	18	19
White, both sexes	1,513	-	-	-	-	-	1	2	9	11	39	78
Male	870	-	-	-	-	-	1	2	9	9	26	66
Female	643	-	-	-	-	-	-	-	-	2	13	12
Black, both sexes	283	-	-	-	-	-	-	-	1	4	11	22
Male	166	-	-	-	-	-	-	-	1	3	6	15
Female	117	-	-	-	-	-	-	-	-	1	5	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Texas												
All races, both sexes	3,909	-	-	-	-	-	2	4	11	46	110	211
Male	2,458	-	-	-	-	-	1	3	9	40	88	159
Female	1,451	-	-	-	-	-	1	1	2	6	22	52
White, both sexes	3,208	-	-	-	-	-	2	2	5	31	83	147
Male	2,000	-	-	-	-	-	1	2	4	28	68	118
Female	1,208	-	-	-	-	-	1	-	1	3	15	29
Black, both sexes	654	-	-	-	-	-	-	2	6	14	25	59
Male	426	-	-	-	-	-	-	1	5	11	19	36
Female	228	-	-	-	-	-	-	1	1	3	6	23
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	2	2
Male	6	-	-	-	-	-	-	-	-	-	1	2
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	-	1	-	3
Male	26	-	-	-	-	-	-	-	-	1	-	3
Female	14	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	105	-	-	-	-	-	-	-	-	1	1	1
Male	65	-	-	-	-	-	-	-	-	-	1	1
Female	40	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	102	-	-	-	-	-	-	-	-	1	1	1
Male	64	-	-	-	-	-	-	-	-	-	1	1
Female	38	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Vermont												
All races, both sexes	269	-	-	-	-	-	-	-	1	1	6	13
Male	151	-	-	-	-	-	-	-	1	-	6	11
Female	118	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	269	-	-	-	-	-	-	-	1	1	6	13
Male	151	-	-	-	-	-	-	-	1	-	6	11
Female	118	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,118	-	-	-	-	-	-	2	4	9	28	66
Male	594	-	-	-	-	-	-	2	4	5	20	51
Female	524	-	-	-	-	-	-	-	-	4	8	15
White, both sexes	869	-	-	-	-	-	-	1	4	6	15	48
Male	451	-	-	-	-	-	-	1	4	4	12	36
Female	418	-	-	-	-	-	-	-	-	2	3	12
Black, both sexes	233	-	-	-	-	-	-	-	-	3	13	16
Male	131	-	-	-	-	-	-	-	-	1	8	14
Female	102	-	-	-	-	-	-	-	-	2	5	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	1	-	-	-	2
Male	11	-	-	-	-	-	-	1	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Washington												
All races, both sexes	1,473	-	-	-	-	-	1	1	9	13	24	43
Male	792	-	-	-	-	-	1	1	8	11	19	32
Female	681	-	-	-	-	-	-	-	1	2	5	11
White, both sexes	1,379	-	-	-	-	-	1	1	7	12	22	40
Male	728	-	-	-	-	-	1	1	6	10	17	29
Female	651	-	-	-	-	-	-	-	1	2	5	11
Black, both sexes	39	-	-	-	-	-	-	-	1	-	-	1
Male	27	-	-	-	-	-	-	-	1	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	45	-	-	-	-	-	-	-	1	1	2	2
Male	30	-	-	-	-	-	-	-	1	1	2	2
Female	15	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	462	-	-	-	-	-	-	-	1	-	5	8
Male	237	-	-	-	-	-	-	-	1	-	3	4
Female	225	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	449	-	-	-	-	-	-	-	1	-	5	7
Male	232	-	-	-	-	-	-	-	1	-	3	4
Female	217	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Wisconsin												
All races, both sexes	842	-	-	-	-	-	-	-	5	3	12	33
Male	512	-	-	-	-	-	-	-	3	3	9	27
Female	330	-	-	-	-	-	-	-	2	-	3	6
White, both sexes	764	-	-	-	-	-	-	-	3	3	10	24
Male	472	-	-	-	-	-	-	-	3	3	9	21
Female	292	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	72	-	-	-	-	-	-	-	2	-	2	8
Male	37	-	-	-	-	-	-	-	-	-	-	5
Female	35	-	-	-	-	-	-	-	2	-	2	3
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	31	-	-	-	-	-	-	-	-	2	-	-
Male	17	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	2	-	-
Male	17	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
United States												
All races, both sexes	3,671	4,749	5,151	4,919	5,501	6,573	8,332	8,618	6,217	2,760	703	16
Male	2,934	3,673	3,754	3,383	3,471	3,678	4,099	3,509	1,937	608	114	10
Female	737	1,076	1,397	1,536	2,030	2,895	4,233	5,109	4,280	2,152	589	6
White, both sexes	2,744	3,591	3,988	3,790	4,324	5,401	7,128	7,670	5,583	2,482	598	13
Male	2,253	2,835	2,929	2,629	2,748	3,051	3,530	3,127	1,758	537	92	9
Female	491	756	1,059	1,161	1,576	2,350	3,598	4,543	3,825	1,945	506	4
Black, both sexes	825	1,008	1,009	976	1,011	997	1,033	793	524	242	91	3
Male	591	715	713	648	620	534	481	307	140	58	16	1
Female	234	293	296	328	391	463	552	486	384	184	75	2
American Indian 1/, both sexes	25	41	36	30	40	33	36	21	11	3	2	-
Male	21	32	23	22	28	19	16	12	4	1	1	-
Female	4	9	13	8	12	14	20	9	7	2	1	-
Asian or Pacific Islander 2/, both sexes	77	109	118	123	126	142	135	134	99	33	12	-
Male	69	91	89	84	75	74	72	63	35	12	5	-
Female	8	18	29	39	51	68	63	71	64	21	7	-
Alabama												
All races, both sexes	41	43	53	39	67	81	112	123	70	31	12	-
Male	32	34	39	31	43	39	46	38	17	5	2	-
Female	9	9	14	8	24	42	66	85	53	26	10	-
White, both sexes	28	24	37	30	50	65	83	107	53	26	3	-
Male	22	18	29	25	34	32	34	35	14	4	-	-
Female	6	6	8	5	16	33	49	72	39	22	3	-
Black, both sexes	11	19	16	9	17	16	29	16	17	5	9	-
Male	8	16	10	6	9	7	12	3	3	1	2	-
Female	3	3	6	3	8	9	17	13	14	4	7	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Alaska												
All races, both sexes	16	9	14	20	15	19	15	7	3	2	-	-
Male	14	7	11	19	11	14	4	3	2	-	-	-
Female	2	2	3	1	4	5	11	4	1	2	-	-
White, both sexes	15	7	11	18	13	16	12	6	2	1	-	-
Male	13	6	9	17	9	12	4	3	1	-	-	-
Female	2	1	2	1	4	4	8	3	1	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	3	2	1	2	3	-	1	1	-	-
Male	1	1	2	2	1	2	-	-	1	-	-	-
Female	-	1	1	-	-	-	3	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	131	147	190	190	199	233	289	269	144	75	15	5
Male	109	113	149	122	126	128	144	114	46	19	3	3
Female	22	34	41	68	73	105	145	155	98	56	12	2
White, both sexes	123	134	173	167	188	220	269	257	141	71	14	3
Male	101	103	138	107	118	122	133	106	45	16	3	2
Female	22	31	35	60	70	98	136	151	96	55	11	1
Black, both sexes	7	8	11	14	10	7	8	6	2	2	1	2
Male	7	6	7	10	7	4	5	4	1	2	-	1
Female	-	2	4	4	3	3	3	2	1	-	1	1
American Indian 1/, both sexes	1	4	5	6	1	4	7	3	-	1	-	-
Male	1	3	4	4	1	2	3	2	-	1	-	-
Female	-	1	1	2	-	2	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	-	2	5	3	1	1	-	-
Male	-	1	-	1	-	-	3	2	-	-	-	-
Female	-	-	1	2	-	2	2	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Arkansas												
All races, both sexes	13	22	12	13	32	28	32	31	25	14	4	-
Male	12	20	9	7	17	14	15	7	8	5	-	-
Female	1	2	3	6	15	14	17	24	17	9	4	-
White, both sexes	8	18	11	9	20	22	29	25	19	10	3	-
Male	7	16	9	6	13	12	14	6	6	3	-	-
Female	1	2	2	3	7	10	15	19	13	7	3	-
Black, both sexes	4	2	1	4	10	6	3	6	6	4	1	-
Male	4	2	-	1	3	2	1	1	2	2	-	-
Female	-	-	1	3	7	4	2	5	4	2	1	-
American Indian 1/, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	489	663	755	723	720	824	1,075	1,105	825	372	107	3
Male	435	547	564	501	461	485	526	433	277	84	12	2
Female	54	116	191	222	259	339	549	672	548	288	95	1
White, both sexes	383	506	598	566	577	661	915	953	710	322	90	3
Male	341	424	452	392	369	390	445	372	244	70	9	2
Female	42	82	146	174	208	271	470	581	466	252	81	1
Black, both sexes	72	100	114	100	89	91	87	80	61	28	9	-
Male	63	74	80	70	57	56	43	29	15	4	1	-
Female	9	26	34	30	32	35	44	51	46	24	8	-
American Indian 1/, both sexes	1	9	1	3	3	3	3	3	-	-	-	-
Male	1	8	-	3	3	2	1	1	-	-	-	-
Female	-	1	1	-	-	1	2	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	48	42	54	51	69	70	69	54	22	8	-
Male	30	41	32	36	32	37	37	31	18	10	2	-
Female	3	7	10	18	19	32	33	38	36	12	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Colorado												
All races, both sexes	46	47	57	46	62	68	84	61	54	25	8	-
Male	37	42	40	28	45	41	40	33	21	5	1	-
Female	9	5	17	18	17	27	44	28	33	20	7	-
White, both sexes	44	43	53	43	59	65	78	61	54	25	8	-
Male	35	38	39	26	42	39	39	33	21	5	1	-
Female	9	5	14	17	17	26	39	28	33	20	7	-
Black, both sexes	1	3	1	3	3	2	1	-	-	-	-	-
Male	1	3	-	2	3	2	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	-	-	4	-	-	-	-	-
Male	-	1	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	3	-	-	-	-	-
Connecticut												
All races, both sexes	33	63	62	40	55	81	95	103	99	56	12	-
Male	27	48	44	30	35	44	41	47	31	12	4	-
Female	6	15	18	10	20	37	54	56	68	44	8	-
White, both sexes	25	52	54	35	55	72	90	99	93	55	12	-
Male	20	40	39	25	35	40	38	45	31	12	4	-
Female	5	12	15	10	20	32	52	54	62	43	8	-
Black, both sexes	7	11	7	5	-	7	3	4	5	-	-	-
Male	6	8	4	5	-	4	2	2	-	-	-	-
Female	1	3	3	-	-	3	1	2	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	1	-	1	1	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Delaware												
All races, both sexes	23	19	34	23	24	28	32	42	33	10	4	1
Male	21	14	24	12	15	18	15	17	7	3	-	1
Female	2	5	10	11	9	10	17	25	26	7	4	-
White, both sexes	19	17	28	19	15	23	27	38	30	10	3	1
Male	18	13	19	9	7	13	13	15	5	3	-	1
Female	1	4	9	10	8	10	14	23	25	7	3	-
Black, both sexes	3	2	6	4	9	5	5	4	3	-	1	-
Male	3	1	5	3	8	5	2	2	2	-	-	-
Female	-	1	1	1	1	-	3	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	31	37	52	76	57	81	96	81	53	17	6	-
Male	18	25	34	50	36	45	44	34	17	4	1	-
Female	13	12	18	26	21	36	52	47	36	13	5	-
White, both sexes	2	5	13	13	11	18	27	23	23	5	1	-
Male	1	4	9	8	7	12	11	12	10	2	-	-
Female	1	1	4	5	4	6	16	11	13	3	1	-
Black, both sexes	29	32	39	63	46	61	68	57	29	12	5	-
Male	17	21	25	42	29	32	32	21	7	2	1	-
Female	12	11	14	21	17	29	36	36	22	10	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Florida												
All races, both sexes	209	231	323	341	381	540	798	926	657	284	73	1
Male	167	181	245	248	244	315	418	415	244	64	14	-
Female	42	50	78	93	137	225	380	511	413	220	59	1
White, both sexes	164	194	284	304	334	494	742	888	619	262	67	1
Male	136	154	218	222	220	293	391	396	235	57	12	-
Female	28	40	66	82	114	201	351	492	384	205	55	1
Black, both sexes	40	37	38	36	44	40	54	36	37	21	5	-
Male	27	27	26	26	22	19	26	18	9	7	1	-
Female	13	10	12	10	22	21	28	18	28	14	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	1	1	3	6	1	2	1	1	1	-
Male	4	-	1	-	2	3	1	1	-	-	1	-
Female	1	-	-	1	1	3	-	1	1	1	-	-
Georgia												
All races, both sexes	71	105	110	82	90	108	122	130	98	43	4	-
Male	52	83	85	63	64	61	60	51	19	12	2	-
Female	19	22	25	19	26	47	62	79	79	31	2	-
White, both sexes	46	70	70	63	57	70	82	96	78	37	1	-
Male	36	60	52	49	42	42	40	40	14	10	-	-
Female	10	10	18	14	15	28	42	56	64	27	1	-
Black, both sexes	24	35	38	19	31	37	40	34	20	6	3	-
Male	15	23	32	14	21	18	20	11	5	2	2	-
Female	9	12	6	5	10	19	20	23	15	4	1	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	1	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hawaii												
All races, both sexes	8	15	28	20	21	20	22	18	14	2	1	-
Male	7	11	20	18	14	15	16	7	8	1	1	-
Female	1	4	8	2	7	5	6	11	6	1	-	-
White, both sexes	6	6	6	7	5	5	7	5	4	1	-	-
Male	6	4	5	7	5	5	5	2	1	-	-	-
Female	-	2	1	-	-	-	2	3	3	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	9	22	13	15	15	15	13	10	1	1	-
Male	1	7	15	11	9	10	11	5	7	1	1	-
Female	1	2	7	2	6	5	4	8	3	-	-	-
Idaho												
All races, both sexes	5	4	3	4	9	7	15	16	10	4	-	-
Male	3	3	3	3	6	4	7	4	1	1	-	-
Female	2	1	-	1	3	3	8	12	9	3	-	-
White, both sexes	5	4	3	4	9	7	15	16	10	4	-	-
Male	3	3	3	3	6	4	7	4	1	1	-	-
Female	2	1	-	1	3	3	8	12	9	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Illinois												
All races, both sexes	155	209	194	183	169	196	205	187	133	63	11	1
Male	130	156	149	134	110	109	101	81	45	12	2	-
Female	25	53	45	49	59	87	104	106	88	51	9	1
White, both sexes	98	125	127	115	110	145	160	161	121	59	10	-
Male	86	96	98	87	70	79	74	69	42	10	2	-
Female	12	29	29	28	40	66	86	92	79	49	8	-
Black, both sexes	57	79	63	63	57	50	42	24	10	4	1	1
Male	44	57	48	45	40	30	24	11	3	2	-	-
Female	13	22	15	18	17	20	18	13	7	2	1	1
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	5	3	5	2	1	3	2	2	-	-	-
Male	-	3	2	2	-	-	3	1	-	-	-	-
Female	-	2	1	3	2	1	-	1	2	-	-	-
Indiana												
All races, both sexes	38	47	39	47	47	45	78	96	62	35	13	-
Male	27	32	28	34	24	25	36	33	17	12	1	-
Female	11	15	11	13	23	20	42	63	45	23	12	-
White, both sexes	33	37	36	36	39	43	72	92	59	33	12	-
Male	24	25	25	25	18	24	31	32	16	11	1	-
Female	9	12	11	11	21	19	41	60	43	22	11	-
Black, both sexes	5	10	3	10	7	1	6	4	3	2	1	-
Male	3	7	3	8	5	1	5	1	1	1	-	-
Female	2	3	-	2	2	-	1	3	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Iowa												
All races, both sexes	54	64	52	80	74	101	155	165	121	79	22	-
Male	44	48	33	55	51	58	84	68	39	27	3	-
Female	10	16	19	25	23	43	71	97	82	52	19	-
White, both sexes	47	64	50	80	71	99	154	162	121	79	22	-
Male	39	48	32	55	50	57	84	65	39	27	3	-
Female	8	16	18	25	21	42	70	97	82	52	19	-
Black, both sexes	7	-	1	-	2	2	1	3	-	-	-	-
Male	5	-	-	-	1	1	-	3	-	-	-	-
Female	2	-	1	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	25	22	36	34	30	32	49	56	32	14	2	-
Male	22	20	27	24	18	20	25	25	9	1	-	-
Female	3	2	9	10	12	12	24	31	23	13	2	-
White, both sexes	24	19	29	31	25	29	48	51	32	14	2	-
Male	21	17	23	22	15	18	25	21	9	1	-	-
Female	3	2	6	9	10	11	23	30	23	13	2	-
Black, both sexes	1	2	5	2	4	3	1	5	-	-	-	-
Male	1	2	4	1	2	2	-	4	-	-	-	-
Female	-	-	1	1	2	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	1	1	1	-	-	-	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Kentucky												
All races, both sexes	21	31	32	26	36	42	66	70	54	20	7	-
Male	12	20	22	13	21	19	29	26	15	3	-	-
Female	9	11	10	13	15	23	37	44	39	17	7	-
White, both sexes	19	26	31	25	33	42	63	67	52	19	7	-
Male	10	18	21	13	19	19	28	25	14	3	-	-
Female	9	8	10	12	14	23	35	42	38	16	7	-
Black, both sexes	2	5	1	1	3	-	3	3	2	1	-	-
Male	2	2	1	-	2	-	1	1	1	-	-	-
Female	-	3	-	1	1	-	2	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	53	65	66	68	96	94	137	132	80	50	7	-
Male	40	45	49	38	58	48	73	47	19	11	1	-
Female	13	20	17	30	38	46	64	85	61	39	6	-
White, both sexes	34	39	47	45	62	70	106	108	60	42	3	-
Male	27	30	37	27	38	37	58	42	13	11	1	-
Female	7	9	10	18	24	33	48	66	47	31	2	-
Black, both sexes	18	26	19	22	32	24	30	24	19	8	4	-
Male	12	15	12	11	18	11	14	5	6	-	-	-
Female	6	11	7	11	14	13	16	19	13	8	4	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Maine												
All races, both sexes	2	12	13	11	13	15	32	33	21	7	5	-
Male	-	10	8	10	11	7	19	14	9	1	-	-
Female	2	2	5	1	2	8	13	19	12	6	5	-
White, both sexes	2	11	13	11	13	15	32	33	21	7	5	-
Male	-	9	8	10	11	7	19	14	9	1	-	-
Female	2	2	5	1	2	8	13	19	12	6	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	153	168	162	151	206	247	341	368	265	132	31	1
Male	114	125	116	94	125	130	159	135	67	29	6	1
Female	39	43	46	57	81	117	182	233	198	103	25	-
White, both sexes	94	102	98	86	128	177	240	285	206	108	22	1
Male	75	78	68	56	74	96	123	114	57	24	5	1
Female	19	24	30	30	54	81	117	171	149	84	17	-
Black, both sexes	57	64	62	61	75	67	97	80	57	22	9	-
Male	37	45	47	34	48	33	35	19	9	5	1	-
Female	20	19	15	27	27	34	62	61	48	17	8	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	2	4	3	3	4	3	2	2	-	-
Male	2	2	1	4	3	1	1	2	1	-	-	-
Female	-	-	1	-	-	2	3	1	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Massachusetts												
All races, both sexes	46	69	80	76	113	116	157	131	93	44	22	-
Male	41	55	62	51	70	60	76	51	30	9	3	-
Female	5	14	18	25	43	56	81	80	63	35	19	-
White, both sexes	45	64	73	72	108	109	150	124	91	43	21	-
Male	40	50	55	47	67	59	72	48	29	9	3	-
Female	5	14	18	25	41	50	78	76	62	34	18	-
Black, both sexes	-	2	6	4	5	5	5	5	1	1	-	-
Male	-	2	6	4	3	1	4	2	1	-	-	-
Female	-	-	-	-	2	4	1	3	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	-	-	2	2	2	1	-	1	-
Male	-	3	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	2	1	1	-	1	-
Michigan												
All races, both sexes	294	410	344	338	364	418	495	412	268	95	22	-
Male	213	302	237	209	227	241	245	174	83	21	5	-
Female	81	108	107	129	137	177	250	238	185	74	17	-
White, both sexes	173	255	207	200	222	296	346	307	210	68	11	-
Male	137	195	144	125	142	171	171	122	62	17	2	-
Female	36	60	63	75	80	125	175	185	148	51	9	-
Black, both sexes	119	149	134	132	132	119	144	100	54	27	11	-
Male	75	102	92	82	78	69	73	47	20	4	3	-
Female	44	47	42	50	54	50	71	53	34	23	8	-
American Indian 1/, both sexes	1	4	1	5	3	2	3	2	2	-	-	-
Male	-	3	-	2	2	-	1	2	1	-	-	-
Female	1	1	1	3	1	2	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	1	7	1	2	3	2	-	-	-
Male	1	2	1	-	5	1	-	3	-	-	-	-
Female	-	-	1	1	2	-	2	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Minnesota												
All races, both sexes	12	23	18	13	25	31	55	55	43	28	4	-
Male	9	16	10	6	16	21	35	34	13	7	-	-
Female	3	7	8	7	9	10	20	21	30	21	4	-
White, both sexes	12	23	17	13	23	29	55	55	42	28	4	-
Male	9	16	10	6	14	20	35	34	13	7	-	-
Female	3	7	7	7	9	9	20	21	29	21	4	-
Black, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Mississippi												
All races, both sexes	16	22	31	31	56	65	93	104	104	43	15	-
Male	12	9	19	21	24	30	33	27	19	7	-	-
Female	4	13	12	10	32	35	60	77	85	36	15	-
White, both sexes	8	11	19	23	35	57	71	81	86	37	10	-
Male	5	5	13	16	14	27	28	17	14	4	-	-
Female	3	6	6	7	21	30	43	64	72	33	10	-
Black, both sexes	8	10	12	8	20	8	22	23	18	6	5	-
Male	7	4	6	5	10	3	5	10	5	3	-	-
Female	1	6	6	3	10	5	17	13	13	3	5	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Missouri												
All races, both sexes	12	30	39	49	52	69	94	104	65	34	12	-
Male	12	23	26	37	31	33	46	39	15	6	1	-
Female	-	7	13	12	21	36	48	65	50	28	11	-
White, both sexes	10	26	31	43	44	61	89	94	60	33	11	-
Male	10	21	20	32	27	30	44	36	14	6	1	-
Female	-	5	11	11	17	31	45	58	46	27	10	-
Black, both sexes	1	4	8	6	7	7	5	9	5	1	1	-
Male	1	2	6	5	4	3	2	2	1	-	-	-
Female	-	2	2	1	3	4	3	7	4	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Montana												
All races, both sexes	8	15	16	24	18	17	19	18	14	7	-	-
Male	8	12	10	14	10	10	12	12	6	1	-	-
Female	-	3	6	10	8	7	7	6	8	6	-	-
White, both sexes	8	15	16	23	18	16	19	18	14	7	-	-
Male	8	12	10	13	10	10	12	12	6	1	-	-
Female	-	3	6	10	8	6	7	6	8	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Nebraska												
All races, both sexes	1	2	5	2	8	8	19	20	27	15	1	-
Male	1	1	4	2	4	5	7	9	9	2	-	-
Female	-	1	1	-	4	3	12	11	18	13	1	-
White, both sexes	1	2	4	2	8	8	19	20	27	15	1	-
Male	1	1	3	2	4	5	7	9	9	2	-	-
Female	-	1	1	-	4	3	12	11	18	13	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	85	115	131	132	137	137	126	104	49	20	3	-
Male	71	93	94	98	94	77	68	45	23	3	-	-
Female	14	22	37	34	43	60	58	59	26	17	3	-
White, both sexes	74	102	118	112	113	123	116	98	45	20	1	-
Male	62	82	84	87	78	69	60	40	22	3	-	-
Female	12	20	34	25	35	54	56	58	23	17	1	-
Black, both sexes	8	7	7	12	18	11	7	3	2	-	2	-
Male	6	7	6	6	13	6	6	2	-	-	-	-
Female	2	-	1	6	5	5	1	1	2	-	2	-
American Indian 1/, both sexes	1	1	2	1	-	1	1	1	-	-	-	-
Male	1	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	4	7	6	2	2	2	2	-	-	-
Male	2	4	3	4	3	1	1	2	1	-	-	-
Female	-	1	1	3	3	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Hampshire												
All races, both sexes	17	28	26	21	29	30	37	41	24	9	1	-
Male	14	25	20	20	17	12	20	16	8	4	1	-
Female	3	3	6	1	12	18	17	25	16	5	-	-
White, both sexes	17	28	26	21	29	30	37	40	24	9	1	-
Male	14	25	20	20	17	12	20	16	8	4	1	-
Female	3	3	6	1	12	18	17	24	16	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
New Jersey												
All races, both sexes	108	132	150	146	164	216	303	313	235	78	19	2
Male	91	101	106	102	100	103	158	112	75	15	2	2
Female	17	31	44	44	64	113	145	201	160	63	17	-
White, both sexes	84	100	126	112	128	184	264	292	219	69	18	2
Male	71	80	89	74	77	85	139	107	69	12	2	2
Female	13	20	37	38	51	99	125	185	150	57	16	-
Black, both sexes	20	27	18	29	32	24	33	17	15	9	1	-
Male	16	17	12	24	22	14	17	3	6	3	-	-
Female	4	10	6	5	10	10	16	14	9	6	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	5	6	4	4	8	5	4	1	-	-	-
Male	4	4	5	4	1	4	2	2	-	-	-	-
Female	-	1	1	-	3	4	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Mexico												
All races, both sexes	50	67	57	57	82	82	81	84	58	29	5	-
Male	42	49	43	38	47	45	44	41	20	6	2	-
Female	8	18	14	19	35	37	37	43	38	23	3	-
White, both sexes	45	60	52	54	71	82	74	83	55	27	5	-
Male	38	45	39	36	41	45	40	40	19	6	2	-
Female	7	15	13	18	30	37	34	43	36	21	3	-
Black, both sexes	1	2	-	2	2	-	3	1	2	2	-	-
Male	1	1	-	1	-	-	1	1	-	-	-	-
Female	-	1	-	1	2	-	2	-	2	2	-	-
American Indian 1/, both sexes	4	4	5	1	9	-	3	-	1	-	-	-
Male	3	2	4	1	6	-	3	-	1	-	-	-
Female	1	2	1	-	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	199	308	389	373	441	449	508	483	338	155	37	-
Male	161	239	276	235	275	251	244	217	122	36	9	-
Female	38	69	113	138	166	198	264	266	216	119	28	-
White, both sexes	129	219	259	231	310	318	395	411	304	135	28	-
Male	111	178	186	149	202	183	198	186	111	31	6	-
Female	18	41	73	82	108	135	197	225	193	104	22	-
Black, both sexes	61	77	118	132	117	120	105	57	28	18	8	-
Male	41	50	80	80	62	61	43	24	8	4	2	-
Female	20	27	38	52	55	59	62	33	20	14	6	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	12	11	10	13	11	8	14	6	2	1	-
Male	9	11	9	6	10	7	3	6	3	1	1	-
Female	-	1	2	4	3	4	5	8	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
North Carolina												
All races, both sexes	60	67	86	55	102	136	163	186	140	58	10	-
Male	48	56	60	38	65	66	69	68	41	10	1	-
Female	12	11	26	17	37	70	94	118	99	48	9	-
White, both sexes	42	47	67	41	79	115	130	165	125	52	9	-
Male	33	39	46	28	52	57	55	57	36	9	1	-
Female	9	8	21	13	27	58	75	108	89	43	8	-
Black, both sexes	16	19	17	14	22	20	32	19	14	6	1	-
Male	13	16	13	10	13	9	13	10	4	1	-	-
Female	3	3	4	4	9	11	19	9	10	5	1	-
American Indian 1/, both sexes	1	1	1	-	1	-	-	1	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	1	1	1	-	-	-
Male	1	-	1	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	13	6	5	10	11	8	13	28	15	8	3	-
Male	10	5	4	4	6	4	4	11	2	2	-	-
Female	3	1	1	6	5	4	9	17	13	6	3	-
White, both sexes	12	5	5	10	10	6	13	27	15	8	3	-
Male	9	4	4	4	5	3	4	11	2	2	-	-
Female	3	1	1	6	5	3	9	16	13	6	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	1	2	-	1	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Ohio												
All races, both sexes	127	156	149	154	166	210	304	300	220	81	17	-
Male	97	120	103	112	94	112	149	120	62	14	3	-
Female	30	36	46	42	72	98	155	180	158	67	14	-
White, both sexes	91	133	123	127	145	172	275	268	203	78	16	-
Male	73	102	86	93	83	95	132	106	60	13	3	-
Female	18	31	37	34	62	77	143	162	143	65	13	-
Black, both sexes	34	23	25	26	21	38	29	31	14	3	1	-
Male	22	18	17	19	11	17	17	14	1	1	-	-
Female	12	5	8	7	10	21	12	17	13	2	1	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	1	3	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	2	-	-	-
Oklahoma												
All races, both sexes	70	81	104	88	96	112	120	147	100	50	9	-
Male	52	62	71	63	62	65	60	56	35	7	2	-
Female	18	19	33	25	34	47	60	91	65	43	7	-
White, both sexes	55	62	83	75	80	94	107	132	93	45	8	-
Male	43	46	54	51	51	58	54	49	34	5	2	-
Female	12	16	29	24	29	36	53	83	59	40	6	-
Black, both sexes	9	11	13	5	4	10	5	7	1	5	1	-
Male	5	8	10	5	2	3	1	3	-	2	-	-
Female	4	3	3	-	2	7	4	4	1	3	1	-
American Indian 1/, both sexes	6	8	8	7	12	6	8	7	6	-	-	-
Male	4	8	7	6	9	3	5	4	1	-	-	-
Female	2	-	1	1	3	3	3	3	5	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Oregon												
All races, both sexes	15	16	19	20	23	33	39	49	29	13	4	-
Male	11	14	14	16	15	17	19	22	14	-	1	-
Female	4	2	5	4	8	16	20	27	15	13	3	-
White, both sexes	15	16	19	19	23	33	39	49	29	12	4	-
Male	11	14	14	15	15	17	19	22	14	-	1	-
Female	4	2	5	4	8	16	20	27	15	12	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	165	270	251	222	269	356	492	581	482	205	48	2
Male	126	207	180	148	172	199	229	232	121	43	4	1
Female	39	63	71	74	97	157	263	349	361	162	44	1
White, both sexes	101	191	181	164	203	278	442	547	460	192	46	2
Male	80	150	131	107	129	148	206	219	116	38	4	1
Female	21	41	50	57	74	130	236	328	344	154	42	1
Black, both sexes	62	78	66	55	64	75	49	32	20	13	1	-
Male	44	56	46	38	42	49	23	13	5	5	-	-
Female	18	22	20	17	22	26	26	19	15	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	1	4	3	2	2	1	2	2	-	-	-
Male	2	1	3	3	1	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Rhode Island												
All races, both sexes	21	22	20	31	22	54	77	104	72	41	12	-
Male	15	19	15	20	15	25	44	39	14	4	2	-
Female	6	3	5	11	7	29	33	65	58	37	10	-
White, both sexes	20	20	19	30	21	52	75	104	71	40	12	-
Male	15	18	14	19	15	25	42	39	14	4	2	-
Female	5	2	5	11	6	27	33	65	57	36	10	-
Black, both sexes	1	2	-	1	-	2	2	-	1	1	-	-
Male	-	1	-	1	-	-	2	-	-	-	-	-
Female	1	1	-	-	-	2	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	32	38	35	25	34	45	49	57	37	11	5	-
Male	27	27	21	19	21	31	18	26	13	2	-	-
Female	5	11	14	6	13	14	31	31	24	9	5	-
White, both sexes	23	28	25	19	26	34	37	42	26	8	4	-
Male	20	21	14	13	15	22	15	22	10	2	-	-
Female	3	7	11	6	11	12	22	20	16	6	4	-
Black, both sexes	8	10	10	6	8	11	12	15	11	3	1	-
Male	7	6	7	6	6	9	3	4	3	-	-	-
Female	1	4	3	-	2	2	9	11	8	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
South Dakota												
All races, both sexes	2	2	2	2	6	9	11	27	18	13	1	-
Male	2	1	2	-	5	7	5	10	5	-	1	-
Female	-	1	-	2	1	2	6	17	13	13	-	-
White, both sexes	2	1	2	2	5	9	9	26	18	12	1	-
Male	2	1	2	-	4	7	5	9	5	-	1	-
Female	-	-	-	2	1	2	4	17	13	12	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	2	1	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	138	171	173	125	157	202	216	211	158	57	18	-
Male	105	121	126	80	105	117	107	82	44	13	4	-
Female	33	50	47	45	52	85	109	129	114	44	14	-
White, both sexes	99	133	139	105	128	176	195	197	134	52	15	-
Male	74	96	104	68	85	104	99	77	37	10	3	-
Female	25	37	35	37	43	72	96	120	97	42	12	-
Black, both sexes	37	37	33	19	27	25	21	14	24	5	3	-
Male	29	25	21	11	19	12	8	5	7	3	1	-
Female	8	12	12	8	8	13	13	9	17	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	1	2	-	-	-	-	-	-	-
Male	2	-	1	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Texas												
All races, both sexes	323	391	428	385	376	406	408	391	272	114	31	-
Male	253	314	322	282	233	249	216	169	90	24	6	-
Female	70	77	106	103	143	157	192	222	182	90	25	-
White, both sexes	250	309	341	313	305	344	356	349	246	99	26	-
Male	198	247	256	234	190	213	183	150	81	23	4	-
Female	52	62	85	79	115	131	173	199	165	76	22	-
Black, both sexes	68	73	83	69	68	57	48	40	23	14	5	-
Male	51	60	62	46	43	33	30	18	8	1	2	-
Female	17	13	21	23	25	24	18	22	15	13	3	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	9	3	3	3	4	3	2	3	1	-	-
Male	4	7	3	2	-	2	2	1	1	-	-	-
Female	1	2	-	1	3	2	1	1	2	1	-	-
Utah												
All races, both sexes	4	4	9	8	6	14	15	20	8	13	1	-
Male	3	2	8	4	4	10	9	11	6	6	-	-
Female	1	2	1	4	2	4	6	9	2	7	1	-
White, both sexes	4	4	9	6	6	13	15	20	8	13	1	-
Male	3	2	8	4	4	9	9	11	6	6	-	-
Female	1	2	1	2	2	4	6	9	2	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Vermont												
All races, both sexes	17	28	15	23	14	35	31	43	30	7	5	-
Male	14	22	12	14	9	20	18	16	6	1	1	-
Female	3	6	3	9	5	15	13	27	24	6	4	-
White, both sexes	17	28	15	23	14	35	31	43	30	7	5	-
Male	14	22	12	14	9	20	18	16	6	1	1	-
Female	3	6	3	9	5	15	13	27	24	6	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	67	68	72	82	88	129	153	169	120	50	11	-
Male	46	51	52	60	50	67	67	65	38	14	2	-
Female	21	17	20	22	38	62	86	104	82	36	9	-
White, both sexes	52	43	53	54	70	97	130	141	105	40	10	-
Male	38	33	36	40	38	53	57	53	33	11	2	-
Female	14	10	17	14	32	44	73	88	72	29	8	-
Black, both sexes	15	24	18	25	16	30	21	27	14	10	1	-
Male	8	17	15	17	11	12	8	12	5	3	-	-
Female	7	7	3	8	5	18	13	15	9	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	2	1	2	1	1	-	-	-
Male	-	1	1	3	1	1	2	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Washington												
All races, both sexes	73	85	118	103	119	142	215	218	201	83	25	-
Male	63	64	86	80	78	77	97	84	58	27	6	-
Female	10	21	32	23	41	65	118	134	143	56	19	-
White, both sexes	66	75	105	92	112	130	209	208	195	80	24	-
Male	57	55	76	70	74	70	95	79	56	27	5	-
Female	9	20	29	22	38	60	114	129	139	53	19	-
Black, both sexes	4	7	6	5	2	3	2	4	2	2	-	-
Male	3	7	4	5	2	2	-	2	-	-	-	-
Female	1	-	2	-	-	1	2	2	2	2	-	-
American Indian 1/, both sexes	1	1	1	1	1	3	-	1	-	-	1	-
Male	1	1	1	1	-	2	-	-	-	-	1	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	6	5	4	6	4	5	4	1	-	-
Male	2	1	5	4	2	3	2	3	2	-	-	-
Female	-	1	1	1	2	3	2	2	2	1	-	-
West Virginia												
All races, both sexes	27	31	26	30	46	54	63	78	64	24	5	-
Male	20	26	21	22	33	26	25	31	19	5	1	-
Female	7	5	5	8	13	28	38	47	45	19	4	-
White, both sexes	27	30	23	30	44	51	61	78	64	23	5	-
Male	20	25	19	22	32	25	25	31	19	5	1	-
Female	7	5	4	8	12	26	36	47	45	18	4	-
Black, both sexes	-	1	3	-	2	3	2	-	-	1	-	-
Male	-	1	2	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	1	2	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Wisconsin												
All races, both sexes	50	66	66	74	70	97	120	118	77	38	13	-
Male	42	53	51	52	52	57	68	56	27	8	4	-
Female	8	13	15	22	18	40	52	62	50	30	9	-
White, both sexes	41	55	59	64	66	88	112	115	73	38	13	-
Male	34	46	47	47	50	54	65	56	25	8	4	-
Female	7	9	12	17	16	34	47	59	48	30	9	-
Black, both sexes	8	10	6	10	3	8	8	3	4	-	-	-
Male	7	6	4	5	2	3	3	-	2	-	-	-
Female	1	4	2	5	1	5	5	3	2	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	3	-	3	4	3	6	3	3	2	-	-
Male	2	-	-	2	2	2	5	2	1	-	-	-
Female	-	3	-	1	2	1	1	1	2	2	-	-
White, both sexes	2	3	-	3	4	3	6	3	3	2	-	-
Male	2	-	-	2	2	2	5	2	1	-	-	-
Female	-	3	-	1	2	1	1	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
United States												
All races, both sexes	219,001	9	1	2	5	13	38	109	292	676	1,814	3,588
Male	112,248	6	1	1	3	10	31	91	233	508	1,383	2,694
Female	106,753	3	-	1	2	3	7	18	59	168	431	894
White, both sexes	194,461	3	1	2	4	9	27	81	220	490	1,447	2,791
Male	100,009	3	1	1	3	8	23	69	177	381	1,140	2,157
Female	94,452	-	-	1	1	1	4	12	43	109	307	634
Black, both sexes	19,858	5	-	-	-	4	11	22	56	157	305	676
Male	9,620	2	-	-	-	2	8	17	41	102	192	445
Female	10,238	3	-	-	-	2	3	5	15	55	113	231
American Indian 1/, both sexes	846	-	-	-	-	-	-	-	3	6	19	30
Male	472	-	-	-	-	-	-	-	3	6	14	23
Female	374	-	-	-	-	-	-	-	-	-	5	7
Asian or Pacific Islander 2/, both sexes	3,836	1	-	-	1	-	-	6	13	23	43	91
Male	2,147	1	-	-	-	-	-	5	12	19	37	69
Female	1,689	-	-	-	1	-	-	1	1	4	6	22
Alabama												
All races, both sexes	2,620	1	-	-	-	1	1	2	4	7	20	50
Male	1,376	1	-	-	-	1	1	1	3	4	12	36
Female	1,244	-	-	-	-	-	-	1	1	3	8	14
White, both sexes	2,152	1	-	-	-	1	1	-	2	5	15	32
Male	1,146	1	-	-	-	1	1	-	1	4	9	27
Female	1,006	-	-	-	-	-	-	-	1	1	6	5
Black, both sexes	462	-	-	-	-	-	-	2	2	2	5	18
Male	227	-	-	-	-	-	-	1	2	-	3	9
Female	235	-	-	-	-	-	-	1	-	2	2	9
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Alaska												
All races, both sexes	133	-	-	-	-	1	-	-	-	1	3	2
Male	84	-	-	-	-	1	-	-	-	1	2	2
Female	49	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	111	-	-	-	-	1	-	-	-	-	2	2
Male	68	-	-	-	-	1	-	-	-	-	2	2
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3,916	-	-	1	-	-	3	1	6	10	28	54
Male	2,134	-	-	-	-	-	3	1	6	7	22	41
Female	1,782	-	-	1	-	-	-	-	-	3	6	13
White, both sexes	3,744	-	-	1	-	-	3	1	5	9	25	51
Male	2,039	-	-	-	-	-	3	1	5	6	20	38
Female	1,705	-	-	1	-	-	-	-	-	3	5	13
Black, both sexes	81	-	-	-	-	-	-	-	1	-	1	2
Male	49	-	-	-	-	-	-	-	1	-	1	2
Female	32	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	60	-	-	-	-	-	-	-	-	1	2	-
Male	31	-	-	-	-	-	-	-	-	1	1	-
Female	29	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Arkansas												
All races, both sexes	1,738	-	-	-	-	-	-	1	3	5	8	27
Male	861	-	-	-	-	-	-	1	1	4	4	17
Female	877	-	-	-	-	-	-	-	2	1	4	10
White, both sexes	1,533	-	-	-	-	-	-	-	2	4	5	20
Male	770	-	-	-	-	-	-	-	1	3	2	13
Female	763	-	-	-	-	-	-	-	1	1	3	7
Black, both sexes	202	-	-	-	-	-	-	1	1	1	3	7
Male	89	-	-	-	-	-	-	1	-	1	2	4
Female	113	-	-	-	-	-	-	-	1	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	25,246	-	-	1	2	3	2	8	24	81	198	418
Male	12,994	-	-	1	1	2	2	8	20	63	156	317
Female	12,252	-	-	-	1	1	-	-	4	18	42	101
White, both sexes	21,651	-	-	1	1	3	2	6	18	55	152	323
Male	11,132	-	-	1	1	2	2	6	14	43	120	250
Female	10,519	-	-	-	-	1	-	-	4	12	32	73
Black, both sexes	1,652	-	-	-	-	-	-	1	2	15	28	61
Male	812	-	-	-	-	-	-	1	2	10	18	39
Female	840	-	-	-	-	-	-	-	-	5	10	22
American Indian 1/, both sexes	116	-	-	-	-	-	-	-	-	1	3	1
Male	57	-	-	-	-	-	-	-	-	1	3	1
Female	59	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1,827	-	-	-	1	-	-	1	4	10	15	33
Male	993	-	-	-	-	-	-	1	4	9	15	27
Female	834	-	-	-	1	-	-	-	-	1	-	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Colorado												
All races, both sexes	2,068	1	1	-	-	1	-	4	4	9	22	48
Male	1,147	1	1	-	-	1	-	3	4	8	19	39
Female	921	-	-	-	-	-	-	1	-	1	3	9
White, both sexes	1,993	1	1	-	-	-	-	4	3	9	21	43
Male	1,102	1	1	-	-	-	-	3	3	8	18	35
Female	891	-	-	-	-	-	-	1	-	1	3	8
Black, both sexes	53	-	-	-	-	1	-	-	1	-	-	5
Male	34	-	-	-	-	1	-	-	1	-	-	4
Female	19	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,598	-	-	-	-	-	1	1	3	13	17	23
Male	1,267	-	-	-	-	-	1	1	3	10	14	19
Female	1,331	-	-	-	-	-	-	-	-	3	3	4
White, both sexes	2,452	-	-	-	-	-	1	1	3	11	14	14
Male	1,198	-	-	-	-	-	1	1	3	8	13	12
Female	1,254	-	-	-	-	-	-	-	-	3	1	2
Black, both sexes	131	-	-	-	-	-	-	-	-	2	3	8
Male	59	-	-	-	-	-	-	-	-	2	1	6
Female	72	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Delaware												
All races, both sexes	647	-	-	-	-	-	-	-	1	2	10	16
Male	317	-	-	-	-	-	-	-	1	1	7	13
Female	330	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	558	-	-	-	-	-	-	-	-	2	9	10
Male	273	-	-	-	-	-	-	-	-	1	6	8
Female	285	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	83	-	-	-	-	-	-	-	1	-	1	6
Male	42	-	-	-	-	-	-	-	1	-	1	5
Female	41	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	293	-	-	-	-	-	-	1	1	-	3	4
Male	134	-	-	-	-	-	-	1	1	-	1	3
Female	159	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	77	-	-	-	-	-	-	-	-	-	1	2
Male	42	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	215	-	-	-	-	-	-	1	1	-	2	2
Male	91	-	-	-	-	-	-	1	1	-	1	1
Female	124	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Florida												
All races, both sexes	19,046	-	-	-	1	1	3	6	17	40	141	269
Male	10,201	-	-	-	1	-	1	3	14	30	106	201
Female	8,845	-	-	-	-	1	2	3	3	10	35	68
White, both sexes	17,403	-	-	-	1	-	2	5	11	28	107	213
Male	9,379	-	-	-	1	-	1	3	10	21	83	169
Female	8,024	-	-	-	-	-	1	2	1	7	24	44
Black, both sexes	1,555	-	-	-	-	1	1	1	5	12	31	50
Male	768	-	-	-	-	-	-	-	3	9	20	27
Female	787	-	-	-	-	1	1	1	2	3	11	23
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	74	-	-	-	-	-	-	-	-	-	3	6
Male	47	-	-	-	-	-	-	-	-	-	3	5
Female	27	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	4,092	-	-	-	-	-	-	7	14	25	66	128
Male	2,218	-	-	-	-	-	-	5	8	17	55	103
Female	1,874	-	-	-	-	-	-	2	6	8	11	25
White, both sexes	3,162	-	-	-	-	-	-	4	8	9	45	75
Male	1,774	-	-	-	-	-	-	2	6	7	39	65
Female	1,388	-	-	-	-	-	-	2	2	2	6	10
Black, both sexes	894	-	-	-	-	-	-	3	6	16	19	52
Male	425	-	-	-	-	-	-	3	2	10	14	38
Female	469	-	-	-	-	-	-	-	4	6	5	14
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	15	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hawaii												
All races, both sexes	675	-	-	-	-	-	-	1	2	5	8	13
Male	438	-	-	-	-	-	-	1	2	3	5	9
Female	237	-	-	-	-	-	-	-	-	2	3	4
White, both sexes	202	-	-	-	-	-	-	-	-	-	-	1
Male	127	-	-	-	-	-	-	-	-	-	-	-
Female	75	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	468	-	-	-	-	-	-	1	2	4	8	11
Male	307	-	-	-	-	-	-	1	2	2	5	8
Female	161	-	-	-	-	-	-	-	-	2	3	3
Idaho												
All races, both sexes	668	-	-	-	-	-	-	1	-	1	4	6
Male	388	-	-	-	-	-	-	1	-	1	3	3
Female	280	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	663	-	-	-	-	-	-	1	-	1	4	6
Male	385	-	-	-	-	-	-	1	-	1	3	3
Female	278	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Illinois												
All races, both sexes	8,963	3	-	-	-	-	-	7	10	33	104	154
Male	4,437	1	-	-	-	-	-	6	10	26	70	114
Female	4,526	2	-	-	-	-	-	1	-	7	34	40
White, both sexes	7,806	-	-	-	-	-	-	4	4	24	76	109
Male	3,843	-	-	-	-	-	-	3	4	21	57	84
Female	3,963	-	-	-	-	-	-	1	-	3	19	25
Black, both sexes	1,063	3	-	-	-	-	-	3	6	9	27	43
Male	537	1	-	-	-	-	-	3	6	5	13	29
Female	526	2	-	-	-	-	-	-	-	4	14	14
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	90	-	-	-	-	-	-	-	-	-	1	2
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	5,001	-	-	-	-	1	1	4	15	21	62	101
Male	2,632	-	-	-	-	-	-	4	9	14	51	71
Female	2,369	-	-	-	-	1	1	-	6	7	11	30
White, both sexes	4,591	-	-	-	-	-	1	4	11	19	54	87
Male	2,409	-	-	-	-	-	-	4	7	14	44	63
Female	2,182	-	-	-	-	-	1	-	4	5	10	24
Black, both sexes	392	-	-	-	-	1	-	-	4	2	7	13
Male	210	-	-	-	-	-	-	-	2	-	6	7
Female	182	-	-	-	-	1	-	-	2	2	1	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Iowa												
All races, both sexes	2,674	-	-	-	-	-	1	2	4	4	17	41
Male	1,316	-	-	-	-	-	1	2	3	3	15	30
Female	1,358	-	-	-	-	-	-	-	1	1	2	11
White, both sexes	2,638	-	-	-	-	-	1	1	4	2	17	37
Male	1,295	-	-	-	-	-	1	1	3	2	15	29
Female	1,343	-	-	-	-	-	-	-	1	-	2	8
Black, both sexes	26	-	-	-	-	-	-	-	-	2	-	2
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	1	-	-	-	2
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	1,951	-	-	-	-	-	-	1	5	2	18	42
Male	1,060	-	-	-	-	-	-	1	4	2	10	27
Female	891	-	-	-	-	-	-	-	1	-	8	15
White, both sexes	1,850	-	-	-	-	-	-	1	4	2	16	36
Male	1,007	-	-	-	-	-	-	1	4	2	10	23
Female	843	-	-	-	-	-	-	-	-	-	6	13
Black, both sexes	81	-	-	-	-	-	-	-	1	-	2	4
Male	40	-	-	-	-	-	-	-	-	-	-	3
Female	41	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Kentucky												
All races, both sexes	2,651	-	-	-	-	-	1	2	5	16	40	56
Male	1,361	-	-	-	-	-	-	2	5	11	34	47
Female	1,290	-	-	-	-	-	1	-	-	5	6	9
White, both sexes	2,498	-	-	-	-	-	1	2	3	13	34	55
Male	1,286	-	-	-	-	-	-	2	3	9	30	46
Female	1,212	-	-	-	-	-	1	-	-	4	4	9
Black, both sexes	145	-	-	-	-	-	-	-	1	3	6	1
Male	71	-	-	-	-	-	-	-	1	2	4	1
Female	74	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2,201	-	-	-	-	-	-	1	6	9	26	58
Male	1,206	-	-	-	-	-	-	1	5	7	14	42
Female	995	-	-	-	-	-	-	-	1	2	12	16
White, both sexes	1,657	-	-	-	-	-	-	1	2	4	17	33
Male	920	-	-	-	-	-	-	1	2	3	9	27
Female	737	-	-	-	-	-	-	-	-	1	8	6
Black, both sexes	536	-	-	-	-	-	-	-	4	5	9	25
Male	282	-	-	-	-	-	-	-	3	4	5	15
Female	254	-	-	-	-	-	-	-	1	1	4	10
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Maine												
All races, both sexes	966	-	-	-	-	-	-	2	3	4	8	19
Male	530	-	-	-	-	-	-	2	2	2	7	16
Female	436	-	-	-	-	-	-	-	1	2	1	3
White, both sexes	961	-	-	-	-	-	-	2	3	4	8	18
Male	525	-	-	-	-	-	-	2	2	2	7	15
Female	436	-	-	-	-	-	-	-	1	2	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,874	-	-	-	-	-	-	-	1	7	8	35
Male	1,417	-	-	-	-	-	-	-	1	5	6	27
Female	1,457	-	-	-	-	-	-	-	-	2	2	8
White, both sexes	2,200	-	-	-	-	-	-	-	-	3	4	18
Male	1,095	-	-	-	-	-	-	-	-	2	3	14
Female	1,105	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	630	-	-	-	-	-	-	-	-	4	4	16
Male	297	-	-	-	-	-	-	-	-	3	3	13
Female	333	-	-	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	41	-	-	-	-	-	-	-	1	-	-	1
Male	24	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Massachusetts												
All races, both sexes	4,429	-	-	-	-	-	-	4	6	9	35	50
Male	2,157	-	-	-	-	-	-	3	6	6	29	38
Female	2,272	-	-	-	-	-	-	1	-	3	6	12
White, both sexes	4,278	-	-	-	-	-	-	4	6	7	31	44
Male	2,075	-	-	-	-	-	-	3	6	5	26	32
Female	2,203	-	-	-	-	-	-	1	-	2	5	12
Black, both sexes	115	-	-	-	-	-	-	-	-	2	4	3
Male	64	-	-	-	-	-	-	-	-	1	3	3
Female	51	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	8,168	-	-	-	-	-	-	2	10	26	50	114
Male	4,023	-	-	-	-	-	-	1	8	16	36	75
Female	4,145	-	-	-	-	-	-	1	2	10	14	39
White, both sexes	7,350	-	-	-	-	-	-	2	9	20	43	93
Male	3,657	-	-	-	-	-	-	1	7	12	34	67
Female	3,693	-	-	-	-	-	-	1	2	8	9	26
Black, both sexes	762	-	-	-	-	-	-	-	1	5	6	19
Male	344	-	-	-	-	-	-	-	1	3	2	7
Female	418	-	-	-	-	-	-	-	-	2	4	12
American Indian 1/, both sexes	35	-	-	-	-	-	-	-	-	1	1	2
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	23	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Minnesota												
All races, both sexes	2,760	-	-	-	-	-	2	2	6	14	40	68
Male	1,617	-	-	-	-	-	2	2	3	11	32	60
Female	1,143	-	-	-	-	-	-	-	3	3	8	8
White, both sexes	2,661	-	-	-	-	-	1	2	5	14	36	62
Male	1,549	-	-	-	-	-	1	2	3	11	28	54
Female	1,112	-	-	-	-	-	-	-	2	3	8	8
Black, both sexes	53	-	-	-	-	-	1	-	-	-	2	3
Male	34	-	-	-	-	-	1	-	-	-	2	3
Female	19	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	1	2
Male	18	-	-	-	-	-	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	1	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	1	-	-	-
Mississippi												
All races, both sexes	1,805	-	-	-	-	-	-	1	13	20	46	61
Male	964	-	-	-	-	-	-	1	10	15	33	44
Female	841	-	-	-	-	-	-	-	3	5	13	17
White, both sexes	1,345	-	-	-	-	-	-	-	10	14	32	37
Male	714	-	-	-	-	-	-	-	8	10	21	26
Female	631	-	-	-	-	-	-	-	2	4	11	11
Black, both sexes	452	-	-	-	-	-	-	1	3	5	14	21
Male	246	-	-	-	-	-	-	1	2	4	12	18
Female	206	-	-	-	-	-	-	-	1	1	2	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	2
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Missouri												
All races, both sexes	5,006	-	-	-	-	1	3	2	13	12	49	108
Male	2,464	-	-	-	-	1	2	2	8	7	41	79
Female	2,542	-	-	-	-	-	1	-	5	5	8	29
White, both sexes	4,537	-	-	-	-	1	2	2	10	10	40	85
Male	2,230	-	-	-	-	1	1	2	6	5	34	60
Female	2,307	-	-	-	-	-	1	-	4	5	6	25
Black, both sexes	443	-	-	-	-	-	1	-	3	2	9	21
Male	220	-	-	-	-	-	1	-	2	2	7	17
Female	223	-	-	-	-	-	-	-	1	-	2	4
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	562	-	-	-	-	-	-	-	3	1	12	17
Male	333	-	-	-	-	-	-	-	3	-	9	13
Female	229	-	-	-	-	-	-	-	-	1	3	4
White, both sexes	537	-	-	-	-	-	-	-	2	-	10	17
Male	319	-	-	-	-	-	-	-	2	-	8	13
Female	218	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	1	-	2	-
Male	13	-	-	-	-	-	-	-	1	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Nebraska												
All races, both sexes	1,196	-	-	-	-	-	1	-	-	3	7	25
Male	628	-	-	-	-	-	1	-	-	3	7	18
Female	568	-	-	-	-	-	-	-	-	-	-	7
White, both sexes	1,152	-	-	-	-	-	1	-	-	1	7	21
Male	605	-	-	-	-	-	1	-	-	1	7	15
Female	547	-	-	-	-	-	-	-	-	-	-	6
Black, both sexes	33	-	-	-	-	-	-	-	-	2	-	4
Male	18	-	-	-	-	-	-	-	-	2	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	961	-	-	-	-	-	2	1	2	5	17	22
Male	573	-	-	-	-	-	2	1	2	5	14	17
Female	388	-	-	-	-	-	-	-	-	-	3	5
White, both sexes	864	-	-	-	-	-	2	-	2	2	14	19
Male	520	-	-	-	-	-	2	-	2	2	11	16
Female	344	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	52	-	-	-	-	-	-	1	-	3	2	1
Male	27	-	-	-	-	-	-	1	-	3	2	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	-	-	1	1
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Hampshire												
All races, both sexes	893	-	-	-	-	-	-	-	1	2	9	15
Male	448	-	-	-	-	-	-	-	1	1	7	13
Female	445	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	887	-	-	-	-	-	-	-	1	2	9	15
Male	444	-	-	-	-	-	-	-	1	1	7	13
Female	443	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7,459	-	-	-	-	-	-	1	5	22	56	80
Male	3,509	-	-	-	-	-	-	1	5	18	45	58
Female	3,950	-	-	-	-	-	-	-	-	4	11	22
White, both sexes	6,788	-	-	-	-	-	-	1	4	15	46	66
Male	3,198	-	-	-	-	-	-	1	4	12	41	51
Female	3,590	-	-	-	-	-	-	-	-	3	5	15
Black, both sexes	571	-	-	-	-	-	-	-	1	5	9	13
Male	262	-	-	-	-	-	-	-	1	4	3	6
Female	309	-	-	-	-	-	-	-	-	1	6	7
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	93	-	-	-	-	-	-	-	-	2	1	1
Male	45	-	-	-	-	-	-	-	-	2	1	1
Female	48	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Mexico												
All races, both sexes	1,064	-	-	-	-	1	-	-	3	1	5	13
Male	554	-	-	-	-	1	-	-	3	1	5	9
Female	510	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	1,014	-	-	-	-	1	-	-	3	1	5	10
Male	524	-	-	-	-	1	-	-	3	1	5	6
Female	490	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	25,809	-	-	-	1	-	8	13	10	44	95	238
Male	11,393	-	-	-	-	-	8	6	8	39	70	179
Female	14,416	-	-	-	1	-	-	7	2	5	25	59
White, both sexes	22,137	-	-	-	1	-	5	7	8	35	73	166
Male	9,871	-	-	-	-	-	5	4	6	32	56	134
Female	12,266	-	-	-	1	-	-	3	2	3	17	32
Black, both sexes	3,045	-	-	-	-	-	3	4	1	8	18	59
Male	1,207	-	-	-	-	-	3	1	1	6	11	35
Female	1,838	-	-	-	-	-	-	3	-	2	7	24
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	606	-	-	-	-	-	-	2	1	1	4	13
Male	311	-	-	-	-	-	-	1	1	1	3	10
Female	295	-	-	-	-	-	-	1	-	-	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
North Carolina												
All races, both sexes	6,145	-	-	-	-	-	1	5	13	33	91	160
Male	3,514	-	-	-	-	-	-	5	12	24	65	129
Female	2,631	-	-	-	-	-	1	-	1	9	26	31
White, both sexes	4,858	-	-	-	-	-	-	2	11	21	57	109
Male	2,819	-	-	-	-	-	-	2	10	15	45	90
Female	2,039	-	-	-	-	-	-	-	1	6	12	19
Black, both sexes	1,193	-	-	-	-	-	1	3	2	12	31	47
Male	641	-	-	-	-	-	-	3	2	9	18	37
Female	552	-	-	-	-	-	1	-	-	3	13	10
American Indian 1/, both sexes	76	-	-	-	-	-	-	-	-	-	2	4
Male	47	-	-	-	-	-	-	-	-	-	1	2
Female	29	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	514	-	-	-	-	-	-	1	-	1	2	5
Male	276	-	-	-	-	-	-	1	-	1	1	4
Female	238	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	498	-	-	-	-	-	-	1	-	-	2	4
Male	266	-	-	-	-	-	-	1	-	-	1	3
Female	232	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Ohio												
All races, both sexes	11,297	-	-	-	-	-	1	5	10	35	77	175
Male	5,740	-	-	-	-	-	1	5	9	28	63	134
Female	5,557	-	-	-	-	-	-	-	1	7	14	41
White, both sexes	10,194	-	-	-	-	-	1	5	9	30	64	140
Male	5,158	-	-	-	-	-	1	5	8	25	54	109
Female	5,036	-	-	-	-	-	-	-	1	5	10	31
Black, both sexes	1,068	-	-	-	-	-	-	-	1	5	13	35
Male	563	-	-	-	-	-	-	-	1	3	9	25
Female	505	-	-	-	-	-	-	-	-	2	4	10
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3,116	-	-	-	-	-	-	2	1	15	29	48
Male	1,669	-	-	-	-	-	-	2	1	11	17	36
Female	1,447	-	-	-	-	-	-	-	-	4	12	12
White, both sexes	2,781	-	-	-	-	-	-	1	-	12	22	44
Male	1,479	-	-	-	-	-	-	1	-	10	12	34
Female	1,302	-	-	-	-	-	-	-	-	2	10	10
Black, both sexes	153	-	-	-	-	-	-	1	-	3	3	1
Male	82	-	-	-	-	-	-	1	-	1	2	1
Female	71	-	-	-	-	-	-	-	-	2	1	-
American Indian 1/, both sexes	172	-	-	-	-	-	-	-	1	-	4	3
Male	101	-	-	-	-	-	-	-	1	-	3	1
Female	71	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Oregon												
All races, both sexes	2,457	-	-	-	-	1	-	-	2	9	23	52
Male	1,500	-	-	-	-	1	-	-	1	7	20	42
Female	957	-	-	-	-	-	-	-	1	2	3	10
White, both sexes	2,391	-	-	-	-	1	-	-	2	9	23	48
Male	1,464	-	-	-	-	1	-	-	1	7	20	39
Female	927	-	-	-	-	-	-	-	1	2	3	9
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	10,617	-	-	-	1	-	2	1	9	20	46	122
Male	5,204	-	-	-	1	-	2	1	7	16	33	91
Female	5,413	-	-	-	-	-	-	-	2	4	13	31
White, both sexes	9,911	-	-	-	1	-	1	1	7	9	39	95
Male	4,879	-	-	-	1	-	1	1	6	9	28	74
Female	5,032	-	-	-	-	-	-	-	1	-	11	21
Black, both sexes	672	-	-	-	-	-	1	-	2	9	7	23
Male	304	-	-	-	-	-	1	-	1	5	5	13
Female	368	-	-	-	-	-	-	-	1	4	2	10
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	2	-	4
Male	19	-	-	-	-	-	-	-	-	2	-	4
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Rhode Island												
All races, both sexes	933	-	-	-	-	-	-	-	-	1	9	6
Male	413	-	-	-	-	-	-	-	-	1	8	6
Female	520	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	915	-	-	-	-	-	-	-	-	-	9	6
Male	405	-	-	-	-	-	-	-	-	-	8	6
Female	510	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,475	-	-	-	-	-	1	2	-	16	35	68
Male	1,464	-	-	-	-	-	-	2	-	11	29	45
Female	1,011	-	-	-	-	-	1	-	-	5	6	23
White, both sexes	1,948	-	-	-	-	-	-	2	-	12	26	53
Male	1,182	-	-	-	-	-	-	2	-	9	24	38
Female	766	-	-	-	-	-	-	-	-	3	2	15
Black, both sexes	522	-	-	-	-	-	1	-	-	4	9	14
Male	277	-	-	-	-	-	-	-	-	2	5	6
Female	245	-	-	-	-	-	1	-	-	2	4	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
South Dakota												
All races, both sexes	659	-	-	-	-	-	-	-	1	-	4	15
Male	341	-	-	-	-	-	-	-	1	-	4	13
Female	318	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	634	-	-	-	-	-	-	-	1	-	3	13
Male	326	-	-	-	-	-	-	-	1	-	3	11
Female	308	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4,368	2	-	-	-	-	-	-	5	17	54	96
Male	2,274	1	-	-	-	-	-	-	4	13	37	75
Female	2,094	1	-	-	-	-	-	-	1	4	17	21
White, both sexes	3,835	-	-	-	-	-	-	-	4	9	44	73
Male	2,012	-	-	-	-	-	-	-	3	8	30	58
Female	1,823	-	-	-	-	-	-	-	1	1	14	15
Black, both sexes	520	2	-	-	-	-	-	-	1	8	9	22
Male	256	1	-	-	-	-	-	-	1	5	6	16
Female	264	1	-	-	-	-	-	-	-	3	3	6
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Texas												
All races, both sexes	11,158	1	-	-	-	1	1	5	21	23	90	221
Male	5,869	1	-	-	-	1	1	4	16	17	68	153
Female	5,289	-	-	-	-	-	-	1	5	6	22	68
White, both sexes	9,986	1	-	-	-	1	-	5	18	19	78	172
Male	5,282	1	-	-	-	1	-	4	13	16	61	124
Female	4,704	-	-	-	-	-	-	1	5	3	17	48
Black, both sexes	1,072	-	-	-	-	-	1	-	2	4	12	45
Male	535	-	-	-	-	-	1	-	2	1	7	26
Female	537	-	-	-	-	-	-	-	-	3	5	19
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	79	-	-	-	-	-	-	-	1	-	-	2
Male	42	-	-	-	-	-	-	-	1	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	809	-	-	-	-	-	-	-	-	6	9	21
Male	526	-	-	-	-	-	-	-	-	6	7	17
Female	283	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	795	-	-	-	-	-	-	-	-	4	8	21
Male	514	-	-	-	-	-	-	-	-	4	6	17
Female	281	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Vermont												
All races, both sexes	353	-	-	-	-	-	-	-	-	-	3	4
Male	183	-	-	-	-	-	-	-	-	-	2	4
Female	170	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	352	-	-	-	-	-	-	-	-	-	3	4
Male	182	-	-	-	-	-	-	-	-	-	2	4
Female	170	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,785	-	-	-	-	1	1	4	8	16	40	78
Male	2,111	-	-	-	-	1	1	4	4	12	36	59
Female	1,674	-	-	-	-	-	-	-	4	4	4	19
White, both sexes	3,102	-	-	-	-	-	-	3	6	11	33	56
Male	1,747	-	-	-	-	-	-	3	3	10	31	41
Female	1,355	-	-	-	-	-	-	-	3	1	2	15
Black, both sexes	638	-	-	-	-	1	1	-	1	5	6	21
Male	338	-	-	-	-	1	1	-	-	2	4	17
Female	300	-	-	-	-	-	-	-	1	3	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	1	1	-	1	1
Male	26	-	-	-	-	-	-	1	1	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Washington												
All races, both sexes	3,645	-	-	-	-	-	1	-	8	8	23	49
Male	2,017	-	-	-	-	-	1	-	8	6	18	40
Female	1,628	-	-	-	-	-	-	-	-	2	5	9
White, both sexes	3,449	-	-	-	-	-	1	-	7	7	19	41
Male	1,902	-	-	-	-	-	1	-	7	5	14	32
Female	1,547	-	-	-	-	-	-	-	-	2	5	9
Black, both sexes	64	-	-	-	-	-	-	-	-	1	1	4
Male	33	-	-	-	-	-	-	-	-	1	1	4
Female	31	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	40	-	-	-	-	-	-	-	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	2	2
Female	20	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	92	-	-	-	-	-	-	-	1	-	1	2
Male	62	-	-	-	-	-	-	-	1	-	1	2
Female	30	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,896	-	-	-	-	-	1	1	7	15	29	52
Male	1,036	-	-	-	-	-	1	1	4	10	20	39
Female	860	-	-	-	-	-	-	-	3	5	9	13
White, both sexes	1,842	-	-	-	-	-	1	1	6	15	29	52
Male	1,008	-	-	-	-	-	1	1	3	10	20	39
Female	834	-	-	-	-	-	-	-	3	5	9	13
Black, both sexes	53	-	-	-	-	-	-	-	1	-	-	-
Male	28	-	-	-	-	-	-	-	1	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Wisconsin												
All races, both sexes	3,367	1	-	-	-	-	-	4	7	6	15	36
Male	1,764	1	-	-	-	-	-	4	7	4	13	26
Female	1,603	-	-	-	-	-	-	-	-	2	2	10
White, both sexes	3,247	-	-	-	-	-	-	4	6	6	14	35
Male	1,699	-	-	-	-	-	-	4	6	4	12	25
Female	1,548	-	-	-	-	-	-	-	-	2	2	10
Black, both sexes	100	-	-	-	-	-	-	-	1	-	1	-
Male	53	-	-	-	-	-	-	-	1	-	1	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	1	-	-	-	-	-	-	-	-	-	1
Male	4	1	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	274	-	-	-	-	-	-	1	-	1	3	5
Male	166	-	-	-	-	-	-	1	-	-	1	1
Female	108	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	271	-	-	-	-	-	-	1	-	1	2	5
Male	164	-	-	-	-	-	-	1	-	-	1	1
Female	107	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
United States												
All races, both sexes	5,482	8,006	9,831	12,644	18,150	27,973	39,682	42,293	31,996	13,337	3,053	7
Male	4,134	5,887	6,848	8,459	11,366	16,565	20,603	18,648	11,079	3,196	497	5
Female	1,348	2,119	2,983	4,185	6,784	11,408	19,079	23,645	20,917	10,141	2,556	2
White, both sexes	4,338	6,421	8,094	10,483	15,511	24,755	35,854	39,127	29,739	12,329	2,728	7
Male	3,359	4,832	5,778	7,150	9,887	14,866	18,907	17,460	10,396	2,961	445	5
Female	979	1,589	2,316	3,333	5,624	9,889	16,947	21,667	19,343	9,368	2,283	2
Black, both sexes	1,007	1,352	1,452	1,761	2,194	2,596	3,005	2,437	1,739	808	271	-
Male	669	873	868	1,029	1,225	1,340	1,292	869	461	151	34	-
Female	338	479	584	732	969	1,256	1,713	1,568	1,278	657	237	-
American Indian 1/, both sexes	42	57	83	99	101	111	123	93	58	18	3	-
Male	31	44	51	63	64	63	54	36	18	2	-	-
Female	11	13	32	36	37	48	69	57	40	16	3	-
Asian or Pacific Islander 2/, both sexes	95	176	202	301	344	511	700	636	460	182	51	-
Male	75	138	151	217	190	296	350	283	204	82	18	-
Female	20	38	51	84	154	215	350	353	256	100	33	-
Alabama												
All races, both sexes	77	123	127	190	307	389	455	427	303	112	24	-
Male	59	89	95	121	200	218	218	178	108	28	3	-
Female	18	34	32	69	107	171	237	249	195	84	21	-
White, both sexes	51	94	93	159	256	310	380	369	270	95	18	-
Male	38	68	73	101	171	181	187	155	99	27	2	-
Female	13	26	20	58	85	129	193	214	171	68	16	-
Black, both sexes	26	28	34	31	49	79	74	57	32	17	6	-
Male	21	20	22	20	28	37	31	23	8	1	1	-
Female	5	8	12	11	21	42	43	34	24	16	5	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Alaska												
All races, both sexes	9	12	5	10	16	20	23	17	7	5	2	-
Male	6	8	3	6	11	18	12	9	3	2	-	-
Female	3	4	2	4	5	2	11	8	4	3	2	-
White, both sexes	7	10	4	8	12	18	19	16	6	4	2	-
Male	4	7	2	4	8	16	10	9	2	1	-	-
Female	3	3	2	4	4	2	9	7	4	3	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	2	4	2	3	1	1	-	-	-
Male	1	-	1	2	3	2	2	-	1	-	-	-
Female	-	1	-	-	1	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	96	161	206	235	319	539	691	770	537	213	43	3
Male	76	117	149	159	205	320	381	374	212	45	13	3
Female	20	44	57	76	114	219	310	396	325	168	30	-
White, both sexes	87	153	190	222	302	517	668	740	521	206	40	3
Male	68	110	135	153	195	311	371	359	206	43	12	3
Female	19	43	55	69	107	206	297	381	315	163	28	-
Black, both sexes	6	3	9	4	9	8	9	16	6	5	2	-
Male	5	3	9	3	7	2	5	7	3	1	-	-
Female	1	-	-	1	2	6	4	9	3	4	2	-
American Indian 1/, both sexes	2	5	6	7	2	7	11	11	6	-	-	-
Male	2	4	4	2	-	5	4	7	1	-	-	-
Female	-	1	2	5	2	2	7	4	5	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	6	7	3	3	4	2	1	-
Male	1	-	1	1	3	2	1	1	2	1	1	-
Female	-	-	-	1	3	5	2	2	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Arkansas												
All races, both sexes	32	53	98	118	174	222	305	323	245	105	19	-
Male	20	38	67	72	108	124	156	138	84	24	3	-
Female	12	15	31	46	66	98	149	185	161	81	16	-
White, both sexes	19	41	78	98	150	203	276	304	216	98	19	-
Male	14	32	54	62	97	117	142	131	77	22	3	-
Female	5	9	24	36	53	86	134	173	139	76	16	-
Black, both sexes	13	12	20	20	24	18	28	19	29	6	-	-
Male	6	6	13	10	11	7	13	7	7	1	-	-
Female	7	6	7	10	13	11	15	12	22	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
California												
All races, both sexes	542	877	953	1,318	1,978	3,063	4,724	5,100	3,899	1,657	396	2
Male	411	605	674	852	1,220	1,841	2,470	2,391	1,454	425	80	1
Female	131	272	279	466	758	1,222	2,254	2,709	2,445	1,232	316	1
White, both sexes	435	676	780	1,045	1,631	2,574	4,093	4,554	3,477	1,486	337	2
Male	335	478	562	683	1,024	1,559	2,150	2,174	1,298	365	64	1
Female	100	198	218	362	607	1,015	1,943	2,380	2,179	1,121	273	1
Black, both sexes	71	123	87	137	191	222	252	202	160	72	28	-
Male	47	72	53	73	109	131	126	67	44	17	3	-
Female	24	51	34	64	82	91	126	135	116	55	25	-
American Indian 1/, both sexes	4	4	9	19	12	18	21	14	7	3	-	-
Male	4	2	4	12	8	10	9	2	1	-	-	-
Female	-	2	5	7	4	8	12	12	6	3	-	-
Asian or Pacific Islander 2/, both sexes	32	74	77	117	144	249	358	330	255	96	31	-
Male	25	53	55	84	79	141	185	148	111	43	13	-
Female	7	21	22	33	65	108	173	182	144	53	18	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Colorado												
All races, both sexes	71	101	114	113	171	257	333	376	287	136	19	-
Male	55	84	85	73	108	163	176	182	108	35	2	-
Female	16	17	29	40	63	94	157	194	179	101	17	-
White, both sexes	68	94	108	107	160	245	326	371	282	132	18	-
Male	52	79	82	69	103	156	173	179	105	33	2	-
Female	16	15	26	38	57	89	153	192	177	99	16	-
Black, both sexes	2	6	6	6	6	7	5	2	2	3	1	-
Male	2	4	3	4	4	4	2	2	2	1	-	-
Female	-	2	3	2	2	3	3	-	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	1	3	1	1	1	-	-	-
Male	-	-	-	-	1	3	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	4	2	1	2	2	1	-	-
Male	1	1	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	4	2	-	2	2	-	-	-
Connecticut												
All races, both sexes	40	64	85	121	179	305	484	579	428	201	54	-
Male	31	50	56	83	115	173	256	260	147	43	5	-
Female	9	14	29	38	64	132	228	319	281	158	49	-
White, both sexes	36	53	72	103	164	279	467	562	423	198	51	-
Male	29	43	51	75	104	164	247	253	146	43	5	-
Female	7	10	21	28	60	115	220	309	277	155	46	-
Black, both sexes	4	11	12	15	14	22	16	15	3	3	3	-
Male	2	7	4	6	10	7	8	5	1	-	-	-
Female	2	4	8	9	4	15	8	10	2	3	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	1	4	1	2	-	-	-	-
Male	-	-	-	2	1	2	1	2	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Delaware												
All races, both sexes	14	16	26	40	57	82	128	131	87	26	11	-
Male	7	12	17	27	34	47	55	65	20	10	1	-
Female	7	4	9	13	23	35	73	66	67	16	10	-
White, both sexes	9	12	19	32	47	76	116	120	76	21	9	-
Male	4	9	13	24	27	44	49	59	19	9	1	-
Female	5	3	6	8	20	32	67	61	57	12	8	-
Black, both sexes	5	4	6	6	9	5	12	11	10	5	2	-
Male	3	3	3	3	6	3	6	6	1	1	-	-
Female	2	1	3	3	3	2	6	5	9	4	2	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	1	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	9	9	15	14	25	33	49	60	42	19	9	-
Male	7	4	6	10	15	20	22	19	19	5	1	-
Female	2	5	9	4	10	13	27	41	23	14	8	-
White, both sexes	1	2	1	2	2	10	14	17	16	7	2	-
Male	1	2	1	1	1	8	6	9	7	4	-	-
Female	-	-	-	1	1	2	8	8	9	3	2	-
Black, both sexes	8	7	14	12	23	23	35	43	25	12	7	-
Male	6	2	5	9	14	12	16	10	11	1	1	-
Female	2	5	9	3	9	11	19	33	14	11	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Florida												
All races, both sexes	420	651	788	1,036	1,505	2,484	3,634	3,747	2,859	1,185	259	-
Male	333	481	565	723	970	1,531	1,967	1,733	1,144	337	61	-
Female	87	170	223	313	535	953	1,667	2,014	1,715	848	198	-
White, both sexes	332	536	670	882	1,318	2,283	3,372	3,569	2,712	1,128	234	-
Male	269	420	492	619	855	1,428	1,848	1,673	1,104	325	58	-
Female	63	116	178	263	463	855	1,524	1,896	1,608	803	176	-
Black, both sexes	83	110	112	144	183	192	244	168	140	53	25	-
Male	59	59	70	96	112	95	109	57	38	11	3	-
Female	24	51	42	48	71	97	135	111	102	42	22	-
American Indian 1/, both sexes	1	2	1	3	-	-	3	1	1	1	-	-
Male	1	-	1	2	-	-	2	-	-	-	-	-
Female	-	2	-	1	-	-	1	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	4	3	5	7	4	9	15	9	6	3	-	-
Male	4	2	2	6	3	8	8	3	2	1	-	-
Female	-	1	3	1	1	1	7	6	4	2	-	-
Georgia												
All races, both sexes	164	215	278	325	392	532	710	616	429	163	27	1
Male	110	165	199	216	256	288	365	267	124	36	3	1
Female	54	50	79	109	136	244	345	349	305	127	24	-
White, both sexes	95	143	198	239	302	416	597	518	356	135	21	1
Male	72	113	156	169	208	236	319	237	109	34	1	1
Female	23	30	42	70	94	180	278	281	247	101	20	-
Black, both sexes	69	70	77	82	87	106	108	95	71	28	5	-
Male	38	50	42	44	46	46	44	30	15	2	1	-
Female	31	20	35	38	41	60	64	65	56	26	4	-
American Indian 1/, both sexes	-	-	1	-	1	3	1	-	1	-	-	-
Male	-	-	-	-	1	3	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	4	2	7	4	3	1	-	1	-
Male	-	2	1	3	1	3	1	-	-	-	1	-
Female	-	-	1	1	1	4	3	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hawaii												
All races, both sexes	30	38	40	56	56	90	112	109	75	31	9	-
Male	24	30	32	42	31	61	76	62	39	18	3	-
Female	6	8	8	14	25	29	36	47	36	13	6	-
White, both sexes	12	14	14	14	14	34	28	35	22	11	3	-
Male	9	11	13	7	5	24	21	19	11	6	1	-
Female	3	3	1	7	9	10	7	16	11	5	2	-
Black, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	24	26	42	41	55	83	74	53	20	6	-
Male	15	19	19	35	25	36	55	43	28	12	2	-
Female	3	5	7	7	16	19	28	31	25	8	4	-
Idaho												
All races, both sexes	8	14	25	35	49	92	130	141	117	40	5	-
Male	4	13	21	23	33	64	78	77	51	14	2	-
Female	4	1	4	12	16	28	52	64	66	26	3	-
White, both sexes	8	14	25	35	49	91	129	139	117	39	5	-
Male	4	13	21	23	33	64	78	75	51	13	2	-
Female	4	1	4	12	16	27	51	64	66	26	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Illinois												
All races, both sexes	247	299	380	494	677	1,109	1,606	1,716	1,422	571	131	-
Male	199	231	257	325	414	656	790	754	439	130	15	-
Female	48	68	123	169	263	453	816	962	983	441	116	-
White, both sexes	191	229	292	392	565	955	1,431	1,576	1,323	519	116	-
Male	156	179	203	257	360	569	717	694	409	117	13	-
Female	35	50	89	135	205	386	714	882	914	402	103	-
Black, both sexes	54	66	80	89	101	142	164	123	93	46	14	-
Male	42	49	48	58	46	80	66	52	27	10	2	-
Female	12	17	32	31	55	62	98	71	66	36	12	-
American Indian 1/, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	7	12	11	11	11	16	6	6	1	-
Male	1	3	6	9	8	7	7	7	3	3	-	-
Female	1	1	1	3	3	4	4	9	3	3	1	-
Indiana												
All races, both sexes	138	193	254	331	460	690	903	880	631	250	66	-
Male	103	135	177	224	293	412	485	383	204	58	9	-
Female	35	58	77	107	167	278	418	497	427	192	57	-
White, both sexes	118	156	219	284	426	642	834	835	599	239	63	-
Male	85	114	150	195	270	386	451	364	197	56	9	-
Female	33	42	69	89	156	256	383	471	402	183	54	-
Black, both sexes	20	35	33	43	32	47	67	42	32	11	3	-
Male	18	20	25	26	21	26	32	18	7	2	-	-
Female	2	15	8	17	11	21	35	24	25	9	3	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	3	1	1	2	3	-	-	-	-
Male	-	1	2	3	1	-	2	1	-	-	-	-
Female	-	1	-	-	-	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Iowa												
All races, both sexes	49	67	96	130	185	312	454	566	476	211	59	-
Male	31	41	64	95	120	181	248	269	158	49	6	-
Female	18	26	32	35	65	131	206	297	318	162	53	-
White, both sexes	46	63	94	125	183	306	449	565	475	211	59	-
Male	29	39	64	91	119	175	245	269	158	49	6	-
Female	17	24	30	34	64	131	204	296	317	162	53	-
Black, both sexes	2	4	2	4	1	4	3	1	1	-	-	-
Male	1	2	-	3	-	4	2	-	-	-	-	-
Female	1	2	2	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	2	1	-	-	-	-	-
Male	1	-	-	1	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	61	91	95	124	161	227	328	377	280	107	32	-
Male	51	72	69	87	97	139	184	179	108	26	4	-
Female	10	19	26	37	64	88	144	198	172	81	28	-
White, both sexes	56	81	89	116	150	208	315	366	276	103	31	-
Male	46	68	65	84	90	131	175	173	107	24	4	-
Female	10	13	24	32	60	77	140	193	169	79	27	-
Black, both sexes	4	10	5	4	9	14	10	9	4	4	1	-
Male	4	4	3	1	6	5	6	5	1	2	-	-
Female	-	6	2	3	3	9	4	4	3	2	1	-
American Indian 1/, both sexes	1	-	-	3	1	3	3	1	-	-	-	-
Male	1	-	-	1	1	3	3	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	2	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Kentucky												
All races, both sexes	105	95	144	187	282	359	460	448	310	116	25	-
Male	82	64	91	125	164	220	222	177	90	24	3	-
Female	23	31	53	62	118	139	238	271	220	92	22	-
White, both sexes	93	88	136	171	260	339	442	429	301	106	25	-
Male	72	60	87	118	151	210	211	173	88	23	3	-
Female	21	28	49	53	109	129	231	256	213	83	22	-
Black, both sexes	11	7	7	15	20	19	18	19	8	10	-	-
Male	10	4	4	7	12	9	11	4	1	1	-	-
Female	1	3	3	8	8	10	7	15	7	9	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	82	117	147	180	225	286	389	349	220	81	25	-
Male	60	85	96	125	145	174	207	151	73	17	4	-
Female	22	32	51	55	80	112	182	198	147	64	21	-
White, both sexes	47	78	101	116	175	221	309	288	184	64	17	-
Male	38	58	70	88	117	137	168	123	62	15	2	-
Female	9	20	31	28	58	84	141	165	122	49	15	-
Black, both sexes	35	38	45	64	48	65	78	59	36	17	8	-
Male	22	26	25	37	28	37	38	27	11	2	2	-
Female	13	12	20	27	20	28	40	32	25	15	6	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	1	2	-	-	-	-
Male	-	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Maine												
All races, both sexes	22	36	48	69	73	119	187	177	142	43	14	-
Male	19	27	38	45	47	73	113	73	53	12	1	-
Female	3	9	10	24	26	46	74	104	89	31	13	-
White, both sexes	19	36	48	69	73	119	187	177	142	42	14	-
Male	16	27	38	45	47	73	113	73	53	11	1	-
Female	3	9	10	24	26	46	74	104	89	31	13	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	71	100	159	195	259	388	520	547	404	148	32	-
Male	43	60	105	117	159	228	246	239	132	47	2	-
Female	28	40	54	78	100	160	274	308	272	101	30	-
White, both sexes	37	61	104	122	170	291	420	470	349	126	25	-
Male	27	37	73	77	111	166	212	214	116	42	1	-
Female	10	24	31	45	59	125	208	256	233	84	24	-
Black, both sexes	33	38	50	67	87	92	92	71	50	20	6	-
Male	16	22	28	34	47	58	30	23	14	5	1	-
Female	17	16	22	33	40	34	62	48	36	15	5	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	6	2	5	7	6	4	2	1	-
Male	-	1	3	6	1	4	4	2	2	-	-	-
Female	1	-	1	-	1	1	3	4	2	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Massachusetts												
All races, both sexes	101	124	177	177	323	504	794	963	727	350	85	-
Male	80	101	117	134	201	301	428	395	239	69	10	-
Female	21	23	60	43	122	203	366	568	488	281	75	-
White, both sexes	90	118	165	165	299	491	769	943	718	344	84	-
Male	71	95	109	127	194	292	416	386	237	66	10	-
Female	19	23	56	38	105	199	353	557	481	278	74	-
Black, both sexes	10	6	9	10	18	7	21	14	7	4	-	-
Male	8	6	6	6	5	4	11	7	2	2	-	-
Female	2	-	3	4	13	3	10	7	5	2	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	1	-	-	1	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	6	6	3	6	2	1	1	-
Male	-	-	1	1	2	5	-	2	-	1	-	-
Female	-	-	1	1	4	1	3	4	2	-	1	-
Michigan												
All races, both sexes	188	277	330	420	644	1,081	1,546	1,649	1,220	501	110	-
Male	139	198	221	275	388	606	777	721	427	115	20	-
Female	49	79	109	145	256	475	769	928	793	386	90	-
White, both sexes	160	219	282	368	568	964	1,388	1,526	1,139	468	101	-
Male	120	158	196	247	352	560	712	668	400	105	18	-
Female	40	61	86	121	216	404	676	858	739	363	83	-
Black, both sexes	25	57	46	47	68	108	147	116	75	33	9	-
Male	17	39	24	26	32	42	62	52	25	10	2	-
Female	8	18	22	21	36	66	85	64	50	23	7	-
American Indian 1/, both sexes	1	-	2	4	4	7	5	5	3	-	-	-
Male	-	-	1	1	2	3	1	1	1	-	-	-
Female	1	-	1	3	2	4	4	4	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	4	2	6	2	3	-	-	-
Male	2	1	-	1	2	1	2	-	1	-	-	-
Female	-	-	-	-	2	1	4	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Minnesota												
All races, both sexes	111	118	129	140	211	299	481	507	424	164	44	-
Male	97	93	93	105	149	198	300	257	173	37	5	-
Female	14	25	36	35	62	101	181	250	251	127	39	-
White, both sexes	97	99	113	131	202	292	472	503	424	164	44	-
Male	87	79	82	99	143	196	294	255	173	37	5	-
Female	10	20	31	32	59	96	178	248	251	127	39	-
Black, both sexes	8	12	8	3	5	4	4	3	-	-	-	-
Male	6	8	5	-	3	2	3	1	-	-	-	-
Female	2	4	3	3	2	2	1	2	-	-	-	-
American Indian 1/, both sexes	3	3	5	5	1	2	3	1	-	-	-	-
Male	1	2	3	5	-	-	3	1	-	-	-	-
Female	2	1	2	-	1	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	3	1	3	1	2	-	-	-	-	-
Male	3	4	3	1	3	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Mississippi												
All races, both sexes	103	102	116	151	186	247	239	250	188	71	11	-
Male	82	70	68	102	124	137	114	97	50	15	2	-
Female	21	32	48	49	62	110	125	153	138	56	9	-
White, both sexes	59	70	83	115	130	186	197	205	144	57	6	-
Male	46	49	47	80	87	109	96	82	40	12	1	-
Female	13	21	36	35	43	77	101	123	104	45	5	-
Black, both sexes	44	30	33	35	56	60	42	45	44	14	5	-
Male	36	19	21	21	37	28	18	15	10	3	1	-
Female	8	11	12	14	19	32	24	30	34	11	4	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Missouri												
All races, both sexes	160	223	235	298	401	550	835	986	749	312	69	-
Male	115	163	154	193	250	308	425	404	239	62	11	-
Female	45	60	81	105	151	242	410	582	510	250	58	-
White, both sexes	129	184	200	252	350	502	771	936	708	290	65	-
Male	94	136	131	167	218	283	404	389	230	59	10	-
Female	35	48	69	85	132	219	367	547	478	231	55	-
Black, both sexes	29	38	34	45	47	42	61	47	39	21	4	-
Male	19	26	23	26	29	20	20	15	9	3	1	-
Female	10	12	11	19	18	22	41	32	30	18	3	-
American Indian 1/, both sexes	1	-	1	-	2	2	3	2	-	-	-	-
Male	1	-	-	-	2	2	1	-	-	-	-	-
Female	-	-	1	-	-	-	2	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	2	4	-	1	2	1	-	-
Male	1	1	-	-	1	3	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	1	2	1	-	-
Montana												
All races, both sexes	14	28	29	35	48	74	83	113	71	31	3	-
Male	10	22	22	26	32	51	43	57	36	8	1	-
Female	4	6	7	9	16	23	40	56	35	23	2	-
White, both sexes	12	26	28	31	44	72	83	110	68	31	3	-
Male	8	20	21	23	29	50	43	57	36	8	1	-
Female	4	6	7	8	15	22	40	53	32	23	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	1	4	4	2	-	3	2	-	-	-
Male	2	1	1	3	3	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Nebraska												
All races, both sexes	30	39	50	57	75	124	236	227	196	102	24	-
Male	22	31	35	46	48	73	147	96	66	30	5	-
Female	8	8	15	11	27	51	89	131	130	72	19	-
White, both sexes	27	35	44	56	73	120	232	221	192	98	24	-
Male	19	28	31	46	47	72	145	93	65	30	5	-
Female	8	7	13	10	26	48	87	128	127	68	19	-
Black, both sexes	3	4	2	-	2	4	2	3	4	3	-	-
Male	3	3	2	-	1	1	1	1	1	-	-	-
Female	-	1	-	-	1	3	1	2	3	3	-	-
American Indian 1/, both sexes	-	-	4	1	-	-	1	-	-	1	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	25	51	45	73	112	145	172	146	107	28	8	-
Male	21	34	37	52	78	90	99	75	36	9	1	-
Female	4	17	8	21	34	55	73	71	71	19	7	-
White, both sexes	20	40	37	62	103	129	160	141	102	25	6	-
Male	18	30	32	43	72	82	93	73	36	7	1	-
Female	2	10	5	19	31	47	67	68	66	18	5	-
Black, both sexes	4	7	5	7	2	8	3	3	4	1	1	-
Male	2	2	2	7	2	3	1	1	-	1	-	-
Female	2	5	3	-	-	5	2	2	4	-	1	-
American Indian 1/, both sexes	-	2	1	1	2	1	1	-	-	1	1	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	2	-	1	1	1	1	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	1	2	2	3	5	7	8	2	1	1	-	-
Male	1	2	2	2	3	5	5	1	-	1	-	-
Female	-	-	-	1	2	2	3	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Hampshire												
All races, both sexes	21	30	26	43	72	117	163	169	149	63	13	-
Male	17	24	22	34	44	70	75	72	48	17	3	-
Female	4	6	4	9	28	47	88	97	101	46	10	-
White, both sexes	21	29	25	43	72	115	162	168	149	63	13	-
Male	17	23	21	34	44	69	75	71	48	17	3	-
Female	4	6	4	9	28	46	87	97	101	46	10	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New Jersey												
All races, both sexes	129	181	252	333	537	945	1,402	1,610	1,217	557	132	-
Male	96	133	168	221	335	543	688	678	381	126	13	-
Female	33	48	84	112	202	402	714	932	836	431	119	-
White, both sexes	108	142	197	279	448	859	1,300	1,520	1,148	533	122	-
Male	84	102	144	190	284	495	645	647	363	123	12	-
Female	24	40	53	89	164	364	655	873	785	410	110	-
Black, both sexes	18	35	46	42	79	76	85	76	59	20	7	-
Male	10	28	17	23	46	42	38	27	15	2	-	-
Female	8	7	29	19	33	34	47	49	44	18	7	-
American Indian 1/, both sexes	-	1	-	1	-	1	2	-	2	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	9	11	10	9	15	14	8	4	3	-
Male	2	2	7	8	5	5	4	4	2	1	1	-
Female	1	1	2	3	5	4	11	10	6	3	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Mexico												
All races, both sexes	19	29	34	55	96	138	188	220	178	68	16	-
Male	15	23	23	42	68	78	103	100	69	14	-	-
Female	4	6	11	13	28	60	85	120	109	54	16	-
White, both sexes	17	27	32	49	89	130	182	214	174	65	15	-
Male	13	22	21	36	65	74	99	99	66	13	-	-
Female	4	5	11	13	24	56	83	115	108	52	15	-
Black, both sexes	1	1	-	4	3	3	1	2	-	2	-	-
Male	1	-	-	4	1	1	-	-	-	-	-	-
Female	-	1	-	-	2	2	1	2	-	2	-	-
American Indian 1/, both sexes	1	1	2	1	3	4	3	4	4	1	1	-
Male	1	1	2	1	2	3	2	1	3	1	-	-
Female	-	-	-	-	1	1	1	3	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	2	-	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
New York												
All races, both sexes	433	669	863	1,253	1,849	3,101	4,675	5,434	4,509	2,065	549	-
Male	287	487	582	776	1,051	1,691	2,167	2,140	1,380	434	88	-
Female	146	182	281	477	798	1,410	2,508	3,294	3,129	1,631	461	-
White, both sexes	332	515	657	948	1,471	2,612	4,021	4,842	4,101	1,851	492	-
Male	243	392	454	612	864	1,463	1,927	1,935	1,267	394	83	-
Female	89	123	203	336	607	1,149	2,094	2,907	2,834	1,457	409	-
Black, both sexes	85	130	174	265	319	412	528	481	324	181	53	-
Male	34	76	102	140	155	185	187	160	74	32	5	-
Female	51	54	72	125	164	227	341	321	250	149	48	-
American Indian 1/, both sexes	-	-	2	1	4	6	2	4	-	2	-	-
Male	-	-	1	-	-	2	-	1	-	-	-	-
Female	-	-	1	1	4	4	2	3	-	2	-	-
Asian or Pacific Islander 2/, both sexes	16	24	30	39	55	71	124	107	84	31	4	-
Male	10	19	25	24	32	41	53	44	39	8	-	-
Female	6	5	5	15	23	30	71	63	45	23	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
North Carolina												
All races, both sexes	264	377	469	514	620	785	943	950	641	223	56	-
Male	210	296	355	352	435	474	515	403	185	47	7	-
Female	54	81	114	162	185	311	428	547	456	176	49	-
White, both sexes	183	269	349	394	485	625	780	799	538	193	43	-
Male	150	216	268	267	353	392	451	349	165	42	4	-
Female	33	53	81	127	132	233	329	450	373	151	39	-
Black, both sexes	76	95	112	110	126	151	146	144	97	27	13	-
Male	55	68	81	77	76	78	59	53	17	5	3	-
Female	21	27	31	33	50	73	87	91	80	22	10	-
American Indian 1/, both sexes	5	11	5	8	7	7	15	4	6	2	-	-
Male	5	10	5	7	6	3	5	-	3	-	-	-
Female	-	1	-	1	1	4	10	4	3	2	-	-
Asian or Pacific Islander 2/, both sexes	-	2	3	2	2	2	2	3	-	1	-	-
Male	-	2	1	1	-	1	-	1	-	-	-	-
Female	-	-	2	1	2	1	2	2	-	1	-	-
North Dakota												
All races, both sexes	10	15	14	17	39	51	97	103	112	39	8	-
Male	9	12	11	12	23	35	57	44	47	16	3	-
Female	1	3	3	5	16	16	40	59	65	23	5	-
White, both sexes	9	13	12	15	37	49	96	102	111	39	8	-
Male	9	10	10	10	23	34	56	43	47	16	3	-
Female	-	3	2	5	14	15	40	59	64	23	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	2	2	2	2	1	1	1	-	-	-
Male	-	2	1	2	-	1	1	1	-	-	-	-
Female	1	-	1	-	2	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Ohio												
All races, both sexes	342	472	514	657	951	1,484	2,154	2,138	1,558	606	118	-
Male	260	345	353	417	566	848	1,115	914	520	146	16	-
Female	82	127	161	240	385	636	1,039	1,224	1,038	460	102	-
White, both sexes	272	380	430	560	813	1,338	1,987	2,025	1,474	563	103	-
Male	212	282	303	359	483	767	1,035	874	494	133	14	-
Female	60	98	127	201	330	571	952	1,151	980	430	89	-
Black, both sexes	68	90	82	92	134	138	162	111	82	40	15	-
Male	48	61	48	53	81	77	79	40	25	11	2	-
Female	20	29	34	39	53	61	83	71	57	29	13	-
American Indian 1/, both sexes	1	1	1	1	-	2	-	-	-	-	-	-
Male	-	1	1	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	4	4	6	5	2	2	3	-	-
Male	-	1	1	4	2	3	1	-	1	2	-	-
Female	1	-	-	-	2	3	4	2	1	1	-	-
Oklahoma												
All races, both sexes	88	128	169	222	309	408	543	563	387	168	36	-
Male	62	106	118	159	195	259	299	235	131	32	6	-
Female	26	22	51	63	114	149	244	328	256	136	30	-
White, both sexes	69	111	141	181	258	366	504	523	357	158	34	-
Male	48	93	101	130	161	235	280	216	122	31	5	-
Female	21	18	40	51	97	131	224	307	235	127	29	-
Black, both sexes	12	8	10	21	18	18	18	16	17	6	1	-
Male	10	4	5	14	11	11	8	6	6	1	1	-
Female	2	4	5	7	7	7	10	10	11	5	-	-
American Indian 1/, both sexes	7	8	17	17	31	24	19	24	12	4	1	-
Male	4	8	11	12	22	13	10	13	3	-	-	-
Female	3	-	6	5	9	11	9	11	9	4	1	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	2	-	2	-	1	-	-	-
Male	-	1	1	3	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Oregon												
All races, both sexes	94	122	132	179	198	288	417	464	337	126	13	-
Male	80	97	97	132	131	197	250	262	139	41	3	-
Female	14	25	35	47	67	91	167	202	198	85	10	-
White, both sexes	90	119	120	172	192	284	409	452	333	124	13	-
Male	76	95	90	126	128	194	248	257	139	40	3	-
Female	14	24	30	46	64	90	161	195	194	84	10	-
Black, both sexes	3	3	5	-	2	1	2	3	1	2	-	-
Male	3	2	2	-	1	1	1	-	-	1	-	-
Female	-	1	3	-	1	-	1	3	1	1	-	-
American Indian 1/, both sexes	1	-	4	3	1	-	-	5	1	-	-	-
Male	1	-	2	2	1	-	-	3	-	-	-	-
Female	-	-	2	1	-	-	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	4	3	3	6	4	2	-	-	-
Male	-	-	3	4	1	2	1	2	-	-	-	-
Female	-	-	-	-	2	1	5	2	2	-	-	-
Pennsylvania												
All races, both sexes	161	269	388	515	831	1,433	2,074	2,297	1,662	664	122	-
Male	119	185	275	343	518	829	1,083	981	538	166	16	-
Female	42	84	113	172	313	604	991	1,316	1,124	498	106	-
White, both sexes	142	228	347	459	753	1,339	1,953	2,197	1,593	637	110	-
Male	109	158	249	313	479	781	1,023	949	521	163	14	-
Female	33	70	98	146	274	558	930	1,248	1,072	474	96	-
Black, both sexes	19	38	39	53	75	89	117	94	67	27	12	-
Male	10	25	25	28	38	44	59	28	17	3	2	-
Female	9	13	14	25	37	45	58	66	50	24	10	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	3	2	5	4	6	2	-	-	-
Male	-	1	1	2	-	4	1	4	-	-	-	-
Female	-	1	1	1	2	1	3	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Rhode Island												
All races, both sexes	12	17	29	29	55	104	154	224	189	84	20	-
Male	9	11	15	19	30	65	77	93	58	16	5	-
Female	3	6	14	10	25	39	77	131	131	68	15	-
White, both sexes	12	17	28	27	53	100	151	221	188	83	20	-
Male	9	11	15	19	28	63	76	92	58	15	5	-
Female	3	6	13	8	25	37	75	129	130	68	15	-
Black, both sexes	-	-	1	2	2	3	2	3	1	-	-	-
Male	-	-	-	-	2	2	-	1	-	-	-	-
Female	-	-	1	2	-	1	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	101	155	161	203	258	368	438	365	212	77	15	-
Male	76	115	108	139	181	246	241	166	85	17	3	-
Female	25	40	53	64	77	122	197	199	127	60	12	-
White, both sexes	67	109	125	150	190	298	354	303	185	68	6	-
Male	51	78	88	110	139	205	202	142	77	14	3	-
Female	16	31	37	40	51	93	152	161	108	54	3	-
Black, both sexes	34	43	36	53	68	69	84	62	27	9	9	-
Male	25	34	20	29	42	40	39	24	8	3	-	-
Female	9	9	16	24	26	29	45	38	19	6	9	-
American Indian 1/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
South Dakota												
All races, both sexes	18	28	29	27	43	76	112	122	114	54	16	-
Male	14	22	18	20	25	50	62	57	40	12	3	-
Female	4	6	11	7	18	26	50	65	74	42	13	-
White, both sexes	16	25	26	23	40	73	109	121	114	54	16	-
Male	12	20	17	18	24	47	61	57	40	12	3	-
Female	4	5	9	5	16	26	48	64	74	42	13	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	3	4	3	1	3	1	-	-	-	-
Male	2	2	1	2	1	1	1	-	-	-	-	-
Female	-	1	2	2	2	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	121	201	266	295	454	570	772	756	522	195	42	-
Male	85	163	177	193	286	346	372	311	165	38	8	-
Female	36	38	89	102	168	224	400	445	357	157	34	-
White, both sexes	93	167	225	244	390	497	698	705	475	179	32	-
Male	69	137	151	161	252	301	348	298	152	36	8	-
Female	24	30	74	83	138	196	350	407	323	143	24	-
Black, both sexes	28	33	40	49	63	73	72	49	45	16	10	-
Male	16	25	25	31	34	45	23	13	13	2	-	-
Female	12	8	15	18	29	28	49	36	32	14	10	-
American Indian 1/, both sexes	-	1	1	1	-	-	1	-	-	-	-	-
Male	-	1	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	2	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Texas												
All races, both sexes	299	501	582	758	1,021	1,600	2,055	1,930	1,386	541	121	1
Male	225	348	390	486	633	932	1,052	844	526	151	21	-
Female	74	153	192	272	388	668	1,003	1,086	860	390	100	1
White, both sexes	244	406	480	639	884	1,444	1,882	1,815	1,303	487	107	1
Male	185	289	335	420	556	849	970	804	494	141	19	-
Female	59	117	145	219	328	595	912	1,011	809	346	88	1
Black, both sexes	53	85	94	104	125	146	156	102	75	54	14	-
Male	38	50	51	58	71	78	76	36	28	10	2	-
Female	15	35	43	46	54	68	80	66	47	44	12	-
American Indian 1/, both sexes	-	2	1	3	3	1	4	3	2	-	-	-
Male	-	2	-	2	2	1	1	-	-	-	-	-
Female	-	-	1	1	1	-	3	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	8	7	12	9	9	13	10	6	-	-	-
Male	2	7	4	6	4	4	5	4	4	-	-	-
Female	-	1	3	6	5	5	8	6	2	-	-	-
Utah												
All races, both sexes	30	38	46	62	64	95	137	155	107	33	6	-
Male	27	33	34	56	42	67	86	89	48	13	1	-
Female	3	5	12	6	22	28	51	66	59	20	5	-
White, both sexes	27	38	46	60	63	92	136	155	106	33	6	-
Male	24	33	34	54	41	65	86	89	47	13	1	-
Female	3	5	12	6	22	27	50	66	59	20	5	-
Black, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	1	-	-	1	-	-	-
Male	1	-	-	1	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Vermont												
All races, both sexes	2	8	13	15	33	51	61	74	54	28	7	-
Male	-	8	11	12	20	28	33	33	25	5	2	-
Female	2	-	2	3	13	23	28	41	29	23	5	-
White, both sexes	2	8	13	15	32	51	61	74	54	28	7	-
Male	-	8	11	12	19	28	33	33	25	5	2	-
Female	2	-	2	3	13	23	28	41	29	23	5	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	89	202	239	263	366	515	670	673	412	180	28	-
Male	67	156	187	184	238	345	357	286	139	32	3	-
Female	22	46	52	79	128	170	313	387	273	148	25	-
White, both sexes	65	146	184	202	296	428	562	576	360	151	23	-
Male	50	116	147	143	200	291	310	246	124	30	2	-
Female	15	30	37	59	96	137	252	330	236	121	21	-
Black, both sexes	22	49	52	58	66	78	103	90	51	29	5	-
Male	15	34	38	40	35	49	46	39	14	2	1	-
Female	7	15	14	18	31	29	57	51	37	27	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	7	3	3	4	9	3	7	1	-	-	-
Male	2	6	2	1	3	5	1	1	1	-	-	-
Female	-	1	1	2	1	4	2	6	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Washington												
All races, both sexes	76	109	177	214	301	477	630	725	581	227	39	-
Male	61	83	124	145	204	289	366	355	236	70	11	-
Female	15	26	53	69	97	188	264	370	345	157	28	-
White, both sexes	68	98	156	194	282	445	605	704	564	220	38	-
Male	55	76	111	132	192	274	354	345	227	66	11	-
Female	13	22	45	62	90	171	251	359	337	154	27	-
Black, both sexes	4	6	8	9	3	12	6	4	5	1	-	-
Male	3	3	4	6	2	4	3	-	2	-	-	-
Female	1	3	4	3	1	8	3	4	3	1	-	-
American Indian 1/, both sexes	1	2	6	2	7	5	7	3	3	-	-	-
Male	-	1	2	1	4	2	3	1	2	-	-	-
Female	1	1	4	1	3	3	4	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	7	9	9	15	12	14	9	6	1	-
Male	3	3	7	6	6	9	6	9	5	4	-	-
Female	-	-	-	3	3	6	6	5	4	2	1	-
West Virginia												
All races, both sexes	80	78	101	150	180	264	341	290	211	83	13	-
Male	63	56	72	106	105	161	180	134	69	15	-	-
Female	17	22	29	44	75	103	161	156	142	68	13	-
White, both sexes	76	77	96	148	176	258	327	284	204	81	11	-
Male	60	55	70	104	102	155	174	133	66	15	-	-
Female	16	22	26	44	74	103	153	151	138	66	11	-
Black, both sexes	4	1	5	2	4	6	13	6	7	2	2	-
Male	3	1	2	2	3	6	6	1	3	-	-	-
Female	1	-	3	-	1	-	7	5	4	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Wisconsin												
All races, both sexes	67	74	118	146	289	434	611	690	543	259	67	-
Male	48	55	83	113	197	262	338	349	201	54	9	-
Female	19	19	35	33	92	172	273	341	342	205	58	-
White, both sexes	57	63	105	138	275	412	597	678	537	254	66	-
Male	45	48	74	108	188	250	331	343	200	52	9	-
Female	12	15	31	30	87	162	266	335	337	202	57	-
Black, both sexes	8	7	11	7	14	19	12	11	6	3	-	-
Male	2	4	7	4	9	10	7	6	1	1	-	-
Female	6	3	4	3	5	9	5	5	5	2	-	-
American Indian 1/, both sexes	2	3	1	1	-	2	2	-	-	2	-	-
Male	1	2	1	1	-	2	-	-	-	1	-	-
Female	1	1	-	-	-	-	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	-	1	-	-	1	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	1	-
Wyoming												
All races, both sexes	7	9	15	22	34	47	34	44	33	18	1	-
Male	7	6	12	15	24	35	27	20	13	4	-	-
Female	-	3	3	7	10	12	7	24	20	14	1	-
White, both sexes	7	9	13	22	34	47	34	44	33	18	1	-
Male	7	6	10	15	24	35	27	20	13	4	-	-
Female	-	3	3	7	10	12	7	24	20	14	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
United States												
All races, both sexes	170,248	320	154	85	146	311	562	803	1,100	1,566	2,569	4,039
Male	75,221	169	84	46	84	200	355	522	695	977	1,563	2,546
Female	95,027	151	70	39	62	111	207	281	405	589	1,006	1,493
White, both sexes	147,421	192	105	59	99	204	383	517	683	993	1,794	2,849
Male	64,243	100	57	33	59	134	254	343	444	624	1,104	1,823
Female	83,178	92	48	26	40	70	129	174	239	369	690	1,026
Black, both sexes	19,805	109	42	20	40	98	155	255	382	520	702	1,083
Male	9,467	59	25	9	24	60	86	157	229	316	410	653
Female	10,338	50	17	11	16	38	69	98	153	204	292	430
American Indian 1/, both sexes	686	7	-	-	2	4	8	7	9	27	31	42
Male	356	3	-	-	-	1	3	5	4	19	22	29
Female	330	4	-	-	2	3	5	2	5	8	9	13
Asian or Pacific Islander 2/, both sexes	2,336	12	7	6	5	5	16	24	26	26	42	65
Male	1,155	7	2	4	1	5	12	17	18	18	27	41
Female	1,181	5	5	2	4	-	4	7	8	8	15	24
Alabama												
All races, both sexes	6,057	8	4	5	4	8	16	39	53	77	114	194
Male	2,757	4	3	1	1	6	10	28	33	48	69	119
Female	3,300	4	1	4	3	2	6	11	20	29	45	75
White, both sexes	4,585	6	2	2	3	6	11	22	33	37	74	116
Male	2,061	4	2	-	-	4	9	17	19	24	47	70
Female	2,524	2	-	2	3	2	2	5	14	13	27	46
Black, both sexes	1,451	2	2	3	1	2	5	15	19	39	40	78
Male	686	-	1	1	1	2	1	9	13	23	22	49
Female	765	2	1	2	-	-	4	6	6	16	18	29
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	2	1	-	-	-
Male	8	-	-	-	-	-	-	2	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Alaska												
All races, both sexes	241	1	-	-	1	-	1	2	1	5	3	6
Male	120	-	-	-	-	-	1	-	-	2	2	2
Female	121	1	-	-	1	-	-	2	1	3	1	4
White, both sexes	176	-	-	-	-	-	-	1	-	1	2	3
Male	91	-	-	-	-	-	-	-	-	-	1	1
Female	85	-	-	-	-	-	-	1	-	1	1	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	45	1	-	-	1	-	1	1	-	3	-	2
Male	18	-	-	-	-	-	1	-	-	1	-	-
Female	27	1	-	-	1	-	-	1	-	2	-	2
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	1	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	-
Arizona												
All races, both sexes	2,199	10	5	2	5	6	10	13	14	31	29	62
Male	1,088	7	3	1	2	3	8	11	12	25	21	37
Female	1,111	3	2	1	3	3	2	2	2	6	8	25
White, both sexes	2,073	8	5	2	4	3	8	9	11	23	23	56
Male	1,008	5	3	1	2	1	6	9	9	17	16	34
Female	1,065	3	2	1	2	2	2	-	2	6	7	22
Black, both sexes	57	1	-	-	-	3	-	2	1	3	2	4
Male	40	1	-	-	-	2	-	1	1	3	2	2
Female	17	-	-	-	-	1	-	1	-	-	-	2
American Indian 1/, both sexes	61	1	-	-	-	-	2	1	1	5	4	2
Male	36	1	-	-	-	-	2	1	1	5	3	1
Female	25	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	1	-	-	1	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Arkansas												
All races, both sexes	1,958	7	5	1	4	4	9	10	9	19	29	59
Male	875	4	3	1	2	1	5	5	9	10	14	38
Female	1,083	3	2	-	2	3	4	5	-	9	15	21
White, both sexes	1,689	4	3	1	2	4	5	6	6	10	16	38
Male	746	2	3	1	2	1	2	4	6	8	7	24
Female	943	2	-	-	-	3	3	2	-	2	9	14
Black, both sexes	263	3	2	-	2	-	4	4	3	7	13	21
Male	126	2	-	-	-	-	3	1	3	1	7	14
Female	137	1	2	-	2	-	1	3	-	6	6	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	2	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	13,462	27	14	8	19	21	48	71	114	153	269	374
Male	6,274	14	9	6	9	19	31	56	81	101	187	255
Female	7,188	13	5	2	10	2	17	15	33	52	82	119
White, both sexes	11,617	18	12	5	16	15	37	52	84	112	211	295
Male	5,387	10	8	4	8	14	24	40	66	71	150	208
Female	6,230	8	4	1	8	1	13	12	18	41	61	87
Black, both sexes	1,004	2	-	1	2	5	8	11	21	28	42	54
Male	492	1	-	1	1	4	5	9	10	22	28	29
Female	512	1	-	-	1	1	3	2	11	6	14	25
American Indian 1/, both sexes	72	-	-	-	-	-	-	1	-	4	2	7
Male	42	-	-	-	-	-	-	1	-	3	-	7
Female	30	-	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	769	7	2	2	1	1	3	7	9	9	14	18
Male	353	3	1	1	-	1	2	6	5	5	9	11
Female	416	4	1	1	1	-	1	1	4	4	5	7

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Colorado												
All races, both sexes	1,864	2	2	2	3	4	14	8	9	11	28	35
Male	765	-	1	2	2	1	8	3	5	8	18	23
Female	1,099	2	1	-	1	3	6	5	4	3	10	12
White, both sexes	1,774	2	2	1	3	3	12	8	8	10	26	31
Male	725	-	1	1	2	1	8	3	4	7	17	20
Female	1,049	2	1	-	1	2	4	5	4	3	9	11
Black, both sexes	53	-	-	-	-	1	2	-	1	1	-	3
Male	23	-	-	-	-	-	-	-	1	1	-	3
Female	30	-	-	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	24	-	-	1	-	-	-	-	-	-	-	1
Male	10	-	-	1	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Connecticut												
All races, both sexes	2,575	2	1	1	-	5	6	14	7	18	24	53
Male	1,119	2	-	1	-	2	5	12	5	13	12	30
Female	1,456	-	1	-	-	3	1	2	2	5	12	23
White, both sexes	2,427	2	1	1	-	5	5	9	4	14	17	44
Male	1,048	2	-	1	-	2	4	7	4	10	10	24
Female	1,379	-	1	-	-	3	1	2	-	4	7	20
Black, both sexes	138	-	-	-	-	-	1	5	3	4	7	9
Male	67	-	-	-	-	-	1	5	1	3	2	6
Female	71	-	-	-	-	-	-	-	2	1	5	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Delaware												
All races, both sexes	460	2	-	-	-	1	1	2	6	7	8	11
Male	208	1	-	-	-	-	1	1	3	6	6	5
Female	252	1	-	-	-	1	-	1	3	1	2	6
White, both sexes	384	-	-	-	-	-	1	2	3	3	3	8
Male	176	-	-	-	-	-	1	1	2	3	2	5
Female	208	-	-	-	-	-	-	1	1	-	1	3
Black, both sexes	73	1	-	-	-	1	-	-	3	4	5	3
Male	31	-	-	-	-	-	-	-	1	3	4	-
Female	42	1	-	-	-	1	-	-	2	1	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	206	1	1	-	1	2	2	3	-	3	5	5
Male	84	-	1	-	-	1	-	2	-	1	4	2
Female	122	1	-	-	1	1	2	1	-	2	1	3
White, both sexes	48	-	-	-	-	-	1	-	-	1	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	157	1	1	-	1	2	1	3	-	2	5	5
Male	62	-	1	-	-	1	-	2	-	1	4	2
Female	95	1	-	-	1	1	1	1	-	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Florida												
All races, both sexes	8,831	19	13	6	10	13	30	39	48	74	116	205
Male	4,213	11	7	4	5	8	19	25	35	49	73	125
Female	4,618	8	6	2	5	5	11	14	13	25	43	80
White, both sexes	7,845	9	8	5	7	10	16	22	27	45	73	134
Male	3,709	4	4	3	5	7	13	15	20	30	49	81
Female	4,136	5	4	2	2	3	3	7	7	15	24	53
Black, both sexes	949	10	3	1	3	3	14	16	21	29	42	69
Male	487	7	2	1	-	1	6	10	15	19	23	43
Female	462	3	1	-	3	2	8	6	6	10	19	26
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	2	-	-	-	-	1	-	-	1	1
Male	14	-	1	-	-	-	-	-	-	-	1	-
Female	18	-	1	-	-	-	-	1	-	-	-	1
Georgia												
All races, both sexes	6,843	18	5	5	6	16	23	40	48	78	117	189
Male	3,167	12	3	3	2	11	8	26	32	53	69	124
Female	3,676	6	2	2	4	5	15	14	16	25	48	65
White, both sexes	5,003	9	4	3	4	6	9	19	21	32	65	107
Male	2,290	6	3	2	1	4	2	13	13	25	40	75
Female	2,713	3	1	1	3	2	7	6	8	7	25	32
Black, both sexes	1,790	9	1	2	2	9	12	21	25	46	51	82
Male	856	6	-	1	1	7	5	13	17	28	29	49
Female	934	3	1	1	1	2	7	8	8	18	22	33
American Indian 1/, both sexes	8	-	-	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	42	-	-	-	-	-	2	-	2	-	1	-
Male	16	-	-	-	-	-	1	-	2	-	-	-
Female	26	-	-	-	-	-	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Hawaii												
All races, both sexes	849	-	1	1	-	1	3	4	4	9	14	30
Male	458	-	-	1	-	1	2	3	3	5	8	16
Female	391	-	1	-	-	-	1	1	1	4	6	14
White, both sexes	201	-	-	-	-	-	-	-	1	3	2	6
Male	105	-	-	-	-	-	-	-	1	1	2	1
Female	96	-	-	-	-	-	-	-	-	2	-	5
Black, both sexes	5	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	641	-	1	1	-	1	2	4	3	5	12	23
Male	351	-	-	1	-	1	2	3	2	4	6	15
Female	290	-	1	-	-	-	-	1	1	1	6	8
Idaho												
All races, both sexes	699	1	2	1	-	-	3	4	5	2	9	14
Male	303	1	1	-	-	-	2	3	3	1	3	8
Female	396	-	1	1	-	-	1	1	2	1	6	6
White, both sexes	691	1	2	1	-	-	3	4	5	2	9	13
Male	300	1	1	-	-	-	2	3	3	1	3	8
Female	391	-	1	1	-	-	1	1	2	1	6	5
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Illinois												
All races, both sexes	7,699	10	4	1	4	21	26	49	53	72	95	160
Male	3,255	6	2	-	2	18	19	32	30	44	56	100
Female	4,444	4	2	1	2	3	7	17	23	28	39	60
White, both sexes	6,646	6	3	1	2	12	18	33	30	38	61	108
Male	2,764	3	1	-	1	10	13	23	19	22	35	70
Female	3,882	3	2	1	1	2	5	10	11	16	26	38
Black, both sexes	992	4	1	-	1	8	8	14	22	33	33	52
Male	461	3	1	-	1	7	6	8	11	22	20	30
Female	531	1	-	-	-	1	2	6	11	11	13	22
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	1	1	-	2	1	1	1	-
Male	29	-	-	-	-	1	-	1	-	-	1	-
Female	30	-	-	-	1	-	-	1	1	1	-	-
Indiana												
All races, both sexes	4,589	15	3	1	2	10	16	19	33	35	85	112
Male	1,960	6	2	-	-	7	9	9	17	22	53	70
Female	2,629	9	1	1	2	3	7	10	16	13	32	42
White, both sexes	4,241	7	3	1	2	7	13	17	24	29	74	91
Male	1,791	3	2	-	-	6	9	9	12	19	44	57
Female	2,450	4	1	1	2	1	4	8	12	10	30	34
Black, both sexes	344	8	-	-	-	3	3	2	9	6	11	21
Male	168	3	-	-	-	1	-	-	5	3	9	13
Female	176	5	-	-	-	2	3	2	4	3	2	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Iowa												
All races, both sexes	1,321	1	1	-	1	2	3	3	2	18	17	24
Male	548	1	-	-	-	1	1	3	1	11	12	13
Female	773	-	1	-	1	1	2	-	1	7	5	11
White, both sexes	1,292	-	1	-	1	2	3	3	2	14	16	24
Male	534	-	-	-	-	1	1	3	1	8	11	13
Female	758	-	1	-	1	1	2	-	1	6	5	11
Black, both sexes	27	1	-	-	-	-	-	-	-	3	1	-
Male	13	1	-	-	-	-	-	-	-	2	1	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,081	5	1	1	-	6	4	11	5	14	26	28
Male	835	3	1	-	-	2	4	5	4	5	15	18
Female	1,246	2	-	1	-	4	-	6	1	9	11	10
White, both sexes	1,953	3	-	1	-	5	4	9	3	12	20	23
Male	774	2	-	-	-	1	4	4	2	4	10	15
Female	1,179	1	-	1	-	4	-	5	1	8	10	8
Black, both sexes	108	2	1	-	-	-	-	2	1	2	5	4
Male	48	1	1	-	-	-	-	1	1	1	4	2
Female	60	1	-	-	-	-	-	1	-	1	1	2
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	1	-	-	1	-	-	-
Male	5	-	-	-	-	1	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Kentucky												
All races, both sexes	3,367	-	3	2	2	7	12	16	20	29	51	92
Male	1,429	-	1	1	2	3	7	13	9	16	32	57
Female	1,938	-	2	1	-	4	5	3	11	13	19	35
White, both sexes	3,143	-	3	2	2	4	9	12	14	22	44	79
Male	1,332	-	1	1	2	2	5	9	7	13	27	50
Female	1,811	-	2	1	-	2	4	3	7	9	17	29
Black, both sexes	212	-	-	-	-	3	3	4	6	7	5	12
Male	92	-	-	-	-	1	2	4	2	3	4	7
Female	120	-	-	-	-	2	1	-	4	4	1	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Louisiana												
All races, both sexes	3,293	12	6	2	2	2	17	18	31	34	53	81
Male	1,483	5	5	-	1	1	9	8	13	24	34	52
Female	1,810	7	1	2	1	1	8	10	18	10	19	29
White, both sexes	2,411	4	4	-	1	1	12	7	13	19	26	40
Male	1,075	2	3	-	-	1	7	5	6	14	17	25
Female	1,336	2	1	-	1	-	5	2	7	5	9	15
Black, both sexes	873	8	2	2	1	1	5	11	18	15	26	41
Male	404	3	2	-	1	-	2	3	7	10	16	27
Female	469	5	-	2	-	1	3	8	11	5	10	14
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Maine												
All races, both sexes	907	-	1	-	1	3	4	3	3	6	10	18
Male	437	-	1	-	1	2	2	3	3	3	7	15
Female	470	-	-	-	-	1	2	-	-	3	3	3
White, both sexes	899	-	1	-	1	3	4	3	3	5	9	18
Male	431	-	1	-	1	2	2	3	3	2	6	15
Female	468	-	-	-	-	1	2	-	-	3	3	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,572	5	3	1	5	6	10	18	35	33	63	67
Male	1,117	2	2	-	3	4	5	16	24	19	35	36
Female	1,455	3	1	1	2	2	5	2	11	14	28	31
White, both sexes	1,891	2	1	1	2	3	7	10	15	15	31	38
Male	811	1	1	-	2	2	4	10	9	9	21	20
Female	1,080	1	-	1	-	1	3	-	6	6	10	18
Black, both sexes	645	3	1	-	3	3	3	8	19	18	32	29
Male	289	1	1	-	1	2	1	6	14	10	14	16
Female	356	2	-	-	2	1	2	2	5	8	18	13
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	1	-	-	-	-	-	1	-	-	-
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	16	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Massachusetts												
All races, both sexes	4,296	3	2	1	2	3	10	16	10	22	43	59
Male	1,752	1	1	1	2	3	9	11	7	15	28	43
Female	2,544	2	1	-	-	-	1	5	3	7	15	16
White, both sexes	4,129	2	2	1	1	2	10	12	9	18	37	51
Male	1,680	1	1	1	1	2	9	9	7	13	23	39
Female	2,449	1	1	-	-	-	1	3	2	5	14	12
Black, both sexes	137	1	-	-	1	1	-	4	1	4	5	7
Male	64	-	-	-	1	1	-	2	-	2	5	3
Female	73	1	-	-	-	-	-	2	1	2	-	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	5,973	10	5	6	3	7	18	34	41	63	96	143
Male	2,579	5	2	3	2	5	14	24	32	43	68	82
Female	3,394	5	3	3	1	2	4	10	9	20	28	61
White, both sexes	5,205	4	3	6	3	3	9	17	19	36	59	85
Male	2,204	2	1	3	2	2	7	13	16	27	44	51
Female	3,001	2	2	3	1	1	2	4	3	9	15	34
Black, both sexes	723	4	2	-	-	4	8	17	22	27	36	55
Male	345	2	1	-	-	3	7	11	16	16	23	28
Female	378	2	1	-	-	1	1	6	6	11	13	27
American Indian 1/, both sexes	24	1	-	-	-	-	-	-	-	-	1	3
Male	16	-	-	-	-	-	-	-	-	-	1	3
Female	8	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	1	-	-	-	-	1	-	-	-	-	-
Male	14	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Minnesota												
All races, both sexes	2,816	3	2	3	1	4	9	8	16	18	28	46
Male	1,218	-	1	-	1	2	4	2	10	8	15	32
Female	1,598	3	1	3	-	2	5	6	6	10	13	14
White, both sexes	2,722	2	2	3	-	1	8	8	12	13	25	39
Male	1,169	-	1	-	-	1	3	2	8	5	13	26
Female	1,553	2	1	3	-	-	5	6	4	8	12	13
Black, both sexes	56	-	-	-	-	3	1	-	3	3	2	5
Male	29	-	-	-	-	1	1	-	2	2	1	4
Female	27	-	-	-	-	2	-	-	1	1	1	1
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	24	1	-	-	1	-	-	-	-	2	-	1
Male	13	-	-	-	1	-	-	-	-	1	-	1
Female	11	1	-	-	-	-	-	-	-	1	-	-
Mississippi												
All races, both sexes	2,975	4	-	2	4	2	13	15	21	20	42	73
Male	1,278	2	-	-	4	1	6	5	10	14	24	44
Female	1,697	2	-	2	-	1	7	10	11	6	18	29
White, both sexes	2,019	4	-	-	1	-	4	4	6	7	17	30
Male	855	2	-	-	1	-	2	2	3	5	10	17
Female	1,164	2	-	-	-	-	2	2	3	2	7	13
Black, both sexes	948	-	-	2	3	2	9	11	15	13	25	43
Male	417	-	-	-	3	1	4	3	7	9	14	27
Female	531	-	-	2	-	1	5	8	8	4	11	16
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Missouri												
All races, both sexes	3,944	2	2	1	5	8	14	7	21	38	59	89
Male	1,640	-	1	-	2	4	9	6	14	22	37	56
Female	2,304	2	1	1	3	4	5	1	7	16	22	33
White, both sexes	3,582	2	1	1	5	7	9	4	13	26	50	65
Male	1,483	-	-	-	2	4	6	4	9	16	33	42
Female	2,099	2	1	1	3	3	3	-	4	10	17	23
Black, both sexes	352	-	1	-	-	1	4	2	8	12	9	24
Male	153	-	1	-	-	-	3	1	5	6	4	14
Female	199	-	-	-	-	1	1	1	3	6	5	10
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	1	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	704	-	-	-	1	-	1	3	3	7	3	10
Male	331	-	-	-	1	-	1	2	3	4	2	6
Female	373	-	-	-	-	-	-	1	-	3	1	4
White, both sexes	677	-	-	-	1	-	1	2	3	6	3	6
Male	316	-	-	-	1	-	1	1	3	3	2	4
Female	361	-	-	-	-	-	-	1	-	3	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	1	-	4
Male	14	-	-	-	-	-	-	-	-	1	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Nebraska												
All races, both sexes	1,427	3	-	-	2	-	4	5	4	12	14	26
Male	592	1	-	-	1	-	1	3	3	10	6	15
Female	835	2	-	-	1	-	3	2	1	2	8	11
White, both sexes	1,371	2	-	-	1	-	3	3	4	11	12	21
Male	564	1	-	-	1	-	1	2	3	9	5	11
Female	807	1	-	-	-	-	2	1	1	2	7	10
Black, both sexes	47	1	-	-	1	-	1	2	-	1	1	4
Male	24	-	-	-	-	-	-	1	-	1	1	3
Female	23	1	-	-	1	-	1	1	-	-	-	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,893	2	1	1	-	1	6	14	25	19	49	64
Male	996	1	1	1	-	1	4	8	17	10	23	41
Female	897	1	-	-	-	-	2	6	8	9	26	23
White, both sexes	1,638	2	-	1	-	1	3	10	18	12	38	51
Male	867	1	-	1	-	1	2	5	11	6	20	32
Female	771	1	-	-	-	-	1	5	7	6	18	19
Black, both sexes	177	-	1	-	-	-	1	3	4	7	10	11
Male	88	-	1	-	-	-	1	2	3	4	2	8
Female	89	-	-	-	-	-	-	1	1	3	8	3
American Indian 1/, both sexes	21	-	-	-	-	-	1	1	1	-	-	1
Male	11	-	-	-	-	-	-	1	1	-	-	1
Female	10	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	1	-	2	-	1	1
Male	30	-	-	-	-	-	1	-	2	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
New Hampshire												
All races, both sexes	732	-	1	-	2	4	2	-	2	6	9	12
Male	312	-	1	-	1	1	2	-	1	4	3	8
Female	420	-	-	-	1	3	-	-	1	2	6	4
White, both sexes	730	-	1	-	2	4	2	-	2	6	8	12
Male	310	-	1	-	1	1	2	-	1	4	2	8
Female	420	-	-	-	1	3	-	-	1	2	6	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,897	8	3	1	4	3	15	21	30	40	73	94
Male	2,134	6	1	1	3	2	9	20	19	27	49	64
Female	2,763	2	2	-	1	1	6	1	11	13	24	30
White, both sexes	4,281	3	2	1	4	3	8	13	21	23	49	56
Male	1,827	2	-	1	3	2	6	12	13	16	36	37
Female	2,454	1	2	-	1	1	2	1	8	7	13	19
Black, both sexes	546	5	1	-	-	-	7	8	8	15	24	36
Male	277	4	1	-	-	-	3	8	5	9	13	26
Female	269	1	-	-	-	-	4	-	3	6	11	10
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	67	-	-	-	-	-	-	-	1	2	-	2
Male	30	-	-	-	-	-	-	-	1	2	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
New Mexico												
All races, both sexes	945	2	-	-	-	1	3	5	9	8	14	24
Male	406	2	-	-	-	1	1	3	3	4	9	16
Female	539	-	-	-	-	-	2	2	6	4	5	8
White, both sexes	892	2	-	-	-	1	3	5	8	6	9	19
Male	377	2	-	-	-	1	1	3	3	2	4	12
Female	515	-	-	-	-	-	2	2	5	4	5	7
Black, both sexes	20	-	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	2	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	1	2	3	4
Male	14	-	-	-	-	-	-	-	-	2	3	3
Female	12	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	8,089	12	10	6	8	20	16	39	49	74	120	193
Male	3,554	8	5	4	5	14	14	26	29	47	65	121
Female	4,535	4	5	2	3	6	2	13	20	27	55	72
White, both sexes	7,275	8	7	3	6	15	8	28	35	52	85	137
Male	3,167	6	3	3	4	10	7	17	19	35	46	89
Female	4,108	2	4	-	2	5	1	11	16	17	39	48
Black, both sexes	710	4	3	2	2	5	6	11	14	21	31	51
Male	332	2	2	1	1	4	5	9	10	12	16	30
Female	378	2	1	1	1	1	1	2	4	9	15	21
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	92	-	-	1	-	-	2	-	-	-	4	4
Male	51	-	-	-	-	-	2	-	-	-	3	2
Female	41	-	-	1	-	-	-	-	-	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
North Carolina												
All races, both sexes	5,309	16	10	5	6	11	23	27	33	54	90	119
Male	2,349	11	2	4	5	8	12	15	20	32	55	78
Female	2,960	5	8	1	1	3	11	12	13	22	35	41
White, both sexes	4,113	7	6	3	1	5	12	16	13	25	49	70
Male	1,781	4	2	2	1	5	6	10	8	18	31	46
Female	2,332	3	4	1	-	-	6	6	5	7	18	24
Black, both sexes	1,137	7	3	2	5	6	10	11	20	29	39	47
Male	531	6	-	2	4	3	6	5	12	14	22	31
Female	606	1	3	-	1	3	4	6	8	15	17	16
American Indian 1/, both sexes	37	1	-	-	-	-	1	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	16	1	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	22	1	1	-	-	-	-	-	-	-	1	1
Male	16	1	-	-	-	-	-	-	-	-	1	1
Female	6	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	339	-	-	-	1	-	2	1	2	2	4	2
Male	136	-	-	-	1	-	1	-	1	1	3	1
Female	203	-	-	-	-	-	1	1	1	1	1	1
White, both sexes	321	-	-	-	1	-	2	1	-	2	2	-
Male	128	-	-	-	1	-	1	-	-	1	1	-
Female	193	-	-	-	-	-	1	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	2	-	2	2
Male	8	-	-	-	-	-	-	-	1	-	2	1
Female	10	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Ohio												
All races, both sexes	6,154	16	4	1	1	12	22	22	42	50	82	155
Male	2,603	8	2	1	1	11	12	14	27	35	45	96
Female	3,551	8	2	-	-	1	10	8	15	15	37	59
White, both sexes	5,510	11	3	1	1	7	20	13	32	33	66	118
Male	2,289	5	2	1	1	7	11	7	22	23	38	79
Female	3,221	6	1	-	-	-	9	6	10	10	28	39
Black, both sexes	627	5	1	-	-	5	2	9	10	17	16	37
Male	305	3	-	-	-	4	1	7	5	12	7	17
Female	322	2	1	-	-	1	1	2	5	5	9	20
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,534	3	2	-	3	7	6	17	13	21	41	55
Male	1,089	3	-	-	1	4	4	13	10	12	23	38
Female	1,445	-	2	-	2	3	2	4	3	9	18	17
White, both sexes	2,255	1	2	-	2	5	5	10	7	11	34	41
Male	953	1	-	-	-	3	3	8	6	7	17	28
Female	1,302	-	2	-	2	2	2	2	1	4	17	13
Black, both sexes	154	-	-	-	1	1	1	4	4	5	1	10
Male	73	-	-	-	1	-	1	4	3	2	-	8
Female	81	-	-	-	-	1	-	-	1	3	1	2
American Indian 1/, both sexes	118	2	-	-	-	1	-	2	2	5	6	4
Male	63	2	-	-	-	1	-	1	1	3	6	2
Female	55	-	-	-	-	-	-	1	1	2	-	2
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Oregon												
All races, both sexes	2,179	2	-	2	2	1	6	9	11	15	13	33
Male	939	-	-	1	1	1	1	5	6	11	9	23
Female	1,240	2	-	1	1	-	5	4	5	4	4	10
White, both sexes	2,102	1	-	2	2	1	6	7	7	15	12	28
Male	904	-	-	1	1	1	1	3	3	11	9	18
Female	1,198	1	-	1	1	-	5	4	4	4	3	10
Black, both sexes	30	-	-	-	-	-	-	1	4	-	-	3
Male	17	-	-	-	-	-	-	1	3	-	-	3
Female	13	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	20	1	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	12	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	1	-	-	-	1
Male	10	-	-	-	-	-	-	1	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10,654	12	5	1	8	7	18	37	42	60	119	197
Male	4,508	5	4	1	6	2	13	19	23	37	77	131
Female	6,146	7	1	-	2	5	5	18	19	23	42	66
White, both sexes	9,738	11	2	-	5	5	11	26	27	40	86	149
Male	4,051	5	2	-	3	2	7	13	14	24	53	99
Female	5,687	6	-	-	2	3	4	13	13	16	33	50
Black, both sexes	885	1	3	1	3	2	7	11	15	20	32	47
Male	442	-	2	1	3	-	6	6	9	13	23	31
Female	443	1	1	-	-	2	1	5	6	7	9	16
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Rhode Island												
All races, both sexes	465	1	-	-	1	2	1	4	3	4	3	11
Male	183	-	-	-	-	2	1	3	3	1	2	6
Female	282	1	-	-	1	-	-	1	-	3	1	5
White, both sexes	449	1	-	-	-	2	1	3	3	4	3	10
Male	175	-	-	-	-	2	1	2	3	1	2	5
Female	274	1	-	-	-	-	-	1	-	3	1	5
Black, both sexes	12	-	-	-	1	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	7	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3,148	11	4	1	-	6	12	23	27	34	66	110
Male	1,516	5	2	-	-	4	8	18	14	21	41	68
Female	1,632	6	2	1	-	2	4	5	13	13	25	42
White, both sexes	2,184	4	2	1	-	2	6	11	10	13	34	60
Male	1,031	2	1	-	-	2	4	8	4	8	24	39
Female	1,153	2	1	1	-	-	2	3	6	5	10	21
Black, both sexes	951	7	2	-	-	4	6	12	17	21	32	50
Male	482	3	1	-	-	2	4	10	10	13	17	29
Female	469	4	1	-	-	2	2	2	7	8	15	21
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
South Dakota												
All races, both sexes	285	1	-	-	1	-	3	1	1	2	8	10
Male	126	1	-	-	1	-	-	1	-	1	7	8
Female	159	-	-	-	-	-	3	-	1	1	1	2
White, both sexes	274	-	-	-	1	-	-	1	1	2	7	9
Male	119	-	-	-	1	-	-	1	-	1	6	7
Female	155	-	-	-	-	-	-	-	1	1	1	2
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	3	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3,043	12	2	1	2	11	12	18	34	46	74	93
Male	1,369	6	1	-	1	5	8	12	21	23	41	54
Female	1,674	6	1	1	1	6	4	6	13	23	33	39
White, both sexes	2,516	8	1	1	1	7	5	12	23	32	52	69
Male	1,102	4	-	-	-	2	4	10	13	17	25	43
Female	1,414	4	1	1	1	5	1	2	10	15	27	26
Black, both sexes	516	4	1	-	1	3	7	5	10	14	22	24
Male	261	2	1	-	1	2	4	1	7	6	16	11
Female	255	2	-	-	-	1	3	4	3	8	6	13
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	1	-	1	1	-	-	-
Male	5	-	-	-	-	1	-	1	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Texas												
All races, both sexes	10,651	30	16	7	11	31	43	50	92	108	185	291
Male	4,707	18	9	5	7	18	32	28	59	64	100	167
Female	5,944	12	7	2	4	13	11	22	33	44	85	124
White, both sexes	9,197	24	11	5	7	22	34	31	61	69	131	219
Male	4,011	14	6	4	5	12	27	16	43	39	73	121
Female	5,186	10	5	1	2	10	7	15	18	30	58	98
Black, both sexes	1,353	5	5	2	3	9	9	19	31	39	53	66
Male	647	3	3	1	2	6	5	12	16	25	27	42
Female	706	2	2	1	1	3	4	7	15	14	26	24
American Indian 1/, both sexes	17	-	-	-	1	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	84	1	-	-	-	-	-	-	-	-	1	6
Male	39	1	-	-	-	-	-	-	-	-	-	4
Female	45	-	-	-	-	-	-	-	-	-	1	2
Utah												
All races, both sexes	1,311	1	1	-	2	3	5	7	7	10	8	25
Male	562	-	-	-	2	2	4	1	7	3	5	15
Female	749	1	1	-	-	1	1	6	-	7	3	10
White, both sexes	1,278	1	1	-	2	3	4	6	7	10	6	24
Male	540	-	-	-	2	2	3	1	7	3	3	14
Female	738	1	1	-	-	1	1	5	-	7	3	10
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	1	1	-	-	-	-
Male	13	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Vermont												
All races, both sexes	284	1	-	1	1	1	-	-	-	2	3	1
Male	113	-	-	-	1	-	-	-	-	2	1	-
Female	171	1	-	1	-	1	-	-	-	-	2	1
White, both sexes	283	1	-	1	1	1	-	-	-	2	3	1
Male	112	-	-	-	1	-	-	-	-	2	1	-
Female	171	1	-	1	-	1	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5,006	12	7	3	3	17	17	10	29	43	79	133
Male	2,253	4	5	2	2	12	14	6	18	28	53	92
Female	2,753	8	2	1	1	5	3	4	11	15	26	41
White, both sexes	3,942	5	3	1	-	11	11	7	11	27	47	68
Male	1,726	1	2	1	-	9	10	5	6	18	30	49
Female	2,216	4	1	-	-	2	1	2	5	9	17	19
Black, both sexes	1,011	7	4	1	3	6	5	3	17	15	32	63
Male	492	3	3	-	2	3	3	1	11	9	23	41
Female	519	4	1	1	1	3	2	2	6	6	9	22
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	50	-	-	1	-	-	1	-	1	1	-	2
Male	34	-	-	1	-	-	1	-	1	1	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Washington												
All races, both sexes	2,397	2	1	-	-	3	8	11	15	32	23	56
Male	1,041	1	-	-	-	2	5	8	5	16	12	37
Female	1,356	1	1	-	-	1	3	3	10	16	11	19
White, both sexes	2,253	1	-	-	-	2	6	8	13	20	20	50
Male	973	-	-	-	-	2	4	6	5	8	11	33
Female	1,280	1	-	-	-	-	2	2	8	12	9	17
Black, both sexes	64	1	1	-	-	-	1	1	1	9	1	4
Male	32	1	-	-	-	-	-	-	-	5	-	2
Female	32	-	1	-	-	-	1	1	1	4	1	2
American Indian 1/, both sexes	29	-	-	-	-	1	-	1	1	1	1	2
Male	14	-	-	-	-	-	-	1	-	1	-	2
Female	15	-	-	-	-	1	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	1	1	-	2	1	-
Male	22	-	-	-	-	-	1	1	-	2	1	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,613	3	1	-	1	2	4	4	5	10	9	30
Male	663	1	1	-	1	1	2	3	4	2	6	19
Female	950	2	-	-	-	1	2	1	1	8	3	11
White, both sexes	1,553	2	1	-	1	2	4	4	5	10	8	29
Male	638	1	1	-	1	1	2	3	4	2	5	18
Female	915	1	-	-	-	1	2	1	1	8	3	11
Black, both sexes	59	1	-	-	-	-	-	-	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	1	1
Female	34	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Wisconsin												
All races, both sexes	3,728	2	-	1	1	4	9	7	12	17	43	54
Male	1,603	1	-	-	-	3	7	2	6	13	22	31
Female	2,125	1	-	1	1	1	2	5	6	4	21	23
White, both sexes	3,569	2	-	-	-	2	8	5	5	14	37	48
Male	1,542	1	-	-	-	1	6	2	3	11	20	31
Female	2,027	1	-	-	-	1	2	3	2	3	17	17
Black, both sexes	125	-	-	1	-	2	-	2	6	1	5	5
Male	47	-	-	-	-	2	-	-	3	1	1	-
Female	78	-	-	1	-	-	-	2	3	-	4	5
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	2	1	1
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	20	-	-	-	1	-	1	-	1	-	-	-
Male	9	-	-	-	-	-	1	-	-	-	-	-
Female	11	-	-	-	1	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	384	-	-	1	-	2	2	2	1	1	4	12
Male	174	-	-	1	-	-	1	-	1	1	2	9
Female	210	-	-	-	-	2	1	2	-	-	2	3
White, both sexes	374	-	-	1	-	1	2	2	1	1	3	12
Male	169	-	-	1	-	-	1	-	1	1	2	9
Female	205	-	-	-	-	1	1	2	-	-	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	1	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
United States												
All races, both sexes	5,125	6,617	7,413	8,893	12,347	19,115	27,950	32,205	25,755	10,781	2,385	7
Male	3,247	4,118	4,432	5,171	6,635	9,445	12,335	12,143	7,692	2,381	375	6
Female	1,878	2,499	2,981	3,722	5,712	9,670	15,615	20,062	18,063	8,400	2,010	1
White, both sexes	3,633	4,913	5,808	7,098	10,259	16,717	25,129	29,791	24,042	10,038	2,111	4
Male	2,329	3,117	3,505	4,186	5,560	8,359	11,209	11,308	7,178	2,191	323	3
Female	1,304	1,796	2,303	2,912	4,699	8,358	13,920	18,483	16,864	7,847	1,788	1
Black, both sexes	1,361	1,543	1,441	1,565	1,831	2,042	2,395	1,973	1,389	622	234	3
Male	831	894	832	850	939	914	943	650	394	149	40	3
Female	530	649	609	715	892	1,128	1,452	1,323	995	473	194	-
American Indian 1/, both sexes	41	47	60	66	57	71	78	64	44	14	7	-
Male	27	31	34	40	31	31	34	26	14	-	2	-
Female	14	16	26	26	26	40	44	38	30	14	5	-
Asian or Pacific Islander 2/, both sexes	90	114	104	164	200	285	348	377	280	107	33	-
Male	60	76	61	95	105	141	149	159	106	41	10	-
Female	30	38	43	69	95	144	199	218	174	66	23	-
Alabama												
All races, both sexes	274	334	362	411	517	734	936	945	689	267	66	-
Male	181	206	230	243	289	338	378	311	199	52	8	-
Female	93	128	132	168	228	396	558	634	490	215	58	-
White, both sexes	171	229	247	296	375	590	746	786	570	223	40	-
Male	115	144	157	186	214	273	308	262	160	42	4	-
Female	56	85	90	110	161	317	438	524	410	181	36	-
Black, both sexes	102	105	112	114	140	142	187	156	117	44	26	-
Male	65	62	71	57	75	65	68	48	39	10	4	-
Female	37	43	41	57	65	77	119	108	78	34	22	-
American Indian 1/, both sexes	-	-	1	1	1	2	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	1	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	1	-	2	2	1	-	-	-
Male	1	-	2	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Alaska												
All races, both sexes	16	12	13	12	31	22	32	41	29	11	2	-
Male	12	8	10	8	19	15	12	18	9	1	1	-
Female	4	4	3	4	12	7	20	23	20	10	1	-
White, both sexes	11	9	8	8	25	14	26	29	27	10	2	-
Male	9	6	6	5	15	12	12	13	9	1	1	-
Female	2	3	2	3	10	2	14	16	18	9	1	-
Black, both sexes	-	2	1	-	1	1	-	-	-	-	-	-
Male	-	2	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	1	4	1	4	6	3	9	2	1	-	-
Male	3	-	3	1	3	2	-	4	-	-	-	-
Female	2	1	1	-	1	4	3	5	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	1	1	3	3	-	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	1	3	2	-	-	-	-
Arizona												
All races, both sexes	71	83	107	138	178	242	363	395	307	106	22	-
Male	48	46	66	91	100	126	165	171	111	27	7	-
Female	23	37	41	47	78	116	198	224	196	79	15	-
White, both sexes	61	75	99	125	172	229	350	387	301	104	18	-
Male	40	40	59	84	97	120	155	169	110	27	4	-
Female	21	35	40	41	75	109	195	218	191	77	14	-
Black, both sexes	5	4	2	8	1	5	8	4	2	1	1	-
Male	4	4	2	5	-	3	8	1	-	-	1	-
Female	1	-	-	3	1	2	-	3	2	1	-	-
American Indian 1/, both sexes	4	4	6	5	4	8	4	3	4	1	2	-
Male	3	2	5	2	3	3	2	-	1	-	1	-
Female	1	2	1	3	1	5	2	3	3	1	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	1	-	-	1	-
Male	1	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Arkansas												
All races, both sexes	50	81	96	118	168	231	308	363	254	107	26	-
Male	35	51	56	67	90	120	132	126	77	25	4	-
Female	15	30	40	51	78	111	176	237	177	82	22	-
White, both sexes	39	62	72	100	139	215	286	340	229	90	22	-
Male	28	36	40	56	72	115	125	120	71	19	4	-
Female	11	26	32	44	67	100	161	220	158	71	18	-
Black, both sexes	11	18	24	17	29	16	21	22	25	17	4	-
Male	7	14	16	11	18	5	6	6	6	6	-	-
Female	4	4	8	6	11	11	15	16	19	11	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
California												
All races, both sexes	441	536	474	569	855	1,296	2,242	2,659	2,157	918	196	1
Male	292	343	293	342	481	654	1,052	1,076	721	218	34	-
Female	149	193	181	227	374	642	1,190	1,583	1,436	700	162	1
White, both sexes	324	413	377	435	709	1,103	1,984	2,413	1,978	848	175	1
Male	215	273	237	270	404	572	949	987	657	192	28	-
Female	109	140	140	165	305	531	1,035	1,426	1,321	656	147	1
Black, both sexes	88	83	65	85	90	98	118	99	69	27	8	-
Male	56	44	37	41	48	40	46	34	24	10	2	-
Female	32	39	28	44	42	58	72	65	45	17	6	-
American Indian 1/, both sexes	4	2	4	5	6	13	11	8	2	2	1	-
Male	3	2	3	4	3	4	8	3	1	-	-	-
Female	1	-	1	1	3	9	3	5	1	2	1	-
Asian or Pacific Islander 2/, both sexes	25	38	28	44	50	82	129	139	108	41	12	-
Male	18	24	16	27	26	38	49	52	39	16	4	-
Female	7	14	12	17	24	44	80	87	69	25	8	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Colorado												
All races, both sexes	52	59	78	89	127	188	302	381	304	139	27	-
Male	29	37	36	52	57	100	128	139	81	29	6	-
Female	23	22	42	37	70	88	174	242	223	110	21	-
White, both sexes	48	54	70	83	116	177	290	370	297	136	27	-
Male	28	33	31	48	52	94	124	136	80	29	6	-
Female	20	21	39	35	64	83	166	234	217	107	21	-
Black, both sexes	2	4	5	4	7	2	7	10	3	1	-	-
Male	-	3	2	3	2	2	2	3	1	-	-	-
Female	2	1	3	1	5	-	5	7	2	1	-	-
American Indian 1/, both sexes	1	-	2	1	1	2	1	-	1	2	-	-
Male	1	-	2	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	3	7	4	1	3	-	-	-
Male	-	1	1	-	2	3	2	-	-	-	-	-
Female	1	-	-	1	1	4	2	1	3	-	-	-
Connecticut												
All races, both sexes	46	80	91	117	158	289	452	535	416	204	56	-
Male	38	49	59	65	88	148	195	214	143	32	6	-
Female	8	31	32	52	70	141	257	321	273	172	50	-
White, both sexes	38	71	84	100	144	276	431	522	408	198	53	-
Male	31	43	53	57	82	143	188	209	143	31	4	-
Female	7	28	31	43	62	133	243	313	265	167	49	-
Black, both sexes	7	8	7	16	12	12	21	12	7	5	2	-
Male	6	5	6	7	6	5	7	4	-	1	2	-
Female	1	3	1	9	6	7	14	8	7	4	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	1	-	1	1	1	1	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	2	1	-	-	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Delaware												
All races, both sexes	19	26	36	21	36	54	67	69	59	28	7	-
Male	10	14	21	10	20	24	31	22	26	5	2	-
Female	9	12	15	11	16	30	36	47	33	23	5	-
White, both sexes	13	21	27	15	27	49	62	62	55	26	7	-
Male	7	12	17	7	16	22	29	20	25	5	2	-
Female	6	9	10	8	11	27	33	42	30	21	5	-
Black, both sexes	6	5	9	6	9	5	5	6	4	1	-	-
Male	3	2	4	3	4	2	2	2	1	-	-	-
Female	3	3	5	3	5	3	3	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	15	5	6	14	18	17	41	25	26	13	3	-
Male	8	3	4	7	9	7	15	6	8	5	1	-
Female	7	2	2	7	9	10	26	19	18	8	2	-
White, both sexes	1	2	2	-	-	7	9	9	10	5	1	-
Male	1	2	-	-	-	5	3	4	4	3	-	-
Female	-	-	2	-	-	2	6	5	6	2	1	-
Black, both sexes	14	3	4	14	18	10	32	16	15	8	2	-
Male	7	1	4	7	9	2	12	2	4	2	1	-
Female	7	2	-	7	9	8	20	14	11	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Florida												
All races, both sexes	234	305	328	446	610	1,040	1,475	1,732	1,389	595	103	1
Male	151	187	211	262	348	536	731	722	508	172	23	1
Female	83	118	117	184	262	504	744	1,010	881	423	80	-
White, both sexes	155	233	268	374	528	929	1,377	1,636	1,318	573	98	-
Male	104	154	170	216	303	483	686	694	481	164	23	-
Female	51	79	98	158	225	446	691	942	837	409	75	-
Black, both sexes	78	66	59	70	79	107	97	86	69	21	5	1
Male	46	31	40	44	43	53	44	24	27	7	-	1
Female	32	35	19	26	36	54	53	62	42	14	5	-
American Indian 1/, both sexes	-	1	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	-	1	3	4	1	9	2	1	-	-
Male	1	2	-	1	2	-	1	4	-	1	-	-
Female	-	3	-	-	1	4	-	5	2	-	-	-
Georgia												
All races, both sexes	307	397	444	473	566	767	1,050	1,121	784	324	65	-
Male	176	275	265	265	309	367	460	395	232	72	8	-
Female	131	122	179	208	257	400	590	726	552	252	57	-
White, both sexes	170	226	300	327	402	585	836	906	659	267	46	-
Male	104	169	179	189	221	287	380	323	190	57	7	-
Female	66	57	121	138	181	298	456	583	469	210	39	-
Black, both sexes	133	169	142	144	158	174	207	206	122	56	19	-
Male	71	104	86	75	83	77	79	67	42	15	1	-
Female	62	65	56	69	75	97	128	139	80	41	18	-
American Indian 1/, both sexes	2	1	-	-	2	1	-	1	-	-	-	-
Male	1	1	-	-	1	1	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	2	4	7	7	8	3	1	-	-
Male	-	1	-	1	4	2	1	4	-	-	-	-
Female	2	-	2	1	-	5	6	4	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Hawaii												
All races, both sexes	36	32	39	52	69	96	126	145	120	53	14	-
Male	26	24	26	32	40	55	68	72	52	20	4	-
Female	10	8	13	20	29	41	58	73	68	33	10	-
White, both sexes	12	10	5	12	12	28	30	39	28	11	2	-
Male	10	8	5	7	9	11	18	19	11	1	1	-
Female	2	2	-	5	3	17	12	20	17	10	1	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	1	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	22	33	40	56	68	95	106	92	42	11	-
Male	16	16	20	25	30	44	50	53	41	19	3	-
Female	8	6	13	15	26	24	45	53	51	23	8	-
Idaho												
All races, both sexes	21	18	26	31	48	77	97	156	126	43	15	-
Male	11	12	17	25	25	38	43	60	38	9	3	-
Female	10	6	9	6	23	39	54	96	88	34	12	-
White, both sexes	20	17	26	31	47	76	96	155	126	42	15	-
Male	11	11	17	25	24	37	43	60	38	9	3	-
Female	9	6	9	6	23	39	53	95	88	33	12	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	1	1	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Illinois												
All races, both sexes	214	286	319	391	534	866	1,281	1,410	1,250	520	133	-
Male	144	166	185	230	266	393	560	523	338	117	24	-
Female	70	120	134	161	268	473	721	887	912	403	109	-
White, both sexes	149	203	254	316	407	745	1,128	1,323	1,195	490	124	-
Male	102	124	151	190	207	339	496	495	329	111	23	-
Female	47	79	103	126	200	406	632	828	866	379	101	-
Black, both sexes	64	80	62	69	118	113	144	78	51	29	8	-
Male	41	42	33	36	53	50	59	23	8	6	1	-
Female	23	38	29	33	65	63	85	55	43	23	7	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	3	5	9	8	9	8	4	1	1	-
Male	1	-	1	4	6	4	5	4	1	-	-	-
Female	-	3	2	1	3	4	4	4	3	1	1	-
Indiana												
All races, both sexes	136	186	201	290	352	487	735	812	715	273	71	-
Male	80	113	131	179	186	248	305	274	188	53	8	-
Female	56	73	70	111	166	239	430	538	527	220	63	-
White, both sexes	112	157	177	256	320	453	691	788	688	263	68	-
Male	66	93	113	157	169	235	290	269	181	49	8	-
Female	46	64	64	99	151	218	401	519	507	214	60	-
Black, both sexes	23	29	23	34	32	33	44	23	27	10	3	-
Male	14	20	17	22	17	13	15	5	7	4	-	-
Female	9	9	6	12	15	20	29	18	20	6	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Iowa												
All races, both sexes	35	57	54	53	78	143	197	244	226	128	34	-
Male	22	36	31	36	35	72	87	91	66	27	2	-
Female	13	21	23	17	43	71	110	153	160	101	32	-
White, both sexes	31	55	53	52	75	141	191	241	226	128	33	-
Male	19	35	30	35	33	72	86	91	66	27	2	-
Female	12	20	23	17	42	69	105	150	160	101	31	-
Black, both sexes	4	2	1	1	3	1	6	3	-	-	1	-
Male	3	1	1	1	2	-	1	-	-	-	-	-
Female	1	1	-	-	1	1	5	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Kansas												
All races, both sexes	45	59	72	72	136	188	318	403	419	221	47	-
Male	29	45	44	41	65	82	132	160	119	54	7	-
Female	16	14	28	31	71	106	186	243	300	167	40	-
White, both sexes	36	55	62	60	125	174	308	384	410	214	45	-
Male	21	41	40	35	61	76	128	153	118	52	7	-
Female	15	14	22	25	64	98	180	231	292	162	38	-
Black, both sexes	8	3	9	10	8	13	8	16	7	7	2	-
Male	7	3	3	5	3	5	3	6	-	2	-	-
Female	1	-	6	5	5	8	5	10	7	5	2	-
American Indian 1/, both sexes	1	-	-	2	2	1	1	3	1	-	-	-
Male	1	-	-	1	1	1	1	1	-	-	-	-
Female	-	-	-	1	1	-	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	1	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Kentucky												
All races, both sexes	104	144	166	250	251	410	546	602	436	184	40	-
Male	75	91	92	143	125	190	216	211	105	35	5	-
Female	29	53	74	107	126	220	330	391	331	149	35	-
White, both sexes	91	123	147	233	241	385	521	574	425	174	38	-
Male	67	80	85	131	121	179	205	205	103	35	4	-
Female	24	43	62	102	120	206	316	369	322	139	34	-
Black, both sexes	12	20	17	17	9	24	24	26	11	10	2	-
Male	7	11	5	12	4	11	10	6	2	-	1	-
Female	5	9	12	5	5	13	14	20	9	10	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	1	1	1	2	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	1	-	2	-	-	-	-
Louisiana												
All races, both sexes	120	159	188	202	279	422	530	535	402	174	24	-
Male	72	88	109	111	139	217	231	190	134	34	6	-
Female	48	71	79	91	140	205	299	345	268	140	18	-
White, both sexes	65	92	119	127	191	312	442	439	340	137	20	-
Male	40	49	69	74	100	166	191	160	117	25	4	-
Female	25	43	50	53	91	146	251	279	223	112	16	-
Black, both sexes	55	66	69	74	85	109	87	96	62	36	4	-
Male	32	38	40	36	38	51	40	30	17	9	2	-
Female	23	28	29	38	47	58	47	66	45	27	2	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	3	-	1	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Maine												
All races, both sexes	25	37	34	47	68	102	152	185	130	62	16	-
Male	18	28	23	29	42	62	71	83	30	11	3	-
Female	7	9	11	18	26	40	81	102	100	51	13	-
White, both sexes	23	35	34	47	68	102	151	184	130	62	16	-
Male	16	26	23	29	42	62	71	83	30	11	3	-
Female	7	9	11	18	26	40	80	101	100	51	13	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	71	110	111	139	197	282	410	469	370	141	25	1
Male	33	65	61	73	107	131	180	183	105	28	4	1
Female	38	45	50	66	90	151	230	286	265	113	21	-
White, both sexes	36	60	67	82	126	206	309	401	330	128	20	1
Male	13	35	37	45	74	101	145	159	92	27	3	1
Female	23	25	30	37	52	105	164	242	238	101	17	-
Black, both sexes	34	48	43	50	69	74	88	65	39	11	5	-
Male	19	29	23	24	32	29	30	24	12	-	1	-
Female	15	19	20	26	37	45	58	41	27	11	4	-
American Indian 1/, both sexes	-	1	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	6	2	2	12	3	1	2	-	-
Male	1	1	-	4	1	1	5	-	1	1	-	-
Female	-	-	-	2	1	1	7	3	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Massachusetts												
All races, both sexes	71	112	131	159	237	458	770	945	804	349	89	-
Male	44	80	72	93	125	246	331	339	220	69	12	-
Female	27	32	59	66	112	212	439	606	584	280	77	-
White, both sexes	60	106	124	149	225	430	751	922	788	343	86	-
Male	37	74	69	88	122	232	325	329	218	68	12	-
Female	23	32	55	61	103	198	426	593	570	275	74	-
Black, both sexes	8	4	7	8	9	20	13	22	15	5	2	-
Male	6	4	3	4	2	13	6	9	2	1	-	-
Female	2	-	4	4	7	7	7	13	13	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	-	2	3	7	5	1	1	1	1	-
Male	1	2	-	1	1	1	-	1	-	-	-	-
Female	2	-	-	1	2	6	5	-	1	1	1	-
Michigan												
All races, both sexes	150	199	218	257	398	660	976	1,214	987	399	89	-
Male	95	123	135	137	204	340	437	446	276	93	13	-
Female	55	76	83	120	194	320	539	768	711	306	76	-
White, both sexes	94	139	169	213	329	579	890	1,148	943	378	79	-
Male	58	92	108	118	174	305	395	424	263	88	11	-
Female	36	47	61	95	155	274	495	724	680	290	68	-
Black, both sexes	53	56	48	40	64	74	81	63	40	21	8	-
Male	35	28	26	16	28	29	39	21	10	5	1	-
Female	18	28	22	24	36	45	42	42	30	16	7	-
American Indian 1/, both sexes	2	3	1	3	2	3	3	1	1	-	-	-
Male	1	2	1	3	-	3	1	1	-	-	-	-
Female	1	1	-	-	2	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	3	4	2	2	3	-	2	-
Male	1	1	-	-	2	3	2	-	3	-	1	-
Female	-	-	-	1	1	1	-	2	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Minnesota												
All races, both sexes	49	70	87	90	151	298	443	617	557	253	63	-
Male	35	44	57	53	83	157	211	253	172	64	14	-
Female	14	26	30	37	68	141	232	364	385	189	49	-
White, both sexes	42	62	82	84	142	291	431	609	551	253	62	-
Male	30	39	55	51	78	152	208	248	171	64	14	-
Female	12	23	27	33	64	139	223	361	380	189	48	-
Black, both sexes	5	7	2	3	6	2	6	4	3	-	1	-
Male	4	4	1	1	3	2	1	2	-	-	-	-
Female	1	3	1	2	3	-	5	2	3	-	1	-
American Indian 1/, both sexes	1	1	2	1	2	2	2	-	-	-	-	-
Male	-	1	-	-	2	1	1	-	-	-	-	-
Female	1	-	2	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	1	3	4	4	3	-	-	-
Male	1	-	1	1	-	2	1	3	1	-	-	-
Female	-	-	-	1	1	1	3	1	2	-	-	-
Mississippi												
All races, both sexes	103	134	137	202	239	366	455	505	409	183	46	-
Male	68	84	79	121	128	172	178	170	121	36	11	-
Female	35	50	58	81	111	194	277	335	288	147	35	-
White, both sexes	46	64	79	134	166	256	341	397	310	129	24	-
Male	31	41	49	80	95	123	142	134	88	25	5	-
Female	15	23	30	54	71	133	199	263	222	104	19	-
Black, both sexes	57	67	58	68	73	108	113	106	99	54	22	-
Male	37	40	30	41	33	47	35	36	33	11	6	-
Female	20	27	28	27	40	61	78	70	66	43	16	-
American Indian 1/, both sexes	-	3	-	-	-	-	-	-	-	-	-	-
Male	-	3	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	2	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Missouri												
All races, both sexes	124	154	167	205	278	424	664	748	601	262	71	-
Male	70	92	89	111	155	195	287	253	173	51	13	-
Female	54	62	78	94	123	229	377	495	428	211	58	-
White, both sexes	98	122	147	170	250	377	619	716	581	253	66	-
Male	56	74	81	97	138	174	273	246	166	49	13	-
Female	42	48	66	73	112	203	346	470	415	204	53	-
Black, both sexes	26	32	18	35	27	45	43	31	20	9	5	-
Male	14	18	6	14	16	21	14	7	7	2	-	-
Female	12	14	12	21	11	24	29	24	13	7	5	-
American Indian 1/, both sexes	-	-	1	-	-	1	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Montana												
All races, both sexes	21	16	28	18	37	68	120	153	149	59	7	-
Male	17	9	20	13	20	37	63	69	48	16	-	-
Female	4	7	8	5	17	31	57	84	101	43	7	-
White, both sexes	19	15	24	17	33	67	117	151	147	59	6	-
Male	15	8	19	12	18	36	61	68	48	16	-	-
Female	4	7	5	5	15	31	56	83	99	43	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	4	1	4	1	3	2	2	-	1	-
Male	2	1	1	1	2	1	2	1	-	-	-	-
Female	-	-	3	-	2	-	1	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Nebraska												
All races, both sexes	31	39	47	60	101	152	226	260	259	141	41	-
Male	23	26	26	34	53	80	110	80	77	36	7	-
Female	8	13	21	26	48	72	116	180	182	105	34	-
White, both sexes	30	35	40	53	100	144	220	255	257	141	39	-
Male	22	24	21	30	52	77	107	78	77	36	7	-
Female	8	11	19	23	48	67	113	177	180	105	32	-
Black, both sexes	1	3	6	5	-	7	6	4	2	-	2	-
Male	1	2	4	3	-	3	3	2	-	-	-	-
Female	-	1	2	2	-	4	3	2	2	-	2	-
American Indian 1/, both sexes	-	1	1	1	1	1	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	75	112	156	165	201	204	317	269	154	47	11	-
Male	46	75	100	104	109	126	155	113	50	9	2	-
Female	29	37	56	61	92	78	162	156	104	38	9	-
White, both sexes	57	91	133	141	171	180	286	246	145	41	11	-
Male	34	63	90	94	95	110	145	100	47	8	2	-
Female	23	28	43	47	76	70	141	146	98	33	9	-
Black, both sexes	14	18	19	12	22	14	19	14	5	3	-	-
Male	8	10	9	6	10	8	7	8	1	-	-	-
Female	6	8	10	6	12	6	12	6	4	3	-	-
American Indian 1/, both sexes	-	1	2	2	2	-	2	4	3	1	-	-
Male	-	1	-	1	2	-	-	2	2	-	-	-
Female	-	-	2	1	-	-	2	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	4	2	2	10	6	10	10	5	1	2	-	-
Male	4	1	1	3	2	8	3	3	-	1	-	-
Female	-	1	1	7	4	2	7	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
New Hampshire												
All races, both sexes	15	22	26	22	46	80	124	156	142	46	15	-
Male	11	15	15	15	28	44	48	63	44	8	-	-
Female	4	7	11	7	18	36	76	93	98	38	15	-
White, both sexes	15	22	26	22	45	80	124	156	142	46	15	-
Male	11	15	15	15	27	44	48	63	44	8	-	-
Female	4	7	11	7	18	36	76	93	98	38	15	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	111	156	183	234	347	573	813	1,005	785	337	61	-
Male	74	100	107	133	191	286	350	385	226	74	7	-
Female	37	56	76	101	156	287	463	620	559	263	54	-
White, both sexes	75	114	135	185	289	516	726	937	743	321	57	-
Male	54	75	76	106	157	262	320	362	212	69	6	-
Female	21	39	59	79	132	254	406	575	531	252	51	-
Black, both sexes	33	39	44	40	53	48	78	58	31	15	3	-
Male	18	23	27	23	31	20	28	21	11	5	1	-
Female	15	16	17	17	22	28	50	37	20	10	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	4	9	5	8	8	9	11	1	1	-
Male	2	2	4	4	3	4	2	2	3	-	-	-
Female	1	1	-	5	2	4	6	7	8	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
New Mexico												
All races, both sexes	18	31	39	44	54	117	155	185	148	75	13	-
Male	8	22	24	21	33	58	67	69	45	18	2	-
Female	10	9	15	23	21	59	88	116	103	57	11	-
White, both sexes	15	28	35	40	49	111	152	180	144	74	11	-
Male	7	20	22	20	30	56	65	68	43	17	1	-
Female	8	8	13	20	19	55	87	112	101	57	10	-
Black, both sexes	1	2	1	-	3	4	1	2	1	1	1	-
Male	-	1	1	-	1	2	1	-	1	1	-	-
Female	1	1	-	-	2	2	-	2	-	-	1	-
American Indian 1/, both sexes	-	-	3	3	2	2	1	2	2	-	1	-
Male	-	-	1	1	2	-	1	-	-	-	1	-
Female	-	-	2	2	-	2	-	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	-	-	1	1	1	-	-	-
Male	1	1	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	1	-	-	1	-	-	-	-	-
New York												
All races, both sexes	183	261	287	331	518	943	1,348	1,653	1,327	570	121	-
Male	118	158	164	206	270	502	615	645	389	134	15	-
Female	65	103	123	125	248	441	733	1,008	938	436	106	-
White, both sexes	129	201	235	265	458	857	1,245	1,568	1,274	546	113	-
Male	86	129	140	164	245	461	566	619	373	130	15	-
Female	43	72	95	101	213	396	679	949	901	416	98	-
Black, both sexes	51	53	45	56	51	74	86	77	40	20	7	-
Male	29	24	21	35	21	35	41	21	10	3	-	-
Female	22	29	24	21	30	39	45	56	30	17	7	-
American Indian 1/, both sexes	1	1	2	2	1	-	2	-	1	-	-	-
Male	1	1	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	1	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	6	5	8	8	12	15	8	12	4	1	-
Male	2	4	1	7	4	6	8	5	6	1	-	-
Female	-	2	4	1	4	6	7	3	6	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
North Carolina												
All races, both sexes	173	229	284	332	446	632	885	935	701	239	59	-
Male	110	134	161	195	243	306	393	318	196	44	7	-
Female	63	95	123	137	203	326	492	617	505	195	52	-
White, both sexes	97	142	177	223	340	528	725	805	623	202	44	-
Male	65	83	101	126	182	259	334	275	183	36	4	-
Female	32	59	76	97	158	269	391	530	440	166	40	-
Black, both sexes	74	84	100	99	99	97	153	126	75	36	15	-
Male	43	50	56	60	56	43	53	43	11	8	3	-
Female	31	34	44	39	43	54	100	83	64	28	12	-
American Indian 1/, both sexes	2	2	6	7	1	5	4	2	3	1	-	-
Male	2	1	4	6	-	2	3	-	2	-	-	-
Female	-	1	2	1	1	3	1	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	6	2	3	2	-	-	-	-
Male	-	-	-	3	5	2	3	-	-	-	-	-
Female	-	1	1	-	1	-	-	2	-	-	-	-
North Dakota												
All races, both sexes	9	10	9	14	24	30	48	76	60	40	5	-
Male	5	6	5	7	17	14	25	32	11	6	-	-
Female	4	4	4	7	7	16	23	44	49	34	5	-
White, both sexes	7	9	9	12	22	28	46	75	60	40	5	-
Male	4	5	5	7	16	13	25	32	11	6	-	-
Female	3	4	4	5	6	15	21	43	49	34	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	2	2	2	2	1	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	2	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Ohio												
All races, both sexes	189	243	235	323	475	734	1,025	1,226	897	343	57	-
Male	110	140	128	169	247	346	419	470	246	68	8	-
Female	79	103	107	154	228	388	606	756	651	275	49	-
White, both sexes	136	192	195	276	403	664	958	1,155	853	325	48	-
Male	75	111	109	145	202	312	396	438	233	64	8	-
Female	61	81	86	131	201	352	562	717	620	261	40	-
Black, both sexes	52	50	38	44	71	68	65	70	42	16	9	-
Male	35	28	18	22	45	33	22	31	11	4	-	-
Female	17	22	20	22	26	35	43	39	31	12	9	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	1	1	2	2	1	2	2	-	-
Male	-	1	1	1	-	1	1	1	2	-	-	-
Female	1	-	1	-	1	1	1	-	-	2	-	-
Oklahoma												
All races, both sexes	67	84	115	124	210	285	458	464	370	145	44	-
Male	48	49	63	69	113	145	186	167	106	28	7	-
Female	19	35	52	55	97	140	272	297	264	117	37	-
White, both sexes	51	65	99	99	177	260	424	443	340	137	42	-
Male	39	37	54	55	100	131	173	160	99	25	7	-
Female	12	28	45	44	77	129	251	283	241	112	35	-
Black, both sexes	12	11	6	14	21	17	18	9	12	6	1	-
Male	8	6	5	5	8	8	6	2	3	3	-	-
Female	4	5	1	9	13	9	12	7	9	3	1	-
American Indian 1/, both sexes	4	8	9	11	11	7	16	12	15	2	1	-
Male	1	6	4	9	5	6	7	5	4	-	-	-
Female	3	2	5	2	6	1	9	7	11	2	1	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	-	3	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Oregon												
All races, both sexes	62	56	67	78	120	230	366	457	434	177	38	-
Male	31	36	44	47	73	104	151	202	138	48	7	-
Female	31	20	23	31	47	126	215	255	296	129	31	-
White, both sexes	58	51	58	73	117	221	354	446	429	177	37	-
Male	28	33	40	46	73	102	147	197	136	48	6	-
Female	30	18	18	27	44	119	207	249	293	129	31	-
Black, both sexes	2	2	6	-	2	1	5	3	1	-	-	-
Male	2	2	4	-	-	-	2	-	-	-	-	-
Female	-	-	2	-	2	1	3	3	1	-	-	-
American Indian 1/, both sexes	2	1	3	4	-	2	2	1	2	-	-	-
Male	1	-	-	1	-	1	1	1	2	-	-	-
Female	1	1	3	3	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	1	6	5	7	2	-	1	-
Male	-	1	-	-	-	1	1	4	-	-	1	-
Female	-	1	-	1	1	5	4	3	2	-	-	-
Pennsylvania												
All races, both sexes	249	343	440	525	739	1,260	1,883	2,202	1,715	646	145	1
Male	154	215	283	306	382	583	834	796	494	120	22	1
Female	95	128	157	219	357	677	1,049	1,406	1,221	526	123	-
White, both sexes	197	278	369	445	656	1,141	1,765	2,113	1,652	625	135	-
Male	123	175	238	261	340	532	790	762	471	116	21	-
Female	74	103	131	184	316	609	975	1,351	1,181	509	114	-
Black, both sexes	49	63	69	75	83	114	115	83	61	20	10	1
Male	29	39	44	43	42	50	42	31	22	4	1	1
Female	20	24	25	32	41	64	73	52	39	16	9	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	2	4	-	5	2	6	2	1	-	-
Male	2	1	1	1	-	1	1	3	1	-	-	-
Female	1	1	1	3	-	4	1	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Rhode Island												
All races, both sexes	8	6	13	26	35	53	72	86	91	37	8	-
Male	6	4	7	18	21	23	28	25	22	10	1	-
Female	2	2	6	8	14	30	44	61	69	27	7	-
White, both sexes	7	6	12	25	33	50	70	84	91	36	8	-
Male	6	4	7	17	20	20	28	24	22	10	1	-
Female	1	2	5	8	13	30	42	60	69	26	7	-
Black, both sexes	-	-	1	-	2	1	2	2	-	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	1	-	1	-	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	146	171	214	235	290	352	472	499	335	118	21	1
Male	95	117	133	133	151	185	226	165	98	28	3	1
Female	51	54	81	102	139	167	246	334	237	90	18	-
White, both sexes	75	95	122	146	202	265	354	411	261	97	13	-
Male	48	63	78	89	104	140	170	148	74	24	1	-
Female	27	32	44	57	98	125	184	263	187	73	12	-
Black, both sexes	70	75	92	87	87	85	114	87	73	21	8	1
Male	47	53	55	44	47	44	55	17	24	4	2	1
Female	23	22	37	43	40	41	59	70	49	17	6	-
American Indian 1/, both sexes	-	1	-	2	-	2	3	-	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	2	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
South Dakota												
All races, both sexes	8	6	5	16	13	21	41	53	60	23	12	-
Male	4	5	4	9	6	11	20	19	20	6	3	-
Female	4	1	1	7	7	10	21	34	40	17	9	-
White, both sexes	7	5	5	15	12	21	40	53	60	23	12	-
Male	3	4	4	8	6	11	19	19	20	6	3	-
Female	4	1	1	7	6	10	21	34	40	17	9	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	1	1	-	1	-	-	-	-	-
Male	1	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	124	163	191	201	258	322	520	464	339	134	21	1
Male	76	96	111	120	136	141	197	182	103	32	2	1
Female	48	67	80	81	122	181	323	282	236	102	19	-
White, both sexes	89	122	141	147	212	276	459	421	303	120	14	1
Male	56	75	79	85	110	125	173	159	90	30	1	1
Female	33	47	62	62	102	151	286	262	213	90	13	-
Black, both sexes	35	41	50	53	44	46	60	42	33	14	7	-
Male	20	21	32	35	25	16	23	22	13	2	1	-
Female	15	20	18	18	19	30	37	20	20	12	6	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	1	3	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Texas												
All races, both sexes	367	454	510	589	796	1,255	1,676	1,916	1,426	670	128	-
Male	234	273	304	322	440	594	725	736	412	138	22	-
Female	133	181	206	267	356	661	951	1,180	1,014	532	106	-
White, both sexes	270	344	404	494	677	1,092	1,511	1,768	1,306	611	106	-
Male	173	216	242	268	370	527	657	680	380	122	16	-
Female	97	128	162	226	307	565	854	1,088	926	489	90	-
Black, both sexes	90	101	97	87	112	147	156	135	110	55	22	-
Male	55	52	56	49	66	62	67	51	27	14	6	-
Female	35	49	41	38	46	85	89	84	83	41	16	-
American Indian 1/, both sexes	2	3	3	1	1	2	-	2	1	1	-	-
Male	2	1	2	1	1	-	-	2	1	-	-	-
Female	-	2	1	-	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	6	6	7	6	14	9	11	9	3	-	-
Male	4	4	4	4	3	5	1	3	4	2	-	-
Female	1	2	2	3	3	9	8	8	5	1	-	-
Utah												
All races, both sexes	32	38	42	56	107	129	243	294	211	80	9	1
Male	22	18	28	31	55	59	124	101	61	22	1	1
Female	10	20	14	25	52	70	119	193	150	58	8	-
White, both sexes	30	37	42	55	100	124	241	287	210	80	7	1
Male	20	17	28	30	50	55	122	99	60	22	1	1
Female	10	20	14	25	50	69	119	188	150	58	6	-
Black, both sexes	1	-	-	-	2	2	2	-	-	-	1	-
Male	1	-	-	-	1	2	2	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	5	3	-	6	1	-	-	-
Male	1	1	-	1	4	2	-	2	1	-	-	-
Female	-	-	-	-	1	1	-	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Vermont												
All races, both sexes	6	8	10	10	19	28	48	61	52	27	5	-
Male	3	4	5	8	11	10	23	22	16	6	1	-
Female	3	4	5	2	8	18	25	39	36	21	4	-
White, both sexes	6	7	10	10	19	28	48	61	52	27	5	-
Male	3	3	5	8	11	10	23	22	16	6	1	-
Female	3	4	5	2	8	18	25	39	36	21	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	183	218	229	279	417	583	793	895	706	282	68	-
Male	122	145	130	162	227	312	352	307	201	50	9	-
Female	61	73	99	117	190	271	441	588	505	232	59	-
White, both sexes	124	129	160	183	308	468	663	793	624	248	51	-
Male	83	87	94	106	172	254	295	269	184	45	6	-
Female	41	42	66	77	136	214	368	524	440	203	45	-
Black, both sexes	58	86	64	90	104	108	124	94	79	32	16	-
Male	38	55	32	51	53	54	51	35	16	5	3	-
Female	20	31	32	39	51	54	73	59	63	27	13	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	3	5	5	5	7	6	7	3	1	1	-
Male	1	3	4	4	2	4	6	3	1	-	-	-
Female	-	-	1	1	3	3	-	4	2	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Washington												
All races, both sexes	80	89	102	112	144	245	390	480	413	149	42	-
Male	43	47	55	65	79	118	161	193	143	46	5	-
Female	37	42	47	47	65	127	229	287	270	103	37	-
White, both sexes	72	72	93	102	129	229	378	463	406	147	42	-
Male	41	38	49	59	72	111	157	185	142	45	5	-
Female	31	34	44	43	57	118	221	278	264	102	37	-
Black, both sexes	3	9	6	6	7	4	6	2	1	1	-	-
Male	-	6	5	6	2	1	2	1	-	1	-	-
Female	3	3	1	-	5	3	4	1	1	-	-	-
American Indian 1/, both sexes	3	4	2	1	3	1	2	4	2	-	-	-
Male	2	1	1	-	1	-	1	3	1	-	-	-
Female	1	3	1	1	2	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	1	3	5	11	4	11	4	1	-	-
Male	-	2	-	-	4	6	1	4	-	-	-	-
Female	2	2	1	3	1	5	3	7	4	1	-	-
West Virginia												
All races, both sexes	39	61	71	84	138	212	294	292	232	98	23	-
Male	24	36	37	54	68	89	132	103	62	16	2	-
Female	15	25	34	30	70	123	162	189	170	82	21	-
White, both sexes	33	58	69	83	132	204	284	285	226	91	22	-
Male	20	33	35	54	65	86	128	102	62	13	2	-
Female	13	25	34	29	67	118	156	183	164	78	20	-
Black, both sexes	6	3	2	1	6	8	10	7	6	6	1	-
Male	4	3	2	-	3	3	4	1	-	3	-	-
Female	2	-	-	1	3	5	6	6	6	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Wisconsin												
All races, both sexes	91	108	113	144	207	409	597	796	718	320	75	-
Male	55	68	69	91	114	214	265	344	211	77	10	-
Female	36	40	44	53	93	195	332	452	507	243	65	-
White, both sexes	79	84	104	135	191	396	576	783	707	318	75	-
Male	48	56	66	88	109	211	256	338	209	76	10	-
Female	31	28	38	47	82	185	320	445	498	242	65	-
Black, both sexes	12	19	7	9	12	9	15	8	10	2	-	-
Male	7	7	3	3	4	3	7	3	2	1	-	-
Female	5	12	4	6	8	6	8	5	8	1	-	-
American Indian 1/, both sexes	-	1	1	-	2	1	4	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	2	1	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	1	-	2	3	2	4	1	-	-	-
Male	-	4	-	-	1	-	1	2	-	-	-	-
Female	-	-	1	-	1	3	1	2	1	-	-	-
Wyoming												
All races, both sexes	9	18	12	23	26	39	68	72	65	21	6	-
Male	6	10	7	13	13	25	30	29	21	2	3	-
Female	3	8	5	10	13	14	38	43	44	19	3	-
White, both sexes	9	16	12	23	23	38	67	72	64	21	6	-
Male	6	9	7	13	11	24	29	29	21	2	3	-
Female	3	7	5	10	12	14	38	43	43	19	3	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	1	1	-	1	-	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
United States												
All races, both sexes	12,387	81	24	12	13	37	71	128	180	267	392	559
Male	4,790	42	12	5	6	15	31	57	65	116	168	250
Female	7,597	39	12	7	7	22	40	71	115	151	224	309
White, both sexes	10,138	43	14	10	7	23	49	85	106	166	268	393
Male	3,929	21	5	4	3	9	21	37	35	77	115	181
Female	6,209	22	9	6	4	14	28	48	71	89	153	212
Black, both sexes	2,050	35	9	2	4	13	22	41	70	88	113	153
Male	787	19	6	1	2	6	10	19	30	34	50	64
Female	1,263	16	3	1	2	7	12	22	40	54	63	89
American Indian 1/, both sexes	63	2	-	-	-	1	-	-	2	8	4	5
Male	24	1	-	-	-	-	-	-	-	3	1	3
Female	39	1	-	-	-	1	-	-	2	5	3	2
Asian or Pacific Islander 2/, both sexes	136	1	1	-	2	-	-	2	2	5	7	8
Male	50	1	1	-	1	-	-	1	-	2	2	2
Female	86	-	-	-	1	-	-	1	2	3	5	6
Alabama												
All races, both sexes	252	1	1	1	-	1	3	4	5	3	13	11
Male	112	1	1	-	-	1	1	1	1	2	7	6
Female	140	-	-	1	-	-	2	3	4	1	6	5
White, both sexes	179	1	1	1	-	1	1	2	2	-	8	5
Male	82	1	1	-	-	1	1	-	-	-	5	3
Female	97	-	-	1	-	-	-	2	2	-	3	2
Black, both sexes	72	-	-	-	-	-	2	2	3	3	5	6
Male	30	-	-	-	-	-	-	1	1	2	2	3
Female	42	-	-	-	-	-	2	1	2	1	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Alaska												
All races, both sexes	22	-	-	-	-	-	-	-	1	3	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	3	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	1	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	2	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Arizona												
All races, both sexes	169	2	-	-	1	1	-	2	2	6	6	11
Male	73	1	-	-	-	1	-	1	1	3	4	3
Female	96	1	-	-	1	-	-	1	1	3	2	8
White, both sexes	157	2	-	-	-	-	-	1	2	4	5	11
Male	70	1	-	-	-	-	-	1	1	1	4	3
Female	87	1	-	-	-	-	-	-	1	3	1	8
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	2	1	-
Male	2	-	-	-	-	-	-	-	-	2	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Arkansas												
All races, both sexes	198	-	-	-	-	1	-	3	1	9	6	10
Male	81	-	-	-	-	-	-	1	1	6	2	8
Female	117	-	-	-	-	1	-	2	-	3	4	2
White, both sexes	165	-	-	-	-	1	-	-	1	6	6	7
Male	68	-	-	-	-	-	-	-	1	5	2	5
Female	97	-	-	-	-	1	-	-	-	1	4	2
Black, both sexes	31	-	-	-	-	-	-	3	-	1	-	3
Male	12	-	-	-	-	-	-	1	-	-	-	3
Female	19	-	-	-	-	-	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	2	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	748	5	4	1	2	3	7	4	10	15	34	32
Male	302	3	2	1	1	3	4	3	3	6	13	13
Female	446	2	2	-	1	-	3	1	7	9	21	19
White, both sexes	619	5	4	1	1	1	5	2	8	11	22	29
Male	255	3	2	1	1	1	3	1	3	5	9	13
Female	364	2	2	-	-	-	2	1	5	6	13	16
Black, both sexes	83	-	-	-	1	2	2	1	1	2	8	2
Male	34	-	-	-	-	2	1	1	-	1	4	-
Female	49	-	-	-	1	-	1	-	1	1	4	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	41	-	-	-	-	-	-	1	1	1	3	1
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	29	-	-	-	-	-	-	-	1	1	3	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Colorado												
All races, both sexes	218	-	-	-	-	-	4	2	2	3	3	3
Male	77	-	-	-	-	-	1	1	-	3	2	1
Female	141	-	-	-	-	-	3	1	2	-	1	2
White, both sexes	197	-	-	-	-	-	3	2	2	3	2	2
Male	70	-	-	-	-	-	1	1	-	3	2	1
Female	127	-	-	-	-	-	2	1	2	-	-	1
Black, both sexes	15	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Connecticut												
All races, both sexes	182	1	-	1	-	-	1	3	1	3	4	10
Male	64	1	-	1	-	-	1	2	-	1	2	5
Female	118	-	-	-	-	-	-	1	1	2	2	5
White, both sexes	164	1	-	1	-	-	-	2	-	2	3	8
Male	58	1	-	1	-	-	-	1	-	1	2	4
Female	106	-	-	-	-	-	-	1	-	1	1	4
Black, both sexes	18	-	-	-	-	-	1	1	1	1	1	2
Male	6	-	-	-	-	-	1	1	-	-	-	1
Female	12	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Delaware												
All races, both sexes	40	-	-	-	-	1	-	-	-	1	2	4
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	1	-	-	-	1	1	4
White, both sexes	30	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	10	-	-	-	-	1	-	-	-	1	1	3
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	1	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	24	-	-	-	-	1	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	1	-	-	-	1	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	1	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	1	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Florida												
All races, both sexes	647	7	2	1	-	1	2	7	4	14	23	33
Male	275	5	1	-	-	-	-	4	2	6	14	17
Female	372	2	1	1	-	1	2	3	2	8	9	16
White, both sexes	526	1	1	1	-	-	1	6	1	10	12	22
Male	219	-	-	-	-	-	-	3	1	5	6	10
Female	307	1	1	1	-	-	1	3	-	5	6	12
Black, both sexes	118	6	-	-	-	1	1	1	3	4	11	10
Male	55	5	-	-	-	-	-	1	1	1	8	7
Female	63	1	-	-	-	1	1	-	2	3	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	-	-	-	-	-	-	-	1
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	417	6	-	-	-	2	7	9	9	11	13	15
Male	167	4	-	-	-	2	2	5	5	4	7	6
Female	250	2	-	-	-	-	5	4	4	7	6	9
White, both sexes	275	2	-	-	-	1	4	7	3	4	4	6
Male	107	1	-	-	-	1	-	4	1	2	2	2
Female	168	1	-	-	-	-	4	3	2	2	2	4
Black, both sexes	141	4	-	-	-	1	3	2	6	7	9	9
Male	60	3	-	-	-	1	2	1	4	2	5	4
Female	81	1	-	-	-	-	1	1	2	5	4	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Hawaii												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	1	-	1
Idaho												
All races, both sexes	77	-	-	-	-	-	2	1	3	1	2	3
Male	34	-	-	-	-	-	1	-	1	-	-	1
Female	43	-	-	-	-	-	1	1	2	1	2	2
White, both sexes	76	-	-	-	-	-	2	1	3	1	2	3
Male	34	-	-	-	-	-	1	-	1	-	-	1
Female	42	-	-	-	-	-	1	1	2	1	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Illinois												
All races, both sexes	604	3	-	-	-	1	2	4	6	10	11	22
Male	220	2	-	-	-	1	2	1	3	4	2	8
Female	384	1	-	-	-	-	-	3	3	6	9	14
White, both sexes	491	2	-	-	-	1	1	3	3	4	9	17
Male	180	1	-	-	-	1	1	1	2	1	1	6
Female	311	1	-	-	-	-	-	2	1	3	8	11
Black, both sexes	112	1	-	-	-	-	1	1	3	6	2	5
Male	39	1	-	-	-	-	1	-	1	3	1	2
Female	73	-	-	-	-	-	-	1	2	3	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	360	2	-	-	-	1	2	5	7	7	15	23
Male	126	-	-	-	-	-	1	2	2	4	7	10
Female	234	2	-	-	-	1	1	3	5	3	8	13
White, both sexes	325	2	-	-	-	1	1	5	6	6	13	18
Male	112	-	-	-	-	-	1	2	2	4	5	8
Female	213	2	-	-	-	1	-	3	4	2	8	10
Black, both sexes	35	-	-	-	-	-	1	-	1	1	2	5
Male	14	-	-	-	-	-	-	-	-	-	2	2
Female	21	-	-	-	-	-	1	-	1	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Iowa												
All races, both sexes	173	-	1	-	-	-	-	2	1	8	4	10
Male	63	-	-	-	-	-	-	2	1	5	3	4
Female	110	-	1	-	-	-	-	-	-	3	1	6
White, both sexes	168	-	1	-	-	-	-	2	1	6	4	10
Male	60	-	-	-	-	-	-	2	1	4	3	4
Female	108	-	1	-	-	-	-	-	-	2	1	6
Black, both sexes	5	-	-	-	-	-	-	-	-	2	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	161	2	-	-	-	2	-	2	2	3	4	6
Male	68	2	-	-	-	-	-	1	1	1	1	4
Female	93	-	-	-	-	2	-	1	1	2	3	2
White, both sexes	148	1	-	-	-	2	-	2	2	3	3	5
Male	58	1	-	-	-	-	-	1	1	1	-	3
Female	90	-	-	-	-	2	-	1	1	2	3	2
Black, both sexes	10	1	-	-	-	-	-	-	-	-	1	1
Male	8	1	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Kentucky												
All races, both sexes	224	-	1	1	1	2	2	4	4	8	9	10
Male	90	-	-	-	1	-	2	3	1	5	4	4
Female	134	-	1	1	-	2	-	1	3	3	5	6
White, both sexes	203	-	1	1	1	1	1	3	3	5	9	10
Male	79	-	-	-	1	-	1	2	-	3	4	4
Female	124	-	1	1	-	1	-	1	3	2	5	6
Black, both sexes	21	-	-	-	-	1	1	1	1	3	-	-
Male	11	-	-	-	-	-	1	1	1	2	-	-
Female	10	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	245	2	-	-	-	-	2	3	6	8	9	12
Male	103	-	-	-	-	-	1	-	1	3	3	6
Female	142	2	-	-	-	-	1	3	5	5	6	6
White, both sexes	169	-	-	-	-	-	2	-	1	4	4	8
Male	71	-	-	-	-	-	1	-	-	1	1	4
Female	98	-	-	-	-	-	1	-	1	3	3	4
Black, both sexes	75	2	-	-	-	-	-	3	5	4	5	4
Male	32	-	-	-	-	-	-	-	1	2	2	2
Female	43	2	-	-	-	-	-	3	4	2	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Maine												
All races, both sexes	60	-	-	-	-	-	2	-	-	-	-	3
Male	25	-	-	-	-	-	1	-	-	-	-	2
Female	35	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	60	-	-	-	-	-	2	-	-	-	-	3
Male	25	-	-	-	-	-	1	-	-	-	-	2
Female	35	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	246	2	1	-	-	-	2	5	8	3	13	9
Male	85	-	1	-	-	-	-	4	2	1	4	2
Female	161	2	-	-	-	-	2	1	6	2	9	7
White, both sexes	152	1	-	-	-	-	2	2	3	1	6	3
Male	57	-	-	-	-	-	-	2	-	-	1	1
Female	95	1	-	-	-	-	2	-	3	1	5	2
Black, both sexes	93	1	1	-	-	-	-	3	5	2	7	6
Male	27	-	1	-	-	-	-	2	2	1	3	1
Female	66	1	-	-	-	-	-	1	3	1	4	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Massachusetts												
All races, both sexes	285	2	-	-	-	-	-	1	1	6	11	8
Male	98	-	-	-	-	-	-	1	-	4	6	2
Female	187	2	-	-	-	-	-	-	1	2	5	6
White, both sexes	265	1	-	-	-	-	-	1	1	5	10	5
Male	91	-	-	-	-	-	-	1	-	4	6	2
Female	174	1	-	-	-	-	-	-	1	1	4	3
Black, both sexes	15	1	-	-	-	-	-	-	-	1	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	397	4	1	1	1	-	1	4	6	3	6	21
Male	160	1	-	-	1	-	1	1	4	2	3	8
Female	237	3	1	1	-	-	-	3	2	1	3	13
White, both sexes	320	1	1	1	1	-	1	2	4	2	3	11
Male	130	-	-	-	1	-	1	1	3	1	1	3
Female	190	1	1	1	-	-	-	1	1	1	2	8
Black, both sexes	73	2	-	-	-	-	-	2	2	1	3	10
Male	28	1	-	-	-	-	-	-	1	1	2	5
Female	45	1	-	-	-	-	-	2	1	-	1	5
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Minnesota												
All races, both sexes	200	-	-	-	1	-	-	3	3	3	6	13
Male	81	-	-	-	1	-	-	1	1	1	3	8
Female	119	-	-	-	-	-	-	2	2	2	3	5
White, both sexes	186	-	-	-	-	-	-	3	2	3	6	13
Male	77	-	-	-	-	-	-	1	-	1	3	8
Female	109	-	-	-	-	-	-	2	2	2	3	5
Black, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	164	2	-	-	-	-	3	4	5	3	5	8
Male	66	1	-	-	-	-	1	1	1	1	3	5
Female	98	1	-	-	-	-	2	3	4	2	2	3
White, both sexes	88	2	-	-	-	-	1	-	1	-	1	2
Male	33	1	-	-	-	-	-	-	-	-	1	1
Female	55	1	-	-	-	-	1	-	1	-	-	1
Black, both sexes	76	-	-	-	-	-	2	4	4	3	4	6
Male	33	-	-	-	-	-	1	1	1	1	2	4
Female	43	-	-	-	-	-	1	3	3	2	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Missouri												
All races, both sexes	306	-	-	1	2	1	1	1	5	8	10	17
Male	118	-	-	-	-	-	1	1	4	2	6	7
Female	188	-	-	1	2	1	-	-	1	6	4	10
White, both sexes	257	-	-	1	2	-	1	-	4	7	8	10
Male	101	-	-	-	-	-	1	-	3	2	5	3
Female	156	-	-	1	2	-	-	-	1	5	3	7
Black, both sexes	48	-	-	-	-	1	-	1	1	1	2	7
Male	17	-	-	-	-	-	-	1	1	-	1	4
Female	31	-	-	-	-	1	-	-	-	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	38	-	-	-	-	-	-	1	-	1	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	37	-	-	-	-	-	-	1	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Nebraska												
All races, both sexes	72	3	-	-	-	-	3	1	-	2	4	3
Male	18	1	-	-	-	-	-	1	-	2	1	1
Female	54	2	-	-	-	-	3	-	-	-	3	2
White, both sexes	69	2	-	-	-	-	2	1	-	2	4	3
Male	18	1	-	-	-	-	-	1	-	2	1	1
Female	51	1	-	-	-	-	2	-	-	-	3	2
Black, both sexes	3	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	111	-	-	-	-	-	-	1	2	-	3	10
Male	46	-	-	-	-	-	-	-	-	-	-	4
Female	65	-	-	-	-	-	-	1	2	-	3	6
White, both sexes	96	-	-	-	-	-	-	1	2	-	2	9
Male	42	-	-	-	-	-	-	-	-	-	-	4
Female	54	-	-	-	-	-	-	1	2	-	2	5
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Hampshire												
All races, both sexes	56	-	-	-	1	-	-	-	-	-	1	4
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	33	-	-	-	1	-	-	-	-	-	1	2
White, both sexes	55	-	-	-	1	-	-	-	-	-	1	4
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	33	-	-	-	1	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	348	2	-	-	-	2	4	2	6	7	14	16
Male	114	2	-	-	-	1	2	2	-	3	8	9
Female	234	-	-	-	-	1	2	-	6	4	6	7
White, both sexes	271	1	-	-	-	2	2	1	5	1	9	10
Male	87	1	-	-	-	1	1	1	-	-	7	5
Female	184	-	-	-	-	1	1	-	5	1	2	5
Black, both sexes	67	1	-	-	-	-	2	1	1	6	5	5
Male	25	1	-	-	-	-	1	1	-	3	1	3
Female	42	-	-	-	-	-	1	-	1	3	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Mexico												
All races, both sexes	78	1	-	-	-	-	1	2	1	3	2	6
Male	30	1	-	-	-	-	-	2	-	-	-	4
Female	48	-	-	-	-	-	1	-	1	3	2	2
White, both sexes	73	1	-	-	-	-	1	2	1	3	2	4
Male	27	1	-	-	-	-	-	2	-	-	-	2
Female	46	-	-	-	-	-	1	-	1	3	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	561	5	2	1	1	-	-	5	6	12	17	19
Male	214	2	-	1	-	-	-	3	1	4	6	7
Female	347	3	2	-	1	-	-	2	5	8	11	12
White, both sexes	459	3	1	1	1	-	-	3	4	6	13	10
Male	182	1	-	1	-	-	-	2	1	2	6	6
Female	277	2	1	-	1	-	-	1	3	4	7	4
Black, both sexes	91	2	1	-	-	-	-	2	2	5	4	8
Male	27	1	-	-	-	-	-	1	-	2	-	1
Female	64	1	1	-	-	-	-	1	2	3	4	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
North Carolina												
All races, both sexes	497	5	2	1	-	2	4	4	9	14	21	29
Male	182	5	-	1	-	1	2	2	5	4	11	16
Female	315	-	2	-	-	1	2	2	4	10	10	13
White, both sexes	344	1	1	-	-	-	3	2	2	6	11	12
Male	125	1	-	-	-	-	2	1	1	3	6	7
Female	219	-	1	-	-	-	1	1	1	3	5	5
Black, both sexes	146	3	1	1	-	2	1	2	7	8	10	16
Male	54	3	-	1	-	1	-	1	4	1	5	9
Female	92	-	1	-	-	1	1	1	3	7	5	7
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	29	-	-	-	-	-	1	-	1	1	2	-
Male	16	-	-	-	-	-	1	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	27	-	-	-	-	-	1	-	-	1	1	-
Male	15	-	-	-	-	-	1	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Ohio												
All races, both sexes	455	2	-	-	-	-	2	3	8	10	16	24
Male	178	2	-	-	-	-	1	1	3	5	7	8
Female	277	-	-	-	-	-	1	2	5	5	9	16
White, both sexes	384	1	-	-	-	-	1	2	6	9	11	20
Male	145	1	-	-	-	-	-	1	3	5	5	7
Female	239	-	-	-	-	-	1	1	3	4	6	13
Black, both sexes	68	1	-	-	-	-	1	1	2	1	5	4
Male	31	1	-	-	-	-	1	-	-	-	2	1
Female	37	-	-	-	-	-	-	1	2	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	137	1	-	-	-	1	1	1	-	4	4	6
Male	56	1	-	-	-	1	1	-	-	2	1	4
Female	81	-	-	-	-	-	-	1	-	2	3	2
White, both sexes	114	-	-	-	-	1	1	1	-	1	3	6
Male	50	-	-	-	-	1	1	-	-	1	1	4
Female	64	-	-	-	-	-	-	1	-	-	2	2
Black, both sexes	12	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	9	1	-	-	-	-	-	-	-	2	-	-
Male	4	1	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Oregon												
All races, both sexes	147	-	-	-	-	-	1	2	2	2	1	8
Male	56	-	-	-	-	-	-	-	-	1	-	6
Female	91	-	-	-	-	-	1	2	2	1	1	2
White, both sexes	136	-	-	-	-	-	1	2	1	2	1	7
Male	52	-	-	-	-	-	-	-	-	1	-	5
Female	84	-	-	-	-	-	1	2	1	1	1	2
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	690	2	2	-	1	-	-	7	9	10	17	24
Male	243	-	2	-	1	-	-	2	2	6	9	12
Female	447	2	-	-	-	-	-	5	7	4	8	12
White, both sexes	586	2	-	-	-	-	-	4	6	7	12	17
Male	205	-	-	-	-	-	-	1	1	5	6	9
Female	381	2	-	-	-	-	-	3	5	2	6	8
Black, both sexes	101	-	2	-	1	-	-	3	3	3	4	7
Male	37	-	2	-	1	-	-	1	1	1	2	3
Female	64	-	-	-	-	-	-	2	2	2	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Rhode Island												
All races, both sexes	46	1	-	-	-	-	-	1	1	-	1	2
Male	17	-	-	-	-	-	-	-	1	-	-	1
Female	29	1	-	-	-	-	-	1	-	-	1	1
White, both sexes	42	1	-	-	-	-	-	1	1	-	1	2
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	26	1	-	-	-	-	-	1	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	252	3	1	-	-	1	2	4	5	7	8	11
Male	97	1	1	-	-	1	-	2	4	3	1	4
Female	155	2	-	-	-	-	2	2	1	4	7	7
White, both sexes	166	1	1	-	-	1	2	2	1	2	3	6
Male	62	1	1	-	-	1	-	1	1	-	-	3
Female	104	-	-	-	-	-	2	1	-	2	3	3
Black, both sexes	85	2	-	-	-	-	-	2	4	5	5	5
Male	35	-	-	-	-	-	-	1	3	3	1	1
Female	50	2	-	-	-	-	-	1	1	2	4	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
South Dakota												
All races, both sexes	24	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	23	-	-	-	-	-	-	-	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	311	3	1	1	-	1	1	2	5	7	9	9
Male	116	1	1	-	-	-	1	1	1	2	5	2
Female	195	2	-	1	-	1	-	1	4	5	4	7
White, both sexes	249	2	-	1	-	1	-	1	5	6	8	4
Male	86	1	-	-	-	-	-	-	1	1	4	-
Female	163	1	-	1	-	1	-	1	4	5	4	4
Black, both sexes	62	1	1	-	-	-	1	1	-	1	1	5
Male	30	-	1	-	-	-	1	1	-	1	1	2
Female	32	1	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Texas												
All races, both sexes	882	4	3	1	2	7	1	10	12	26	32	43
Male	364	3	2	1	1	3	-	5	6	12	10	16
Female	518	1	1	-	1	4	1	5	6	14	22	27
White, both sexes	710	3	2	1	-	5	1	7	6	17	20	35
Male	297	3	1	1	-	2	-	3	2	7	7	14
Female	413	-	1	-	-	3	1	4	4	10	13	21
Black, both sexes	166	1	1	-	2	2	-	3	6	9	11	7
Male	65	-	1	-	1	1	-	2	4	5	3	2
Female	101	1	-	-	1	1	-	1	2	4	8	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	1
Utah												
All races, both sexes	100	-	-	-	-	1	-	3	1	-	2	5
Male	41	-	-	-	-	-	-	-	1	-	-	2
Female	59	-	-	-	-	1	-	3	-	-	2	3
White, both sexes	98	-	-	-	-	1	-	3	1	-	2	5
Male	39	-	-	-	-	-	-	-	1	-	-	2
Female	59	-	-	-	-	1	-	3	-	-	2	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Vermont												
All races, both sexes	18	1	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	1	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	394	6	2	-	-	1	3	1	9	9	11	17
Male	162	2	1	-	-	-	2	-	3	5	7	8
Female	232	4	1	-	-	1	1	1	6	4	4	9
White, both sexes	295	1	-	-	-	1	2	-	5	5	10	8
Male	128	-	-	-	-	-	2	-	1	3	7	5
Female	167	1	-	-	-	1	-	-	4	2	3	3
Black, both sexes	96	5	2	-	-	-	1	1	4	3	1	8
Male	31	2	1	-	-	-	-	-	2	1	-	2
Female	65	3	1	-	-	-	1	1	2	2	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Washington												
All races, both sexes	251	-	-	-	-	-	2	2	7	6	5	7
Male	82	-	-	-	-	-	1	-	1	-	1	4
Female	169	-	-	-	-	-	1	2	6	6	4	3
White, both sexes	234	-	-	-	-	-	2	2	6	4	5	6
Male	78	-	-	-	-	-	1	-	1	-	1	4
Female	156	-	-	-	-	-	1	2	5	4	4	2
Black, both sexes	11	-	-	-	-	-	-	-	-	2	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	110	1	-	-	-	1	-	1	1	1	1	5
Male	39	-	-	-	-	-	-	-	1	1	1	2
Female	71	1	-	-	-	1	-	1	-	-	-	3
White, both sexes	105	1	-	-	-	1	-	1	1	1	1	5
Male	36	-	-	-	-	-	-	-	1	1	1	2
Female	69	1	-	-	-	1	-	1	-	-	-	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Wisconsin												
All races, both sexes	288	-	-	1	-	1	2	-	3	3	9	11
Male	114	-	-	-	-	-	-	-	1	2	1	5
Female	174	-	-	1	-	1	2	-	2	1	8	6
White, both sexes	261	-	-	-	-	1	2	-	-	3	5	8
Male	108	-	-	-	-	-	-	-	-	2	-	5
Female	153	-	-	-	-	1	2	-	-	1	5	3
Black, both sexes	26	-	-	1	-	-	-	-	3	-	4	2
Male	6	-	-	-	-	-	-	-	1	-	1	-
Female	20	-	-	1	-	-	-	-	2	-	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	41	-	-	-	-	1	-	2	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	1	-	2	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	2	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
United States												
All races, both sexes	741	867	889	983	1,187	1,568	1,724	1,532	822	264	46	-
Male	352	387	385	453	512	619	588	466	207	48	6	-
Female	389	480	504	530	675	949	1,136	1,066	615	216	40	-
White, both sexes	574	653	704	776	977	1,350	1,517	1,388	760	239	36	-
Male	283	307	312	360	439	540	526	421	187	41	5	-
Female	291	346	392	416	538	810	991	967	573	198	31	-
Black, both sexes	152	202	174	190	198	187	184	130	52	21	10	-
Male	64	77	69	87	69	60	56	42	16	5	1	-
Female	88	125	105	103	129	127	128	88	36	16	9	-
American Indian 1/, both sexes	5	4	7	4	3	9	3	2	3	1	-	-
Male	2	-	3	1	3	4	2	-	1	-	-	-
Female	3	4	4	3	-	5	1	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	10	8	4	13	9	22	20	12	7	3	-	-
Male	3	3	1	5	1	15	4	3	3	2	-	-
Female	7	5	3	8	8	7	16	9	4	1	-	-
Alabama												
All races, both sexes	21	13	22	21	27	23	44	27	7	4	-	-
Male	13	3	12	8	15	8	18	9	3	2	-	-
Female	8	10	10	13	12	15	26	18	4	2	-	-
White, both sexes	15	6	14	16	19	20	34	24	5	4	-	-
Male	10	2	8	7	9	8	16	7	1	2	-	-
Female	5	4	6	9	10	12	18	17	4	2	-	-
Black, both sexes	6	7	8	4	8	3	10	3	2	-	-	-
Male	3	1	4	1	6	-	2	2	2	-	-	-
Female	3	6	4	3	2	3	8	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Alaska												
All races, both sexes	1	1	1	1	2	3	3	2	3	-	-	-
Male	1	-	-	1	1	3	3	-	1	-	-	-
Female	-	1	1	-	1	-	-	2	2	-	-	-
White, both sexes	1	1	1	1	1	2	3	2	3	-	-	-
Male	1	-	-	1	-	2	3	-	1	-	-	-
Female	-	1	1	-	1	-	-	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	11	10	13	23	22	21	17	15	1	-	-
Male	2	4	3	7	12	11	10	5	5	-	-	-
Female	3	7	7	6	11	11	11	12	10	1	-	-
White, both sexes	5	11	10	10	22	21	21	16	15	1	-	-
Male	2	4	3	7	12	11	10	5	5	-	-	-
Female	3	7	7	3	10	10	11	11	10	1	-	-
Black, both sexes	-	-	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Arkansas												
All races, both sexes	16	18	12	18	19	23	24	21	10	6	1	-
Male	10	9	4	6	13	8	6	2	4	1	-	-
Female	6	9	8	12	6	15	18	19	6	5	1	-
White, both sexes	15	16	10	16	13	20	19	20	9	6	-	-
Male	10	8	2	6	9	8	5	2	4	1	-	-
Female	5	8	8	10	4	12	14	18	5	5	-	-
Black, both sexes	1	2	2	2	6	3	5	1	1	-	1	-
Male	-	1	2	-	4	-	1	-	-	-	-	-
Female	1	1	-	2	2	3	4	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	52	59	48	52	55	92	128	81	53	9	2	-
Male	21	30	21	28	23	39	39	31	15	3	-	-
Female	31	29	27	24	32	53	89	50	38	6	2	-
White, both sexes	44	44	40	41	44	77	109	72	49	8	2	-
Male	20	24	17	21	19	33	35	28	14	2	-	-
Female	24	20	23	20	25	44	74	44	35	6	2	-
Black, both sexes	5	11	6	10	7	8	12	3	2	-	-	-
Male	1	5	4	6	3	2	2	1	1	-	-	-
Female	4	6	2	4	4	6	10	2	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	2	1	4	6	6	6	2	1	-	-
Male	-	1	-	1	1	4	1	2	-	1	-	-
Female	2	3	2	-	3	2	5	4	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Colorado												
All races, both sexes	16	18	20	17	19	23	31	31	21	4	1	-
Male	8	7	5	8	7	13	10	6	4	-	1	-
Female	8	11	15	9	12	10	21	25	17	4	-	-
White, both sexes	14	17	17	15	17	20	27	31	20	4	1	-
Male	8	7	5	6	6	10	9	6	4	-	1	-
Female	6	10	12	9	11	10	18	25	16	4	-	-
Black, both sexes	1	1	3	2	2	1	3	-	1	-	-	-
Male	-	-	-	2	1	1	1	-	-	-	-	-
Female	1	1	3	-	1	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	6	6	13	8	16	34	34	26	9	6	-	-
Male	2	4	6	4	6	10	9	8	2	-	-	-
Female	4	2	7	4	10	24	25	18	7	6	-	-
White, both sexes	4	4	12	7	15	33	31	26	9	6	-	-
Male	1	3	5	4	6	10	9	8	2	-	-	-
Female	3	1	7	3	9	23	22	18	7	6	-	-
Black, both sexes	2	2	1	1	1	1	3	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	1	1	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Delaware												
All races, both sexes	2	2	5	1	3	10	5	3	1	-	-	-
Male	-	1	1	-	1	4	4	1	-	-	-	-
Female	2	1	4	1	2	6	1	2	1	-	-	-
White, both sexes	2	2	3	-	2	10	5	3	1	-	-	-
Male	-	1	1	-	-	4	4	1	-	-	-	-
Female	2	1	2	-	2	6	1	2	1	-	-	-
Black, both sexes	-	-	2	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	2	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	2	2	-	2	3	6	2	2	1	-	-
Male	-	-	1	-	-	-	3	1	1	-	-	-
Female	1	2	1	-	2	3	3	1	1	1	-	-
White, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	1	2	1	-	2	3	5	1	1	1	-	-
Male	-	-	1	-	-	-	3	-	-	-	-	-
Female	1	2	-	-	2	3	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Florida												
All races, both sexes	31	42	46	58	54	95	85	78	48	15	1	-
Male	17	14	26	25	24	37	36	27	16	4	-	-
Female	14	28	20	33	30	58	49	51	32	11	1	-
White, both sexes	26	29	40	47	43	80	78	70	43	14	1	-
Male	15	14	22	18	21	29	34	24	14	3	-	-
Female	11	15	18	29	22	51	44	46	29	11	1	-
Black, both sexes	5	12	6	11	11	15	7	8	5	1	-	-
Male	2	-	4	7	3	8	2	3	2	1	-	-
Female	3	12	2	4	8	7	5	5	3	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	24	27	35	39	43	41	59	41	25	9	2	-
Male	8	15	13	17	15	20	17	19	8	-	-	-
Female	16	12	22	22	28	21	42	22	17	9	2	-
White, both sexes	14	12	25	21	32	31	47	31	22	7	2	-
Male	5	7	8	11	13	17	13	14	6	-	-	-
Female	9	5	17	10	19	14	34	17	16	7	2	-
Black, both sexes	9	15	10	18	11	10	12	10	3	2	-	-
Male	3	8	5	6	2	3	4	5	2	-	-	-
Female	6	7	5	12	9	7	8	5	1	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Hawaii												
All races, both sexes	4	1	2	3	-	4	5	7	2	1	-	-
Male	2	-	2	1	-	2	2	3	-	1	-	-
Female	2	1	-	2	-	2	3	4	2	-	-	-
White, both sexes	1	-	1	-	-	1	3	3	1	-	-	-
Male	1	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	3	-	3	2	4	1	1	-	-
Male	1	-	1	1	-	2	1	1	-	1	-	-
Female	2	1	-	2	-	1	1	3	1	-	-	-
Idaho												
All races, both sexes	9	7	2	8	11	14	1	8	3	2	-	-
Male	6	3	1	6	7	6	-	2	-	-	-	-
Female	3	4	1	2	4	8	1	6	3	2	-	-
White, both sexes	8	7	2	8	11	14	1	8	3	2	-	-
Male	6	3	1	6	7	6	-	2	-	-	-	-
Female	2	4	1	2	4	8	1	6	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Illinois												
All races, both sexes	33	50	51	46	63	79	78	83	40	17	5	-
Male	19	16	18	28	20	24	29	23	14	5	1	-
Female	14	34	33	18	43	55	49	60	26	12	4	-
White, both sexes	26	36	38	37	50	64	71	76	37	12	4	-
Male	16	13	14	23	17	21	25	21	13	2	1	-
Female	10	23	24	14	33	43	46	55	24	10	3	-
Black, both sexes	6	14	13	9	13	15	7	7	3	5	1	-
Male	2	3	4	5	3	3	4	2	1	3	-	-
Female	4	11	9	4	10	12	3	5	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	26	27	28	30	44	39	45	40	12	6	1	-
Male	11	12	8	16	18	11	12	10	1	1	-	-
Female	15	15	20	14	26	28	33	30	11	5	1	-
White, both sexes	23	24	27	26	39	37	41	38	11	6	1	-
Male	9	11	8	13	16	10	11	10	1	1	-	-
Female	14	13	19	13	23	27	30	28	10	5	1	-
Black, both sexes	3	3	1	4	5	2	4	2	1	-	-	-
Male	2	1	-	3	2	1	1	-	-	-	-	-
Female	1	2	1	1	3	1	3	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Iowa												
All races, both sexes	11	15	12	10	8	20	26	25	12	8	-	-
Male	7	7	5	4	3	9	6	6	1	-	-	-
Female	4	8	7	6	5	11	20	19	11	8	-	-
White, both sexes	10	15	11	10	8	20	25	25	12	8	-	-
Male	6	7	4	4	3	9	6	6	1	-	-	-
Female	4	8	7	6	5	11	19	19	11	8	-	-
Black, both sexes	1	-	1	-	-	-	1	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	11	7	12	12	16	18	20	21	16	7	-	-
Male	6	4	6	5	9	8	8	7	4	1	-	-
Female	5	3	6	7	7	10	12	14	12	6	-	-
White, both sexes	10	6	11	9	15	17	20	19	16	7	-	-
Male	5	3	5	3	8	7	8	7	4	1	-	-
Female	5	3	6	6	7	10	12	12	12	6	-	-
Black, both sexes	1	-	1	3	-	1	-	1	-	-	-	-
Male	1	-	1	2	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Kentucky												
All races, both sexes	18	13	21	24	15	28	24	24	12	2	1	-
Male	10	5	11	10	7	12	7	4	4	-	-	-
Female	8	8	10	14	8	16	17	20	8	2	1	-
White, both sexes	15	9	19	21	15	27	23	24	12	2	1	-
Male	9	4	11	7	7	12	6	4	4	-	-	-
Female	6	5	8	14	8	15	17	20	8	2	1	-
Black, both sexes	3	4	2	3	-	1	1	-	-	-	-	-
Male	1	1	-	3	-	-	1	-	-	-	-	-
Female	2	3	2	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	20	25	20	16	20	30	35	26	5	5	1	-
Male	10	9	11	10	10	14	16	7	1	1	-	-
Female	10	16	9	6	10	16	19	19	4	4	1	-
White, both sexes	13	16	16	7	16	23	33	17	3	5	1	-
Male	7	6	9	3	8	9	15	5	1	1	-	-
Female	6	10	7	4	8	14	18	12	2	4	1	-
Black, both sexes	7	9	4	9	4	6	2	9	2	-	-	-
Male	3	3	2	7	2	5	1	2	-	-	-	-
Female	4	6	2	2	2	1	1	7	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Maine												
All races, both sexes	5	4	2	8	7	9	7	3	6	4	-	-
Male	2	3	-	4	3	1	5	2	1	1	-	-
Female	3	1	2	4	4	8	2	1	5	3	-	-
White, both sexes	5	4	2	8	7	9	7	3	6	4	-	-
Male	2	3	-	4	3	1	5	2	1	1	-	-
Female	3	1	2	4	4	8	2	1	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	13	16	15	14	31	33	37	26	12	6	-	-
Male	4	7	2	9	11	14	12	11	1	-	-	-
Female	9	9	13	5	20	19	25	15	11	6	-	-
White, both sexes	9	7	9	10	19	22	25	19	10	4	-	-
Male	3	3	2	6	8	12	9	9	1	-	-	-
Female	6	4	7	4	11	10	16	10	9	4	-	-
Black, both sexes	4	9	6	4	12	11	11	7	2	2	-	-
Male	1	4	-	3	3	2	2	2	-	-	-	-
Female	3	5	6	1	9	9	9	5	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Massachusetts												
All races, both sexes	9	19	16	22	19	43	55	44	17	11	1	-
Male	3	8	8	10	7	22	10	11	1	4	1	-
Female	6	11	8	12	12	21	45	33	16	7	-	-
White, both sexes	7	17	16	20	17	41	53	43	17	10	1	-
Male	2	6	8	8	7	21	10	11	1	3	1	-
Female	5	11	8	12	10	20	43	32	16	7	-	-
Black, both sexes	1	2	-	1	2	1	1	1	-	1	-	-
Male	1	2	-	1	-	1	-	-	-	1	-	-
Female	-	-	-	-	2	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
Michigan												
All races, both sexes	21	29	24	27	38	63	61	47	33	5	1	-
Male	10	10	14	13	17	31	24	11	9	-	-	-
Female	11	19	10	14	21	32	37	36	24	5	1	-
White, both sexes	18	18	17	20	32	57	52	42	32	5	-	-
Male	9	9	10	10	13	28	22	10	8	-	-	-
Female	9	9	7	10	19	29	30	32	24	5	-	-
Black, both sexes	3	11	7	7	6	3	9	5	1	-	1	-
Male	1	1	4	3	4	1	2	1	1	-	-	-
Female	2	10	3	4	2	2	7	4	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Minnesota												
All races, both sexes	8	14	8	14	11	23	33	29	23	7	1	-
Male	5	6	3	6	3	8	11	9	10	5	-	-
Female	3	8	5	8	8	15	22	20	13	2	1	-
White, both sexes	8	12	6	12	11	22	28	29	23	7	1	-
Male	5	6	3	6	3	7	10	9	10	5	-	-
Female	3	6	3	6	8	15	18	20	13	2	1	-
Black, both sexes	-	2	1	1	-	-	5	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	2	1	1	-	-	4	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	11	11	18	12	21	17	18	11	11	4	-	-
Male	10	6	6	4	9	7	4	4	2	1	-	-
Female	1	5	12	8	12	10	14	7	9	3	-	-
White, both sexes	3	5	8	8	14	9	12	9	10	3	-	-
Male	3	3	3	3	8	3	2	3	1	1	-	-
Female	-	2	5	5	6	6	10	6	9	2	-	-
Black, both sexes	8	6	10	4	7	8	6	2	1	1	-	-
Male	7	3	3	1	1	4	2	1	1	-	-	-
Female	1	3	7	3	6	4	4	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Missouri												
All races, both sexes	14	25	32	32	29	31	46	34	11	5	1	-
Male	6	16	17	13	7	13	12	11	2	-	-	-
Female	8	9	15	19	22	18	34	23	9	5	1	-
White, both sexes	12	18	26	25	27	28	43	29	10	5	1	-
Male	5	12	16	12	7	12	11	10	2	-	-	-
Female	7	6	10	13	20	16	32	19	8	5	1	-
Black, both sexes	2	7	6	7	2	3	3	4	1	-	-	-
Male	1	4	1	1	-	1	1	1	-	-	-	-
Female	1	3	5	6	2	2	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	5	4	2	2	2	4	7	6	2	-	-	-
Male	3	2	2	1	1	1	1	2	1	-	-	-
Female	2	2	-	1	1	3	6	4	1	-	-	-
White, both sexes	5	4	2	2	2	4	7	6	2	-	-	-
Male	3	2	2	1	1	1	1	2	1	-	-	-
Female	2	2	-	1	1	3	6	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Nebraska												
All races, both sexes	3	6	4	5	5	5	8	9	5	4	2	-
Male	1	3	1	1	1	1	2	2	-	-	-	-
Female	2	3	3	4	4	4	6	7	5	4	2	-
White, both sexes	3	6	4	4	5	5	8	9	5	4	2	-
Male	1	3	1	1	1	1	2	2	-	-	-	-
Female	2	3	3	3	4	4	6	7	5	4	2	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	12	6	10	13	10	17	11	6	1	-	-
Male	4	6	2	8	5	4	7	4	2	-	-	-
Female	5	6	4	2	8	6	10	7	4	1	-	-
White, both sexes	8	10	5	8	11	10	14	11	5	-	-	-
Male	4	6	2	7	4	4	6	4	1	-	-	-
Female	4	4	3	1	7	6	8	7	4	-	-	-
Black, both sexes	1	2	1	1	1	-	2	-	-	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	1	2	1	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Hampshire												
All races, both sexes	3	5	5	7	6	7	7	5	4	-	1	-
Male	2	1	1	4	4	3	3	2	1	-	-	-
Female	1	4	4	3	2	4	4	3	3	-	1	-
White, both sexes	3	5	5	7	5	7	7	5	4	-	1	-
Male	2	1	1	4	3	3	3	2	1	-	-	-
Female	1	4	4	3	2	4	4	3	3	-	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	19	14	29	38	35	42	42	43	18	13	2	-
Male	10	8	9	17	13	11	7	10	1	1	-	-
Female	9	6	20	21	22	31	35	33	17	12	2	-
White, both sexes	14	10	20	31	28	38	30	38	17	13	1	-
Male	9	4	4	14	11	11	7	9	1	1	-	-
Female	5	6	16	17	17	27	23	29	16	12	1	-
Black, both sexes	4	4	9	6	7	2	9	4	-	-	1	-
Male	-	4	5	3	2	-	-	1	-	-	-	-
Female	4	-	4	3	5	2	9	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	3	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	3	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Mexico												
All races, both sexes	4	1	5	2	9	17	8	7	7	2	-	-
Male	1	1	4	-	6	7	-	2	1	1	-	-
Female	3	-	1	2	3	10	8	5	6	1	-	-
White, both sexes	4	1	3	2	9	16	8	7	7	2	-	-
Male	1	1	3	-	6	7	-	2	1	1	-	-
Female	3	-	-	2	3	9	8	5	6	1	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	30	37	34	41	57	68	82	83	38	21	2	-
Male	14	19	18	20	23	27	28	22	12	6	1	-
Female	16	18	16	21	34	41	54	61	26	15	1	-
White, both sexes	21	32	25	31	49	57	70	75	34	21	2	-
Male	11	17	15	15	22	24	21	20	11	6	1	-
Female	10	15	10	16	27	33	49	55	23	15	1	-
Black, both sexes	9	4	8	10	7	9	9	8	3	-	-	-
Male	3	1	3	5	1	1	6	2	-	-	-	-
Female	6	3	5	5	6	8	3	6	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	2	3	-	1	-	-	-
Male	-	1	-	-	-	2	1	-	1	-	-	-
Female	-	-	1	-	1	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
North Carolina												
All races, both sexes	28	35	34	50	52	56	56	59	28	6	2	-
Male	10	17	15	17	21	18	16	15	5	1	-	-
Female	18	18	19	33	31	38	40	44	23	5	2	-
White, both sexes	15	25	22	31	37	46	46	52	26	5	1	-
Male	7	11	11	10	17	17	13	12	5	1	-	-
Female	8	14	11	21	20	29	33	40	21	4	1	-
Black, both sexes	13	9	11	18	15	9	10	6	2	1	1	-
Male	3	6	3	6	4	1	3	3	-	-	-	-
Female	10	3	8	12	11	8	7	3	2	1	1	-
American Indian 1/, both sexes	-	1	1	1	-	1	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	4	1	2	-	3	4	5	3	-	-	-
Male	1	3	-	1	-	2	2	3	2	-	-	-
Female	1	1	1	1	-	1	2	2	1	-	-	-
White, both sexes	2	4	1	2	-	3	4	5	3	-	-	-
Male	1	3	-	1	-	2	2	3	2	-	-	-
Female	1	1	1	1	-	1	2	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Ohio												
All races, both sexes	19	31	26	42	51	68	63	51	30	9	-	-
Male	8	14	8	19	24	28	24	16	8	2	-	-
Female	11	17	18	23	27	40	39	35	22	7	-	-
White, both sexes	14	23	25	36	37	60	56	45	29	9	-	-
Male	4	10	8	16	17	23	23	13	7	2	-	-
Female	10	13	17	20	20	37	33	32	22	7	-	-
Black, both sexes	5	8	1	5	14	8	6	6	-	-	-	-
Male	4	4	-	2	7	5	1	3	-	-	-	-
Female	1	4	1	3	7	3	5	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	7	9	16	8	13	12	20	20	11	1	2	-
Male	6	1	8	4	7	3	6	10	-	-	1	-
Female	1	8	8	4	6	9	14	10	11	1	1	-
White, both sexes	5	7	13	6	12	8	19	19	10	-	2	-
Male	4	1	7	4	7	2	6	10	-	-	1	-
Female	1	6	6	2	5	6	13	9	10	-	1	-
Black, both sexes	1	1	1	2	-	3	1	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	2	-	2	1	1	-	-	-	-
American Indian 1/, both sexes	1	1	2	-	-	-	-	-	1	1	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Oregon												
All races, both sexes	9	9	11	10	14	20	21	18	18	1	-	-
Male	3	4	3	6	8	6	8	8	3	-	-	-
Female	6	5	8	4	6	14	13	10	15	1	-	-
White, both sexes	8	9	7	10	13	18	20	18	18	1	-	-
Male	2	4	2	6	8	5	8	8	3	-	-	-
Female	6	5	5	4	5	13	12	10	15	1	-	-
Black, both sexes	1	-	3	-	1	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	33	44	52	51	56	93	104	114	53	12	6	-
Male	14	21	20	17	18	33	40	32	14	-	-	-
Female	19	23	32	34	38	60	64	82	39	12	6	-
White, both sexes	27	36	42	42	47	80	97	101	49	11	6	-
Male	10	18	18	14	17	25	38	29	14	-	-	-
Female	17	18	24	28	30	55	59	72	35	11	6	-
Black, both sexes	6	8	10	8	9	13	7	13	4	-	-	-
Male	4	3	2	3	1	8	2	3	-	-	-	-
Female	2	5	8	5	8	5	5	10	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Rhode Island												
All races, both sexes	3	1	2	4	5	7	6	5	6	1	-	-
Male	1	1	-	3	2	3	1	2	2	-	-	-
Female	2	-	2	1	3	4	5	3	4	1	-	-
White, both sexes	2	1	1	4	5	6	5	5	6	1	-	-
Male	1	1	-	3	2	2	1	2	2	-	-	-
Female	1	-	1	1	3	4	4	3	4	1	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	19	14	15	19	41	30	22	28	16	6	-	-
Male	8	6	6	9	16	13	9	8	4	1	-	-
Female	11	8	9	10	25	17	13	20	12	5	-	-
White, both sexes	11	9	8	10	28	22	15	23	15	6	-	-
Male	5	3	3	6	10	11	6	6	3	1	-	-
Female	6	6	5	4	18	11	9	17	12	5	-	-
Black, both sexes	8	5	7	8	13	8	7	5	1	-	-	-
Male	3	3	3	3	6	2	3	2	1	-	-	-
Female	5	2	4	5	7	6	4	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
South Dakota												
All races, both sexes	3	1	1	4	2	2	3	2	1	2	-	-
Male	1	-	1	2	1	1	-	-	-	2	-	-
Female	2	1	-	2	1	1	3	2	1	-	-	-
White, both sexes	2	1	1	4	2	2	3	2	1	2	-	-
Male	-	-	1	2	1	1	-	-	-	2	-	-
Female	2	1	-	2	1	1	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	17	32	22	28	28	39	45	34	22	4	1	-
Male	5	11	10	16	12	11	18	12	7	-	-	-
Female	12	21	12	12	16	28	27	22	15	4	1	-
White, both sexes	10	22	18	23	22	37	39	28	18	4	-	-
Male	3	7	7	12	10	11	14	9	6	-	-	-
Female	7	15	11	11	12	26	25	19	12	4	-	-
Black, both sexes	7	10	4	5	6	2	6	6	4	-	1	-
Male	2	4	3	4	2	-	4	3	1	-	-	-
Female	5	6	1	1	4	2	2	3	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Texas												
All races, both sexes	59	66	65	71	83	109	97	110	60	17	4	-
Male	31	34	35	25	45	46	40	32	14	2	1	-
Female	28	32	30	46	38	63	57	78	46	15	3	-
White, both sexes	44	51	45	59	73	86	90	100	51	13	1	-
Male	25	28	27	21	40	39	36	29	10	2	-	-
Female	19	23	18	38	33	47	54	71	41	11	1	-
Black, both sexes	15	15	19	11	10	22	7	10	8	4	3	-
Male	6	6	7	4	5	7	4	3	3	-	1	-
Female	9	9	12	7	5	15	3	7	5	4	2	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	7	8	11	7	12	12	15	10	5	1	-	-
Male	4	3	4	4	6	5	7	2	3	-	-	-
Female	3	5	7	3	6	7	8	8	2	1	-	-
White, both sexes	7	8	11	6	12	11	15	10	5	1	-	-
Male	4	3	4	3	6	4	7	2	3	-	-	-
Female	3	5	7	3	6	7	8	8	2	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Vermont												
All races, both sexes	1	3	1	1	1	2	3	3	2	-	-	-
Male	1	1	1	-	-	-	1	1	-	-	-	-
Female	-	2	-	1	1	2	2	2	2	-	-	-
White, both sexes	1	3	1	1	1	2	3	3	2	-	-	-
Male	1	1	1	-	-	-	1	1	-	-	-	-
Female	-	2	-	1	1	2	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	30	27	31	27	34	50	63	45	24	3	1	-
Male	14	17	14	11	14	19	19	17	9	-	-	-
Female	16	10	17	16	20	31	44	28	15	3	1	-
White, both sexes	23	16	25	17	25	38	54	42	22	1	-	-
Male	11	11	10	8	10	18	18	16	8	-	-	-
Female	12	5	15	9	15	20	36	26	14	1	-	-
Black, both sexes	7	11	6	10	9	11	9	3	2	2	1	-
Male	3	6	4	3	4	-	1	1	1	-	-	-
Female	4	5	2	7	5	11	8	2	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Washington												
All races, both sexes	13	17	13	22	28	30	37	36	22	4	-	-
Male	2	6	6	10	13	13	9	11	5	-	-	-
Female	11	11	7	12	15	17	28	25	17	4	-	-
White, both sexes	12	14	13	19	25	29	36	36	21	4	-	-
Male	2	5	6	8	13	12	9	11	5	-	-	-
Female	10	9	7	11	12	17	27	25	16	4	-	-
Black, both sexes	1	1	-	2	3	-	-	-	1	-	-	-
Male	-	1	-	2	-	-	-	-	-	-	-	-
Female	1	-	-	-	3	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
West Virginia												
All races, both sexes	7	11	10	10	9	17	15	12	4	3	1	-
Male	3	3	3	5	5	5	5	4	-	1	-	-
Female	4	8	7	5	4	12	10	8	4	2	1	-
White, both sexes	6	11	10	10	9	16	13	11	4	3	1	-
Male	2	3	3	5	5	4	4	4	-	1	-	-
Female	4	8	7	5	4	12	9	7	4	2	1	-
Black, both sexes	1	-	-	-	-	1	2	1	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Wisconsin												
All races, both sexes	20	13	14	12	30	39	42	52	26	7	3	-
Male	9	5	8	7	15	19	19	19	3	1	-	-
Female	11	8	6	5	15	20	23	33	23	6	3	-
White, both sexes	16	8	13	12	28	39	41	50	25	7	3	-
Male	7	4	8	7	15	19	19	18	3	1	-	-
Female	9	4	5	5	13	20	22	32	22	6	3	-
Black, both sexes	4	5	1	-	2	-	1	2	1	-	-	-
Male	2	1	-	-	-	-	-	1	-	-	-	-
Female	2	4	1	-	2	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	1	2	4	5	6	6	7	2	1	-	-
Male	3	1	2	3	4	5	3	-	-	-	-	-
Female	-	-	-	1	1	1	3	7	2	1	-	-
White, both sexes	3	1	2	4	3	5	5	7	1	1	-	-
Male	3	1	2	3	3	4	2	-	-	-	-	-
Female	-	-	-	1	-	1	3	7	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
United States												
All races, both sexes	6,924	2	-	2	4	17	43	83	129	176	276	339
Male	2,946	-	-	1	3	5	18	35	49	81	124	157
Female	3,978	2	-	1	1	12	25	48	80	95	152	182
White, both sexes	5,644	2	-	2	2	11	28	57	76	109	186	242
Male	2,397	-	-	1	2	2	11	23	24	54	83	112
Female	3,247	2	-	1	-	9	17	34	52	55	103	130
Black, both sexes	1,193	-	-	-	2	6	15	26	50	60	84	90
Male	515	-	-	-	1	3	7	12	25	24	39	41
Female	678	-	-	-	1	3	8	14	25	36	45	49
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	1	4	2	4
Male	14	-	-	-	-	-	-	-	-	1	1	3
Female	19	-	-	-	-	-	-	-	1	3	1	1
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	-	-	2	3	4	3
Male	20	-	-	-	-	-	-	-	-	2	1	1
Female	34	-	-	-	-	-	-	-	2	1	3	2
Alabama												
All races, both sexes	169	-	-	-	-	-	2	3	5	2	9	7
Male	81	-	-	-	-	-	1	1	1	1	6	3
Female	88	-	-	-	-	-	1	2	4	1	3	4
White, both sexes	123	-	-	-	-	-	1	2	2	-	7	3
Male	59	-	-	-	-	-	1	-	-	-	5	1
Female	64	-	-	-	-	-	-	2	2	-	2	2
Black, both sexes	45	-	-	-	-	-	1	1	3	2	2	4
Male	22	-	-	-	-	-	-	1	1	1	1	2
Female	23	-	-	-	-	-	1	-	2	1	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Alaska												
All races, both sexes	9	-	-	-	-	-	-	-	1	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Arizona												
All races, both sexes	98	-	-	-	-	1	-	1	1	2	4	7
Male	47	-	-	-	-	1	-	1	1	1	3	1
Female	51	-	-	-	-	-	-	-	-	1	1	6
White, both sexes	94	-	-	-	-	-	-	1	1	1	4	7
Male	45	-	-	-	-	-	-	1	1	-	3	1
Female	49	-	-	-	-	-	-	-	-	1	1	6
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Arkansas												
All races, both sexes	142	-	-	-	-	1	-	1	1	8	6	6
Male	61	-	-	-	-	-	-	-	1	6	2	5
Female	81	-	-	-	-	1	-	1	-	2	4	1
White, both sexes	116	-	-	-	-	1	-	-	1	6	6	5
Male	51	-	-	-	-	-	-	-	1	5	2	4
Female	65	-	-	-	-	1	-	-	-	1	4	1
Black, both sexes	24	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	2	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	282	-	-	-	-	1	-	-	3	7	15	14
Male	114	-	-	-	-	1	-	-	-	2	6	5
Female	168	-	-	-	-	-	-	-	3	5	9	9
White, both sexes	249	-	-	-	-	-	-	-	2	5	10	13
Male	108	-	-	-	-	-	-	-	-	2	5	5
Female	141	-	-	-	-	-	-	-	2	3	5	8
Black, both sexes	23	-	-	-	-	1	-	-	-	1	3	1
Male	6	-	-	-	-	1	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Colorado												
All races, both sexes	99	-	-	-	-	-	1	2	2	1	1	3
Male	35	-	-	-	-	-	-	1	-	1	1	1
Female	64	-	-	-	-	-	1	1	2	-	-	2
White, both sexes	90	-	-	-	-	-	1	2	2	1	1	2
Male	32	-	-	-	-	-	-	1	-	1	1	1
Female	58	-	-	-	-	-	1	1	2	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Connecticut												
All races, both sexes	88	-	-	-	-	-	1	3	1	1	3	5
Male	35	-	-	-	-	-	1	2	-	1	2	3
Female	53	-	-	-	-	-	-	1	1	-	1	2
White, both sexes	79	-	-	-	-	-	-	2	-	1	2	4
Male	32	-	-	-	-	-	-	1	-	1	2	3
Female	47	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	9	-	-	-	-	-	1	1	1	-	1	1
Male	3	-	-	-	-	-	1	1	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Delaware												
All races, both sexes	26	-	-	-	-	1	-	-	-	-	1	3
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	1	-	-	-	-	1	3
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Florida												
All races, both sexes	369	-	-	-	-	1	1	4	1	10	17	20
Male	163	-	-	-	-	-	-	2	-	5	12	8
Female	206	-	-	-	-	1	1	2	1	5	5	12
White, both sexes	298	-	-	-	-	-	-	4	-	8	7	13
Male	130	-	-	-	-	-	-	2	-	5	4	4
Female	168	-	-	-	-	-	-	2	-	3	3	9
Black, both sexes	70	-	-	-	-	1	1	-	1	2	10	6
Male	33	-	-	-	-	-	-	-	-	-	8	4
Female	37	-	-	-	-	1	1	-	1	2	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	196	-	-	-	-	-	4	5	8	5	10	8
Male	86	-	-	-	-	-	-	2	5	2	4	4
Female	110	-	-	-	-	-	4	3	3	3	6	4
White, both sexes	127	-	-	-	-	-	3	4	3	2	2	4
Male	52	-	-	-	-	-	-	2	1	1	-	2
Female	75	-	-	-	-	-	3	2	2	1	2	2
Black, both sexes	69	-	-	-	-	-	1	1	5	3	8	4
Male	34	-	-	-	-	-	-	-	4	1	4	2
Female	35	-	-	-	-	-	1	1	1	2	4	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Hawaii												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	46	-	-	-	-	-	2	1	2	1	2	3
Male	22	-	-	-	-	-	1	-	1	-	-	1
Female	24	-	-	-	-	-	1	1	1	1	2	2
White, both sexes	45	-	-	-	-	-	2	1	2	1	2	3
Male	22	-	-	-	-	-	1	-	1	-	-	1
Female	23	-	-	-	-	-	1	1	1	1	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Illinois												
All races, both sexes	374	-	-	-	-	-	2	3	5	7	9	14
Male	150	-	-	-	-	-	2	-	3	3	2	5
Female	224	-	-	-	-	-	-	3	2	4	7	9
White, both sexes	292	-	-	-	-	-	1	2	3	3	7	11
Male	119	-	-	-	-	-	1	-	2	1	1	3
Female	173	-	-	-	-	-	-	2	1	2	6	8
Black, both sexes	82	-	-	-	-	-	1	1	2	4	2	3
Male	31	-	-	-	-	-	1	-	1	2	1	2
Female	51	-	-	-	-	-	-	1	1	2	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	238	1	-	-	-	-	1	4	4	6	12	19
Male	91	-	-	-	-	-	-	2	1	3	6	8
Female	147	1	-	-	-	-	1	2	3	3	6	11
White, both sexes	215	1	-	-	-	-	-	4	4	5	10	15
Male	79	-	-	-	-	-	-	2	1	3	4	6
Female	136	1	-	-	-	-	-	2	3	2	6	9
Black, both sexes	23	-	-	-	-	-	1	-	-	1	2	4
Male	12	-	-	-	-	-	-	-	-	-	2	2
Female	11	-	-	-	-	-	1	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Iowa												
All races, both sexes	101	-	-	-	-	-	-	2	1	7	1	7
Male	45	-	-	-	-	-	-	2	1	4	1	3
Female	56	-	-	-	-	-	-	-	-	3	-	4
White, both sexes	98	-	-	-	-	-	-	2	1	6	1	7
Male	44	-	-	-	-	-	-	2	1	4	1	3
Female	54	-	-	-	-	-	-	-	-	2	-	4
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	89	-	-	-	-	1	-	1	1	2	2	6
Male	42	-	-	-	-	-	-	1	1	-	-	4
Female	47	-	-	-	-	1	-	-	-	2	2	2
White, both sexes	83	-	-	-	-	1	-	1	1	2	2	5
Male	38	-	-	-	-	-	-	1	1	-	-	3
Female	45	-	-	-	-	1	-	-	-	2	2	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Kentucky												
All races, both sexes	147	-	-	1	1	2	2	3	4	6	7	6
Male	64	-	-	-	1	-	2	2	1	4	3	2
Female	83	-	-	1	-	2	-	1	3	2	4	4
White, both sexes	134	-	-	1	1	1	1	2	3	3	7	6
Male	54	-	-	-	1	-	1	1	-	2	3	2
Female	80	-	-	1	-	1	-	1	3	1	4	4
Black, both sexes	13	-	-	-	-	1	1	1	1	3	-	-
Male	10	-	-	-	-	-	1	1	1	2	-	-
Female	3	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	173	-	-	-	-	-	2	3	5	6	7	8
Male	79	-	-	-	-	-	1	-	1	3	3	4
Female	94	-	-	-	-	-	1	3	4	3	4	4
White, both sexes	125	-	-	-	-	-	2	-	1	3	4	6
Male	53	-	-	-	-	-	1	-	-	1	1	2
Female	72	-	-	-	-	-	1	-	1	2	3	4
Black, both sexes	47	-	-	-	-	-	-	3	4	3	3	2
Male	26	-	-	-	-	-	-	-	1	2	2	2
Female	21	-	-	-	-	-	-	3	3	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Maine												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	139	1	-	-	-	-	1	3	6	2	9	6
Male	45	-	-	-	-	-	-	2	1	1	2	1
Female	94	1	-	-	-	-	1	1	5	1	7	5
White, both sexes	82	1	-	-	-	-	1	1	3	-	4	2
Male	25	-	-	-	-	-	-	1	-	-	-	-
Female	57	1	-	-	-	-	1	-	3	-	4	2
Black, both sexes	56	-	-	-	-	-	-	2	3	2	5	4
Male	19	-	-	-	-	-	-	1	1	1	2	1
Female	37	-	-	-	-	-	-	1	2	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Massachusetts												
All races, both sexes	162	-	-	-	-	-	-	-	1	4	8	4
Male	60	-	-	-	-	-	-	-	-	3	4	1
Female	102	-	-	-	-	-	-	-	1	1	4	3
White, both sexes	151	-	-	-	-	-	-	-	1	3	7	2
Male	54	-	-	-	-	-	-	-	-	3	4	1
Female	97	-	-	-	-	-	-	-	1	-	3	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	213	-	-	-	1	-	-	2	3	1	6	12
Male	101	-	-	-	1	-	-	1	3	-	3	6
Female	112	-	-	-	-	-	-	1	-	1	3	6
White, both sexes	173	-	-	-	1	-	-	1	2	1	3	7
Male	84	-	-	-	1	-	-	1	2	-	1	3
Female	89	-	-	-	-	-	-	-	-	1	2	4
Black, both sexes	37	-	-	-	-	-	-	1	1	-	3	5
Male	15	-	-	-	-	-	-	-	1	-	2	3
Female	22	-	-	-	-	-	-	1	-	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Minnesota												
All races, both sexes	76	-	-	-	-	-	-	1	1	2	3	9
Male	38	-	-	-	-	-	-	-	1	-	1	6
Female	38	-	-	-	-	-	-	1	-	2	2	3
White, both sexes	71	-	-	-	-	-	-	1	-	2	3	9
Male	37	-	-	-	-	-	-	-	-	-	1	6
Female	34	-	-	-	-	-	-	1	-	2	2	3
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	109	-	-	-	-	-	3	3	5	3	5	7
Male	52	-	-	-	-	-	1	1	1	1	3	5
Female	57	-	-	-	-	-	2	2	4	2	2	2
White, both sexes	50	-	-	-	-	-	1	-	1	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	1	-	1	-	-	1
Black, both sexes	59	-	-	-	-	-	2	3	4	3	4	5
Male	29	-	-	-	-	-	1	1	1	1	2	4
Female	30	-	-	-	-	-	1	2	3	2	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Missouri												
All races, both sexes	186	-	-	-	-	-	-	1	3	6	6	6
Male	83	-	-	-	-	-	-	1	2	1	5	3
Female	103	-	-	-	-	-	-	-	1	5	1	3
White, both sexes	155	-	-	-	-	-	-	-	3	5	5	3
Male	72	-	-	-	-	-	-	-	2	1	4	1
Female	83	-	-	-	-	-	-	-	1	4	1	2
Black, both sexes	30	-	-	-	-	-	-	1	-	1	1	3
Male	11	-	-	-	-	-	-	1	-	-	1	2
Female	19	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Nebraska												
All races, both sexes	32	-	-	-	-	-	2	-	-	1	2	3
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	23	-	-	-	-	-	2	-	-	-	2	2
White, both sexes	31	-	-	-	-	-	1	-	-	1	2	3
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	22	-	-	-	-	-	1	-	-	-	2	2
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	68	-	-	-	-	-	-	1	1	-	3	6
Male	29	-	-	-	-	-	-	-	-	-	-	3
Female	39	-	-	-	-	-	-	1	1	-	3	3
White, both sexes	59	-	-	-	-	-	-	1	1	-	2	5
Male	26	-	-	-	-	-	-	-	-	-	-	3
Female	33	-	-	-	-	-	-	1	1	-	2	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
New Hampshire												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	210	-	-	-	-	1	3	-	6	5	10	9
Male	77	-	-	-	-	-	2	-	-	3	7	6
Female	133	-	-	-	-	1	1	-	6	2	3	3
White, both sexes	166	-	-	-	-	1	2	-	5	-	7	7
Male	59	-	-	-	-	-	1	-	-	-	6	5
Female	107	-	-	-	-	1	1	-	5	-	1	2
Black, both sexes	37	-	-	-	-	-	1	-	1	5	3	2
Male	17	-	-	-	-	-	1	-	-	3	1	1
Female	20	-	-	-	-	-	-	-	1	2	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
New Mexico												
All races, both sexes	35	-	-	-	-	-	-	2	1	1	1	4
Male	18	-	-	-	-	-	-	2	-	-	-	3
Female	17	-	-	-	-	-	-	-	1	1	1	1
White, both sexes	31	-	-	-	-	-	-	2	1	1	1	2
Male	16	-	-	-	-	-	-	2	-	-	-	1
Female	15	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	263	-	-	-	-	-	-	1	1	9	8	11
Male	114	-	-	-	-	-	-	1	-	3	3	4
Female	149	-	-	-	-	-	-	-	1	6	5	7
White, both sexes	215	-	-	-	-	-	-	1	1	4	7	7
Male	97	-	-	-	-	-	-	1	-	2	3	4
Female	118	-	-	-	-	-	-	-	1	2	4	3
Black, both sexes	43	-	-	-	-	-	-	-	-	4	1	4
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	29	-	-	-	-	-	-	-	-	3	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
North Carolina												
All races, both sexes	289	-	-	-	-	1	4	3	7	13	14	17
Male	114	-	-	-	-	1	2	2	4	4	8	10
Female	175	-	-	-	-	-	2	1	3	9	6	7
White, both sexes	197	-	-	-	-	-	3	2	2	6	6	7
Male	75	-	-	-	-	-	2	1	1	3	4	5
Female	122	-	-	-	-	-	1	1	1	3	2	2
Black, both sexes	89	-	-	-	-	1	1	1	5	7	8	9
Male	38	-	-	-	-	1	-	1	3	1	4	5
Female	51	-	-	-	-	-	1	-	2	6	4	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	20	-	-	-	-	-	1	-	1	1	2	-
Male	11	-	-	-	-	-	1	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	18	-	-	-	-	-	1	-	-	1	1	-
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Ohio												
All races, both sexes	237	-	-	-	-	-	1	3	6	8	12	18
Male	109	-	-	-	-	-	1	1	3	3	7	7
Female	128	-	-	-	-	-	-	2	3	5	5	11
White, both sexes	196	-	-	-	-	-	-	2	5	7	8	15
Male	88	-	-	-	-	-	-	1	3	3	5	6
Female	108	-	-	-	-	-	-	1	2	4	3	9
Black, both sexes	39	-	-	-	-	-	1	1	1	1	4	3
Male	20	-	-	-	-	-	1	-	-	-	2	1
Female	19	-	-	-	-	-	-	1	1	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	91	-	-	-	-	-	1	-	-	3	3	6
Male	41	-	-	-	-	-	1	-	-	1	1	4
Female	50	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	78	-	-	-	-	-	1	-	-	1	3	6
Male	38	-	-	-	-	-	1	-	-	1	1	4
Female	40	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Oregon												
All races, both sexes	68	-	-	-	-	-	1	2	2	2	-	3
Male	32	-	-	-	-	-	-	-	-	1	-	2
Female	36	-	-	-	-	-	1	2	2	1	-	1
White, both sexes	63	-	-	-	-	-	1	2	1	2	-	2
Male	29	-	-	-	-	-	-	-	-	1	-	1
Female	34	-	-	-	-	-	1	2	1	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	389	-	-	-	-	-	-	5	7	6	14	16
Male	154	-	-	-	-	-	-	2	1	4	8	8
Female	235	-	-	-	-	-	-	3	6	2	6	8
White, both sexes	329	-	-	-	-	-	-	3	4	4	11	10
Male	128	-	-	-	-	-	-	1	-	3	6	5
Female	201	-	-	-	-	-	-	2	4	1	5	5
Black, both sexes	58	-	-	-	-	-	-	2	3	2	2	6
Male	25	-	-	-	-	-	-	1	1	1	1	3
Female	33	-	-	-	-	-	-	1	2	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Rhode Island												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	148	-	-	-	-	1	1	3	5	4	7	6
Male	55	-	-	-	-	1	-	1	4	2	1	2
Female	93	-	-	-	-	-	1	2	1	2	6	4
White, both sexes	91	-	-	-	-	1	1	1	1	1	2	3
Male	33	-	-	-	-	1	-	-	1	-	-	2
Female	58	-	-	-	-	-	1	1	-	1	2	1
Black, both sexes	57	-	-	-	-	-	-	2	4	3	5	3
Male	22	-	-	-	-	-	-	1	3	2	1	-
Female	35	-	-	-	-	-	-	1	1	1	4	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
South Dakota												
All races, both sexes	16	-	-	-	-	-	-	-	-	1	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	15	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	204	-	-	-	-	1	1	1	4	4	6	6
Male	83	-	-	-	-	-	1	1	1	2	3	2
Female	121	-	-	-	-	1	-	-	3	2	3	4
White, both sexes	160	-	-	-	-	1	-	-	4	3	5	2
Male	59	-	-	-	-	-	-	-	1	1	2	-
Female	101	-	-	-	-	1	-	-	3	2	3	2
Black, both sexes	44	-	-	-	-	-	1	1	-	1	1	4
Male	24	-	-	-	-	-	1	1	-	1	1	2
Female	20	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Texas												
All races, both sexes	547	-	-	1	2	2	1	8	7	15	26	25
Male	248	-	-	1	1	1	-	4	5	8	8	11
Female	299	-	-	-	1	1	1	4	2	7	18	14
White, both sexes	445	-	-	1	-	2	1	6	4	9	15	21
Male	200	-	-	1	-	1	-	2	2	4	6	9
Female	245	-	-	-	-	1	1	4	2	5	9	12
Black, both sexes	98	-	-	-	2	-	-	2	3	6	10	4
Male	46	-	-	-	1	-	-	2	3	4	2	2
Female	52	-	-	-	1	-	-	-	-	2	8	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	57	-	-	-	-	1	-	3	-	-	2	2
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	1	-	3	-	-	2	2
White, both sexes	55	-	-	-	-	1	-	3	-	-	2	2
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	1	-	3	-	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	227	-	-	-	-	-	2	1	8	6	9	10
Male	102	-	-	-	-	-	1	-	3	4	6	4
Female	125	-	-	-	-	-	1	1	5	2	3	6
White, both sexes	173	-	-	-	-	-	1	-	4	3	8	6
Male	83	-	-	-	-	-	1	-	1	2	6	3
Female	90	-	-	-	-	-	-	-	3	1	2	3
Black, both sexes	52	-	-	-	-	-	1	1	4	2	1	3
Male	17	-	-	-	-	-	-	-	2	1	-	-
Female	35	-	-	-	-	-	1	1	2	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Washington												
All races, both sexes	128	-	-	-	-	-	1	2	6	3	2	4
Male	45	-	-	-	-	-	-	-	1	-	-	2
Female	83	-	-	-	-	-	1	2	5	3	2	2
White, both sexes	119	-	-	-	-	-	1	2	6	2	2	3
Male	42	-	-	-	-	-	-	-	1	-	-	2
Female	77	-	-	-	-	-	1	2	5	2	2	1
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	61	-	-	-	-	1	-	1	1	1	1	3
Male	25	-	-	-	-	-	-	-	1	1	1	2
Female	36	-	-	-	-	1	-	1	-	-	-	1
White, both sexes	57	-	-	-	-	1	-	1	1	1	1	3
Male	23	-	-	-	-	-	-	-	1	1	1	2
Female	34	-	-	-	-	1	-	1	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Wisconsin												
All races, both sexes	157	-	-	-	-	1	2	-	2	2	8	6
Male	63	-	-	-	-	-	-	-	1	2	1	3
Female	94	-	-	-	-	1	2	-	1	-	7	3
White, both sexes	141	-	-	-	-	1	2	-	-	2	4	5
Male	59	-	-	-	-	-	-	-	-	2	-	3
Female	82	-	-	-	-	1	2	-	-	-	4	2
Black, both sexes	16	-	-	-	-	-	-	-	2	-	4	1
Male	4	-	-	-	-	-	-	-	1	-	1	-
Female	12	-	-	-	-	-	-	-	1	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	24	-	-	-	-	-	-	1	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	1	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
United States												
All races, both sexes	460	508	521	526	643	862	920	811	437	139	26	-
Male	241	259	252	272	309	370	360	258	121	29	2	-
Female	219	249	269	254	334	492	560	553	316	110	24	-
White, both sexes	368	402	408	414	538	728	798	728	401	124	20	-
Male	193	211	202	211	268	319	317	229	108	26	1	-
Female	175	191	206	203	270	409	481	499	293	98	19	-
Black, both sexes	85	101	108	104	103	119	112	79	30	13	6	-
Male	44	47	47	58	40	43	41	29	10	3	1	-
Female	41	54	61	46	63	76	71	50	20	10	5	-
American Indian 1/, both sexes	3	1	5	2	1	5	1	1	2	1	-	-
Male	2	-	3	-	1	2	-	-	1	-	-	-
Female	1	1	2	2	-	3	1	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	4	4	-	6	1	10	9	3	4	1	-	-
Male	2	1	-	3	-	6	2	-	2	-	-	-
Female	2	3	-	3	1	4	7	3	2	1	-	-
Alabama												
All races, both sexes	13	9	16	14	17	16	28	19	7	2	-	-
Male	9	3	10	7	10	6	13	6	3	1	-	-
Female	4	6	6	7	7	10	15	13	4	1	-	-
White, both sexes	10	5	11	11	11	14	21	18	5	2	-	-
Male	7	2	7	6	5	6	11	6	1	1	-	-
Female	3	3	4	5	6	8	10	12	4	1	-	-
Black, both sexes	3	4	5	2	6	2	7	1	2	-	-	-
Male	2	1	3	1	5	-	2	-	2	-	-	-
Female	1	3	2	1	1	2	5	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Alaska												
All races, both sexes	1	-	-	1	-	2	-	-	2	-	-	-
Male	1	-	-	1	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	1	-	2	-	-	2	-	-	-
Male	1	-	-	1	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	5	6	7	15	12	14	10	11	-	-	-
Male	1	2	2	4	7	7	9	4	3	-	-	-
Female	1	3	4	3	8	5	5	6	8	-	-	-
White, both sexes	2	5	6	6	15	11	14	10	11	-	-	-
Male	1	2	2	4	7	7	9	4	3	-	-	-
Female	1	3	4	2	8	4	5	6	8	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Arkansas												
All races, both sexes	12	14	7	13	13	16	21	13	7	2	1	-
Male	8	8	3	5	9	6	5	-	3	-	-	-
Female	4	6	4	8	4	10	16	13	4	2	1	-
White, both sexes	11	13	5	11	7	13	16	13	6	2	-	-
Male	8	7	1	5	5	6	4	-	3	-	-	-
Female	3	6	4	6	2	7	12	13	3	2	-	-
Black, both sexes	1	1	2	2	6	3	5	-	1	-	1	-
Male	-	1	2	-	4	-	1	-	-	-	-	-
Female	1	-	-	2	2	3	4	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	21	21	22	18	26	35	37	28	28	4	2	-
Male	10	11	8	11	12	13	16	10	9	-	-	-
Female	11	10	14	7	14	22	21	18	19	4	2	-
White, both sexes	19	18	20	16	23	33	33	25	26	4	2	-
Male	10	11	8	9	11	13	15	10	9	-	-	-
Female	9	7	12	7	12	20	18	15	17	4	2	-
Black, both sexes	2	2	2	2	2	1	3	2	1	-	-	-
Male	-	-	-	2	1	-	1	-	-	-	-	-
Female	2	2	2	-	1	1	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Colorado												
All races, both sexes	11	9	9	7	11	6	13	10	10	3	-	-
Male	5	3	3	3	4	2	7	2	2	-	-	-
Female	6	6	6	4	7	4	6	8	8	3	-	-
White, both sexes	10	9	8	5	10	6	11	10	9	3	-	-
Male	5	3	3	1	4	2	6	2	2	-	-	-
Female	5	6	5	4	6	4	5	8	7	3	-	-
Black, both sexes	-	-	1	2	1	-	1	-	1	-	-	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	5	4	5	3	5	15	15	14	6	2	-	-
Male	2	3	2	2	3	3	6	3	2	-	-	-
Female	3	1	3	1	2	12	9	11	4	2	-	-
White, both sexes	4	4	5	3	4	15	13	14	6	2	-	-
Male	1	3	2	2	3	3	6	3	2	-	-	-
Female	3	1	3	1	1	12	7	11	4	2	-	-
Black, both sexes	1	-	-	-	1	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Delaware												
All races, both sexes	1	2	4	-	-	7	4	2	1	-	-	-
Male	-	1	1	-	-	4	3	1	-	-	-	-
Female	1	1	3	-	-	3	1	1	1	-	-	-
White, both sexes	1	2	2	-	-	7	4	2	1	-	-	-
Male	-	1	1	-	-	4	3	1	-	-	-	-
Female	1	1	1	-	-	3	1	1	1	-	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	2	1	2	-	1	1	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	1	-	2	1	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	2	1	2	-	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	2	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Florida												
All races, both sexes	13	27	23	29	29	57	51	50	24	11	1	-
Male	6	12	15	13	13	23	20	20	11	3	-	-
Female	7	15	8	16	16	34	31	30	13	8	1	-
White, both sexes	10	20	19	22	24	46	46	46	21	11	1	-
Male	5	12	12	8	11	18	18	18	10	3	-	-
Female	5	8	7	14	13	28	28	28	11	8	1	-
Black, both sexes	3	7	4	7	5	11	5	4	3	-	-	-
Male	1	-	3	5	2	5	2	2	1	-	-	-
Female	2	7	1	2	3	6	3	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	16	10	20	11	25	19	27	17	7	4	-	-
Male	7	9	8	4	9	10	10	8	4	-	-	-
Female	9	1	12	7	16	9	17	9	3	4	-	-
White, both sexes	12	4	16	4	17	13	21	13	5	4	-	-
Male	5	3	4	2	8	8	8	6	2	-	-	-
Female	7	1	12	2	9	5	13	7	3	4	-	-
Black, both sexes	4	6	4	7	8	6	6	4	2	-	-	-
Male	2	6	4	2	1	2	2	2	2	-	-	-
Female	2	-	-	5	7	4	4	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Hawaii												
All races, both sexes	2	1	1	2	-	4	2	-	1	-	-	-
Male	1	-	1	1	-	2	-	-	-	-	-	-
Female	1	1	-	1	-	2	2	-	1	-	-	-
White, both sexes	-	-	1	-	-	1	2	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	2	-	3	-	-	-	-	-	-
Male	1	-	-	1	-	2	-	-	-	-	-	-
Female	1	1	-	1	-	1	-	-	-	-	-	-
Idaho												
All races, both sexes	8	3	2	6	3	9	1	2	-	1	-	-
Male	5	-	1	5	2	5	-	1	-	-	-	-
Female	3	3	1	1	1	4	1	1	-	1	-	-
White, both sexes	7	3	2	6	3	9	1	2	-	1	-	-
Male	5	-	1	5	2	5	-	1	-	-	-	-
Female	2	3	1	1	1	4	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Illinois												
All races, both sexes	29	30	30	31	38	46	43	54	21	9	3	-
Male	18	15	10	21	13	17	15	16	6	3	1	-
Female	11	15	20	10	25	29	28	38	15	6	2	-
White, both sexes	24	22	19	24	28	35	39	47	20	5	2	-
Male	16	12	7	17	11	14	12	14	6	1	1	-
Female	8	10	12	7	17	21	27	33	14	4	1	-
Black, both sexes	5	8	11	7	10	11	4	7	1	4	1	-
Male	2	3	3	4	2	3	3	2	-	2	-	-
Female	3	5	8	3	8	8	1	5	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	20	18	19	21	17	26	26	31	6	6	1	-
Male	10	10	6	11	7	10	7	8	1	1	-	-
Female	10	8	13	10	10	16	19	23	5	5	1	-
White, both sexes	18	16	18	18	16	24	24	29	6	6	1	-
Male	8	9	6	8	7	9	6	8	1	1	-	-
Female	10	7	12	10	9	15	18	21	5	5	1	-
Black, both sexes	2	2	1	3	1	2	2	2	-	-	-	-
Male	2	1	-	3	-	1	1	-	-	-	-	-
Female	-	1	1	-	1	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Iowa												
All races, both sexes	9	9	9	7	5	7	16	13	5	3	-	-
Male	6	5	4	3	3	4	4	5	-	-	-	-
Female	3	4	5	4	2	3	12	8	5	3	-	-
White, both sexes	9	9	8	7	5	7	15	13	5	3	-	-
Male	6	5	3	3	3	4	4	5	-	-	-	-
Female	3	4	5	4	2	3	11	8	5	3	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	5	8	4	12	11	13	11	7	2	-	-
Male	1	4	6	1	8	4	6	4	1	1	-	-
Female	2	1	2	3	4	7	7	7	6	1	-	-
White, both sexes	3	4	7	3	12	11	13	9	7	2	-	-
Male	1	3	5	-	8	4	6	4	1	1	-	-
Female	2	1	2	3	4	7	7	5	6	1	-	-
Black, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Kentucky												
All races, both sexes	14	7	11	15	11	16	16	14	8	2	1	-
Male	8	4	5	8	6	8	5	3	2	-	-	-
Female	6	3	6	7	5	8	11	11	6	2	1	-
White, both sexes	12	7	11	12	11	16	15	14	8	2	1	-
Male	7	4	5	5	6	8	4	3	2	-	-	-
Female	5	3	6	7	5	8	11	11	6	2	1	-
Black, both sexes	2	-	-	3	-	-	1	-	-	-	-	-
Male	1	-	-	3	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	16	19	17	11	11	24	23	12	3	5	1	-
Male	9	5	10	8	4	12	13	4	1	1	-	-
Female	7	14	7	3	7	12	10	8	2	4	1	-
White, both sexes	11	13	15	5	10	17	21	9	2	5	1	-
Male	6	4	8	2	4	7	12	3	1	1	-	-
Female	5	9	7	3	6	10	9	6	1	4	1	-
Black, both sexes	5	6	2	6	1	6	2	3	1	-	-	-
Male	3	1	2	6	-	5	1	1	-	-	-	-
Female	2	5	-	-	1	1	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Maine												
All races, both sexes	-	3	-	5	4	5	6	3	2	1	-	-
Male	-	2	-	3	3	-	5	2	1	-	-	-
Female	-	1	-	2	1	5	1	1	1	1	-	-
White, both sexes	-	3	-	5	4	5	6	3	2	1	-	-
Male	-	2	-	3	3	-	5	2	1	-	-	-
Female	-	1	-	2	1	5	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	8	9	7	6	12	22	21	14	6	6	-	-
Male	2	5	1	3	5	8	8	5	1	-	-	-
Female	6	4	6	3	7	14	13	9	5	6	-	-
White, both sexes	5	3	4	3	9	15	12	9	6	4	-	-
Male	1	1	1	1	4	7	5	3	1	-	-	-
Female	4	2	3	2	5	8	7	6	5	4	-	-
Black, both sexes	3	6	3	3	3	7	8	5	-	2	-	-
Male	1	4	-	2	1	1	2	2	-	-	-	-
Female	2	2	3	1	2	6	6	3	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Massachusetts												
All races, both sexes	5	13	9	12	12	24	28	25	10	7	-	-
Male	3	5	5	8	5	13	4	5	1	3	-	-
Female	2	8	4	4	7	11	24	20	9	4	-	-
White, both sexes	4	12	9	10	12	23	27	25	10	6	-	-
Male	2	4	5	6	5	12	4	5	1	2	-	-
Female	2	8	4	4	7	11	23	20	9	4	-	-
Black, both sexes	1	1	-	1	-	1	-	-	-	1	-	-
Male	1	1	-	1	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	14	15	14	17	25	36	32	17	15	2	1	-
Male	7	5	9	10	15	21	11	4	5	-	-	-
Female	7	10	5	7	10	15	21	13	10	2	1	-
White, both sexes	12	11	11	13	22	32	27	13	15	2	-	-
Male	6	4	8	8	12	19	11	3	5	-	-	-
Female	6	7	3	5	10	13	16	10	10	2	-	-
Black, both sexes	2	4	3	4	3	1	5	4	-	-	1	-
Male	1	1	1	2	3	-	-	1	-	-	-	-
Female	1	3	2	2	-	1	5	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Minnesota												
All races, both sexes	6	9	4	6	5	7	7	10	5	1	-	-
Male	5	4	2	3	3	2	3	4	3	1	-	-
Female	1	5	2	3	2	5	4	6	2	-	-	-
White, both sexes	6	8	4	5	5	7	5	10	5	1	-	-
Male	5	4	2	3	3	2	3	4	3	1	-	-
Female	1	4	2	2	2	5	2	6	2	-	-	-
Black, both sexes	-	1	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	8	11	7	11	11	13	7	6	3	-	-
Male	6	5	3	4	6	6	3	4	2	1	-	-
Female	-	3	8	3	5	5	10	3	4	2	-	-
White, both sexes	-	3	4	4	9	4	8	6	5	2	-	-
Male	-	2	1	3	6	2	2	3	1	1	-	-
Female	-	1	3	1	3	2	6	3	4	1	-	-
Black, both sexes	6	5	7	3	2	7	5	1	1	1	-	-
Male	6	3	2	1	-	4	1	1	1	-	-	-
Female	-	2	5	2	2	3	4	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Missouri												
All races, both sexes	12	19	22	15	17	23	25	23	5	2	1	-
Male	6	13	13	4	6	12	9	7	1	-	-	-
Female	6	6	9	11	11	11	16	16	4	2	1	-
White, both sexes	10	14	18	10	16	20	23	20	5	2	1	-
Male	5	10	12	4	6	11	8	7	1	-	-	-
Female	5	4	6	6	10	9	15	13	4	2	1	-
Black, both sexes	2	5	4	5	1	3	2	2	-	-	-	-
Male	1	3	1	-	-	1	1	-	-	-	-	-
Female	1	2	3	5	1	2	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	3	3	2	1	1	2	4	2	2	-	-	-
Male	2	2	2	-	1	-	1	-	1	-	-	-
Female	1	1	-	1	-	2	3	2	1	-	-	-
White, both sexes	3	3	2	1	1	2	4	2	2	-	-	-
Male	2	2	2	-	1	-	1	-	1	-	-	-
Female	1	1	-	1	-	2	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Nebraska												
All races, both sexes	2	5	2	1	2	4	3	2	2	-	1	-
Male	1	2	1	-	1	1	-	1	-	-	-	-
Female	1	3	1	1	1	3	3	1	2	-	1	-
White, both sexes	2	5	2	1	2	4	3	2	2	-	1	-
Male	1	2	1	-	1	1	-	1	-	-	-	-
Female	1	3	1	1	1	3	3	1	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	5	4	8	9	7	10	5	5	-	-	-
Male	3	3	1	7	3	2	4	1	2	-	-	-
Female	1	2	3	1	6	5	6	4	3	-	-	-
White, both sexes	4	4	3	7	9	7	7	5	4	-	-	-
Male	3	3	1	6	3	2	3	1	1	-	-	-
Female	1	1	2	1	6	5	4	4	3	-	-	-
Black, both sexes	-	1	1	1	-	-	2	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
New Hampshire												
All races, both sexes	3	2	-	3	3	1	5	5	2	-	1	-
Male	2	-	-	3	2	-	3	2	-	-	-	-
Female	1	2	-	-	1	1	2	3	2	-	1	-
White, both sexes	3	2	-	3	2	1	5	5	2	-	1	-
Male	2	-	-	3	1	-	3	2	-	-	-	-
Female	1	2	-	-	1	1	2	3	2	-	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	12	8	17	24	18	27	19	30	11	9	1	-
Male	9	5	7	11	7	7	5	7	-	1	-	-
Female	3	3	10	13	11	20	14	23	11	8	1	-
White, both sexes	11	6	11	19	15	24	13	25	10	9	1	-
Male	8	3	3	8	6	7	5	6	-	1	-	-
Female	3	3	8	11	9	17	8	19	10	8	1	-
Black, both sexes	-	2	6	4	3	2	4	4	-	-	-	-
Male	-	2	4	3	1	-	-	1	-	-	-	-
Female	-	-	2	1	2	2	4	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	2	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
New Mexico												
All races, both sexes	2	-	2	1	4	10	1	2	3	1	-	-
Male	1	-	1	-	4	5	-	1	-	1	-	-
Female	1	-	1	1	-	5	1	1	3	-	-	-
White, both sexes	2	-	1	1	4	9	1	2	3	1	-	-
Male	1	-	1	-	4	5	-	1	-	1	-	-
Female	1	-	-	1	-	4	1	1	3	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	17	17	15	20	28	37	39	37	13	10	-	-
Male	8	11	8	9	16	15	20	9	4	3	-	-
Female	9	6	7	11	12	22	19	28	9	7	-	-
White, both sexes	10	16	10	18	26	30	30	34	11	10	-	-
Male	6	11	6	9	15	12	13	8	4	3	-	-
Female	4	5	4	9	11	18	17	26	7	7	-	-
Black, both sexes	7	1	5	2	2	5	7	3	2	-	-	-
Male	2	-	2	-	1	1	6	1	-	-	-	-
Female	5	1	3	2	1	4	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
North Carolina												
All races, both sexes	15	23	24	31	24	34	27	32	19	1	-	-
Male	8	11	12	14	8	12	7	7	3	1	-	-
Female	7	12	12	17	16	22	20	25	16	-	-	-
White, both sexes	9	18	16	19	16	27	20	26	19	1	-	-
Male	5	8	9	8	6	11	4	4	3	1	-	-
Female	4	10	7	11	10	16	16	22	16	-	-	-
Black, both sexes	6	5	7	12	8	6	7	6	-	-	-	-
Male	3	3	2	6	2	1	3	3	-	-	-	-
Female	3	2	5	6	6	5	4	3	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	2	-	2	-	2	1	3	3	-	-	-
Male	1	2	-	1	-	1	-	2	2	-	-	-
Female	1	-	-	1	-	1	1	1	1	-	-	-
White, both sexes	2	2	-	2	-	2	1	3	3	-	-	-
Male	1	2	-	1	-	1	-	2	2	-	-	-
Female	1	-	-	1	-	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Ohio												
All races, both sexes	11	14	10	21	28	30	30	27	12	6	-	-
Male	3	8	3	11	14	17	12	12	5	2	-	-
Female	8	6	7	10	14	13	18	15	7	4	-	-
White, both sexes	8	14	9	18	20	24	26	23	11	6	-	-
Male	1	8	3	10	10	12	11	9	4	2	-	-
Female	7	6	6	8	10	12	15	14	7	4	-	-
Black, both sexes	3	-	1	3	8	6	3	4	-	-	-	-
Male	2	-	-	1	4	5	1	3	-	-	-	-
Female	1	-	1	2	4	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	5	5	14	3	7	10	12	16	4	1	1	-
Male	4	1	7	3	5	2	4	8	-	-	-	-
Female	1	4	7	-	2	8	8	8	4	1	1	-
White, both sexes	3	5	11	3	7	7	11	15	4	-	1	-
Male	2	1	6	3	5	2	4	8	-	-	-	-
Female	1	4	5	-	2	5	7	7	4	-	1	-
Black, both sexes	1	-	1	-	-	2	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	2	1	1	-	-	-	-
American Indian 1/, both sexes	1	-	2	-	-	-	-	-	-	1	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Oregon												
All races, both sexes	6	4	3	3	8	8	13	4	9	-	-	-
Male	2	1	3	2	6	5	6	2	2	-	-	-
Female	4	3	-	1	2	3	7	2	7	-	-	-
White, both sexes	5	4	2	3	8	7	13	4	9	-	-	-
Male	1	1	2	2	6	5	6	2	2	-	-	-
Female	4	3	-	1	2	2	7	2	7	-	-	-
Black, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	22	26	36	24	28	51	59	57	29	5	4	-
Male	12	15	15	10	8	18	28	19	6	-	-	-
Female	10	11	21	14	20	33	31	38	23	5	4	-
White, both sexes	19	22	29	20	24	44	56	49	26	4	4	-
Male	9	13	13	8	8	13	27	16	6	-	-	-
Female	10	9	16	12	16	31	29	33	20	4	4	-
Black, both sexes	3	4	7	4	4	7	3	8	3	-	-	-
Male	3	2	2	2	-	5	1	3	-	-	-	-
Female	-	2	5	2	4	2	2	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Rhode Island												
All races, both sexes	1	-	1	3	4	4	-	3	3	-	-	-
Male	1	-	-	3	2	1	-	1	2	-	-	-
Female	-	-	1	-	2	3	-	2	1	-	-	-
White, both sexes	1	-	-	3	4	4	-	3	3	-	-	-
Male	1	-	-	3	2	1	-	1	2	-	-	-
Female	-	-	-	-	2	3	-	2	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	8	11	9	24	19	12	14	6	6	-	-
Male	5	3	5	4	8	7	5	4	2	1	-	-
Female	7	5	6	5	16	12	7	10	4	5	-	-
White, both sexes	6	4	5	6	15	13	9	12	5	6	-	-
Male	3	1	2	4	4	6	4	3	1	1	-	-
Female	3	3	3	2	11	7	5	9	4	5	-	-
Black, both sexes	6	4	6	3	9	6	3	2	1	-	-	-
Male	2	2	3	-	4	1	1	1	1	-	-	-
Female	4	2	3	3	5	5	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
South Dakota												
All races, both sexes	2	1	-	4	-	-	3	1	-	2	-	-
Male	1	-	-	2	-	-	-	-	-	2	-	-
Female	1	1	-	2	-	-	3	1	-	-	-	-
White, both sexes	1	1	-	4	-	-	3	1	-	2	-	-
Male	-	-	-	2	-	-	-	-	-	2	-	-
Female	1	1	-	2	-	-	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	11	24	16	18	19	19	28	25	18	3	-	-
Male	3	9	7	14	7	5	13	9	6	-	-	-
Female	8	15	9	4	12	14	15	16	12	3	-	-
White, both sexes	7	15	14	14	15	18	24	20	15	3	-	-
Male	2	6	5	10	6	5	10	6	5	-	-	-
Female	5	9	9	4	9	13	14	14	10	3	-	-
Black, both sexes	4	9	2	4	4	1	4	5	3	-	-	-
Male	1	3	2	4	1	-	3	3	1	-	-	-
Female	3	6	-	-	3	1	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Texas												
All races, both sexes	34	38	39	37	51	64	71	69	45	9	3	-
Male	19	22	26	14	32	31	32	21	10	1	1	-
Female	15	16	13	23	19	33	39	48	35	8	2	-
White, both sexes	28	32	26	30	44	51	67	63	38	6	1	-
Male	16	18	20	11	27	26	30	19	7	1	-	-
Female	12	14	6	19	17	25	37	44	31	5	1	-
Black, both sexes	6	6	12	6	7	13	4	6	6	3	2	-
Male	3	4	5	3	5	5	2	2	2	-	1	-
Female	3	2	7	3	2	8	2	4	4	3	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	4	6	7	4	5	8	7	4	4	-	-	-
Male	2	3	3	3	3	5	3	1	3	-	-	-
Female	2	3	4	1	2	3	4	3	1	-	-	-
White, both sexes	4	6	7	3	5	7	7	4	4	-	-	-
Male	2	3	3	2	3	4	3	1	3	-	-	-
Female	2	3	4	1	2	3	4	3	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Vermont												
All races, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	19	22	17	15	18	20	39	22	17	1	1	-
Male	8	15	11	7	7	11	10	9	6	-	-	-
Female	11	7	6	8	11	9	29	13	11	1	1	-
White, both sexes	16	14	13	10	14	16	31	20	17	-	-	-
Male	7	11	8	5	5	11	9	8	6	-	-	-
Female	9	3	5	5	9	5	22	12	11	-	-	-
Black, both sexes	3	8	4	5	4	4	8	2	-	1	1	-
Male	1	4	3	2	2	-	1	1	-	-	-	-
Female	2	4	1	3	2	4	7	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Washington												
All races, both sexes	8	10	6	12	12	14	19	15	13	1	-	-
Male	1	4	3	4	9	7	7	5	2	-	-	-
Female	7	6	3	8	3	7	12	10	11	1	-	-
White, both sexes	7	7	6	11	12	13	19	15	12	1	-	-
Male	1	3	3	3	9	6	7	5	2	-	-	-
Female	6	4	3	8	3	7	12	10	10	1	-	-
Black, both sexes	1	1	-	1	-	-	-	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	7	6	6	6	9	8	6	-	1	1	-
Male	1	3	2	3	2	3	3	2	-	1	-	-
Female	2	4	4	3	4	6	5	4	-	-	1	-
White, both sexes	3	7	6	6	6	8	6	5	-	1	1	-
Male	1	3	2	3	2	2	2	2	-	1	-	-
Female	2	4	4	3	4	6	4	3	-	-	1	-
Black, both sexes	-	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Wisconsin												
All races, both sexes	13	8	10	5	13	21	22	27	12	4	1	-
Male	6	4	5	2	7	11	11	8	1	1	-	-
Female	7	4	5	3	6	10	11	19	11	3	1	-
White, both sexes	11	6	9	5	12	21	21	26	11	4	1	-
Male	5	3	5	2	7	11	11	8	1	1	-	-
Female	6	3	4	3	5	10	10	18	10	3	1	-
Black, both sexes	2	2	1	-	1	-	1	1	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	1	2	3	4	4	3	2	1	-	-	-
Male	2	1	2	2	4	4	2	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	1	-	-	-
White, both sexes	2	1	2	3	3	3	3	2	-	-	-	-
Male	2	1	2	2	3	3	2	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
United States												
All races, both sexes	5,463	79	24	10	9	20	28	45	51	91	116	220
Male	1,844	42	12	4	3	10	13	22	16	35	44	93
Female	3,619	37	12	6	6	10	15	23	35	56	72	127
White, both sexes	4,494	41	14	8	5	12	21	28	30	57	82	151
Male	1,532	21	5	3	1	7	10	14	11	23	32	69
Female	2,962	20	9	5	4	5	11	14	19	34	50	82
Black, both sexes	857	35	9	2	2	7	7	15	20	28	29	63
Male	272	19	6	1	1	3	3	7	5	10	11	23
Female	585	16	3	1	1	4	4	8	15	18	18	40
American Indian 1/, both sexes	30	2	-	-	-	1	-	-	1	4	2	1
Male	10	1	-	-	-	-	-	-	-	2	-	-
Female	20	1	-	-	-	1	-	-	1	2	2	1
Asian or Pacific Islander 2/, both sexes	82	1	1	-	2	-	-	2	-	2	3	5
Male	30	1	1	-	1	-	-	1	-	-	1	1
Female	52	-	-	-	1	-	-	1	-	2	2	4
Alabama												
All races, both sexes	83	1	1	1	-	1	1	1	-	1	4	4
Male	31	1	1	-	-	1	-	-	-	1	1	3
Female	52	-	-	1	-	-	1	1	-	-	3	1
White, both sexes	56	1	1	1	-	1	-	-	-	-	1	2
Male	23	1	1	-	-	1	-	-	-	-	-	2
Female	33	-	-	1	-	-	-	-	-	-	1	-
Black, both sexes	27	-	-	-	-	-	1	1	-	1	3	2
Male	8	-	-	-	-	-	-	-	-	1	1	1
Female	19	-	-	-	-	-	1	1	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Alaska												
All races, both sexes	13	-	-	-	-	-	-	-	-	2	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	2	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	71	2	-	-	1	-	-	1	1	4	2	4
Male	26	1	-	-	-	-	-	-	-	2	1	2
Female	45	1	-	-	1	-	-	1	1	2	1	2
White, both sexes	63	2	-	-	-	-	-	-	1	3	1	4
Male	25	1	-	-	-	-	-	-	-	1	1	2
Female	38	1	-	-	-	-	-	-	1	2	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Arkansas												
All races, both sexes	56	-	-	-	-	-	-	2	-	1	-	4
Male	20	-	-	-	-	-	-	1	-	-	-	3
Female	36	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	49	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	2	-	1	-	2
Male	3	-	-	-	-	-	-	1	-	-	-	2
Female	4	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	466	5	4	1	2	2	7	4	7	8	19	18
Male	188	3	2	1	1	2	4	3	3	4	7	8
Female	278	2	2	-	1	-	3	1	4	4	12	10
White, both sexes	370	5	4	1	1	1	5	2	6	6	12	16
Male	147	3	2	1	1	1	3	1	3	3	4	8
Female	223	2	2	-	-	-	2	1	3	3	8	8
Black, both sexes	60	-	-	-	1	1	2	1	1	1	5	1
Male	28	-	-	-	-	1	1	1	-	1	3	-
Female	32	-	-	-	1	-	1	-	1	-	2	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	1	-	1	2	1
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	1	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Colorado												
All races, both sexes	119	-	-	-	-	-	3	-	-	2	2	-
Male	42	-	-	-	-	-	1	-	-	2	1	-
Female	77	-	-	-	-	-	2	-	-	-	1	-
White, both sexes	107	-	-	-	-	-	2	-	-	2	1	-
Male	38	-	-	-	-	-	1	-	-	2	1	-
Female	69	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	94	1	-	1	-	-	-	-	-	2	1	5
Male	29	1	-	1	-	-	-	-	-	-	-	2
Female	65	-	-	-	-	-	-	-	-	2	1	3
White, both sexes	85	1	-	1	-	-	-	-	-	1	1	4
Male	26	1	-	1	-	-	-	-	-	-	-	1
Female	59	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Delaware												
All races, both sexes	14	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	16	-	-	-	-	1	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	1	-	-	-	1	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	1	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	1	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Florida												
All races, both sexes	278	7	2	1	-	-	1	3	3	4	6	13
Male	112	5	1	-	-	-	-	2	2	1	2	9
Female	166	2	1	1	-	-	1	1	1	3	4	4
White, both sexes	228	1	1	1	-	-	1	2	1	2	5	9
Male	89	-	-	-	-	-	-	1	1	-	2	6
Female	139	1	1	1	-	-	1	1	-	2	3	3
Black, both sexes	48	6	-	-	-	-	-	1	2	2	1	4
Male	22	5	-	-	-	-	-	1	1	1	-	3
Female	26	1	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	221	6	-	-	-	2	3	4	1	6	3	7
Male	81	4	-	-	-	2	2	3	-	2	3	2
Female	140	2	-	-	-	-	1	1	1	4	-	5
White, both sexes	148	2	-	-	-	1	1	3	-	2	2	2
Male	55	1	-	-	-	1	-	2	-	1	2	-
Female	93	1	-	-	-	-	1	1	-	1	-	2
Black, both sexes	72	4	-	-	-	1	2	1	1	4	1	5
Male	26	3	-	-	-	1	2	1	-	1	1	2
Female	46	1	-	-	-	-	-	-	1	3	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Hawaii												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	1	-	1
Idaho												
All races, both sexes	31	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Illinois												
All races, both sexes	230	3	-	-	-	1	-	1	1	3	2	8
Male	70	2	-	-	-	1	-	1	-	1	-	3
Female	160	1	-	-	-	-	-	-	1	2	2	5
White, both sexes	199	2	-	-	-	1	-	1	-	1	2	6
Male	61	1	-	-	-	1	-	1	-	-	-	3
Female	138	1	-	-	-	-	-	-	-	1	2	3
Black, both sexes	30	1	-	-	-	-	-	-	1	2	-	2
Male	8	1	-	-	-	-	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	1	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	122	1	-	-	-	1	1	1	3	1	3	4
Male	35	-	-	-	-	-	1	-	1	1	1	2
Female	87	1	-	-	-	1	-	1	2	-	2	2
White, both sexes	110	1	-	-	-	1	1	1	2	1	3	3
Male	33	-	-	-	-	-	1	-	1	1	1	2
Female	77	1	-	-	-	1	-	1	1	-	2	1
Black, both sexes	12	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Iowa												
All races, both sexes	72	-	1	-	-	-	-	-	-	1	3	3
Male	18	-	-	-	-	-	-	-	-	1	2	1
Female	54	-	1	-	-	-	-	-	-	-	1	2
White, both sexes	70	-	1	-	-	-	-	-	-	-	3	3
Male	16	-	-	-	-	-	-	-	-	-	2	1
Female	54	-	1	-	-	-	-	-	-	-	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	72	2	-	-	-	1	-	1	1	1	2	-
Male	26	2	-	-	-	-	-	-	-	1	1	-
Female	46	-	-	-	-	1	-	1	1	-	1	-
White, both sexes	65	1	-	-	-	1	-	1	1	1	1	-
Male	20	1	-	-	-	-	-	-	-	1	-	-
Female	45	-	-	-	-	1	-	1	1	-	1	-
Black, both sexes	6	1	-	-	-	-	-	-	-	-	1	-
Male	5	1	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Kentucky												
All races, both sexes	77	-	1	-	-	-	-	1	-	2	2	4
Male	26	-	-	-	-	-	-	1	-	1	1	2
Female	51	-	1	-	-	-	-	-	-	1	1	2
White, both sexes	69	-	1	-	-	-	-	1	-	2	2	4
Male	25	-	-	-	-	-	-	1	-	1	1	2
Female	44	-	1	-	-	-	-	-	-	1	1	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	72	2	-	-	-	-	-	-	1	2	2	4
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	48	2	-	-	-	-	-	-	1	2	2	2
White, both sexes	44	-	-	-	-	-	-	-	-	1	-	2
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	28	2	-	-	-	-	-	-	1	1	2	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	2	-	-	-	-	-	-	1	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Maine												
All races, both sexes	30	-	-	-	-	-	2	-	-	-	-	2
Male	8	-	-	-	-	-	1	-	-	-	-	1
Female	22	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	30	-	-	-	-	-	2	-	-	-	-	2
Male	8	-	-	-	-	-	1	-	-	-	-	1
Female	22	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	107	1	1	-	-	-	1	2	2	1	4	3
Male	40	-	1	-	-	-	-	2	1	-	2	1
Female	67	1	-	-	-	-	1	-	1	1	2	2
White, both sexes	70	-	-	-	-	-	1	1	-	1	2	1
Male	32	-	-	-	-	-	-	1	-	-	1	1
Female	38	-	-	-	-	-	1	-	-	1	1	-
Black, both sexes	37	1	1	-	-	-	-	1	2	-	2	2
Male	8	-	1	-	-	-	-	1	1	-	1	-
Female	29	1	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Massachusetts												
All races, both sexes	123	2	-	-	-	-	-	1	-	2	3	4
Male	38	-	-	-	-	-	-	1	-	1	2	1
Female	85	2	-	-	-	-	-	-	-	1	1	3
White, both sexes	114	1	-	-	-	-	-	1	-	2	3	3
Male	37	-	-	-	-	-	-	1	-	1	2	1
Female	77	1	-	-	-	-	-	-	-	1	1	2
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	184	4	1	1	-	-	1	2	3	2	-	9
Male	59	1	-	-	-	-	1	-	1	2	-	2
Female	125	3	1	1	-	-	-	2	2	-	-	7
White, both sexes	147	1	1	1	-	-	1	1	2	1	-	4
Male	46	-	-	-	-	-	1	-	1	1	-	-
Female	101	1	1	1	-	-	-	1	1	-	-	4
Black, both sexes	36	2	-	-	-	-	-	1	1	1	-	5
Male	13	1	-	-	-	-	-	-	-	1	-	2
Female	23	1	-	-	-	-	-	1	1	-	-	3
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Minnesota												
All races, both sexes	124	-	-	-	1	-	-	2	2	1	3	4
Male	43	-	-	-	1	-	-	1	-	1	2	2
Female	81	-	-	-	-	-	-	1	2	-	1	2
White, both sexes	115	-	-	-	-	-	-	2	2	1	3	4
Male	40	-	-	-	-	-	-	1	-	1	2	2
Female	75	-	-	-	-	-	-	1	2	-	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	55	2	-	-	-	-	-	1	-	-	-	1
Male	14	1	-	-	-	-	-	-	-	-	-	-
Female	41	1	-	-	-	-	-	1	-	-	-	1
White, both sexes	38	2	-	-	-	-	-	-	-	-	-	-
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	28	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Missouri												
All races, both sexes	120	-	-	1	2	1	1	-	2	2	4	11
Male	35	-	-	-	-	-	1	-	2	1	1	4
Female	85	-	-	1	2	1	-	-	-	1	3	7
White, both sexes	102	-	-	1	2	-	1	-	1	2	3	7
Male	29	-	-	-	-	-	1	-	1	1	1	2
Female	73	-	-	1	2	-	-	-	-	1	2	5
Black, both sexes	18	-	-	-	-	1	-	-	1	-	1	4
Male	6	-	-	-	-	-	-	-	1	-	-	2
Female	12	-	-	-	-	1	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Nebraska												
All races, both sexes	40	3	-	-	-	-	1	1	-	1	2	-
Male	9	1	-	-	-	-	-	1	-	1	1	-
Female	31	2	-	-	-	-	1	-	-	-	1	-
White, both sexes	38	2	-	-	-	-	1	1	-	1	2	-
Male	9	1	-	-	-	-	-	1	-	1	1	-
Female	29	1	-	-	-	-	1	-	-	-	1	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	43	-	-	-	-	-	-	-	1	-	-	4
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	37	-	-	-	-	-	-	-	1	-	-	4
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Hampshire												
All races, both sexes	31	-	-	-	1	-	-	-	-	-	1	4
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	1	-	-	-	-	-	1	2
White, both sexes	31	-	-	-	1	-	-	-	-	-	1	4
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	1	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	138	2	-	-	-	1	1	2	-	2	4	7
Male	37	2	-	-	-	1	-	2	-	-	1	3
Female	101	-	-	-	-	-	1	-	-	2	3	4
White, both sexes	105	1	-	-	-	1	-	1	-	1	2	3
Male	28	1	-	-	-	1	-	1	-	-	1	-
Female	77	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	30	1	-	-	-	-	1	1	-	1	2	3
Male	8	1	-	-	-	-	-	1	-	-	-	2
Female	22	-	-	-	-	-	1	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Mexico												
All races, both sexes	43	1	-	-	-	-	1	-	-	2	1	2
Male	12	1	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	1	-	-	2	1	1
White, both sexes	42	1	-	-	-	-	1	-	-	2	1	2
Male	11	1	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	1	-	-	2	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	298	5	2	1	1	-	-	4	5	3	9	8
Male	100	2	-	1	-	-	-	2	1	1	3	3
Female	198	3	2	-	1	-	-	2	4	2	6	5
White, both sexes	244	3	1	1	1	-	-	2	3	2	6	3
Male	85	1	-	1	-	-	-	1	1	-	3	2
Female	159	2	1	-	1	-	-	1	2	2	3	1
Black, both sexes	48	2	1	-	-	-	-	2	2	1	3	4
Male	13	1	-	-	-	-	-	1	-	1	-	1
Female	35	1	1	-	-	-	-	1	2	-	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
North Carolina												
All races, both sexes	208	5	2	1	-	1	-	1	2	1	7	12
Male	68	5	-	1	-	-	-	-	1	-	3	6
Female	140	-	2	-	-	1	-	1	1	1	4	6
White, both sexes	147	1	1	-	-	-	-	-	-	-	5	5
Male	50	1	-	-	-	-	-	-	-	-	2	2
Female	97	-	1	-	-	-	-	-	-	-	3	3
Black, both sexes	57	3	1	1	-	1	-	1	2	1	2	7
Male	16	3	-	1	-	-	-	-	1	-	1	4
Female	41	-	1	-	-	1	-	1	1	1	1	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Ohio												
All races, both sexes	218	2	-	-	-	-	1	-	2	2	4	6
Male	69	2	-	-	-	-	-	-	-	2	-	1
Female	149	-	-	-	-	-	1	-	2	-	4	5
White, both sexes	188	1	-	-	-	-	1	-	1	2	3	5
Male	57	1	-	-	-	-	-	-	-	2	-	1
Female	131	-	-	-	-	-	1	-	1	-	3	4
Black, both sexes	29	1	-	-	-	-	-	-	1	-	1	1
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	46	1	-	-	-	1	-	1	-	1	1	-
Male	15	1	-	-	-	1	-	-	-	1	-	-
Female	31	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	36	-	-	-	-	1	-	1	-	-	-	-
Male	12	-	-	-	-	1	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Oregon												
All races, both sexes	79	-	-	-	-	-	-	-	-	-	1	5
Male	24	-	-	-	-	-	-	-	-	-	-	4
Female	55	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	73	-	-	-	-	-	-	-	-	-	1	5
Male	23	-	-	-	-	-	-	-	-	-	-	4
Female	50	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	301	2	2	-	1	-	-	2	2	4	3	8
Male	89	-	2	-	1	-	-	-	1	2	1	4
Female	212	2	-	-	-	-	-	2	1	2	2	4
White, both sexes	257	2	-	-	-	-	-	1	2	3	1	7
Male	77	-	-	-	-	-	-	-	1	2	-	4
Female	180	2	-	-	-	-	-	1	1	1	1	3
Black, both sexes	43	-	2	-	1	-	-	1	-	1	2	1
Male	12	-	2	-	1	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Rhode Island												
All races, both sexes	26	1	-	-	-	-	-	1	1	-	-	2
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	19	1	-	-	-	-	-	1	-	-	-	1
White, both sexes	23	1	-	-	-	-	-	1	1	-	-	2
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	17	1	-	-	-	-	-	1	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	104	3	1	-	-	-	1	1	-	3	1	5
Male	42	1	1	-	-	-	-	1	-	1	-	2
Female	62	2	-	-	-	-	1	-	-	2	1	3
White, both sexes	75	1	1	-	-	-	1	1	-	1	1	3
Male	29	1	1	-	-	-	-	1	-	-	-	1
Female	46	-	-	-	-	-	1	-	-	1	1	2
Black, both sexes	28	2	-	-	-	-	-	-	-	2	-	2
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	15	2	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
South Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	107	3	1	1	-	-	-	1	1	3	3	3
Male	33	1	1	-	-	-	-	-	-	-	2	-
Female	74	2	-	1	-	-	-	1	1	3	1	3
White, both sexes	89	2	-	1	-	-	-	1	1	3	3	2
Male	27	1	-	-	-	-	-	-	-	-	2	-
Female	62	1	-	1	-	-	-	1	1	3	1	2
Black, both sexes	18	1	1	-	-	-	-	-	-	-	-	1
Male	6	-	1	-	-	-	-	-	-	-	-	-
Female	12	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Texas												
All races, both sexes	335	4	3	-	-	5	-	2	5	11	6	18
Male	116	3	2	-	-	2	-	1	1	4	2	5
Female	219	1	1	-	-	3	-	1	4	7	4	13
White, both sexes	265	3	2	-	-	3	-	1	2	8	5	14
Male	97	3	1	-	-	1	-	1	-	3	1	5
Female	168	-	1	-	-	2	-	-	2	5	4	9
Black, both sexes	68	1	1	-	-	2	-	1	3	3	1	3
Male	19	-	1	-	-	1	-	-	1	1	1	-
Female	49	1	-	-	-	1	-	1	2	2	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	43	-	-	-	-	-	-	-	1	-	-	3
Male	15	-	-	-	-	-	-	-	1	-	-	2
Female	28	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	-	1	-	-	3
Male	15	-	-	-	-	-	-	-	1	-	-	2
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Vermont												
All races, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	167	6	2	-	-	1	1	-	1	3	2	7
Male	60	2	1	-	-	-	1	-	-	1	1	4
Female	107	4	1	-	-	1	-	-	1	2	1	3
White, both sexes	122	1	-	-	-	1	1	-	1	2	2	2
Male	45	-	-	-	-	-	1	-	-	1	1	2
Female	77	1	-	-	-	1	-	-	1	1	1	-
Black, both sexes	44	5	2	-	-	-	-	-	-	1	-	5
Male	14	2	1	-	-	-	-	-	-	-	-	2
Female	30	3	1	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Washington												
All races, both sexes	123	-	-	-	-	-	1	-	1	3	3	3
Male	37	-	-	-	-	-	1	-	-	-	1	2
Female	86	-	-	-	-	-	-	-	1	3	2	1
White, both sexes	115	-	-	-	-	-	1	-	-	2	3	3
Male	36	-	-	-	-	-	1	-	-	-	1	2
Female	79	-	-	-	-	-	-	-	-	2	2	1
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	49	1	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	35	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	48	1	-	-	-	-	-	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	35	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Wisconsin												
All races, both sexes	131	-	-	1	-	-	-	-	1	1	1	5
Male	51	-	-	-	-	-	-	-	-	-	-	2
Female	80	-	-	1	-	-	-	-	1	1	1	3
White, both sexes	120	-	-	-	-	-	-	-	-	1	1	3
Male	49	-	-	-	-	-	-	-	-	-	-	2
Female	71	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	10	-	-	1	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	1	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	17	-	-	-	-	1	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
United States												
All races, both sexes	281	359	368	457	544	706	804	721	385	125	20	-
Male	111	128	133	181	203	249	228	208	86	19	4	-
Female	170	231	235	276	341	457	576	513	299	106	16	-
White, both sexes	206	251	296	362	439	622	719	660	359	115	16	-
Male	90	96	110	149	171	221	209	192	79	15	4	-
Female	116	155	186	213	268	401	510	468	280	100	12	-
Black, both sexes	67	101	66	86	95	68	72	51	22	8	4	-
Male	20	30	22	29	29	17	15	13	6	2	-	-
Female	47	71	44	57	66	51	57	38	16	6	4	-
American Indian 1/, both sexes	2	3	2	2	2	4	2	1	1	-	-	-
Male	-	-	-	1	2	2	2	-	-	-	-	-
Female	2	3	2	1	-	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	4	4	7	8	12	11	9	3	2	-	-
Male	1	2	1	2	1	9	2	3	1	2	-	-
Female	5	2	3	5	7	3	9	6	2	-	-	-
Alabama												
All races, both sexes	8	4	6	7	10	7	16	8	-	2	-	-
Male	4	-	2	1	5	2	5	3	-	1	-	-
Female	4	4	4	6	5	5	11	5	-	1	-	-
White, both sexes	5	1	3	5	8	6	13	6	-	2	-	-
Male	3	-	1	1	4	2	5	1	-	1	-	-
Female	2	1	2	4	4	4	8	5	-	1	-	-
Black, both sexes	3	3	3	2	2	1	3	2	-	-	-	-
Male	1	-	1	-	1	-	-	2	-	-	-	-
Female	2	3	2	2	1	1	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Alaska												
All races, both sexes	-	1	1	-	2	1	3	2	1	-	-	-
Male	-	-	-	-	1	1	3	-	-	-	-	-
Female	-	1	1	-	1	-	-	2	1	-	-	-
White, both sexes	-	1	1	-	1	-	3	2	1	-	-	-
Male	-	-	-	-	-	-	3	-	-	-	-	-
Female	-	1	1	-	1	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	6	4	6	8	10	7	7	4	1	-	-
Male	1	2	1	3	5	4	1	1	2	-	-	-
Female	2	4	3	3	3	6	6	6	2	1	-	-
White, both sexes	3	6	4	4	7	10	7	6	4	1	-	-
Male	1	2	1	3	5	4	1	1	2	-	-	-
Female	2	4	3	1	2	6	6	5	2	1	-	-
Black, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Arkansas												
All races, both sexes	4	4	5	5	6	7	3	8	3	4	-	-
Male	2	1	1	1	4	2	1	2	1	1	-	-
Female	2	3	4	4	2	5	2	6	2	3	-	-
White, both sexes	4	3	5	5	6	7	3	7	3	4	-	-
Male	2	1	1	1	4	2	1	2	1	1	-	-
Female	2	2	4	4	2	5	2	5	2	3	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	31	38	26	34	29	57	91	53	25	5	-	-
Male	11	19	13	17	11	26	23	21	6	3	-	-
Female	20	19	13	17	18	31	68	32	19	2	-	-
White, both sexes	25	26	20	25	21	44	76	47	23	4	-	-
Male	10	13	9	12	8	20	20	18	5	2	-	-
Female	15	13	11	13	13	24	56	29	18	2	-	-
Black, both sexes	3	9	4	8	5	7	9	1	1	-	-	-
Male	1	5	4	4	2	2	1	1	1	-	-	-
Female	2	4	-	4	3	5	8	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	2	1	3	5	5	5	1	1	-	-
Male	-	1	-	1	1	4	1	2	-	1	-	-
Female	2	2	2	-	2	1	4	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Colorado												
All races, both sexes	5	9	11	10	8	17	18	21	11	1	1	-
Male	3	4	2	5	3	11	3	4	2	-	1	-
Female	2	5	9	5	5	6	15	17	9	1	-	-
White, both sexes	4	8	9	10	7	14	16	21	11	1	1	-
Male	3	4	2	5	2	8	3	4	2	-	1	-
Female	1	4	7	5	5	6	13	17	9	1	-	-
Black, both sexes	1	1	2	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	1	2	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	2	8	5	11	19	19	12	3	4	-	-
Male	-	1	4	2	3	7	3	5	-	-	-	-
Female	1	1	4	3	8	12	16	7	3	4	-	-
White, both sexes	-	-	7	4	11	18	18	12	3	4	-	-
Male	-	-	3	2	3	7	3	5	-	-	-	-
Female	-	-	4	2	8	11	15	7	3	4	-	-
Black, both sexes	1	2	1	1	-	1	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Delaware												
All races, both sexes	1	-	1	1	3	3	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	1	-	1	1	2	3	-	1	-	-	-	-
White, both sexes	1	-	1	-	2	3	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	2	3	-	1	-	-	-	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	2	1	-	-	2	4	2	1	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	1	2	-	-	-	2	3	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	2	1	-	-	2	3	1	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	2	-	-	-	2	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Florida												
All races, both sexes	18	15	23	29	25	38	34	28	24	4	-	-
Male	11	2	11	12	11	14	16	7	5	1	-	-
Female	7	13	12	17	14	24	18	21	19	3	-	-
White, both sexes	16	9	21	25	19	34	32	24	22	3	-	-
Male	10	2	10	10	10	11	16	6	4	-	-	-
Female	6	7	11	15	9	23	16	18	18	3	-	-
Black, both sexes	2	5	2	4	6	4	2	4	2	1	-	-
Male	1	-	1	2	1	3	-	1	1	1	-	-
Female	1	5	1	2	5	1	2	3	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	8	17	15	28	18	22	32	24	18	5	2	-
Male	1	6	5	13	6	10	7	11	4	-	-	-
Female	7	11	10	15	12	12	25	13	14	5	2	-
White, both sexes	2	8	9	17	15	18	26	18	17	3	2	-
Male	-	4	4	9	5	9	5	8	4	-	-	-
Female	2	4	5	8	10	9	21	10	13	3	2	-
Black, both sexes	5	9	6	11	3	4	6	6	1	2	-	-
Male	1	2	1	4	1	1	2	3	-	-	-	-
Female	4	7	5	7	2	3	4	3	1	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Hawaii												
All races, both sexes	2	-	1	1	-	-	3	7	1	1	-	-
Male	1	-	1	-	-	-	2	3	-	1	-	-
Female	1	-	-	1	-	-	1	4	1	-	-	-
White, both sexes	1	-	-	-	-	-	1	3	-	-	-	-
Male	1	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	2	4	1	1	-	-
Male	-	-	1	-	-	-	1	1	-	1	-	-
Female	1	-	-	1	-	-	1	3	1	-	-	-
Idaho												
All races, both sexes	1	4	-	2	8	5	-	6	3	1	-	-
Male	1	3	-	1	5	1	-	1	-	-	-	-
Female	-	1	-	1	3	4	-	5	3	1	-	-
White, both sexes	1	4	-	2	8	5	-	6	3	1	-	-
Male	1	3	-	1	5	1	-	1	-	-	-	-
Female	-	1	-	1	3	4	-	5	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Illinois												
All races, both sexes	4	20	21	15	25	33	35	29	19	8	2	-
Male	1	1	8	7	7	7	14	7	8	2	-	-
Female	3	19	13	8	18	26	21	22	11	6	2	-
White, both sexes	2	14	19	13	22	29	32	29	17	7	2	-
Male	-	1	7	6	6	7	13	7	7	1	-	-
Female	2	13	12	7	16	22	19	22	10	6	2	-
Black, both sexes	1	6	2	2	3	4	3	-	2	1	-	-
Male	-	-	1	1	1	-	1	-	1	1	-	-
Female	1	6	1	1	2	4	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	6	9	9	9	27	13	19	9	6	-	-	-
Male	1	2	2	5	11	1	5	2	-	-	-	-
Female	5	7	7	4	16	12	14	7	6	-	-	-
White, both sexes	5	8	9	8	23	13	17	9	5	-	-	-
Male	1	2	2	5	9	1	5	2	-	-	-	-
Female	4	6	7	3	14	12	12	7	5	-	-	-
Black, both sexes	1	1	-	1	4	-	2	-	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	1	1	-	1	2	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Iowa												
All races, both sexes	2	6	3	3	3	13	10	12	7	5	-	-
Male	1	2	1	1	-	5	2	1	1	-	-	-
Female	1	4	2	2	3	8	8	11	6	5	-	-
White, both sexes	1	6	3	3	3	13	10	12	7	5	-	-
Male	-	2	1	1	-	5	2	1	1	-	-	-
Female	1	4	2	2	3	8	8	11	6	5	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	8	2	4	8	4	7	7	10	9	5	-	-
Male	5	-	-	4	1	4	2	3	3	-	-	-
Female	3	2	4	4	3	3	5	7	6	5	-	-
White, both sexes	7	2	4	6	3	6	7	10	9	5	-	-
Male	4	-	-	3	-	3	2	3	3	-	-	-
Female	3	2	4	3	3	3	5	7	6	5	-	-
Black, both sexes	1	-	-	2	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Kentucky												
All races, both sexes	4	6	10	9	4	12	8	10	4	-	-	-
Male	2	1	6	2	1	4	2	1	2	-	-	-
Female	2	5	4	7	3	8	6	9	2	-	-	-
White, both sexes	3	2	8	9	4	11	8	10	4	-	-	-
Male	2	-	6	2	1	4	2	1	2	-	-	-
Female	1	2	2	7	3	7	6	9	2	-	-	-
Black, both sexes	1	4	2	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	3	2	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	6	3	5	9	6	12	14	2	-	-	-
Male	1	4	1	2	6	2	3	3	-	-	-	-
Female	3	2	2	3	3	4	9	11	2	-	-	-
White, both sexes	2	3	1	2	6	6	12	8	1	-	-	-
Male	1	2	1	1	4	2	3	2	-	-	-	-
Female	1	1	-	1	2	4	9	6	1	-	-	-
Black, both sexes	2	3	2	3	3	-	-	6	1	-	-	-
Male	-	2	-	1	2	-	-	1	-	-	-	-
Female	2	1	2	2	1	-	-	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Maine												
All races, both sexes	5	1	2	3	3	4	1	-	4	3	-	-
Male	2	1	-	1	-	1	-	-	-	1	-	-
Female	3	-	2	2	3	3	1	-	4	2	-	-
White, both sexes	5	1	2	3	3	4	1	-	4	3	-	-
Male	2	1	-	1	-	1	-	-	-	1	-	-
Female	3	-	2	2	3	3	1	-	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	7	8	8	19	11	16	12	6	-	-	-
Male	2	2	1	6	6	6	4	6	-	-	-	-
Female	3	5	7	2	13	5	12	6	6	-	-	-
White, both sexes	4	4	5	7	10	7	13	10	4	-	-	-
Male	2	2	1	5	4	5	4	6	-	-	-	-
Female	2	2	4	2	6	2	9	4	4	-	-	-
Black, both sexes	1	3	3	1	9	4	3	2	2	-	-	-
Male	-	-	-	1	2	1	-	-	-	-	-	-
Female	1	3	3	-	7	3	3	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Massachusetts												
All races, both sexes	4	6	7	10	7	19	27	19	7	4	1	-
Male	-	3	3	2	2	9	6	6	-	1	1	-
Female	4	3	4	8	5	10	21	13	7	3	-	-
White, both sexes	3	5	7	10	5	18	26	18	7	4	1	-
Male	-	2	3	2	2	9	6	6	-	1	1	-
Female	3	3	4	8	3	9	20	12	7	3	-	-
Black, both sexes	-	1	-	-	2	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	7	14	10	10	13	27	29	30	18	3	-	-
Male	3	5	5	3	2	10	13	7	4	-	-	-
Female	4	9	5	7	11	17	16	23	14	3	-	-
White, both sexes	6	7	6	7	10	25	25	29	17	3	-	-
Male	3	5	2	2	1	9	11	7	3	-	-	-
Female	3	2	4	5	9	16	14	22	14	3	-	-
Black, both sexes	1	7	4	3	3	2	4	1	1	-	-	-
Male	-	-	3	1	1	1	2	-	1	-	-	-
Female	1	7	1	2	2	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Minnesota												
All races, both sexes	2	5	4	8	6	16	26	19	18	6	1	-
Male	-	2	1	3	-	6	8	5	7	4	-	-
Female	2	3	3	5	6	10	18	14	11	2	1	-
White, both sexes	2	4	2	7	6	15	23	19	18	6	1	-
Male	-	2	1	3	-	5	7	5	7	4	-	-
Female	2	2	1	4	6	10	16	14	11	2	1	-
Black, both sexes	-	1	1	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	5	3	7	5	10	6	5	4	5	1	-	-
Male	4	1	3	-	3	1	1	-	-	-	-	-
Female	1	2	4	5	7	5	4	4	5	1	-	-
White, both sexes	3	2	4	4	5	5	4	3	5	1	-	-
Male	3	1	2	-	2	1	-	-	-	-	-	-
Female	-	1	2	4	3	4	4	3	5	1	-	-
Black, both sexes	2	1	3	1	5	1	1	1	-	-	-	-
Male	1	-	1	-	1	-	1	-	-	-	-	-
Female	1	1	2	1	4	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Missouri												
All races, both sexes	2	6	10	17	12	8	21	11	6	3	-	-
Male	-	3	4	9	1	1	3	4	1	-	-	-
Female	2	3	6	8	11	7	18	7	5	3	-	-
White, both sexes	2	4	8	15	11	8	20	9	5	3	-	-
Male	-	2	4	8	1	1	3	3	1	-	-	-
Female	2	2	4	7	10	7	17	6	4	3	-	-
Black, both sexes	-	2	2	2	1	-	1	2	1	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	-	1	2	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	1	-	1	1	2	3	4	-	-	-	-
Male	1	-	-	1	-	1	-	2	-	-	-	-
Female	1	1	-	-	1	1	3	2	-	-	-	-
White, both sexes	2	1	-	1	1	2	3	4	-	-	-	-
Male	1	-	-	1	-	1	-	2	-	-	-	-
Female	1	1	-	-	1	1	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Nebraska												
All races, both sexes	1	1	2	4	3	1	5	7	3	4	1	-
Male	-	1	-	1	-	-	2	1	-	-	-	-
Female	1	-	2	3	3	1	3	6	3	4	1	-
White, both sexes	1	1	2	3	3	1	5	7	3	4	1	-
Male	-	1	-	1	-	-	2	1	-	-	-	-
Female	1	-	2	2	3	1	3	6	3	4	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	7	2	2	4	3	7	6	1	1	-	-
Male	1	3	1	1	2	2	3	3	-	-	-	-
Female	4	4	1	1	2	1	4	3	1	1	-	-
White, both sexes	4	6	2	1	2	3	7	6	1	-	-	-
Male	1	3	1	1	1	2	3	3	-	-	-	-
Female	3	3	1	-	1	1	4	3	1	-	-	-
Black, both sexes	1	1	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Hampshire												
All races, both sexes	-	3	5	4	3	6	2	-	2	-	-	-
Male	-	1	1	1	2	3	-	-	1	-	-	-
Female	-	2	4	3	1	3	2	-	1	-	-	-
White, both sexes	-	3	5	4	3	6	2	-	2	-	-	-
Male	-	1	1	1	2	3	-	-	1	-	-	-
Female	-	2	4	3	1	3	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	6	12	14	17	15	23	13	7	4	1	-
Male	1	3	2	6	6	4	2	3	1	-	-	-
Female	6	3	10	8	11	11	21	10	6	4	1	-
White, both sexes	3	4	9	12	13	14	17	13	7	4	-	-
Male	1	1	1	6	5	4	2	3	1	-	-	-
Female	2	3	8	6	8	10	15	10	6	4	-	-
Black, both sexes	4	2	3	2	4	-	5	-	-	-	1	-
Male	-	2	1	-	1	-	-	-	-	-	-	-
Female	4	-	2	2	3	-	5	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Mexico												
All races, both sexes	2	1	3	1	5	7	7	5	4	1	-	-
Male	-	1	3	-	2	2	-	1	1	-	-	-
Female	2	-	-	1	3	5	7	4	3	1	-	-
White, both sexes	2	1	2	1	5	7	7	5	4	1	-	-
Male	-	1	2	-	2	2	-	1	1	-	-	-
Female	2	-	-	1	3	5	7	4	3	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	20	19	21	29	31	43	46	25	11	2	-
Male	6	8	10	11	7	12	8	13	8	3	1	-
Female	7	12	9	10	22	19	35	33	17	8	1	-
White, both sexes	11	16	15	13	23	27	40	41	23	11	2	-
Male	5	6	9	6	7	12	8	12	7	3	1	-
Female	6	10	6	7	16	15	32	29	16	8	1	-
Black, both sexes	2	3	3	8	5	4	2	5	1	-	-	-
Male	1	1	1	5	-	-	-	1	-	-	-	-
Female	1	2	2	3	5	4	2	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
North Carolina												
All races, both sexes	13	12	10	19	28	22	29	27	9	5	2	-
Male	2	6	3	3	13	6	9	8	2	-	-	-
Female	11	6	7	16	15	16	20	19	7	5	2	-
White, both sexes	6	7	6	12	21	19	26	26	7	4	1	-
Male	2	3	2	2	11	6	9	8	2	-	-	-
Female	4	4	4	10	10	13	17	18	5	4	1	-
Black, both sexes	7	4	4	6	7	3	3	-	2	1	1	-
Male	-	3	1	-	2	-	-	-	-	-	-	-
Female	7	1	3	6	5	3	3	-	2	1	1	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	2	1	-	-	1	3	2	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	-	1	1	-	-	-	1	1	-	-	-	-
White, both sexes	-	2	1	-	-	1	3	2	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	-	1	1	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Ohio												
All races, both sexes	8	17	16	21	23	38	33	24	18	3	-	-
Male	5	6	5	8	10	11	12	4	3	-	-	-
Female	3	11	11	13	13	27	21	20	15	3	-	-
White, both sexes	6	9	16	18	17	36	30	22	18	3	-	-
Male	3	2	5	6	7	11	12	4	3	-	-	-
Female	3	7	11	12	10	25	18	18	15	3	-	-
Black, both sexes	2	8	-	2	6	2	3	2	-	-	-	-
Male	2	4	-	1	3	-	-	-	-	-	-	-
Female	-	4	-	1	3	2	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	4	2	5	6	2	8	4	7	-	1	-
Male	2	-	1	1	2	1	2	2	-	-	1	-
Female	-	4	1	4	4	1	6	2	7	-	-	-
White, both sexes	2	2	2	3	5	1	8	4	6	-	1	-
Male	2	-	1	1	2	-	2	2	-	-	1	-
Female	-	2	1	2	3	1	6	2	6	-	-	-
Black, both sexes	-	1	-	2	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Oregon												
All races, both sexes	3	5	8	7	6	12	8	14	9	1	-	-
Male	1	3	-	4	2	1	2	6	1	-	-	-
Female	2	2	8	3	4	11	6	8	8	1	-	-
White, both sexes	3	5	5	7	5	11	7	14	9	1	-	-
Male	1	3	-	4	2	-	2	6	1	-	-	-
Female	2	2	5	3	3	11	5	8	8	1	-	-
Black, both sexes	-	-	2	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	11	18	16	27	28	42	45	57	24	7	2	-
Male	2	6	5	7	10	15	12	13	8	-	-	-
Female	9	12	11	20	18	27	33	44	16	7	2	-
White, both sexes	8	14	13	22	23	36	41	52	23	7	2	-
Male	1	5	5	6	9	12	11	13	8	-	-	-
Female	7	9	8	16	14	24	30	39	15	7	2	-
Black, both sexes	3	4	3	4	5	6	4	5	1	-	-	-
Male	1	1	-	1	1	3	1	-	-	-	-	-
Female	2	3	3	3	4	3	3	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Rhode Island												
All races, both sexes	2	1	1	1	1	3	6	2	3	1	-	-
Male	-	1	-	-	-	2	1	1	-	-	-	-
Female	2	-	1	1	1	1	5	1	3	1	-	-
White, both sexes	1	1	1	1	1	2	5	2	3	1	-	-
Male	-	1	-	-	-	1	1	1	-	-	-	-
Female	1	-	1	1	1	1	4	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	7	6	4	10	17	11	10	14	10	-	-	-
Male	3	3	1	5	8	6	4	4	2	-	-	-
Female	4	3	3	5	9	5	6	10	8	-	-	-
White, both sexes	5	5	3	4	13	9	6	11	10	-	-	-
Male	2	2	1	2	6	5	2	3	2	-	-	-
Female	3	3	2	2	7	4	4	8	8	-	-	-
Black, both sexes	2	1	1	5	4	2	4	3	-	-	-	-
Male	1	1	-	3	2	1	2	1	-	-	-	-
Female	1	-	1	2	2	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
South Dakota												
All races, both sexes	1	-	1	-	2	2	-	1	1	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	1	1	-	-	-
White, both sexes	1	-	1	-	2	2	-	1	1	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	8	6	10	9	20	17	9	4	1	1	-
Male	2	2	3	2	5	6	5	3	1	-	-	-
Female	4	6	3	8	4	14	12	6	3	1	1	-
White, both sexes	3	7	4	9	7	19	15	8	3	1	-	-
Male	1	1	2	2	4	6	4	3	1	-	-	-
Female	2	6	2	7	3	13	11	5	2	1	-	-
Black, both sexes	3	1	2	1	2	1	2	1	1	-	1	-
Male	1	1	1	-	1	-	1	-	-	-	-	-
Female	2	-	1	1	1	1	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Texas												
All races, both sexes	25	28	26	34	32	45	26	41	15	8	1	-
Male	12	12	9	11	13	15	8	11	4	1	-	-
Female	13	16	17	23	19	30	18	30	11	7	1	-
White, both sexes	16	19	19	29	29	35	23	37	13	7	-	-
Male	9	10	7	10	13	13	6	10	3	1	-	-
Female	7	9	12	19	16	22	17	27	10	6	-	-
Black, both sexes	9	9	7	5	3	9	3	4	2	1	1	-
Male	3	2	2	1	-	2	2	1	1	-	-	-
Female	6	7	5	4	3	7	1	3	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	2	4	3	7	4	8	6	1	1	-	-
Male	2	-	1	1	3	-	4	1	-	-	-	-
Female	1	2	3	2	4	4	4	5	1	1	-	-
White, both sexes	3	2	4	3	7	4	8	6	1	1	-	-
Male	2	-	1	1	3	-	4	1	-	-	-	-
Female	1	2	3	2	4	4	4	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Vermont												
All races, both sexes	1	3	1	1	-	2	2	1	2	-	-	-
Male	1	1	1	-	-	-	1	-	-	-	-	-
Female	-	2	-	1	-	2	1	1	2	-	-	-
White, both sexes	1	3	1	1	-	2	2	1	2	-	-	-
Male	1	1	1	-	-	-	1	-	-	-	-	-
Female	-	2	-	1	-	2	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	5	14	12	16	30	24	23	7	2	-	-
Male	6	2	3	4	7	8	9	8	3	-	-	-
Female	5	3	11	8	9	22	15	15	4	2	-	-
White, both sexes	7	2	12	7	11	22	23	22	5	1	-	-
Male	4	-	2	3	5	7	9	8	2	-	-	-
Female	3	2	10	4	6	15	14	14	3	1	-	-
Black, both sexes	4	3	2	5	5	7	1	1	2	1	-	-
Male	2	2	1	1	2	-	-	-	1	-	-	-
Female	2	1	1	4	3	7	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Washington												
All races, both sexes	5	7	7	10	16	16	18	21	9	3	-	-
Male	1	2	3	6	4	6	2	6	3	-	-	-
Female	4	5	4	4	12	10	16	15	6	3	-	-
White, both sexes	5	7	7	8	13	16	17	21	9	3	-	-
Male	1	2	3	5	4	6	2	6	3	-	-	-
Female	4	5	4	3	9	10	15	15	6	3	-	-
Black, both sexes	-	-	-	1	3	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	3	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
West Virginia												
All races, both sexes	4	4	4	4	3	8	7	6	4	2	-	-
Male	2	-	1	2	3	2	2	2	-	-	-	-
Female	2	4	3	2	-	6	5	4	4	2	-	-
White, both sexes	3	4	4	4	3	8	7	6	4	2	-	-
Male	1	-	1	2	3	2	2	2	-	-	-	-
Female	2	4	3	2	-	6	5	4	4	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Wisconsin												
All races, both sexes	7	5	4	7	17	18	20	25	14	3	2	-
Male	3	1	3	5	8	8	8	11	2	-	-	-
Female	4	4	1	2	9	10	12	14	12	3	2	-
White, both sexes	5	2	4	7	16	18	20	24	14	3	2	-
Male	2	1	3	5	8	8	8	10	2	-	-	-
Female	3	1	1	2	8	10	12	14	12	3	2	-
Black, both sexes	2	3	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	3	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	1	1	2	3	5	1	1	-	-
Male	1	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	5	1	1	-	-
White, both sexes	1	-	-	1	-	2	2	5	1	1	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
United States												
All races, both sexes	157,861	239	130	73	133	274	491	675	920	1,299	2,177	3,480
Male	70,431	127	72	41	78	185	324	465	630	861	1,395	2,296
Female	87,430	112	58	32	55	89	167	210	290	438	782	1,184
White, both sexes	137,283	149	91	49	92	181	334	432	577	827	1,526	2,456
Male	60,314	79	52	29	56	125	233	306	409	547	989	1,642
Female	76,969	70	39	20	36	56	101	126	168	280	537	814
Black, both sexes	17,755	74	33	18	36	85	133	214	312	432	589	930
Male	8,680	40	19	8	22	54	76	138	199	282	360	589
Female	9,075	34	14	10	14	31	57	76	113	150	229	341
American Indian 1/, both sexes	623	5	-	-	2	3	8	7	7	19	27	37
Male	332	2	-	-	-	1	3	5	4	16	21	26
Female	291	3	-	-	2	2	5	2	3	3	6	11
Asian or Pacific Islander 2/, both sexes	2,200	11	6	6	3	5	16	22	24	21	35	57
Male	1,105	6	1	4	-	5	12	16	18	16	25	39
Female	1,095	5	5	2	3	-	4	6	6	5	10	18
Alabama												
All races, both sexes	5,805	7	3	4	4	7	13	35	48	74	101	183
Male	2,645	3	2	1	1	5	9	27	32	46	62	113
Female	3,160	4	1	3	3	2	4	8	16	28	39	70
White, both sexes	4,406	5	1	1	3	5	10	20	31	37	66	111
Male	1,979	3	1	-	-	3	8	17	19	24	42	67
Female	2,427	2	-	1	3	2	2	3	12	13	24	44
Black, both sexes	1,379	2	2	3	1	2	3	13	16	36	35	72
Male	656	-	1	1	1	2	1	8	12	21	20	46
Female	723	2	1	2	-	-	2	5	4	15	15	26
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	2	1	-	-	-
Male	8	-	-	-	-	-	-	2	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Alaska												
All races, both sexes	219	1	-	-	1	-	1	2	-	2	2	6
Male	110	-	-	-	-	-	1	-	-	2	2	2
Female	109	1	-	-	1	-	-	2	-	-	-	4
White, both sexes	159	-	-	-	-	-	-	1	-	-	1	3
Male	83	-	-	-	-	-	-	-	-	-	1	1
Female	76	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	41	1	-	-	1	-	1	1	-	1	-	2
Male	16	-	-	-	-	-	1	-	-	1	-	-
Female	25	1	-	-	1	-	-	1	-	-	-	2
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,030	8	5	2	4	5	10	11	12	25	23	51
Male	1,015	6	3	1	2	2	8	10	11	22	17	34
Female	1,015	2	2	1	2	3	2	1	1	3	6	17
White, both sexes	1,916	6	5	2	4	3	8	8	9	19	18	45
Male	938	4	3	1	2	1	6	8	8	16	12	31
Female	978	2	2	1	2	2	2	-	1	3	6	14
Black, both sexes	52	1	-	-	-	2	-	2	1	3	2	4
Male	39	1	-	-	-	1	-	1	1	3	2	2
Female	13	-	-	-	-	1	-	1	-	-	-	2
American Indian 1/, both sexes	57	1	-	-	-	-	2	1	1	3	3	2
Male	34	1	-	-	-	-	2	1	1	3	3	1
Female	23	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Arkansas												
All races, both sexes	1,760	7	5	1	4	3	9	7	8	10	23	49
Male	794	4	3	1	2	1	5	4	8	4	12	30
Female	966	3	2	-	2	2	4	3	-	6	11	19
White, both sexes	1,524	4	3	1	2	3	5	6	5	4	10	31
Male	678	2	3	1	2	1	2	4	5	3	5	19
Female	846	2	-	-	-	2	3	2	-	1	5	12
Black, both sexes	232	3	2	-	2	-	4	1	3	6	13	18
Male	114	2	-	-	-	-	3	-	3	1	7	11
Female	118	1	2	-	2	-	1	1	-	5	6	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12,714	22	10	7	17	18	41	67	104	138	235	342
Male	5,972	11	7	5	8	16	27	53	78	95	174	242
Female	6,742	11	3	2	9	2	14	14	26	43	61	100
White, both sexes	10,998	13	8	4	15	14	32	50	76	101	189	266
Male	5,132	7	6	3	7	13	21	39	63	66	141	195
Female	5,866	6	2	1	8	1	11	11	13	35	48	71
Black, both sexes	921	2	-	1	1	3	6	10	20	26	34	52
Male	458	1	-	1	1	2	4	8	10	21	24	29
Female	463	1	-	-	-	1	2	2	10	5	10	23
American Indian 1/, both sexes	67	-	-	-	-	-	-	1	-	3	1	7
Male	41	-	-	-	-	-	-	1	-	3	-	7
Female	26	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	728	7	2	2	1	1	3	6	8	8	11	17
Male	341	3	1	1	-	1	2	5	5	5	9	11
Female	387	4	1	1	1	-	1	1	3	3	2	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Colorado												
All races, both sexes	1,646	2	2	2	3	4	10	6	7	8	25	32
Male	688	-	1	2	2	1	7	2	5	5	16	22
Female	958	2	1	-	1	3	3	4	2	3	9	10
White, both sexes	1,577	2	2	1	3	3	9	6	6	7	24	29
Male	655	-	1	1	2	1	7	2	4	4	15	19
Female	922	2	1	-	1	2	2	4	2	3	9	10
Black, both sexes	38	-	-	-	-	1	1	-	1	1	-	3
Male	18	-	-	-	-	-	-	-	1	1	-	3
Female	20	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	1	-	-	-	-	-	-	-	-
Male	8	-	-	1	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,393	1	1	-	-	5	5	11	6	15	20	43
Male	1,055	1	-	-	-	2	4	10	5	12	10	25
Female	1,338	-	1	-	-	3	1	1	1	3	10	18
White, both sexes	2,263	1	1	-	-	5	5	7	4	12	14	36
Male	990	1	-	-	-	2	4	6	4	9	8	20
Female	1,273	-	1	-	-	3	1	1	-	3	6	16
Black, both sexes	120	-	-	-	-	-	-	4	2	3	6	7
Male	61	-	-	-	-	-	-	4	1	3	2	5
Female	59	-	-	-	-	-	-	-	1	-	4	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Delaware												
All races, both sexes	420	2	-	-	-	-	1	2	6	6	6	7
Male	195	1	-	-	-	-	1	1	3	6	5	5
Female	225	1	-	-	-	-	-	1	3	-	1	2
White, both sexes	354	-	-	-	-	-	1	2	3	3	2	7
Male	165	-	-	-	-	-	1	1	2	3	2	5
Female	189	-	-	-	-	-	-	1	1	-	-	2
Black, both sexes	63	1	-	-	-	-	-	-	3	3	4	-
Male	29	-	-	-	-	-	-	-	1	3	3	-
Female	34	1	-	-	-	-	-	-	2	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	182	1	1	-	1	1	2	3	-	2	5	4
Male	78	-	1	-	-	1	-	2	-	1	4	2
Female	104	1	-	-	1	-	2	1	-	1	1	2
White, both sexes	44	-	-	-	-	-	1	-	-	1	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	137	1	1	-	1	1	1	3	-	1	5	4
Male	58	-	1	-	-	1	-	2	-	1	4	2
Female	79	1	-	-	1	-	1	1	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Florida												
All races, both sexes	8,184	12	11	5	10	12	28	32	44	60	93	172
Male	3,938	6	6	4	5	8	19	21	33	43	59	108
Female	4,246	6	5	1	5	4	9	11	11	17	34	64
White, both sexes	7,319	8	7	4	7	10	15	16	26	35	61	112
Male	3,490	4	4	3	5	7	13	12	19	25	43	71
Female	3,829	4	3	1	2	3	2	4	7	10	18	41
Black, both sexes	831	4	3	1	3	2	13	15	18	25	31	59
Male	432	2	2	1	-	1	6	9	14	18	15	36
Female	399	2	1	-	3	1	7	6	4	7	16	23
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	1	-	-	-	-	1	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	1	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	6,426	12	5	5	6	14	16	31	39	67	104	174
Male	3,000	8	3	3	2	9	6	21	27	49	62	118
Female	3,426	4	2	2	4	5	10	10	12	18	42	56
White, both sexes	4,728	7	4	3	4	5	5	12	18	28	61	101
Male	2,183	5	3	2	1	3	2	9	12	23	38	73
Female	2,545	2	1	1	3	2	3	3	6	5	23	28
Black, both sexes	1,649	5	1	2	2	8	9	19	19	39	42	73
Male	796	3	-	1	1	6	3	12	13	26	24	45
Female	853	2	1	1	1	2	6	7	6	13	18	28
American Indian 1/, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	42	-	-	-	-	-	2	-	2	-	1	-
Male	16	-	-	-	-	-	1	-	2	-	-	-
Female	26	-	-	-	-	-	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Hawaii												
All races, both sexes	817	-	1	1	-	1	3	4	4	8	13	29
Male	444	-	-	1	-	1	2	3	3	5	7	16
Female	373	-	1	-	-	-	1	1	1	3	6	13
White, both sexes	191	-	-	-	-	-	-	-	1	3	2	6
Male	100	-	-	-	-	-	-	-	1	1	2	1
Female	91	-	-	-	-	-	-	-	-	2	-	5
Black, both sexes	5	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	619	-	1	1	-	1	2	4	3	4	11	22
Male	342	-	-	1	-	1	2	3	2	4	5	15
Female	277	-	1	-	-	-	-	1	1	-	6	7
Idaho												
All races, both sexes	622	1	2	1	-	-	1	3	2	1	7	11
Male	269	1	1	-	-	-	1	3	2	1	3	7
Female	353	-	1	1	-	-	-	-	-	-	4	4
White, both sexes	615	1	2	1	-	-	1	3	2	1	7	10
Male	266	1	1	-	-	-	1	3	2	1	3	7
Female	349	-	1	1	-	-	-	-	-	-	4	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Illinois												
All races, both sexes	7,095	7	4	1	4	20	24	45	47	62	84	138
Male	3,035	4	2	-	2	17	17	31	27	40	54	92
Female	4,060	3	2	1	2	3	7	14	20	22	30	46
White, both sexes	6,155	4	3	1	2	11	17	30	27	34	52	91
Male	2,584	2	1	-	1	9	12	22	17	21	34	64
Female	3,571	2	2	1	1	2	5	8	10	13	18	27
Black, both sexes	880	3	1	-	1	8	7	13	19	27	31	47
Male	422	2	1	-	1	7	5	8	10	19	19	28
Female	458	1	-	-	-	1	2	5	9	8	12	19
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	58	-	-	-	1	1	-	2	1	1	1	-
Male	28	-	-	-	-	1	-	1	-	-	1	-
Female	30	-	-	-	1	-	-	1	1	1	-	-
Indiana												
All races, both sexes	4,229	13	3	1	2	9	14	14	26	28	70	89
Male	1,834	6	2	-	-	7	8	7	15	18	46	60
Female	2,395	7	1	1	2	2	6	7	11	10	24	29
White, both sexes	3,916	5	3	1	2	6	12	12	18	23	61	73
Male	1,679	3	2	-	-	6	8	7	10	15	39	49
Female	2,237	2	1	1	2	-	4	5	8	8	22	24
Black, both sexes	309	8	-	-	-	3	2	2	8	5	9	16
Male	154	3	-	-	-	1	-	-	5	3	7	11
Female	155	5	-	-	-	2	2	2	3	2	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Iowa												
All races, both sexes	1,148	1	-	-	1	2	3	1	1	10	13	14
Male	485	1	-	-	-	1	1	1	-	6	9	9
Female	663	-	-	-	1	1	2	-	1	4	4	5
White, both sexes	1,124	-	-	-	1	2	3	1	1	8	12	14
Male	474	-	-	-	-	1	1	1	-	4	8	9
Female	650	-	-	-	1	1	2	-	1	4	4	5
Black, both sexes	22	1	-	-	-	-	-	-	-	1	1	-
Male	10	1	-	-	-	-	-	-	-	1	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,920	3	1	1	-	4	4	9	3	11	22	22
Male	767	1	1	-	-	2	4	4	3	4	14	14
Female	1,153	2	-	1	-	2	-	5	-	7	8	8
White, both sexes	1,805	2	-	1	-	3	4	7	1	9	17	18
Male	716	1	-	-	-	1	4	3	1	3	10	12
Female	1,089	1	-	1	-	2	-	4	-	6	7	6
Black, both sexes	98	1	1	-	-	-	-	2	1	2	4	3
Male	40	-	1	-	-	-	-	1	1	1	3	1
Female	58	1	-	-	-	-	-	1	-	1	1	2
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	1	-	-	1	-	-	-
Male	4	-	-	-	-	1	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Kentucky												
All races, both sexes	3,143	-	2	1	1	5	10	12	16	21	42	82
Male	1,339	-	1	1	1	3	5	10	8	11	28	53
Female	1,804	-	1	-	-	2	5	2	8	10	14	29
White, both sexes	2,940	-	2	1	1	3	8	9	11	17	35	69
Male	1,253	-	1	1	1	2	4	7	7	10	23	46
Female	1,687	-	1	-	-	1	4	2	4	7	12	23
Black, both sexes	191	-	-	-	-	2	2	3	5	4	5	12
Male	81	-	-	-	-	1	1	3	1	1	4	7
Female	110	-	-	-	-	1	1	-	4	3	1	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Louisiana												
All races, both sexes	3,048	10	6	2	2	2	15	15	25	26	44	69
Male	1,380	5	5	-	1	1	8	8	12	21	31	46
Female	1,668	5	1	2	1	1	7	7	13	5	13	23
White, both sexes	2,242	4	4	-	1	1	10	7	12	15	22	32
Male	1,004	2	3	-	-	1	6	5	6	13	16	21
Female	1,238	2	1	-	1	-	4	2	6	2	6	11
Black, both sexes	798	6	2	2	1	1	5	8	13	11	21	37
Male	372	3	2	-	1	-	2	3	6	8	14	25
Female	426	3	-	2	-	1	3	5	7	3	7	12
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Maine												
All races, both sexes	847	-	1	-	1	3	2	3	3	6	10	15
Male	412	-	1	-	1	2	1	3	3	3	7	13
Female	435	-	-	-	-	1	1	-	-	3	3	2
White, both sexes	839	-	1	-	1	3	2	3	3	5	9	15
Male	406	-	1	-	1	2	1	3	3	2	6	13
Female	433	-	-	-	-	1	1	-	-	3	3	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,326	3	2	1	5	6	8	13	27	30	50	58
Male	1,032	2	1	-	3	4	5	12	22	18	31	34
Female	1,294	1	1	1	2	2	3	1	5	12	19	24
White, both sexes	1,739	1	1	1	2	3	5	8	12	14	25	35
Male	754	1	1	-	2	2	4	8	9	9	20	19
Female	985	-	-	1	-	1	1	-	3	5	5	16
Black, both sexes	552	2	-	-	3	3	3	5	14	16	25	23
Male	262	1	-	-	1	2	1	4	12	9	11	15
Female	290	1	-	-	2	1	2	1	2	7	14	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	1	-	-	-	-	-	1	-	-	-
Male	15	-	-	-	-	-	-	-	1	-	-	-
Female	16	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Massachusetts												
All races, both sexes	4,011	1	2	1	2	3	10	15	9	16	32	51
Male	1,654	1	1	1	2	3	9	10	7	11	22	41
Female	2,357	-	1	-	-	-	1	5	2	5	10	10
White, both sexes	3,864	1	2	1	1	2	10	11	8	13	27	46
Male	1,589	1	1	1	1	2	9	8	7	9	17	37
Female	2,275	-	1	-	-	-	1	3	1	4	10	9
Black, both sexes	122	-	-	-	1	1	-	4	1	3	5	4
Male	58	-	-	-	1	1	-	2	-	2	5	3
Female	64	-	-	-	-	-	-	2	1	1	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5,576	6	4	5	2	7	17	30	35	60	90	122
Male	2,419	4	2	3	1	5	13	23	28	41	65	74
Female	3,157	2	2	2	1	2	4	7	7	19	25	48
White, both sexes	4,885	3	2	5	2	3	8	15	15	34	56	74
Male	2,074	2	1	3	1	2	6	12	13	26	43	48
Female	2,811	1	1	2	1	1	2	3	2	8	13	26
Black, both sexes	650	2	2	-	-	4	8	15	20	26	33	45
Male	317	1	1	-	-	3	7	11	15	15	21	23
Female	333	1	1	-	-	1	1	4	5	11	12	22
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	-	1	3
Male	15	-	-	-	-	-	-	-	-	-	1	3
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	1	-	-	-	-	1	-	-	-	-	-
Male	13	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Minnesota												
All races, both sexes	2,616	3	2	3	-	4	9	5	13	15	22	33
Male	1,137	-	1	-	-	2	4	1	9	7	12	24
Female	1,479	3	1	3	-	2	5	4	4	8	10	9
White, both sexes	2,536	2	2	3	-	1	8	5	10	10	19	26
Male	1,092	-	1	-	-	1	3	1	8	4	10	18
Female	1,444	2	1	3	-	-	5	4	2	6	9	8
Black, both sexes	46	-	-	-	-	3	1	-	2	3	2	5
Male	27	-	-	-	-	1	1	-	1	2	1	4
Female	19	-	-	-	-	2	-	-	1	1	1	1
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	21	1	-	-	-	-	-	-	-	2	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	10	1	-	-	-	-	-	-	-	1	-	-
Mississippi												
All races, both sexes	2,811	2	-	2	4	2	10	11	16	17	37	65
Male	1,212	1	-	-	4	1	5	4	9	13	21	39
Female	1,599	1	-	2	-	1	5	7	7	4	16	26
White, both sexes	1,931	2	-	-	1	-	3	4	5	7	16	28
Male	822	1	-	-	1	-	2	2	3	5	9	16
Female	1,109	1	-	-	-	-	1	2	2	2	7	12
Black, both sexes	872	-	-	2	3	2	7	7	11	10	21	37
Male	384	-	-	-	3	1	3	2	6	8	12	23
Female	488	-	-	2	-	1	4	5	5	2	9	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Missouri												
All races, both sexes	3,638	2	2	-	3	7	13	6	16	30	49	72
Male	1,522	-	1	-	2	4	8	5	10	20	31	49
Female	2,116	2	1	-	1	3	5	1	6	10	18	23
White, both sexes	3,325	2	1	-	3	7	8	4	9	19	42	55
Male	1,382	-	-	-	2	4	5	4	6	14	28	39
Female	1,943	2	1	-	1	3	3	-	3	5	14	16
Black, both sexes	304	-	1	-	-	-	4	1	7	11	7	17
Male	136	-	1	-	-	-	3	-	4	6	3	10
Female	168	-	-	-	-	-	1	1	3	5	4	7
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	1	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	666	-	-	-	1	-	1	2	3	6	3	8
Male	316	-	-	-	1	-	1	2	3	4	2	5
Female	350	-	-	-	-	-	-	-	-	2	1	3
White, both sexes	640	-	-	-	1	-	1	1	3	5	3	5
Male	302	-	-	-	1	-	1	1	3	3	2	4
Female	338	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	1	-	3
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Nebraska												
All races, both sexes	1,355	-	-	-	2	-	1	4	4	10	10	23
Male	574	-	-	-	1	-	1	2	3	8	5	14
Female	781	-	-	-	1	-	-	2	1	2	5	9
White, both sexes	1,302	-	-	-	1	-	1	2	4	9	8	18
Male	546	-	-	-	1	-	1	1	3	7	4	10
Female	756	-	-	-	-	-	-	1	1	2	4	8
Black, both sexes	44	-	-	-	1	-	-	2	-	1	1	4
Male	24	-	-	-	-	-	-	1	-	1	1	3
Female	20	-	-	-	1	-	-	1	-	-	-	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,782	2	1	1	-	1	6	13	23	19	46	54
Male	950	1	1	1	-	1	4	8	17	10	23	37
Female	832	1	-	-	-	-	2	5	6	9	23	17
White, both sexes	1,542	2	-	1	-	1	3	9	16	12	36	42
Male	825	1	-	1	-	1	2	5	11	6	20	28
Female	717	1	-	-	-	-	1	4	5	6	16	14
Black, both sexes	166	-	1	-	-	-	1	3	4	7	9	10
Male	85	-	1	-	-	-	1	2	3	4	2	8
Female	81	-	-	-	-	-	-	1	1	3	7	2
American Indian 1/, both sexes	20	-	-	-	-	-	1	1	1	-	-	1
Male	10	-	-	-	-	-	-	1	1	-	-	1
Female	10	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	1	-	2	-	1	1
Male	30	-	-	-	-	-	1	-	2	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
New Hampshire												
All races, both sexes	676	-	1	-	1	4	2	-	2	6	8	8
Male	289	-	1	-	1	1	2	-	1	4	3	6
Female	387	-	-	-	-	3	-	-	1	2	5	2
White, both sexes	675	-	1	-	1	4	2	-	2	6	7	8
Male	288	-	1	-	1	1	2	-	1	4	2	6
Female	387	-	-	-	-	3	-	-	1	2	5	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,549	6	3	1	4	1	11	19	24	33	59	78
Male	2,020	4	1	1	3	1	7	18	19	24	41	55
Female	2,529	2	2	-	1	-	4	1	5	9	18	23
White, both sexes	4,010	2	2	1	4	1	6	12	16	22	40	46
Male	1,740	1	-	1	3	1	5	11	13	16	29	32
Female	2,270	1	2	-	1	-	1	1	3	6	11	14
Black, both sexes	479	4	1	-	-	-	5	7	7	9	19	31
Male	252	3	1	-	-	-	2	7	5	6	12	23
Female	227	1	-	-	-	-	3	-	2	3	7	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	58	-	-	-	-	-	-	-	1	2	-	1
Male	28	-	-	-	-	-	-	-	1	2	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
New Mexico												
All races, both sexes	867	1	-	-	-	1	2	3	8	5	12	18
Male	376	1	-	-	-	1	1	1	3	4	9	12
Female	491	-	-	-	-	-	1	2	5	1	3	6
White, both sexes	819	1	-	-	-	1	2	3	7	3	7	15
Male	350	1	-	-	-	1	1	1	3	2	4	10
Female	469	-	-	-	-	-	1	2	4	1	3	5
Black, both sexes	18	-	-	-	-	-	-	-	-	-	2	1
Male	10	-	-	-	-	-	-	-	-	-	2	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	1	2	3	2
Male	12	-	-	-	-	-	-	-	-	2	3	1
Female	11	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7,528	7	8	5	7	20	16	34	43	62	103	174
Male	3,340	6	5	3	5	14	14	23	28	43	59	114
Female	4,188	1	3	2	2	6	2	11	15	19	44	60
White, both sexes	6,816	5	6	2	5	15	8	25	31	46	72	127
Male	2,985	5	3	2	4	10	7	15	18	33	40	83
Female	3,831	-	3	-	1	5	1	10	13	13	32	44
Black, both sexes	619	2	2	2	2	5	6	9	12	16	27	43
Male	305	1	2	1	1	4	5	8	10	10	16	29
Female	314	1	-	1	1	1	1	1	2	6	11	14
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	82	-	-	1	-	-	2	-	-	-	4	3
Male	46	-	-	-	-	-	2	-	-	-	3	2
Female	36	-	-	1	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
North Carolina												
All races, both sexes	4,812	11	8	4	6	9	19	23	24	40	69	90
Male	2,167	6	2	3	5	7	10	13	15	28	44	62
Female	2,645	5	6	1	1	2	9	10	9	12	25	28
White, both sexes	3,769	6	5	3	1	5	9	14	11	19	38	58
Male	1,656	3	2	2	1	5	4	9	7	15	25	39
Female	2,113	3	3	1	-	-	5	5	4	4	13	19
Black, both sexes	991	4	2	1	5	4	9	9	13	21	29	31
Male	477	3	-	1	4	2	6	4	8	13	17	22
Female	514	1	2	-	1	2	3	5	5	8	12	9
American Indian 1/, both sexes	31	1	-	-	-	-	1	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	12	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	1	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	310	-	-	-	1	-	1	1	1	1	2	2
Male	120	-	-	-	1	-	-	-	1	1	2	1
Female	190	-	-	-	-	-	1	1	-	-	-	1
White, both sexes	294	-	-	-	1	-	1	1	-	1	1	-
Male	113	-	-	-	1	-	-	-	-	1	1	-
Female	181	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	1	-	1	2
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Ohio												
All races, both sexes	5,699	14	4	1	1	12	20	19	34	40	66	131
Male	2,425	6	2	1	1	11	11	13	24	30	38	88
Female	3,274	8	2	-	-	1	9	6	10	10	28	43
White, both sexes	5,126	10	3	1	1	7	19	11	26	24	55	98
Male	2,144	4	2	1	1	7	11	6	19	18	33	72
Female	2,982	6	1	-	-	-	8	5	7	6	22	26
Black, both sexes	559	4	1	-	-	5	1	8	8	16	11	33
Male	274	2	-	-	-	4	-	7	5	12	5	16
Female	285	2	1	-	-	1	1	1	3	4	6	17
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,397	2	2	-	3	6	5	16	13	17	37	49
Male	1,033	2	-	-	1	3	3	13	10	10	22	34
Female	1,364	-	2	-	2	3	2	3	3	7	15	15
White, both sexes	2,141	1	2	-	2	4	4	9	7	10	31	35
Male	903	1	-	-	-	2	2	8	6	6	16	24
Female	1,238	-	2	-	2	2	2	1	1	4	15	11
Black, both sexes	142	-	-	-	1	1	1	4	4	4	-	10
Male	71	-	-	-	1	-	1	4	3	2	-	8
Female	71	-	-	-	-	1	-	-	1	2	-	2
American Indian 1/, both sexes	109	1	-	-	-	1	-	2	2	3	6	4
Male	59	1	-	-	-	1	-	1	1	2	6	2
Female	50	-	-	-	-	-	-	1	1	1	-	2
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Oregon												
All races, both sexes	2,032	2	-	2	2	1	5	7	9	13	12	25
Male	883	-	-	1	1	1	1	5	6	10	9	17
Female	1,149	2	-	1	1	-	4	2	3	3	3	8
White, both sexes	1,966	1	-	2	2	1	5	5	6	13	11	21
Male	852	-	-	1	1	1	1	3	3	10	9	13
Female	1,114	1	-	1	1	-	4	2	3	3	2	8
Black, both sexes	22	-	-	-	-	-	-	1	3	-	-	2
Male	14	-	-	-	-	-	-	1	3	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	1	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	11	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	1	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	9,964	10	3	1	7	7	18	30	33	50	102	173
Male	4,265	5	2	1	5	2	13	17	21	31	68	119
Female	5,699	5	1	-	2	5	5	13	12	19	34	54
White, both sexes	9,152	9	2	-	5	5	11	22	21	33	74	132
Male	3,846	5	2	-	3	2	7	12	13	19	47	90
Female	5,306	4	-	-	2	3	4	10	8	14	27	42
Black, both sexes	784	1	1	1	2	2	7	8	12	17	28	40
Male	405	-	-	1	2	-	6	5	8	12	21	28
Female	379	1	1	-	-	2	1	3	4	5	7	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Rhode Island												
All races, both sexes	419	-	-	-	1	2	1	3	2	4	2	9
Male	166	-	-	-	-	2	1	3	2	1	2	5
Female	253	-	-	-	1	-	-	-	-	3	-	4
White, both sexes	407	-	-	-	-	2	1	2	2	4	2	8
Male	159	-	-	-	-	2	1	2	2	1	2	4
Female	248	-	-	-	-	-	-	-	-	3	-	4
Black, both sexes	10	-	-	-	1	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	5	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,896	8	3	1	-	5	10	19	22	27	58	99
Male	1,419	4	1	-	-	3	8	16	10	18	40	64
Female	1,477	4	2	1	-	2	2	3	12	9	18	35
White, both sexes	2,018	3	1	1	-	1	4	9	9	11	31	54
Male	969	1	-	-	-	1	4	7	3	8	24	36
Female	1,049	2	1	1	-	-	-	2	6	3	7	18
Black, both sexes	866	5	2	-	-	4	6	10	13	16	27	45
Male	447	3	1	-	-	2	4	9	7	10	16	28
Female	419	2	1	-	-	2	2	1	6	6	11	17
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
South Dakota												
All races, both sexes	261	1	-	-	1	-	3	1	1	1	7	9
Male	117	1	-	-	1	-	-	1	-	1	7	7
Female	144	-	-	-	-	-	3	-	1	-	-	2
White, both sexes	251	-	-	-	1	-	-	1	1	1	6	8
Male	111	-	-	-	1	-	-	1	-	1	6	6
Female	140	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	3	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2,732	9	1	-	2	10	11	16	29	39	65	84
Male	1,253	5	-	-	1	5	7	11	20	21	36	52
Female	1,479	4	1	-	1	5	4	5	9	18	29	32
White, both sexes	2,267	6	1	-	1	6	5	11	18	26	44	65
Male	1,016	3	-	-	-	2	4	10	12	16	21	43
Female	1,251	3	1	-	1	4	1	1	6	10	23	22
Black, both sexes	454	3	-	-	1	3	6	4	10	13	21	19
Male	231	2	-	-	1	2	3	-	7	5	15	9
Female	223	1	-	-	-	1	3	4	3	8	6	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	1	-	1	1	-	-	-
Male	5	-	-	-	-	1	-	1	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Texas												
All races, both sexes	9,769	26	13	6	9	24	42	40	80	82	153	248
Male	4,343	15	7	4	6	15	32	23	53	52	90	151
Female	5,426	11	6	2	3	9	10	17	27	30	63	97
White, both sexes	8,487	21	9	4	7	17	33	24	55	52	111	184
Male	3,714	11	5	3	5	10	27	13	41	32	66	107
Female	4,773	10	4	1	2	7	6	11	14	20	45	77
Black, both sexes	1,187	4	4	2	1	7	9	16	25	30	42	59
Male	582	3	2	1	1	5	5	10	12	20	24	40
Female	605	1	2	1	-	2	4	6	13	10	18	19
American Indian 1/, both sexes	15	-	-	-	1	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	80	1	-	-	-	-	-	-	-	-	-	5
Male	38	1	-	-	-	-	-	-	-	-	-	4
Female	42	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	1,211	1	1	-	2	2	5	4	6	10	6	20
Male	521	-	-	-	2	2	4	1	6	3	5	13
Female	690	1	1	-	-	-	1	3	-	7	1	7
White, both sexes	1,180	1	1	-	2	2	4	3	6	10	4	19
Male	501	-	-	-	2	2	3	1	6	3	3	12
Female	679	1	1	-	-	-	1	2	-	7	1	7
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	1	1	-	-	-	-
Male	12	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Vermont												
All races, both sexes	266	-	-	1	1	1	-	-	-	2	3	1
Male	108	-	-	-	1	-	-	-	-	2	1	-
Female	158	-	-	1	-	1	-	-	-	-	2	1
White, both sexes	265	-	-	1	1	1	-	-	-	2	3	1
Male	107	-	-	-	1	-	-	-	-	2	1	-
Female	158	-	-	1	-	1	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4,612	6	5	3	3	16	14	9	20	34	68	116
Male	2,091	2	4	2	2	12	12	6	15	23	46	84
Female	2,521	4	1	1	1	4	2	3	5	11	22	32
White, both sexes	3,647	4	3	1	-	10	9	7	6	22	37	60
Male	1,598	1	2	1	-	9	8	5	5	15	23	44
Female	2,049	3	1	-	-	1	1	2	1	7	14	16
Black, both sexes	915	2	2	1	3	6	4	2	13	12	31	55
Male	461	1	2	-	2	3	3	1	9	8	23	39
Female	454	1	-	1	1	3	1	1	4	4	8	16
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	1	-	-	1	-	1	-	-	1
Male	31	-	-	1	-	-	1	-	1	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Washington												
All races, both sexes	2,146	2	1	-	-	3	6	9	8	26	18	49
Male	959	1	-	-	-	2	4	8	4	16	11	33
Female	1,187	1	1	-	-	1	2	1	4	10	7	16
White, both sexes	2,019	1	-	-	-	2	4	6	7	16	15	44
Male	895	-	-	-	-	2	3	6	4	8	10	29
Female	1,124	1	-	-	-	-	1	-	3	8	5	15
Black, both sexes	53	1	1	-	-	-	1	1	1	7	1	3
Male	29	1	-	-	-	-	-	-	-	5	-	2
Female	24	-	1	-	-	-	1	1	1	2	1	1
American Indian 1/, both sexes	27	-	-	-	-	1	-	1	-	1	1	2
Male	14	-	-	-	-	-	-	1	-	1	-	2
Female	13	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	1	1	-	2	1	-
Male	21	-	-	-	-	-	1	1	-	2	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,503	2	1	-	1	1	4	3	4	9	8	25
Male	624	1	1	-	1	1	2	3	3	1	5	17
Female	879	1	-	-	-	-	2	-	1	8	3	8
White, both sexes	1,448	1	1	-	1	1	4	3	4	9	7	24
Male	602	1	1	-	1	1	2	3	3	1	4	16
Female	846	-	-	-	-	-	2	-	1	8	3	8
Black, both sexes	54	1	-	-	-	-	-	-	-	-	1	1
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	32	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Wisconsin												
All races, both sexes	3,440	2	-	-	1	3	7	7	9	14	34	43
Male	1,489	1	-	-	-	3	7	2	5	11	21	26
Female	1,951	1	-	-	1	-	-	5	4	3	13	17
White, both sexes	3,308	2	-	-	-	1	6	5	5	11	32	40
Male	1,434	1	-	-	-	1	6	2	3	9	20	26
Female	1,874	1	-	-	-	-	-	3	2	2	12	14
Black, both sexes	99	-	-	-	-	2	-	2	3	1	1	3
Male	41	-	-	-	-	2	-	-	2	1	-	-
Female	58	-	-	-	-	-	-	2	1	-	1	3
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	2	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	1	-	1	-	1	-	-	-
Male	9	-	-	-	-	-	1	-	-	-	-	-
Female	11	-	-	-	1	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	343	-	-	1	-	1	2	-	1	1	4	11
Male	152	-	-	1	-	-	1	-	1	1	2	8
Female	191	-	-	-	-	1	1	-	-	-	2	3
White, both sexes	339	-	-	1	-	1	2	-	1	1	3	11
Male	150	-	-	1	-	-	1	-	1	1	2	8
Female	189	-	-	-	-	1	1	-	-	-	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
United States												
All races, both sexes	4,384	5,750	6,524	7,910	11,160	17,547	26,226	30,673	24,933	10,517	2,339	7
Male	2,895	3,731	4,047	4,718	6,123	8,826	11,747	11,677	7,485	2,333	369	6
Female	1,489	2,019	2,477	3,192	5,037	8,721	14,479	18,996	17,448	8,184	1,970	1
White, both sexes	3,059	4,260	5,104	6,322	9,282	15,367	23,612	28,403	23,282	9,799	2,075	4
Male	2,046	2,810	3,193	3,826	5,121	7,819	10,683	10,887	6,991	2,150	318	3
Female	1,013	1,450	1,911	2,496	4,161	7,548	12,929	17,516	16,291	7,649	1,757	1
Black, both sexes	1,209	1,341	1,267	1,375	1,633	1,855	2,211	1,843	1,337	601	224	3
Male	767	817	763	763	870	854	887	608	378	144	39	3
Female	442	524	504	612	763	1,001	1,324	1,235	959	457	185	-
American Indian 1/, both sexes	36	43	53	62	54	62	75	62	41	13	7	-
Male	25	31	31	39	28	27	32	26	13	-	2	-
Female	11	12	22	23	26	35	43	36	28	13	5	-
Asian or Pacific Islander 2/, both sexes	80	106	100	151	191	263	328	365	273	104	33	-
Male	57	73	60	90	104	126	145	156	103	39	10	-
Female	23	33	40	61	87	137	183	209	170	65	23	-
Alabama												
All races, both sexes	253	321	340	390	490	711	892	918	682	263	66	-
Male	168	203	218	235	274	330	360	302	196	50	8	-
Female	85	118	122	155	216	381	532	616	486	213	58	-
White, both sexes	156	223	233	280	356	570	712	762	565	219	40	-
Male	105	142	149	179	205	265	292	255	159	40	4	-
Female	51	81	84	101	151	305	420	507	406	179	36	-
Black, both sexes	96	98	104	110	132	139	177	153	115	44	26	-
Male	62	61	67	56	69	65	66	46	37	10	4	-
Female	34	37	37	54	63	74	111	107	78	34	22	-
American Indian 1/, both sexes	-	-	1	-	1	2	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	1	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	1	-	2	2	1	-	-	-
Male	1	-	2	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Alaska												
All races, both sexes	15	11	12	11	29	19	29	39	26	11	2	-
Male	11	8	10	7	18	12	9	18	8	1	1	-
Female	4	3	2	4	11	7	20	21	18	10	1	-
White, both sexes	10	8	7	7	24	12	23	27	24	10	2	-
Male	8	6	6	4	15	10	9	13	8	1	1	-
Female	2	2	1	3	9	2	14	14	16	9	1	-
Black, both sexes	-	2	1	-	1	1	-	-	-	-	-	-
Male	-	2	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	1	4	1	3	5	3	9	2	1	-	-
Male	3	-	3	1	2	1	-	4	-	-	-	-
Female	2	1	1	-	1	4	3	5	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	1	1	3	3	-	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	1	3	2	-	-	-	-
Arizona												
All races, both sexes	66	72	97	125	155	220	342	378	292	105	22	-
Male	46	42	63	84	88	115	155	166	106	27	7	-
Female	20	30	34	41	67	105	187	212	186	78	15	-
White, both sexes	56	64	89	115	150	208	329	371	286	103	18	-
Male	38	36	56	77	85	109	145	164	105	27	4	-
Female	18	28	33	38	65	99	184	207	181	76	14	-
Black, both sexes	5	4	2	6	1	4	8	3	2	1	1	-
Male	4	4	2	5	-	3	8	1	-	-	1	-
Female	1	-	-	1	1	1	-	2	2	1	-	-
American Indian 1/, both sexes	4	4	6	4	4	8	4	3	4	1	2	-
Male	3	2	5	2	3	3	2	-	1	-	1	-
Female	1	2	1	2	1	5	2	3	3	1	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	1	-	-	1	-
Male	1	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Arkansas												
All races, both sexes	34	63	84	100	149	208	284	342	244	101	25	-
Male	25	42	52	61	77	112	126	124	73	24	4	-
Female	9	21	32	39	72	96	158	218	171	77	21	-
White, both sexes	24	46	62	84	126	195	267	320	220	84	22	-
Male	18	28	38	50	63	107	120	118	67	18	4	-
Female	6	18	24	34	63	88	147	202	153	66	18	-
Black, both sexes	10	16	22	15	23	13	16	21	24	17	3	-
Male	7	13	14	11	14	5	5	6	6	6	-	-
Female	3	3	8	4	9	8	11	15	18	11	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
California												
All races, both sexes	389	477	426	517	800	1,204	2,114	2,578	2,104	909	194	1
Male	271	313	272	314	458	615	1,013	1,045	706	215	34	-
Female	118	164	154	203	342	589	1,101	1,533	1,398	694	160	1
White, both sexes	280	369	337	394	665	1,026	1,875	2,341	1,929	840	173	1
Male	195	249	220	249	385	539	914	959	643	190	28	-
Female	85	120	117	145	280	487	961	1,382	1,286	650	145	1
Black, both sexes	83	72	59	75	83	90	106	96	67	27	8	-
Male	55	39	33	35	45	38	44	33	23	10	2	-
Female	28	33	26	40	38	52	62	63	44	17	6	-
American Indian 1/, both sexes	3	2	4	5	6	12	10	8	2	2	1	-
Male	3	2	3	4	3	4	7	3	1	-	-	-
Female	-	-	1	1	3	8	3	5	1	2	1	-
Asian or Pacific Islander 2/, both sexes	23	34	26	43	46	76	123	133	106	40	12	-
Male	18	23	16	26	25	34	48	50	39	15	4	-
Female	5	11	10	17	21	42	75	83	67	25	8	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Colorado												
All races, both sexes	36	41	58	72	108	165	271	350	283	135	26	-
Male	21	30	31	44	50	87	118	133	77	29	5	-
Female	15	11	27	28	58	78	153	217	206	106	21	-
White, both sexes	34	37	53	68	99	157	263	339	277	132	26	-
Male	20	26	26	42	46	84	115	130	76	29	5	-
Female	14	11	27	26	53	73	148	209	201	103	21	-
Black, both sexes	1	3	2	2	5	1	4	10	2	1	-	-
Male	-	3	2	1	1	1	1	3	1	-	-	-
Female	1	-	-	1	4	-	3	7	1	1	-	-
American Indian 1/, both sexes	1	-	2	1	1	2	-	-	1	2	-	-
Male	1	-	2	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	3	5	4	1	3	-	-	-
Male	-	1	1	-	2	1	2	-	-	-	-	-
Female	-	-	-	1	1	4	2	1	3	-	-	-
Connecticut												
All races, both sexes	40	74	78	109	142	255	418	509	407	198	56	-
Male	36	45	53	61	82	138	186	206	141	32	6	-
Female	4	29	25	48	60	117	232	303	266	166	50	-
White, both sexes	34	67	72	93	129	243	400	496	399	192	53	-
Male	30	40	48	53	76	133	179	201	141	31	4	-
Female	4	27	24	40	53	110	221	295	258	161	49	-
Black, both sexes	5	6	6	15	11	11	18	12	7	5	2	-
Male	5	4	5	7	6	5	7	4	-	1	2	-
Female	-	2	1	8	5	6	11	8	7	4	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	1	-	1	1	1	1	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	2	1	-	-	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Delaware												
All races, both sexes	17	24	31	20	33	44	62	66	58	28	7	-
Male	10	13	20	10	19	20	27	21	26	5	2	-
Female	7	11	11	10	14	24	35	45	32	23	5	-
White, both sexes	11	19	24	15	25	39	57	59	54	26	7	-
Male	7	11	16	7	16	18	25	19	25	5	2	-
Female	4	8	8	8	9	21	32	40	29	21	5	-
Black, both sexes	6	5	7	5	8	5	5	6	4	1	-	-
Male	3	2	4	3	3	2	2	2	1	-	-	-
Female	3	3	3	2	5	3	3	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	14	3	4	14	16	14	35	23	24	12	3	-
Male	8	3	3	7	9	7	12	5	7	5	1	-
Female	6	-	1	7	7	7	23	18	17	7	2	-
White, both sexes	1	2	1	-	-	7	8	8	9	5	1	-
Male	1	2	-	-	-	5	3	3	3	3	-	-
Female	-	-	1	-	-	2	5	5	6	2	1	-
Black, both sexes	13	1	3	14	16	7	27	15	14	7	2	-
Male	7	1	3	7	9	2	9	2	4	2	1	-
Female	6	-	-	7	7	5	18	13	10	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Florida												
All races, both sexes	203	263	282	388	556	945	1,390	1,654	1,341	580	102	1
Male	134	173	185	237	324	499	695	695	492	168	23	1
Female	69	90	97	151	232	446	695	959	849	412	79	-
White, both sexes	129	204	228	327	485	849	1,299	1,566	1,275	559	97	-
Male	89	140	148	198	282	454	652	670	467	161	23	-
Female	40	64	80	129	203	395	647	896	808	398	74	-
Black, both sexes	73	54	53	59	68	92	90	78	64	20	5	1
Male	44	31	36	37	40	45	42	21	25	6	-	1
Female	29	23	17	22	28	47	48	57	39	14	5	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	-	1	3	4	1	9	2	1	-	-
Male	1	2	-	1	2	-	1	4	-	1	-	-
Female	-	3	-	-	1	4	-	5	2	-	-	-
Georgia												
All races, both sexes	283	370	409	434	523	726	991	1,080	759	315	63	-
Male	168	260	252	248	294	347	443	376	224	72	8	-
Female	115	110	157	186	229	379	548	704	535	243	55	-
White, both sexes	156	214	275	306	370	554	789	875	637	260	44	-
Male	99	162	171	178	208	270	367	309	184	57	7	-
Female	57	52	104	128	162	284	422	566	453	203	37	-
Black, both sexes	124	154	132	126	147	164	195	196	119	54	19	-
Male	68	96	81	69	81	74	75	62	40	15	1	-
Female	56	58	51	57	66	90	120	134	79	39	18	-
American Indian 1/, both sexes	1	1	-	-	2	1	-	1	-	-	-	-
Male	1	1	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	2	4	7	7	8	3	1	-	-
Male	-	1	-	1	4	2	1	4	-	-	-	-
Female	2	-	2	1	-	5	6	4	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Hawaii												
All races, both sexes	32	31	37	49	69	92	121	138	118	52	14	-
Male	24	24	24	31	40	53	66	69	52	19	4	-
Female	8	7	13	18	29	39	55	69	66	33	10	-
White, both sexes	11	10	4	12	12	27	27	36	27	11	2	-
Male	9	8	4	7	9	11	17	17	11	1	1	-
Female	2	2	-	5	3	16	10	19	16	10	1	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	1	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	21	32	37	56	65	93	102	91	41	11	-
Male	15	16	19	24	30	42	49	52	41	18	3	-
Female	6	5	13	13	26	23	44	50	50	23	8	-
Idaho												
All races, both sexes	12	11	24	23	37	63	96	148	123	41	15	-
Male	5	9	16	19	18	32	43	58	38	9	3	-
Female	7	2	8	4	19	31	53	90	85	32	12	-
White, both sexes	12	10	24	23	36	62	95	147	123	40	15	-
Male	5	8	16	19	17	31	43	58	38	9	3	-
Female	7	2	8	4	19	31	52	89	85	31	12	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	1	1	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Illinois												
All races, both sexes	181	236	268	345	471	787	1,203	1,327	1,210	503	128	-
Male	125	150	167	202	246	369	531	500	324	112	23	-
Female	56	86	101	143	225	418	672	827	886	391	105	-
White, both sexes	123	167	216	279	357	681	1,057	1,247	1,158	478	120	-
Male	86	111	137	167	190	318	471	474	316	109	22	-
Female	37	56	79	112	167	363	586	773	842	369	98	-
Black, both sexes	58	66	49	60	105	98	137	71	48	24	7	-
Male	39	39	29	31	50	47	55	21	7	3	1	-
Female	19	27	20	29	55	51	82	50	41	21	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	3	5	9	8	9	8	4	1	1	-
Male	-	-	1	4	6	4	5	4	1	-	-	-
Female	-	3	2	1	3	4	4	4	3	1	1	-
Indiana												
All races, both sexes	110	159	173	260	308	448	690	772	703	267	70	-
Male	69	101	123	163	168	237	293	264	187	52	8	-
Female	41	58	50	97	140	211	397	508	516	215	62	-
White, both sexes	89	133	150	230	281	416	650	750	677	257	67	-
Male	57	82	105	144	153	225	279	259	180	48	8	-
Female	32	51	45	86	128	191	371	491	497	209	59	-
Black, both sexes	20	26	22	30	27	31	40	21	26	10	3	-
Male	12	19	17	19	15	12	14	5	7	4	-	-
Female	8	7	5	11	12	19	26	16	19	6	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Iowa												
All races, both sexes	24	42	42	43	70	123	171	219	214	120	34	-
Male	15	29	26	32	32	63	81	85	65	27	2	-
Female	9	13	16	11	38	60	90	134	149	93	32	-
White, both sexes	21	40	42	42	67	121	166	216	214	120	33	-
Male	13	28	26	31	30	63	80	85	65	27	2	-
Female	8	12	16	11	37	58	86	131	149	93	31	-
Black, both sexes	3	2	-	1	3	1	5	3	-	-	1	-
Male	2	1	-	1	2	-	1	-	-	-	-	-
Female	1	1	-	-	1	1	4	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Kansas												
All races, both sexes	34	52	60	60	120	170	298	382	403	214	47	-
Male	23	41	38	36	56	74	124	153	115	53	7	-
Female	11	11	22	24	64	96	174	229	288	161	40	-
White, both sexes	26	49	51	51	110	157	288	365	394	207	45	-
Male	16	38	35	32	53	69	120	146	114	51	7	-
Female	10	11	16	19	57	88	168	219	280	156	38	-
Black, both sexes	7	3	8	7	8	12	8	15	7	7	2	-
Male	6	3	2	3	3	4	3	6	-	2	-	-
Female	1	-	6	4	5	8	5	9	7	5	2	-
American Indian 1/, both sexes	1	-	-	2	1	1	1	2	1	-	-	-
Male	1	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Kentucky												
All races, both sexes	86	131	145	226	236	382	522	578	424	182	39	-
Male	65	86	81	133	118	178	209	207	101	35	5	-
Female	21	45	64	93	118	204	313	371	323	147	34	-
White, both sexes	76	114	128	212	226	358	498	550	413	172	37	-
Male	58	76	74	124	114	167	199	201	99	35	4	-
Female	18	38	54	88	112	191	299	349	314	137	33	-
Black, both sexes	9	16	15	14	9	23	23	26	11	10	2	-
Male	6	10	5	9	4	11	9	6	2	-	1	-
Female	3	6	10	5	5	12	14	20	9	10	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	1	1	1	2	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	1	-	2	-	-	-	-
Louisiana												
All races, both sexes	100	134	168	186	259	392	495	509	397	169	23	-
Male	62	79	98	101	129	203	215	183	133	33	6	-
Female	38	55	70	85	130	189	280	326	264	136	17	-
White, both sexes	52	76	103	120	175	289	409	422	337	132	19	-
Male	33	43	60	71	92	157	176	155	116	24	4	-
Female	19	33	43	49	83	132	233	267	221	108	15	-
Black, both sexes	48	57	65	65	81	103	85	87	60	36	4	-
Male	29	35	38	29	36	46	39	28	17	9	2	-
Female	19	22	27	36	45	57	46	59	43	27	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	3	-	1	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Maine												
All races, both sexes	20	33	32	39	61	93	145	182	124	58	16	-
Male	16	25	23	25	39	61	66	81	29	10	3	-
Female	4	8	9	14	22	32	79	101	95	48	13	-
White, both sexes	18	31	32	39	61	93	144	181	124	58	16	-
Male	14	23	23	25	39	61	66	81	29	10	3	-
Female	4	8	9	14	22	32	78	100	95	48	13	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	58	94	96	125	166	249	373	443	358	135	25	1
Male	29	58	59	64	96	117	168	172	104	28	4	1
Female	29	36	37	61	70	132	205	271	254	107	21	-
White, both sexes	27	53	58	72	107	184	284	382	320	124	20	1
Male	10	32	35	39	66	89	136	150	91	27	3	1
Female	17	21	23	33	41	95	148	232	229	97	17	-
Black, both sexes	30	39	37	46	57	63	77	58	37	9	5	-
Male	18	25	23	21	29	27	28	22	12	-	1	-
Female	12	14	14	25	28	36	49	36	25	9	4	-
American Indian 1/, both sexes	-	1	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	6	2	2	11	3	1	2	-	-
Male	1	1	-	4	1	1	4	-	1	1	-	-
Female	-	-	-	2	1	1	7	3	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Massachusetts												
All races, both sexes	62	93	115	137	218	415	715	901	787	338	88	-
Male	41	72	64	83	118	224	321	328	219	65	11	-
Female	21	21	51	54	100	191	394	573	568	273	77	-
White, both sexes	53	89	108	129	208	389	698	879	771	333	85	-
Male	35	68	61	80	115	211	315	318	217	65	11	-
Female	18	21	47	49	93	178	383	561	554	268	74	-
Black, both sexes	7	2	7	7	7	19	12	21	15	4	2	-
Male	5	2	3	3	2	12	6	9	2	-	-	-
Female	2	-	4	4	5	7	6	12	13	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	1	3	6	4	1	1	1	1	-
Male	1	2	-	-	1	1	-	1	-	-	-	-
Female	1	-	-	1	2	5	4	-	1	1	1	-
Michigan												
All races, both sexes	129	170	194	230	360	597	915	1,167	954	394	88	-
Male	85	113	121	124	187	309	413	435	267	93	13	-
Female	44	57	73	106	173	288	502	732	687	301	75	-
White, both sexes	76	121	152	193	297	522	838	1,106	911	373	79	-
Male	49	83	98	108	161	277	373	414	255	88	11	-
Female	27	38	54	85	136	245	465	692	656	285	68	-
Black, both sexes	50	45	41	33	58	71	72	58	39	21	7	-
Male	34	27	22	13	24	28	37	20	9	5	1	-
Female	16	18	19	20	34	43	35	38	30	16	6	-
American Indian 1/, both sexes	2	3	1	3	2	2	3	1	1	-	-	-
Male	1	2	1	3	-	2	1	1	-	-	-	-
Female	1	1	-	-	2	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	3	2	2	2	3	-	2	-
Male	1	1	-	-	2	2	2	-	3	-	1	-
Female	-	-	-	1	1	-	-	2	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Minnesota												
All races, both sexes	41	56	79	76	140	275	410	588	534	246	62	-
Male	30	38	54	47	80	149	200	244	162	59	14	-
Female	11	18	25	29	60	126	210	344	372	187	48	-
White, both sexes	34	50	76	72	131	269	403	580	528	246	61	-
Male	25	33	52	45	75	145	198	239	161	59	14	-
Female	9	17	24	27	56	124	205	341	367	187	47	-
Black, both sexes	5	5	1	2	6	2	1	4	3	-	1	-
Male	4	4	1	1	3	2	-	2	-	-	-	-
Female	1	1	-	1	3	-	1	2	3	-	1	-
American Indian 1/, both sexes	1	1	1	1	2	2	2	-	-	-	-	-
Male	-	1	-	-	2	1	1	-	-	-	-	-
Female	1	-	1	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	2	4	4	3	-	-	-
Male	1	-	1	1	-	1	1	3	1	-	-	-
Female	-	-	-	-	1	1	3	1	2	-	-	-
Mississippi												
All races, both sexes	92	123	119	190	218	349	437	494	398	179	46	-
Male	58	78	73	117	119	165	174	166	119	35	11	-
Female	34	45	46	73	99	184	263	328	279	144	35	-
White, both sexes	43	59	71	126	152	247	329	388	300	126	24	-
Male	28	38	46	77	87	120	140	131	87	24	5	-
Female	15	21	25	49	65	127	189	257	213	102	19	-
Black, both sexes	49	61	48	64	66	100	107	104	98	53	22	-
Male	30	37	27	40	32	43	33	35	32	11	6	-
Female	19	24	21	24	34	57	74	69	66	42	16	-
American Indian 1/, both sexes	-	3	-	-	-	-	-	-	-	-	-	-
Male	-	3	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	2	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Missouri												
All races, both sexes	110	129	135	173	249	393	618	714	590	257	70	-
Male	64	76	72	98	148	182	275	242	171	51	13	-
Female	46	53	63	75	101	211	343	472	419	206	57	-
White, both sexes	86	104	121	145	223	349	576	687	571	248	65	-
Male	51	62	65	85	131	162	262	236	164	49	13	-
Female	35	42	56	60	92	187	314	451	407	199	52	-
Black, both sexes	24	25	12	28	25	42	40	27	19	9	5	-
Male	13	14	5	13	16	20	13	6	7	2	-	-
Female	11	11	7	15	9	22	27	21	12	7	5	-
American Indian 1/, both sexes	-	-	1	-	-	1	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	16	12	26	16	35	64	113	147	147	59	7	-
Male	14	7	18	12	19	36	62	67	47	16	-	-
Female	2	5	8	4	16	28	51	80	100	43	7	-
White, both sexes	14	11	22	15	31	63	110	145	145	59	6	-
Male	12	6	17	11	17	35	60	66	47	16	-	-
Female	2	5	5	4	14	28	50	79	98	43	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	4	1	4	1	3	2	2	-	1	-
Male	2	1	1	1	2	1	2	1	-	-	-	-
Female	-	-	3	-	2	-	1	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Nebraska												
All races, both sexes	28	33	43	55	96	147	218	251	254	137	39	-
Male	22	23	25	33	52	79	108	78	77	36	7	-
Female	6	10	18	22	44	68	110	173	177	101	32	-
White, both sexes	27	29	36	49	95	139	212	246	252	137	37	-
Male	21	21	20	29	51	76	105	76	77	36	7	-
Female	6	8	16	20	44	63	107	170	175	101	30	-
Black, both sexes	1	3	6	4	-	7	6	4	2	-	2	-
Male	1	2	4	3	-	3	3	2	-	-	-	-
Female	-	1	2	1	-	4	3	2	2	-	2	-
American Indian 1/, both sexes	-	1	1	1	1	1	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	66	100	150	155	188	194	300	258	148	46	11	-
Male	42	69	98	96	104	122	148	109	48	9	2	-
Female	24	31	52	59	84	72	152	149	100	37	9	-
White, both sexes	49	81	128	133	160	170	272	235	140	41	11	-
Male	30	57	88	87	91	106	139	96	46	8	2	-
Female	19	24	40	46	69	64	133	139	94	33	9	-
Black, both sexes	13	16	18	11	21	14	17	14	5	2	-	-
Male	8	10	9	5	9	8	6	8	1	-	-	-
Female	5	6	9	6	12	6	11	6	4	2	-	-
American Indian 1/, both sexes	-	1	2	2	2	-	2	4	2	1	-	-
Male	-	1	-	1	2	-	-	2	1	-	-	-
Female	-	-	2	1	-	-	2	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	4	2	2	9	5	10	9	5	1	2	-	-
Male	4	1	1	3	2	8	3	3	-	1	-	-
Female	-	1	1	6	3	2	6	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
New Hampshire												
All races, both sexes	12	17	21	15	40	73	117	151	138	46	14	-
Male	9	14	14	11	24	41	45	61	43	8	-	-
Female	3	3	7	4	16	32	72	90	95	38	14	-
White, both sexes	12	17	21	15	40	73	117	151	138	46	14	-
Male	9	14	14	11	24	41	45	61	43	8	-	-
Female	3	3	7	4	16	32	72	90	95	38	14	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	92	142	154	196	312	531	771	962	767	324	59	-
Male	64	92	98	116	178	275	343	375	225	73	7	-
Female	28	50	56	80	134	256	428	587	542	251	52	-
White, both sexes	61	104	115	154	261	478	696	899	726	308	56	-
Male	45	71	72	92	146	251	313	353	211	68	6	-
Female	16	33	43	62	115	227	383	546	515	240	50	-
Black, both sexes	29	35	35	34	46	46	69	54	31	15	2	-
Male	18	19	22	20	29	20	28	20	11	5	1	-
Female	11	16	13	14	17	26	41	34	20	10	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	4	8	5	7	5	8	10	1	1	-
Male	1	2	4	4	3	4	2	2	3	-	-	-
Female	1	1	-	4	2	3	3	6	7	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
New Mexico												
All races, both sexes	14	30	34	42	45	100	147	178	141	73	13	-
Male	7	21	20	21	27	51	67	67	44	17	2	-
Female	7	9	14	21	18	49	80	111	97	56	11	-
White, both sexes	11	27	32	38	40	95	144	173	137	72	11	-
Male	6	19	19	20	24	49	65	66	42	16	1	-
Female	5	8	13	18	16	46	79	107	95	56	10	-
Black, both sexes	1	2	-	-	3	3	1	2	1	1	1	-
Male	-	1	-	-	1	2	1	-	1	1	-	-
Female	1	1	-	-	2	1	-	2	-	-	1	-
American Indian 1/, both sexes	-	-	2	3	2	2	1	2	2	-	1	-
Male	-	-	1	1	2	-	1	-	-	-	1	-
Female	-	-	1	2	-	2	-	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	-	-	1	1	1	-	-	-
Male	1	1	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	1	-	-	1	-	-	-	-	-
New York												
All races, both sexes	153	224	253	290	461	875	1,266	1,570	1,289	549	119	-
Male	104	139	146	186	247	475	587	623	377	128	14	-
Female	49	85	107	104	214	400	679	947	912	421	105	-
White, both sexes	108	169	210	234	409	800	1,175	1,493	1,240	525	111	-
Male	75	112	125	149	223	437	545	599	362	124	14	-
Female	33	57	85	85	186	363	630	894	878	401	97	-
Black, both sexes	42	49	37	46	44	65	77	69	37	20	7	-
Male	26	23	18	30	20	34	35	19	10	3	-	-
Female	16	26	19	16	24	31	42	50	27	17	7	-
American Indian 1/, both sexes	1	1	2	2	1	-	2	-	1	-	-	-
Male	1	1	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	1	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	4	8	7	10	12	8	11	4	1	-
Male	2	3	1	7	4	4	7	5	5	1	-	-
Female	-	2	3	1	3	6	5	3	6	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
North Carolina												
All races, both sexes	145	194	250	282	394	576	829	876	673	233	57	-
Male	100	117	146	178	222	288	377	303	191	43	7	-
Female	45	77	104	104	172	288	452	573	482	190	50	-
White, both sexes	82	117	155	192	303	482	679	753	597	197	43	-
Male	58	72	90	116	165	242	321	263	178	35	4	-
Female	24	45	65	76	138	240	358	490	419	162	39	-
Black, both sexes	61	75	89	81	84	88	143	120	73	35	14	-
Male	40	44	53	54	52	42	50	40	11	8	3	-
Female	21	31	36	27	32	46	93	80	62	27	11	-
American Indian 1/, both sexes	2	1	5	6	1	4	4	1	3	1	-	-
Male	2	1	3	5	-	2	3	-	2	-	-	-
Female	-	-	2	1	1	2	1	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	6	2	3	2	-	-	-	-
Male	-	-	-	3	5	2	3	-	-	-	-	-
Female	-	1	1	-	1	-	-	2	-	-	-	-
North Dakota												
All races, both sexes	7	6	8	12	24	27	44	71	57	40	5	-
Male	4	3	5	6	17	12	23	29	9	6	-	-
Female	3	3	3	6	7	15	21	42	48	34	5	-
White, both sexes	5	5	8	10	22	25	42	70	57	40	5	-
Male	3	2	5	6	16	11	23	29	9	6	-	-
Female	2	3	3	4	6	14	19	41	48	34	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	2	2	2	2	1	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	2	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Ohio												
All races, both sexes	170	212	209	281	424	666	962	1,175	867	334	57	-
Male	102	126	120	150	223	318	395	454	238	66	8	-
Female	68	86	89	131	201	348	567	721	629	268	49	-
White, both sexes	122	169	170	240	366	604	902	1,110	824	316	48	-
Male	71	101	101	129	185	289	373	425	226	62	8	-
Female	51	68	69	111	181	315	529	685	598	254	40	-
Black, both sexes	47	42	37	39	57	60	59	64	42	16	9	-
Male	31	24	18	20	38	28	21	28	11	4	-	-
Female	16	18	19	19	19	32	38	36	31	12	9	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	-	1	2	1	1	1	2	-	-
Male	-	1	1	-	-	1	1	1	1	-	-	-
Female	1	-	1	-	1	1	-	-	-	2	-	-
Oklahoma												
All races, both sexes	60	75	99	116	197	273	438	444	359	144	42	-
Male	42	48	55	65	106	142	180	157	106	28	6	-
Female	18	27	44	51	91	131	258	287	253	116	36	-
White, both sexes	46	58	86	93	165	252	405	424	330	137	40	-
Male	35	36	47	51	93	129	167	150	99	25	6	-
Female	11	22	39	42	72	123	238	274	231	112	34	-
Black, both sexes	11	10	5	12	21	14	17	8	12	6	1	-
Male	7	6	5	5	8	7	6	2	3	3	-	-
Female	4	4	-	7	13	7	11	6	9	3	1	-
American Indian 1/, both sexes	3	7	7	11	11	7	16	12	14	1	1	-
Male	-	6	3	9	5	6	7	5	4	-	-	-
Female	3	1	4	2	6	1	9	7	10	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	3	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Oregon												
All races, both sexes	53	47	56	68	106	210	345	439	416	176	38	-
Male	28	32	41	41	65	98	143	194	135	48	7	-
Female	25	15	15	27	41	112	202	245	281	128	31	-
White, both sexes	50	42	51	63	104	203	334	428	411	176	37	-
Male	26	29	38	40	65	97	139	189	133	48	6	-
Female	24	13	13	23	39	106	195	239	278	128	31	-
Black, both sexes	1	2	3	-	1	-	5	3	1	-	-	-
Male	1	2	3	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	3	3	1	-	-	-
American Indian 1/, both sexes	2	1	2	4	-	2	2	1	2	-	-	-
Male	1	-	-	1	-	1	1	1	2	-	-	-
Female	1	1	2	3	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	1	5	4	7	2	-	1	-
Male	-	1	-	-	-	-	1	4	-	-	1	-
Female	-	1	-	1	1	5	3	3	2	-	-	-
Pennsylvania												
All races, both sexes	216	299	388	474	683	1,167	1,779	2,088	1,662	634	139	1
Male	140	194	263	289	364	550	794	764	480	120	22	1
Female	76	105	125	185	319	617	985	1,324	1,182	514	117	-
White, both sexes	170	242	327	403	609	1,061	1,668	2,012	1,603	614	129	-
Male	113	157	220	247	323	507	752	733	457	116	21	-
Female	57	85	107	156	286	554	916	1,279	1,146	498	108	-
Black, both sexes	43	55	59	67	74	101	108	70	57	20	10	1
Male	25	36	42	40	41	42	40	28	22	4	1	1
Female	18	19	17	27	33	59	68	42	35	16	9	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	2	3	-	5	2	6	2	-	-	-
Male	2	1	1	1	-	1	1	3	1	-	-	-
Female	1	1	1	2	-	4	1	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Rhode Island												
All races, both sexes	5	5	11	22	30	46	66	81	85	36	8	-
Male	5	3	7	15	19	20	27	23	20	10	1	-
Female	-	2	4	7	11	26	39	58	65	26	7	-
White, both sexes	5	5	11	21	28	44	65	79	85	35	8	-
Male	5	3	7	14	18	18	27	22	20	10	1	-
Female	-	2	4	7	10	26	38	57	65	25	7	-
Black, both sexes	-	-	-	-	2	1	1	2	-	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	127	157	199	216	249	322	450	471	319	112	21	1
Male	87	111	127	124	135	172	217	157	94	27	3	1
Female	40	46	72	92	114	150	233	314	225	85	18	-
White, both sexes	64	86	114	136	174	243	339	388	246	91	13	-
Male	43	60	75	83	94	129	164	142	71	23	1	-
Female	21	26	39	53	80	114	175	246	175	68	12	-
Black, both sexes	62	70	85	79	74	77	107	82	72	21	8	1
Male	44	50	52	41	41	42	52	15	23	4	2	1
Female	18	20	33	38	33	35	55	67	49	17	6	-
American Indian 1/, both sexes	-	1	-	1	-	2	3	-	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
South Dakota												
All races, both sexes	5	5	4	12	11	19	38	51	59	21	12	-
Male	3	5	3	7	5	10	20	19	20	4	3	-
Female	2	-	1	5	6	9	18	32	39	17	9	-
White, both sexes	5	4	4	11	10	19	37	51	59	21	12	-
Male	3	4	3	6	5	10	19	19	20	4	3	-
Female	2	-	1	5	5	9	18	32	39	17	9	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	107	131	169	173	230	283	475	430	317	130	20	1
Male	71	85	101	104	124	130	179	170	96	32	2	1
Female	36	46	68	69	106	153	296	260	221	98	18	-
White, both sexes	79	100	123	124	190	239	420	393	285	116	14	1
Male	53	68	72	73	100	114	159	150	84	30	1	1
Female	26	32	51	51	90	125	261	243	201	86	13	-
Black, both sexes	28	31	46	48	38	44	54	36	29	14	6	-
Male	18	17	29	31	23	16	19	19	12	2	1	-
Female	10	14	17	17	15	28	35	17	17	12	5	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	1	3	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Texas												
All races, both sexes	308	388	445	518	713	1,146	1,579	1,806	1,366	653	124	-
Male	203	239	269	297	395	548	685	704	398	136	21	-
Female	105	149	176	221	318	598	894	1,102	968	517	103	-
White, both sexes	226	293	359	435	604	1,006	1,421	1,668	1,255	598	105	-
Male	148	188	215	247	330	488	621	651	370	120	16	-
Female	78	105	144	188	274	518	800	1,017	885	478	89	-
Black, both sexes	75	86	78	76	102	125	149	125	102	51	19	-
Male	49	46	49	45	61	55	63	48	24	14	5	-
Female	26	40	29	31	41	70	86	77	78	37	14	-
American Indian 1/, both sexes	2	3	2	1	1	1	-	2	1	1	-	-
Male	2	1	1	1	1	-	-	2	1	-	-	-
Female	-	2	1	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	6	6	6	6	14	9	11	8	3	-	-
Male	4	4	4	4	3	5	1	3	3	2	-	-
Female	1	2	2	2	3	9	8	8	5	1	-	-
Utah												
All races, both sexes	25	30	31	49	95	117	228	284	206	79	9	1
Male	18	15	24	27	49	54	117	99	58	22	1	1
Female	7	15	7	22	46	63	111	185	148	57	8	-
White, both sexes	23	29	31	49	88	113	226	277	205	79	7	1
Male	16	14	24	27	44	51	115	97	57	22	1	1
Female	7	15	7	22	44	62	111	180	148	57	6	-
Black, both sexes	1	-	-	-	2	1	2	-	-	-	1	-
Male	1	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	5	3	-	6	1	-	-	-
Male	1	1	-	-	4	2	-	2	1	-	-	-
Female	-	-	-	-	1	1	-	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Vermont												
All races, both sexes	5	5	9	9	18	26	45	58	50	27	5	-
Male	2	3	4	8	11	10	22	21	16	6	1	-
Female	3	2	5	1	7	16	23	37	34	21	4	-
White, both sexes	5	4	9	9	18	26	45	58	50	27	5	-
Male	2	2	4	8	11	10	22	21	16	6	1	-
Female	3	2	5	1	7	16	23	37	34	21	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	153	191	198	252	383	533	730	850	682	279	67	-
Male	108	128	116	151	213	293	333	290	192	50	9	-
Female	45	63	82	101	170	240	397	560	490	229	58	-
White, both sexes	101	113	135	166	283	430	609	751	602	247	51	-
Male	72	76	84	98	162	236	277	253	176	45	6	-
Female	29	37	51	68	121	194	332	498	426	202	45	-
Black, both sexes	51	75	58	80	95	97	115	91	77	30	15	-
Male	35	49	28	48	49	54	50	34	15	5	3	-
Female	16	26	30	32	46	43	65	57	62	25	12	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	3	5	5	5	6	6	7	3	1	1	-
Male	1	3	4	4	2	3	6	3	1	-	-	-
Female	-	-	1	1	3	3	-	4	2	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Washington												
All races, both sexes	67	72	89	90	116	215	353	444	391	145	42	-
Male	41	41	49	55	66	105	152	182	138	46	5	-
Female	26	31	40	35	50	110	201	262	253	99	37	-
White, both sexes	60	58	80	83	104	200	342	427	385	143	42	-
Male	39	33	43	51	59	99	148	174	137	45	5	-
Female	21	25	37	32	45	101	194	253	248	98	37	-
Black, both sexes	2	8	6	4	4	4	6	2	-	1	-	-
Male	-	5	5	4	2	1	2	1	-	1	-	-
Female	2	3	1	-	2	3	4	1	-	-	-	-
American Indian 1/, both sexes	3	3	2	1	3	1	2	4	2	-	-	-
Male	2	1	1	-	1	-	1	3	1	-	-	-
Female	1	2	1	1	2	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	2	5	10	3	11	4	1	-	-
Male	-	2	-	-	4	5	1	4	-	-	-	-
Female	2	1	1	2	1	5	2	7	4	1	-	-
West Virginia												
All races, both sexes	32	50	61	74	129	195	279	280	228	95	22	-
Male	21	33	34	49	63	84	127	99	62	15	2	-
Female	11	17	27	25	66	111	152	181	166	80	20	-
White, both sexes	27	47	59	73	123	188	271	274	222	88	21	-
Male	18	30	32	49	60	82	124	98	62	12	2	-
Female	9	17	27	24	63	106	147	176	160	76	19	-
Black, both sexes	5	3	2	1	6	7	8	6	6	6	1	-
Male	3	3	2	-	3	2	3	1	-	3	-	-
Female	2	-	-	1	3	5	5	5	6	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Wisconsin												
All races, both sexes	71	95	99	132	177	370	555	744	692	313	72	-
Male	46	63	61	84	99	195	246	325	208	76	10	-
Female	25	32	38	48	78	175	309	419	484	237	62	-
White, both sexes	63	76	91	123	163	357	535	733	682	311	72	-
Male	41	52	58	81	94	192	237	320	206	75	10	-
Female	22	24	33	42	69	165	298	413	476	236	62	-
Black, both sexes	8	14	6	9	10	9	14	6	9	2	-	-
Male	5	6	3	3	4	3	7	2	2	1	-	-
Female	3	8	3	6	6	6	7	4	7	1	-	-
American Indian 1/, both sexes	-	1	1	-	2	1	4	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	2	1	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	1	-	2	3	2	4	1	-	-	-
Male	-	4	-	-	1	-	1	2	-	-	-	-
Female	-	-	1	-	1	3	1	2	1	-	-	-
Wyoming												
All races, both sexes	6	17	10	19	21	33	62	65	63	20	6	-
Male	3	9	5	10	9	20	27	29	21	2	3	-
Female	3	8	5	9	12	13	35	36	42	18	3	-
White, both sexes	6	15	10	19	20	33	62	65	63	20	6	-
Male	3	8	5	10	8	20	27	29	21	2	3	-
Female	3	7	5	9	12	13	35	36	42	18	3	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
United States												
All races, both sexes	1,216	-	1	2	-	4	6	12	21	34	51	101
Male	660	-	-	1	-	1	3	4	14	22	29	71
Female	556	-	1	1	-	3	3	8	7	12	22	30
White, both sexes	958	-	1	2	-	3	6	8	14	22	40	77
Male	518	-	-	1	-	1	3	3	9	13	23	54
Female	440	-	1	1	-	2	3	5	5	9	17	23
Black, both sexes	228	-	-	-	-	1	-	4	7	10	10	20
Male	122	-	-	-	-	-	-	1	5	7	5	13
Female	106	-	-	-	-	1	-	3	2	3	5	7
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	1	1
Male	6	-	-	-	-	-	-	-	-	1	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	1	-	3
Male	14	-	-	-	-	-	-	-	-	1	-	3
Female	7	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	17	-	-	-	-	-	-	1	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Alaska												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	26	-	-	1	-	-	-	-	-	1	4	3
Male	16	-	-	1	-	-	-	-	-	1	2	2
Female	10	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	24	-	-	1	-	-	-	-	-	-	3	3
Male	14	-	-	1	-	-	-	-	-	-	1	2
Female	10	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Arkansas												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	137	-	-	-	-	2	2	2	4	4	8	7
Male	86	-	-	-	-	1	1	2	2	4	4	7
Female	51	-	-	-	-	1	1	-	2	-	4	-
White, both sexes	110	-	-	-	-	1	2	2	3	2	8	5
Male	70	-	-	-	-	1	1	2	1	2	4	5
Female	40	-	-	-	-	-	1	-	2	-	4	-
Black, both sexes	14	-	-	-	-	1	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	1	1	-	-
Female	7	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Colorado												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Delaware												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Florida												
All races, both sexes	72	-	1	-	-	-	-	-	2	2	3	4
Male	37	-	-	-	-	-	-	-	1	1	3	3
Female	35	-	1	-	-	-	-	-	1	1	-	1
White, both sexes	58	-	1	-	-	-	-	-	-	2	2	2
Male	29	-	-	-	-	-	-	-	-	1	2	1
Female	29	-	1	-	-	-	-	-	-	1	-	1
Black, both sexes	14	-	-	-	-	-	-	-	2	-	1	2
Male	8	-	-	-	-	-	-	-	1	-	1	2
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	43	-	-	-	-	-	-	-	-	4	1	5
Male	26	-	-	-	-	-	-	-	-	4	1	4
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	1	1	4
Male	15	-	-	-	-	-	-	-	-	1	1	3
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	17	-	-	-	-	-	-	-	-	3	-	1
Male	11	-	-	-	-	-	-	-	-	3	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Hawaii												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Illinois												
All races, both sexes	48	-	-	-	-	-	1	1	1	-	-	5
Male	30	-	-	-	-	-	1	-	1	-	-	2
Female	18	-	-	-	-	-	-	1	-	-	-	3
White, both sexes	36	-	-	-	-	-	1	1	1	-	-	3
Male	22	-	-	-	-	-	1	-	1	-	-	1
Female	14	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	24	-	-	-	-	-	-	-	-	1	4	1
Male	16	-	-	-	-	-	-	-	-	-	3	1
Female	8	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	3	1
Male	14	-	-	-	-	-	-	-	-	-	2	1
Female	7	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Iowa												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	-	-	-	-	-	-	1	1	1	-	1
Male	5	-	-	-	-	-	-	1	1	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	1	1	-	-	1
Male	4	-	-	-	-	-	-	1	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Kentucky												
All races, both sexes	18	-	-	-	-	-	1	1	-	1	-	2
Male	9	-	-	-	-	-	1	-	-	-	-	1
Female	9	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	18	-	-	-	-	-	1	1	-	1	-	2
Male	9	-	-	-	-	-	1	-	-	-	-	1
Female	9	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	25	-	-	-	-	-	-	-	2	-	-	4
Male	13	-	-	-	-	-	-	-	1	-	-	2
Female	12	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	11	-	-	-	-	-	-	-	1	-	-	2
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Maine												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	18	-	-	-	-	-	-	-	1	-	1	2
Male	8	-	-	-	-	-	-	-	1	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Massachusetts												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	46	-	-	1	-	-	-	-	1	1	5	3
Male	24	-	-	-	-	-	-	-	1	-	1	3
Female	22	-	-	1	-	-	-	-	-	1	4	-
White, both sexes	28	-	-	1	-	-	-	-	-	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	14	-	-	1	-	-	-	-	-	-	1	-
Black, both sexes	17	-	-	-	-	-	-	-	1	1	3	1
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Minnesota												
All races, both sexes	22	-	-	-	-	-	-	1	1	2	-	1
Male	12	-	-	-	-	-	-	-	1	1	-	1
Female	10	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	22	-	-	-	-	-	-	1	1	2	-	1
Male	12	-	-	-	-	-	-	-	1	1	-	1
Female	10	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	19	-	-	-	-	-	-	-	2	-	4	1
Male	8	-	-	-	-	-	-	-	2	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	13	-	-	-	-	-	-	-	1	-	3	1
Male	6	-	-	-	-	-	-	-	1	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	6	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Missouri												
All races, both sexes	31	-	-	-	-	-	-	-	-	3	3	4
Male	16	-	-	-	-	-	-	-	-	1	2	2
Female	15	-	-	-	-	-	-	-	-	2	1	2
White, both sexes	26	-	-	-	-	-	-	-	-	2	3	4
Male	15	-	-	-	-	-	-	-	-	1	2	2
Female	11	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Nebraska												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	31	-	-	-	-	-	-	1	-	1	1	2
Male	16	-	-	-	-	-	-	-	-	1	1	2
Female	15	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	1	-	1	1	2
Male	15	-	-	-	-	-	-	-	-	1	1	2
Female	11	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
New Mexico												
All races, both sexes	9	-	-	-	-	-	-	-	1	1	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	8	-	-	-	-	-	-	-	1	1	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	89	-	-	-	-	1	-	-	-	-	1	6
Male	37	-	-	-	-	-	-	-	-	-	1	3
Female	52	-	-	-	-	1	-	-	-	-	-	3
White, both sexes	69	-	-	-	-	1	-	-	-	-	-	5
Male	27	-	-	-	-	-	-	-	-	-	-	3
Female	42	-	-	-	-	1	-	-	-	-	-	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
North Carolina												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	2	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Ohio												
All races, both sexes	48	-	-	-	-	-	-	1	-	4	-	3
Male	30	-	-	-	-	-	-	1	-	3	-	3
Female	18	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	2	-	2
Male	21	-	-	-	-	-	-	-	-	2	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	1	-	2	-	1
Male	9	-	-	-	-	-	-	1	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	20	-	-	-	-	-	-	-	1	1	-	4
Male	14	-	-	-	-	-	-	-	1	-	-	3
Female	6	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	18	-	-	-	-	-	-	-	1	1	-	3
Male	12	-	-	-	-	-	-	-	1	-	-	2
Female	6	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Oregon												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	64	-	-	-	-	-	-	-	1	2	1	2
Male	33	-	-	-	-	-	-	-	1	2	1	1
Female	31	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	57	-	-	-	-	-	-	-	1	2	1	1
Male	28	-	-	-	-	-	-	-	1	2	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Rhode Island												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	19	-	-	-	-	-	-	-	1	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	15	-	-	-	-	-	-	-	1	-	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	30	-	-	-	-	-	-	3	1	1	2	1
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	16	-	-	-	-	-	-	3	1	-	1	1
White, both sexes	21	-	-	-	-	-	-	-	1	1	1	1
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	9	-	-	-	-	-	-	3	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Texas												
All races, both sexes	79	-	-	-	-	1	1	-	1	2	6	7
Male	40	-	-	-	-	-	-	-	1	-	5	5
Female	39	-	-	-	-	1	1	-	-	2	1	2
White, both sexes	68	-	-	-	-	1	1	-	1	2	5	6
Male	36	-	-	-	-	-	-	-	1	-	4	5
Female	32	-	-	-	-	1	1	-	-	2	1	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	31	-	-	-	-	-	1	-	-	-	-	4
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	13	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	21	-	-	-	-	-	1	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	4
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Washington												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	5
Male	15	-	-	-	-	-	-	-	-	-	-	4
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	5
Male	14	-	-	-	-	-	-	-	-	-	-	4
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	2
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	2
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Wisconsin												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	3
Male	11	-	-	-	-	-	-	-	-	-	-	3
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	3
Male	9	-	-	-	-	-	-	-	-	-	-	3
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
United States												
All races, both sexes	102	123	105	100	119	165	154	74	36	3	3	-
Male	69	84	58	59	61	72	73	27	11	1	-	-
Female	33	39	47	41	58	93	81	47	25	2	3	-
White, both sexes	63	84	75	83	101	136	142	64	31	3	3	-
Male	42	58	43	47	55	63	65	26	11	1	-	-
Female	21	26	32	36	46	73	77	38	20	2	3	-
Black, both sexes	36	36	26	14	16	25	10	9	4	-	-	-
Male	24	23	14	11	4	8	7	-	-	-	-	-
Female	12	13	12	3	12	17	3	9	4	-	-	-
American Indian 1/, both sexes	2	-	1	1	1	1	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	3	2	1	3	2	1	1	-	-	-
Male	1	3	1	1	1	1	1	1	-	-	-	-
Female	-	-	2	1	-	2	1	-	1	-	-	-
Alabama												
All races, both sexes	2	3	1	1	3	2	1	2	1	-	-	-
Male	2	1	1	1	2	1	1	-	-	-	-	-
Female	-	2	-	-	1	1	-	2	1	-	-	-
White, both sexes	-	-	-	1	3	1	1	-	1	-	-	-
Male	-	-	-	1	2	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	2	3	1	-	-	1	-	2	-	-	-	-
Male	2	1	1	-	-	1	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Alaska												
All races, both sexes	1	-	-	-	1	2	1	-	-	-	-	-
Male	1	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	1	2	3	-	2	2	-	2	-	-	-
Male	4	1	1	2	-	1	1	-	-	-	-	-
Female	1	-	1	1	-	1	1	-	2	-	-	-
White, both sexes	5	1	2	3	-	2	2	-	2	-	-	-
Male	4	1	1	2	-	1	1	-	-	-	-	-
Female	1	-	1	1	-	1	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Arkansas												
All races, both sexes	1	3	2	1	2	3	1	1	-	-	-	-
Male	1	2	1	1	1	1	-	1	-	-	-	-
Female	-	1	1	-	1	2	1	-	-	-	-	-
White, both sexes	1	2	1	-	2	3	1	1	-	-	-	-
Male	1	1	-	-	1	1	-	1	-	-	-	-
Female	-	1	1	-	1	2	1	-	-	-	-	-
Black, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	11	16	8	10	9	19	21	8	5	1	-	-
Male	7	15	5	7	6	8	13	3	1	-	-	-
Female	4	1	3	3	3	11	8	5	4	1	-	-
White, both sexes	8	11	6	9	9	14	19	7	3	1	-	-
Male	5	10	4	7	6	7	12	2	1	-	-	-
Female	3	1	2	2	3	7	7	5	2	1	-	-
Black, both sexes	2	4	1	1	-	2	-	-	1	-	-	-
Male	1	4	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	-	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	-	2	2	1	1	-	-	-
Male	1	1	1	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Colorado												
All races, both sexes	-	1	1	1	-	1	1	2	-	-	-	-
Male	-	1	-	-	-	-	1	2	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
White, both sexes	-	1	1	1	-	1	1	2	-	-	-	-
Male	-	1	-	-	-	-	1	2	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	2	1	3	5	7	2	-	-	-	-	-
Male	1	-	1	1	2	3	-	-	-	-	-	-
Female	-	2	-	2	3	4	2	-	-	-	-	-
White, both sexes	-	1	1	2	4	5	2	-	-	-	-	-
Male	-	-	1	-	2	2	-	-	-	-	-	-
Female	-	1	-	2	2	3	2	-	-	-	-	-
Black, both sexes	1	1	-	1	1	2	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Delaware												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Florida												
All races, both sexes	5	8	4	4	10	12	10	4	2	-	1	-
Male	3	4	1	3	5	5	5	1	2	-	-	-
Female	2	4	3	1	5	7	5	3	-	-	1	-
White, both sexes	3	8	4	2	10	10	9	2	2	-	1	-
Male	3	4	1	1	5	4	4	1	2	-	-	-
Female	-	4	3	1	5	6	5	1	-	-	1	-
Black, both sexes	2	-	-	2	-	2	1	2	-	-	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	2	-	-	-	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	6	7	8	2	1	5	2	1	1	-	-	-
Male	4	6	5	1	-	-	-	-	1	-	-	-
Female	2	1	3	1	1	5	2	1	-	-	-	-
White, both sexes	3	2	7	1	1	2	2	1	1	-	-	-
Male	2	2	5	-	-	-	-	-	1	-	-	-
Female	1	-	2	1	1	2	2	1	-	-	-	-
Black, both sexes	3	5	1	1	-	3	-	-	-	-	-	-
Male	2	4	-	1	-	-	-	-	-	-	-	-
Female	1	1	1	-	-	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Hawaii												
All races, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	1	-	-	3	-	-	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	3	-	-	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Illinois												
All races, both sexes	3	3	4	9	3	7	5	3	3	-	-	-
Male	1	2	3	7	3	5	2	2	1	-	-	-
Female	2	1	1	2	-	2	3	1	2	-	-	-
White, both sexes	2	3	3	7	1	6	3	3	2	-	-	-
Male	-	2	3	5	1	4	1	2	1	-	-	-
Female	2	1	-	2	-	2	2	1	1	-	-	-
Black, both sexes	1	-	1	2	1	1	2	-	1	-	-	-
Male	1	-	-	2	1	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	4	2	4	-	4	2	2	-	-	-	-	-
Male	3	2	2	-	3	1	1	-	-	-	-	-
Female	1	-	2	-	1	1	1	-	-	-	-	-
White, both sexes	4	1	4	-	3	2	2	-	-	-	-	-
Male	3	1	2	-	3	1	1	-	-	-	-	-
Female	1	-	2	-	-	1	1	-	-	-	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Iowa												
All races, both sexes	-	2	1	1	1	2	1	1	-	-	-	-
Male	-	1	1	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	2	1	1	1	2	1	1	-	-	-	-
Male	-	1	1	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	1	-	1	1	2	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-
White, both sexes	1	-	1	-	1	1	2	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Kentucky												
All races, both sexes	2	-	2	1	2	2	2	1	1	-	-	-
Male	2	-	1	-	1	1	1	1	-	-	-	-
Female	-	-	1	1	1	1	1	-	1	-	-	-
White, both sexes	2	-	2	1	2	2	2	1	1	-	-	-
Male	2	-	1	-	1	1	1	1	-	-	-	-
Female	-	-	1	1	1	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	1	2	2	3	4	1	2	-	-	-	-
Male	4	1	1	2	1	-	-	1	-	-	-	-
Female	-	-	1	-	2	4	1	1	-	-	-	-
White, both sexes	-	1	1	1	1	1	1	2	-	-	-	-
Male	-	1	-	1	1	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
Black, both sexes	4	-	1	1	2	3	-	-	-	-	-	-
Male	4	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Maine												
All races, both sexes	-	1	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	1	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	2	1	2	1	1	2	3	-	-	-	-
Male	1	1	1	1	-	1	1	-	-	-	-	-
Female	1	1	-	1	1	-	1	3	-	-	-	-
White, both sexes	1	1	1	1	1	-	1	2	-	-	-	-
Male	-	1	1	1	-	-	1	-	-	-	-	-
Female	1	-	-	-	1	-	-	2	-	-	-	-
Black, both sexes	1	1	-	-	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Massachusetts												
All races, both sexes	-	2	2	4	-	2	4	1	2	-	-	-
Male	-	1	1	3	-	1	2	-	2	-	-	-
Female	-	1	1	1	-	1	2	1	-	-	-	-
White, both sexes	-	2	2	4	-	2	4	1	2	-	-	-
Male	-	1	1	3	-	1	2	-	2	-	-	-
Female	-	1	1	1	-	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5	2	5	4	7	8	3	1	-	-	-	-
Male	3	2	4	1	3	4	2	-	-	-	-	-
Female	2	-	1	3	4	4	1	1	-	-	-	-
White, both sexes	3	1	1	4	4	7	2	1	-	-	-	-
Male	2	1	1	1	2	3	1	-	-	-	-	-
Female	1	-	-	3	2	4	1	1	-	-	-	-
Black, both sexes	1	1	4	-	3	1	1	-	-	-	-	-
Male	-	1	3	-	1	1	1	-	-	-	-	-
Female	1	-	1	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Minnesota												
All races, both sexes	-	-	3	2	4	4	2	1	1	-	-	-
Male	-	-	3	1	2	-	2	1	-	-	-	-
Female	-	-	-	1	2	4	-	-	1	-	-	-
White, both sexes	-	-	3	2	4	4	2	1	1	-	-	-
Male	-	-	3	1	2	-	2	1	-	-	-	-
Female	-	-	-	1	2	4	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	2	3	-	1	2	3	-	-	-	-	-
Male	1	-	1	-	-	2	-	-	-	-	-	-
Female	-	2	2	-	1	-	3	-	-	-	-	-
White, both sexes	1	2	1	-	-	2	2	-	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	2	1	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	2	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Missouri												
All races, both sexes	4	2	1	-	3	5	6	-	-	-	-	-
Male	1	1	-	-	2	4	3	-	-	-	-	-
Female	3	1	1	-	1	1	3	-	-	-	-	-
White, both sexes	3	1	-	-	3	4	6	-	-	-	-	-
Male	1	1	-	-	2	3	3	-	-	-	-	-
Female	2	-	-	-	1	1	3	-	-	-	-	-
Black, both sexes	1	1	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Nebraska												
All races, both sexes	-	1	-	1	-	-	1	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	1	-	1	-	-	1	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	3	-	2	-	-	-	-	-	-	-
Male	-	-	2	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	3	-	2	-	-	-	-	-	-	-
Male	-	-	2	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
New Hampshire												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	3	2	4	4	6	3	2	-	-	-
Male	2	-	2	2	3	-	2	1	-	-	-	-
Female	-	-	1	-	1	4	4	2	2	-	-	-
White, both sexes	1	-	3	2	3	2	6	2	2	-	-	-
Male	1	-	2	2	3	-	2	1	-	-	-	-
Female	-	-	1	-	-	2	4	1	2	-	-	-
Black, both sexes	1	-	-	-	1	2	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
New Mexico												
All races, both sexes	-	2	-	1	1	-	1	-	-	-	-	-
Male	-	2	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	2	-	1	-	-	1	-	-	-	-	-
Male	-	2	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	11	5	6	11	17	11	10	6	-	-	-
Male	1	7	2	4	3	5	7	4	-	-	-	-
Female	3	4	3	2	8	12	4	6	6	-	-	-
White, both sexes	1	6	5	5	10	13	10	8	5	-	-	-
Male	-	2	2	3	3	4	6	4	-	-	-	-
Female	1	4	3	2	7	9	4	4	5	-	-	-
Black, both sexes	3	4	-	1	1	4	1	2	1	-	-	-
Male	1	4	-	1	-	1	1	-	-	-	-	-
Female	2	-	-	-	1	3	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
North Carolina												
All races, both sexes	6	3	3	3	3	5	7	1	2	-	-	-
Male	5	1	1	2	2	5	3	-	1	-	-	-
Female	1	2	2	1	1	-	4	1	1	-	-	-
White, both sexes	5	1	1	2	3	5	5	1	2	-	-	-
Male	4	1	-	1	2	5	1	-	1	-	-	-
Female	1	-	1	1	1	-	4	1	1	-	-	-
Black, both sexes	1	2	2	-	-	-	2	-	-	-	-	-
Male	1	-	1	-	-	-	2	-	-	-	-	-
Female	-	2	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Ohio												
All races, both sexes	4	9	3	4	5	4	6	4	-	1	-	-
Male	4	6	2	2	3	1	2	2	-	1	-	-
Female	-	3	1	2	2	3	4	2	-	-	-	-
White, both sexes	1	6	1	3	4	4	6	3	-	1	-	-
Male	1	5	1	1	3	1	2	2	-	1	-	-
Female	-	1	-	2	1	3	4	1	-	-	-	-
Black, both sexes	3	3	2	1	1	-	-	1	-	-	-	-
Male	3	1	1	1	-	-	-	-	-	-	-	-
Female	-	2	1	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	2	2	1	2	1	3	-	-	-	-	-
Male	3	2	2	-	1	1	1	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	-	-
White, both sexes	2	2	2	1	2	1	3	-	-	-	-	-
Male	2	2	2	-	1	1	1	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Oregon												
All races, both sexes	-	-	3	1	1	-	2	1	1	1	-	-
Male	-	-	2	-	1	-	1	-	1	-	-	-
Female	-	-	1	1	-	-	1	1	-	1	-	-
White, both sexes	-	-	3	-	1	-	2	1	1	1	-	-
Male	-	-	2	-	1	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	9	1	6	8	8	13	5	3	-	1	-
Male	3	8	-	4	2	4	4	1	2	-	-	-
Female	1	1	1	2	6	4	9	4	1	-	1	-
White, both sexes	3	5	1	6	8	7	13	5	3	-	1	-
Male	2	4	-	4	2	4	4	1	2	-	-	-
Female	1	1	1	2	6	3	9	4	1	-	1	-
Black, both sexes	1	4	-	-	-	1	-	-	-	-	-	-
Male	1	4	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Rhode Island												
All races, both sexes	-	-	-	-	1	-	2	1	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	-	-	-	-	1	-	2	1	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	3	1	2	1	3	4	-	-	-	-	-
Male	-	2	1	1	-	1	3	-	-	-	-	-
Female	1	1	-	1	1	2	1	-	-	-	-	-
White, both sexes	1	2	-	2	-	3	3	-	-	-	-	-
Male	-	1	-	1	-	1	2	-	-	-	-	-
Female	1	1	-	1	-	2	1	-	-	-	-	-
Black, both sexes	-	1	1	-	1	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
South Dakota												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	2	6	7	1	1	1	1	-	-	-	-
Male	2	1	3	3	1	1	-	1	-	-	-	-
Female	1	1	3	4	-	-	1	-	-	-	-	-
White, both sexes	3	1	4	6	-	1	1	1	-	-	-	-
Male	2	1	2	3	-	1	-	1	-	-	-	-
Female	1	-	2	3	-	-	1	-	-	-	-	-
Black, both sexes	-	1	2	1	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Texas												
All races, both sexes	5	9	5	7	7	10	10	6	1	-	1	-
Male	4	6	3	3	4	2	6	1	-	-	-	-
Female	1	3	2	4	3	8	4	5	1	-	1	-
White, both sexes	3	8	2	6	7	9	10	6	-	-	1	-
Male	3	6	1	3	4	2	6	1	-	-	-	-
Female	-	2	1	3	3	7	4	5	-	-	1	-
Black, both sexes	2	1	3	1	-	1	-	-	1	-	-	-
Male	1	-	2	-	-	-	-	-	-	-	-	-
Female	1	1	1	1	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	1	-	1	3	1	-	-	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
White, both sexes	-	1	1	-	1	3	1	-	-	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	6	3	1	4	4	2	4	-	-	-	-
Male	2	3	1	1	1	4	2	2	-	-	-	-
Female	-	3	2	-	3	-	-	2	-	-	-	-
White, both sexes	1	5	-	-	4	4	2	4	-	-	-	-
Male	1	3	-	-	1	4	2	2	-	-	-	-
Female	-	2	-	-	3	-	-	2	-	-	-	-
Black, both sexes	1	1	2	1	-	-	-	-	-	-	-	-
Male	1	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Washington												
All races, both sexes	-	2	4	5	1	5	1	4	-	-	-	-
Male	-	1	1	3	-	3	1	2	-	-	-	-
Female	-	1	3	2	1	2	-	2	-	-	-	-
White, both sexes	-	1	4	5	1	5	1	4	-	-	-	-
Male	-	-	1	3	-	3	1	2	-	-	-	-
Female	-	1	3	2	1	2	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	-	2	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	2	1	-	-	-	-	-	-	-
White, both sexes	-	1	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	1	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Wisconsin												
All races, both sexes	4	2	4	-	1	1	2	1	-	-	-	-
Male	2	2	1	-	-	1	2	-	-	-	-	-
Female	2	-	3	-	1	-	-	1	-	-	-	-
White, both sexes	3	1	2	-	1	1	2	1	-	-	-	-
Male	1	1	1	-	-	1	2	-	-	-	-	-
Female	2	-	1	-	1	-	-	1	-	-	-	-
Black, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
United States												
All races, both sexes	816	18	24	14	7	11	25	31	17	37	41	52
Male	431	12	14	6	2	8	11	18	12	23	22	35
Female	385	6	10	8	5	3	14	13	5	14	19	17
White, both sexes	675	11	19	8	4	7	20	23	15	24	34	38
Male	364	7	11	5	1	7	8	14	11	13	19	27
Female	311	4	8	3	3	-	12	9	4	11	15	11
Black, both sexes	109	5	4	3	1	4	4	7	2	9	5	14
Male	55	3	3	-	1	1	2	3	1	8	2	8
Female	54	2	1	3	-	3	2	4	1	1	3	6
American Indian 1/, both sexes	10	-	-	-	1	-	-	1	-	2	1	-
Male	4	-	-	-	-	-	-	1	-	1	1	-
Female	6	-	-	-	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	22	2	1	3	1	-	1	-	-	2	1	-
Male	8	2	-	1	-	-	1	-	-	1	-	-
Female	14	-	1	2	1	-	-	-	-	1	1	-
Alabama												
All races, both sexes	8	-	2	-	-	-	-	-	-	-	-	1
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	6	-	1	-	-	-	-	-	-	-	-	1
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Alaska												
All races, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	14	2	1	-	-	-	1	2	-	-	-	1
Male	10	2	1	-	-	-	1	2	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	2	1	-	-	-	1	1	-	-	-	1
Male	9	2	1	-	-	-	1	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Arkansas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	74	2	4	2	3	1	-	4	1	5	2	8
Male	39	1	2	1	-	1	-	3	1	2	1	7
Female	35	1	2	1	3	-	-	1	-	3	1	1
White, both sexes	58	1	3	-	2	1	-	3	1	4	2	8
Male	32	-	2	-	-	1	-	2	1	1	1	7
Female	26	1	1	-	2	-	-	1	-	3	1	1
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	1	1	2	1	-	-	-	-	-	-	-
Male	3	1	-	1	-	-	-	-	-	-	-	-
Female	6	-	1	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Colorado												
All races, both sexes	13	-	1	-	-	-	1	2	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	1	-	-	-	-	2	-	-	-	-
White, both sexes	11	-	1	-	-	-	1	2	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	1	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Florida												
All races, both sexes	49	-	2	2	-	1	-	1	1	1	1	1
Male	31	-	1	1	-	1	-	1	1	1	1	-
Female	18	-	1	1	-	-	-	-	-	-	-	1
White, both sexes	46	-	2	2	-	1	-	1	1	1	1	-
Male	29	-	1	1	-	1	-	1	1	1	1	-
Female	17	-	1	1	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	25	2	1	1	-	-	1	2	-	1	2	1
Male	12	1	1	-	-	-	-	1	-	1	1	1
Female	13	1	-	1	-	-	1	1	-	-	1	-
White, both sexes	16	1	1	-	-	-	1	1	-	-	1	-
Male	9	1	1	-	-	-	-	1	-	-	1	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	9	1	-	1	-	-	-	1	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	6	1	-	1	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hawaii												
All races, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	4	1	1	-	-	-	-	-	-	-	-	-
Male	4	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	1	-	-	-	-	-	-	-	-	-
Male	4	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Illinois												
All races, both sexes	32	-	-	-	-	-	-	1	2	1	1	3
Male	20	-	-	-	-	-	-	-	2	1	1	3
Female	12	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	1	2	-	1	2
Male	16	-	-	-	-	-	-	-	2	-	1	2
Female	8	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	15	-	-	-	-	1	1	1	1	1	-	-
Male	6	-	-	-	-	1	-	-	-	1	-	-
Female	9	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	14	-	-	-	-	1	1	1	1	1	-	-
Male	6	-	-	-	-	1	-	-	-	1	-	-
Female	8	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Iowa												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	16	1	-	-	-	-	-	1	-	-	1	1
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	15	1	-	-	-	-	-	1	-	-	1	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Kentucky												
All races, both sexes	9	-	-	-	-	-	1	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14	-	3	1	-	-	1	-	-	-	-	1
Male	6	-	2	-	-	-	-	-	-	-	-	1
Female	8	-	1	1	-	-	1	-	-	-	-	-
White, both sexes	8	-	2	-	-	-	1	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	6	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	6	-	1	1	-	-	-	-	-	-	-	1
Male	4	-	1	-	-	-	-	-	-	-	-	1
Female	2	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Maine												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	24	1	-	-	-	-	1	-	1	-	2	1
Male	8	1	-	-	-	-	-	-	1	-	1	-
Female	16	-	-	-	-	-	1	-	-	-	1	1
White, both sexes	20	-	-	-	-	-	1	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	1	-	1	-
Female	13	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Massachusetts												
All races, both sexes	26	-	1	1	-	-	3	3	-	2	1	3
Male	16	-	-	1	-	-	3	3	-	2	-	2
Female	10	-	1	-	-	-	-	-	-	-	1	1
White, both sexes	26	-	1	1	-	-	3	3	-	2	1	3
Male	16	-	-	1	-	-	3	3	-	2	-	2
Female	10	-	1	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	26	1	-	1	-	-	1	2	-	1	-	-
Male	17	1	-	1	-	-	1	2	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	1	-	-	-	1	-	-	-	-
Male	12	-	-	1	-	-	-	1	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	1	1	-	1	-	-
Male	4	-	-	-	-	-	1	1	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Minnesota												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Mississippi												
All races, both sexes	7	-	-	-	-	1	-	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	1	-	-	1	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Missouri												
All races, both sexes	12	1	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Nebraska												
All races, both sexes	7	-	-	-	-	-	-	1	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Hampshire												
All races, both sexes	4	-	-	-	-	1	-	-	-	2	-	-
Male	4	-	-	-	-	1	-	-	-	2	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	1	-	-	-	2	-	-
Male	4	-	-	-	-	1	-	-	-	2	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	26	-	-	1	1	-	-	-	1	1	2	-
Male	12	-	-	1	-	-	-	-	-	-	1	-
Female	14	-	-	-	1	-	-	-	1	1	1	-
White, both sexes	23	-	-	1	1	-	-	-	1	1	2	-
Male	10	-	-	1	-	-	-	-	-	-	1	-
Female	13	-	-	-	1	-	-	-	1	1	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Mexico												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	50	1	3	2	-	1	1	1	3	2	2	3
Male	32	1	2	-	-	1	1	1	2	-	1	2
Female	18	-	1	2	-	-	-	-	1	2	1	1
White, both sexes	40	-	2	-	-	1	-	1	3	1	2	2
Male	27	-	1	-	-	1	-	1	2	-	1	1
Female	13	-	1	-	-	-	-	-	1	1	1	1
Black, both sexes	7	1	1	1	-	-	-	-	-	1	-	1
Male	4	1	1	-	-	-	-	-	-	-	-	1
Female	3	-	-	1	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
North Carolina												
All races, both sexes	29	2	1	1	-	1	-	2	1	1	2	2
Male	17	2	1	1	-	1	-	1	1	1	-	1
Female	12	-	-	-	-	-	-	1	-	-	2	1
White, both sexes	24	1	1	1	-	1	-	2	-	1	1	1
Male	14	1	1	1	-	1	-	1	-	1	-	1
Female	10	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	5	1	-	-	-	-	-	-	1	-	1	1
Male	3	1	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Ohio												
All races, both sexes	36	2	-	-	-	2	5	-	1	-	3	3
Male	19	1	-	-	-	2	1	-	1	-	2	2
Female	17	1	-	-	-	-	4	-	-	-	1	1
White, both sexes	32	1	-	-	-	1	5	-	1	-	3	3
Male	17	1	-	-	-	1	1	-	1	-	2	2
Female	15	-	-	-	-	-	4	-	-	-	1	1
Black, both sexes	4	1	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Oregon												
All races, both sexes	10	-	-	1	-	-	1	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	9	-	-	1	-	-	1	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	39	-	-	-	-	-	-	1	-	1	4	2
Male	16	-	-	-	-	-	-	-	-	-	1	2
Female	23	-	-	-	-	-	-	1	-	1	3	-
White, both sexes	36	-	-	-	-	-	-	1	-	1	3	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	22	-	-	-	-	-	-	1	-	1	2	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	15	-	2	1	-	-	1	1	-	1	-	2
Male	7	-	1	-	-	-	-	1	-	1	-	1
Female	8	-	1	1	-	-	1	-	-	-	-	1
White, both sexes	8	-	1	1	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	1	1	-	-	-	-	-	-	-	-
Black, both sexes	7	-	1	-	-	-	1	1	-	1	-	2
Male	5	-	1	-	-	-	-	1	-	1	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	24	-	-	-	-	1	1	1	1	-	1	3
Male	14	-	-	-	-	-	1	1	1	-	1	2
Female	10	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	1	1	1	-	1	2
Male	14	-	-	-	-	-	1	1	1	-	1	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Texas												
All races, both sexes	57	2	1	-	-	1	2	2	1	4	2	5
Male	27	-	-	-	-	-	1	-	1	2	2	3
Female	30	2	1	-	-	1	1	2	-	2	-	2
White, both sexes	44	2	1	-	-	-	1	1	1	3	2	4
Male	21	-	-	-	-	-	-	-	1	1	2	3
Female	23	2	1	-	-	-	1	1	-	2	-	1
Black, both sexes	12	-	-	-	-	1	1	1	-	1	-	1
Male	6	-	-	-	-	-	1	-	-	1	-	-
Female	6	-	-	-	-	1	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	15	-	-	-	1	-	-	-	-	-	2	1
Male	9	-	-	-	1	-	-	-	-	-	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	1	-	-	-	-	-	1	1
Male	4	-	-	-	1	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Washington												
All races, both sexes	17	-	-	-	-	-	-	-	1	2	2	1
Male	7	-	-	-	-	-	-	-	-	1	1	1
Female	10	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	14	-	-	-	-	-	-	-	1	1	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	10	-	1	-	1	-	1	1	-	2	-	-
Male	5	-	1	-	1	-	-	1	-	1	-	-
Female	5	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	10	-	1	-	1	-	1	1	-	2	-	-
Male	5	-	1	-	1	-	-	1	-	1	-	-
Female	5	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Wisconsin												
All races, both sexes	16	-	-	-	-	-	-	2	-	-	2	1
Male	9	-	-	-	-	-	-	1	-	-	1	1
Female	7	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	15	-	-	-	-	-	-	1	-	-	2	1
Male	9	-	-	-	-	-	-	1	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
United States												
All races, both sexes	55	57	65	46	70	79	72	53	34	8	-	-
Male	33	34	37	28	35	34	37	20	8	2	-	-
Female	22	23	28	18	35	45	35	33	26	6	-	-
White, both sexes	44	44	59	40	59	71	67	49	33	6	-	-
Male	26	28	36	24	31	33	35	18	8	2	-	-
Female	18	16	23	16	28	38	32	31	25	4	-	-
Black, both sexes	8	10	5	5	8	7	4	1	1	2	-	-
Male	6	6	-	4	4	1	1	1	-	-	-	-
Female	2	4	5	1	4	6	3	-	1	2	-	-
American Indian 1/, both sexes	2	1	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	-	3	-	1	3	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	1	2	-	-	3	-	-	2	-	-	-	-
Alabama												
All races, both sexes	1	1	-	-	-	1	1	1	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	1	1	-	1	-	1	1	1	-	-	-
Male	1	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	1	1	1	-	1	-	1	1	1	-	-	-
Male	1	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Arkansas												
All races, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	7	3	4	7	3	3	6	3	1	-	-
Male	4	4	1	3	3	-	2	3	-	-	-	-
Female	1	3	2	1	4	3	1	3	3	1	-	-
White, both sexes	3	6	3	4	5	3	3	2	3	1	-	-
Male	3	4	1	3	3	-	2	1	-	-	-	-
Female	-	2	2	1	2	3	1	1	3	1	-	-
Black, both sexes	1	1	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Colorado												
All races, both sexes	-	1	2	-	-	2	4	-	-	-	-	-
Male	-	1	1	-	-	1	4	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	2	3	-	-	-	-	-
Male	-	1	-	-	-	1	3	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	2	-	-	1	4	2	-	1	-	-
Male	1	1	1	-	-	1	3	-	-	-	-	-
Female	-	-	1	-	-	-	1	2	-	1	-	-
White, both sexes	1	1	2	-	-	1	4	2	-	1	-	-
Male	1	1	1	-	-	1	3	-	-	-	-	-
Female	-	-	1	-	-	-	1	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Florida												
All races, both sexes	3	3	3	2	9	7	5	5	2	-	-	-
Male	3	3	1	-	7	3	4	3	-	-	-	-
Female	-	-	2	2	2	4	1	2	2	-	-	-
White, both sexes	3	3	3	2	7	7	5	5	2	-	-	-
Male	3	3	1	-	5	3	4	3	-	-	-	-
Female	-	-	2	2	2	4	1	2	2	-	-	-
Black, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	2	2	1	3	-	-	3	1	-	-
Male	1	1	1	1	1	-	-	-	-	1	-	-
Female	-	-	1	1	-	3	-	-	3	-	-	-
White, both sexes	1	1	2	1	1	1	-	-	3	1	-	-
Male	1	1	1	-	1	-	-	-	-	1	-	-
Female	-	-	1	1	-	1	-	-	3	-	-	-
Black, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hawaii												
All races, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Illinois												
All races, both sexes	2	2	3	3	4	6	2	-	1	1	-	-
Male	2	1	-	3	3	2	1	-	1	-	-	-
Female	-	1	3	-	1	4	1	-	-	1	-	-
White, both sexes	1	2	2	3	4	4	1	-	1	-	-	-
Male	1	1	-	3	3	2	-	-	1	-	-	-
Female	-	1	2	-	1	2	1	-	-	-	-	-
Black, both sexes	1	-	1	-	-	2	1	-	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	2	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	2	1	-	1	-	1	3	-	-	-
Male	1	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	1	1	-	1	-	1	2	-	-	-
White, both sexes	1	1	2	1	-	1	-	1	2	-	-	-
Male	1	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	1	1	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Iowa												
All races, both sexes	-	1	2	-	-	2	1	-	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	-	-	-	-
White, both sexes	-	1	2	-	-	2	1	-	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	1	-	-	2	2	2	2	1	-	-	-
Male	1	1	-	-	2	1	1	-	1	-	-	-
Female	1	-	-	-	-	1	1	2	-	-	-	-
White, both sexes	2	1	-	-	2	2	2	2	1	-	-	-
Male	1	1	-	-	2	1	1	-	1	-	-	-
Female	1	-	-	-	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Kentucky												
All races, both sexes	1	-	1	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	1	-	1	-	1	-	1	-	-	-
White, both sexes	1	-	1	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	1	-	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	2	1	-	3	-	1	1	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	2	-	1	1	-	-	-
White, both sexes	-	-	2	-	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Maine												
All races, both sexes	-	1	1	1	-	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	1	1	1	-	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	3	2	-	1	5	3	3	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	1	2	2	-	1	4	1	2	-	-	-	-
White, both sexes	1	2	2	-	-	4	3	3	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	1	1	2	-	-	3	1	2	-	-	-	-
Black, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Massachusetts												
All races, both sexes	1	-	-	1	3	1	2	2	2	-	-	-
Male	1	-	-	1	2	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	2	2	-	-	-
White, both sexes	1	-	-	1	3	1	2	2	2	-	-	-
Male	1	-	-	1	2	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	-	4	3	1	3	5	1	1	-	-	-
Male	1	-	3	2	-	2	2	-	1	-	-	-
Female	1	-	1	1	1	1	3	1	-	-	-	-
White, both sexes	1	-	4	3	1	3	4	1	1	-	-	-
Male	-	-	3	2	-	2	2	-	1	-	-	-
Female	1	-	1	1	1	1	2	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Minnesota												
All races, both sexes	1	3	-	2	-	2	3	-	-	-	-	-
Male	1	3	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	-	-	-	-	-
White, both sexes	1	2	-	2	-	2	3	-	-	-	-	-
Male	1	2	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	1	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Missouri												
All races, both sexes	-	2	2	-	5	1	-	-	-	-	-	-
Male	-	2	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	5	1	-	-	-	-	-	-
White, both sexes	-	2	2	-	4	1	-	-	-	-	-	-
Male	-	2	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	4	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Nebraska												
All races, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	2	-	1	-	1	-	1	-	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	2	-	1	-	1	-	1	-	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Hampshire												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	3	2	3	3	1	2	2	1	-	-
Male	1	-	2	2	1	2	1	1	-	-	-	-
Female	-	2	1	-	2	1	-	1	2	1	-	-
White, both sexes	-	2	2	1	3	3	1	2	2	1	-	-
Male	-	-	2	1	1	2	1	1	-	-	-	-
Female	-	2	-	-	2	1	-	1	2	1	-	-
Black, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Mexico												
All races, both sexes	-	-	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	3	4	2	4	5	5	3	1	-	-	-
Male	2	2	3	2	2	4	5	1	-	-	-	-
Female	2	1	1	-	2	1	-	2	1	-	-	-
White, both sexes	4	2	4	2	2	5	5	3	1	-	-	-
Male	2	1	3	2	2	4	5	1	-	-	-	-
Female	2	1	1	-	-	1	-	2	1	-	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
North Carolina												
All races, both sexes	2	1	3	1	2	1	3	1	2	-	-	-
Male	1	1	2	1	-	-	1	1	1	-	-	-
Female	1	-	1	-	2	1	2	-	1	-	-	-
White, both sexes	1	1	3	1	2	1	3	1	2	-	-	-
Male	-	1	2	1	-	-	1	1	1	-	-	-
Female	1	-	1	-	2	1	2	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Ohio												
All races, both sexes	3	1	1	4	3	2	3	2	1	-	-	-
Male	2	-	1	2	2	-	2	1	-	-	-	-
Female	1	1	-	2	1	2	1	1	1	-	-	-
White, both sexes	3	-	1	4	2	2	3	2	1	-	-	-
Male	2	-	1	2	1	-	2	1	-	-	-	-
Female	1	-	-	2	1	2	1	1	1	-	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	-	-	-	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	2	-	-	-	-	3	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Oregon												
All races, both sexes	-	-	1	1	1	1	1	1	1	-	-	-
Male	-	-	1	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	1	1	1	1	1	-	-	-
Male	-	-	1	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	4	1	2	2	9	5	2	1	1	-	-
Male	3	3	-	-	1	2	2	1	1	-	-	-
Female	1	1	1	2	1	7	3	1	-	1	-	-
White, both sexes	4	3	1	2	2	9	5	2	1	1	-	-
Male	3	2	-	-	1	2	2	1	1	-	-	-
Female	1	1	1	2	1	7	3	1	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	1	1	-	1	-	-	1	-	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	2	1	1	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
South Dakota												
All races, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	1	4	-	2	3	-	1	3	1	-	-
Male	-	-	2	-	2	3	-	-	1	-	-	-
Female	1	1	2	-	-	-	-	1	2	1	-	-
White, both sexes	-	1	3	-	2	3	-	1	3	-	-	-
Male	-	-	2	-	2	3	-	-	1	-	-	-
Female	-	1	1	-	-	-	-	1	2	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Texas												
All races, both sexes	5	4	5	7	5	2	3	4	1	1	-	-
Male	2	3	4	2	2	2	-	2	-	1	-	-
Female	3	1	1	5	3	-	3	2	1	-	-	-
White, both sexes	4	1	5	5	4	2	2	4	1	1	-	-
Male	2	1	4	1	1	2	-	2	-	1	-	-
Female	2	-	1	4	3	-	2	2	1	-	-	-
Black, both sexes	-	3	-	2	1	-	1	-	-	-	-	-
Male	-	2	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	1	1	-	1	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	2	-	1	1	-	1	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Vermont												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	2	-	2	2	3	-	-	-	-	-
Male	-	1	-	-	1	2	1	-	-	-	-	-
Female	1	-	2	-	1	-	2	-	-	-	-	-
White, both sexes	1	-	1	-	2	2	3	-	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	1	-	1	-	1	-	2	-	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Washington												
All races, both sexes	-	3	1	1	1	1	-	4	-	-	-	-
Male	-	-	1	1	-	-	-	2	-	-	-	-
Female	-	3	-	-	1	1	-	2	-	-	-	-
White, both sexes	-	2	1	1	-	1	-	4	-	-	-	-
Male	-	-	1	1	-	-	-	2	-	-	-	-
Female	-	2	-	-	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	1	-	-	1	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Wisconsin												
All races, both sexes	1	2	2	-	3	-	3	-	-	-	-	-
Male	-	1	1	-	2	-	2	-	-	-	-	-
Female	1	1	1	-	1	-	1	-	-	-	-	-
White, both sexes	1	2	2	-	3	-	3	-	-	-	-	-
Male	-	1	1	-	2	-	2	-	-	-	-	-
Female	1	1	1	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
United States												
All races, both sexes	21,386	1	-	4	4	17	32	50	83	93	190	276
Male	8,432	-	-	2	3	6	17	28	49	55	111	176
Female	12,954	1	-	2	1	11	15	22	34	38	79	100
White, both sexes	19,779	1	-	4	4	14	28	36	58	67	146	208
Male	7,749	-	-	2	3	5	16	20	37	39	88	133
Female	12,030	1	-	2	1	9	12	16	21	28	58	75
Black, both sexes	1,212	-	-	-	-	3	3	11	20	23	39	60
Male	512	-	-	-	-	1	-	6	10	14	21	38
Female	700	-	-	-	-	2	3	5	10	9	18	22
American Indian 1/, both sexes	73	-	-	-	-	-	-	2	-	1	2	5
Male	40	-	-	-	-	-	-	2	-	1	1	5
Female	33	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	322	-	-	-	-	-	1	1	5	2	3	3
Male	131	-	-	-	-	-	1	-	2	1	1	-
Female	191	-	-	-	-	-	-	1	3	1	2	3
Alabama												
All races, both sexes	262	-	-	-	-	-	1	-	3	-	4	3
Male	95	-	-	-	-	-	-	-	1	-	1	3
Female	167	-	-	-	-	-	1	-	2	-	3	-
White, both sexes	215	-	-	-	-	-	1	-	2	-	3	-
Male	79	-	-	-	-	-	-	-	1	-	1	-
Female	136	-	-	-	-	-	1	-	1	-	2	-
Black, both sexes	46	-	-	-	-	-	-	-	1	-	1	3
Male	16	-	-	-	-	-	-	-	-	-	-	3
Female	30	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Alaska												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	328	-	-	-	-	-	1	2	4	1	4	3
Male	150	-	-	-	-	-	1	2	3	1	3	2
Female	178	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	316	-	-	-	-	-	1	1	3	1	3	2
Male	142	-	-	-	-	-	1	1	2	1	2	2
Female	174	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Arkansas												
All races, both sexes	186	-	-	-	-	1	2	-	1	1	-	4
Male	66	-	-	-	-	-	-	-	1	-	-	2
Female	120	-	-	-	-	1	2	-	-	1	-	2
White, both sexes	176	-	-	-	-	1	2	-	1	1	-	4
Male	64	-	-	-	-	-	-	-	1	-	-	2
Female	112	-	-	-	-	1	2	-	-	1	-	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,255	-	-	1	1	-	2	7	7	9	12	19
Male	965	-	-	-	1	-	1	5	3	7	7	13
Female	1,290	-	-	1	-	-	1	2	4	2	5	6
White, both sexes	2,018	-	-	1	1	-	2	4	4	6	10	18
Male	866	-	-	-	1	-	1	3	3	5	7	13
Female	1,152	-	-	1	-	-	1	1	1	1	3	5
Black, both sexes	92	-	-	-	-	-	-	3	1	1	1	-
Male	34	-	-	-	-	-	-	2	-	1	-	-
Female	58	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	136	-	-	-	-	-	-	-	2	2	1	1
Male	57	-	-	-	-	-	-	-	-	1	-	-
Female	79	-	-	-	-	-	-	-	2	1	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Colorado												
All races, both sexes	301	-	-	-	-	-	-	-	1	-	2	5
Male	118	-	-	-	-	-	-	-	1	-	-	4
Female	183	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	293	-	-	-	-	-	-	-	1	-	2	5
Male	116	-	-	-	-	-	-	-	1	-	-	4
Female	177	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	323	-	-	-	-	-	-	1	2	1	2	4
Male	130	-	-	-	-	-	-	1	2	-	1	2
Female	193	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	308	-	-	-	-	-	-	-	1	1	2	4
Male	121	-	-	-	-	-	-	-	1	-	1	2
Female	187	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	15	-	-	-	-	-	-	1	1	-	-	-
Male	9	-	-	-	-	-	-	1	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Delaware												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	30	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Florida												
All races, both sexes	1,489	-	-	-	-	-	-	1	4	7	15	18
Male	634	-	-	-	-	-	-	-	3	5	6	11
Female	855	-	-	-	-	-	-	1	1	2	9	7
White, both sexes	1,392	-	-	-	-	-	-	1	3	6	11	13
Male	590	-	-	-	-	-	-	-	2	4	5	10
Female	802	-	-	-	-	-	-	1	1	2	6	3
Black, both sexes	88	-	-	-	-	-	-	-	1	1	4	5
Male	40	-	-	-	-	-	-	-	1	1	1	1
Female	48	-	-	-	-	-	-	-	-	-	3	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	393	-	-	-	-	-	-	1	-	1	6	8
Male	175	-	-	-	-	-	-	1	-	1	4	6
Female	218	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	324	-	-	-	-	-	-	-	-	1	3	4
Male	140	-	-	-	-	-	-	-	-	1	2	3
Female	184	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	67	-	-	-	-	-	-	1	-	-	2	4
Male	34	-	-	-	-	-	-	1	-	-	2	3
Female	33	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Hawaii												
All races, both sexes	92	-	-	-	-	-	-	-	-	-	-	2
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	74	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	115	-	-	-	-	-	-	-	-	-	3	1
Male	45	-	-	-	-	-	-	-	-	-	3	-
Female	70	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	115	-	-	-	-	-	-	-	-	-	3	1
Male	45	-	-	-	-	-	-	-	-	-	3	-
Female	70	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Illinois												
All races, both sexes	828	-	-	-	-	-	-	3	4	6	8	11
Male	314	-	-	-	-	-	-	2	2	4	5	10
Female	514	-	-	-	-	-	-	1	2	2	3	1
White, both sexes	758	-	-	-	-	-	-	2	4	3	3	9
Male	289	-	-	-	-	-	-	1	2	2	3	8
Female	469	-	-	-	-	-	-	1	2	1	-	1
Black, both sexes	63	-	-	-	-	-	-	1	-	3	4	2
Male	22	-	-	-	-	-	-	1	-	2	1	2
Female	41	-	-	-	-	-	-	-	-	1	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	494	-	-	-	-	-	-	1	4	3	7	5
Male	176	-	-	-	-	-	-	1	2	3	5	5
Female	318	-	-	-	-	-	-	-	2	-	2	-
White, both sexes	470	-	-	-	-	-	-	1	3	3	7	3
Male	166	-	-	-	-	-	-	1	1	3	5	3
Female	304	-	-	-	-	-	-	-	2	-	2	-
Black, both sexes	22	-	-	-	-	-	-	-	1	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Iowa												
All races, both sexes	273	-	-	-	1	-	1	-	-	1	1	-
Male	102	-	-	-	-	-	-	-	-	-	1	-
Female	171	-	-	-	1	-	1	-	-	1	-	-
White, both sexes	272	-	-	-	1	-	1	-	-	1	1	-
Male	102	-	-	-	-	-	-	-	-	-	1	-
Female	170	-	-	-	1	-	1	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	233	-	-	-	-	-	-	1	-	1	3	2
Male	85	-	-	-	-	-	-	-	-	1	1	2
Female	148	-	-	-	-	-	-	1	-	-	2	-
White, both sexes	225	-	-	-	-	-	-	1	-	1	3	1
Male	81	-	-	-	-	-	-	-	-	1	1	1
Female	144	-	-	-	-	-	-	1	-	-	2	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Kentucky												
All races, both sexes	243	-	-	-	-	3	3	1	2	1	1	8
Male	100	-	-	-	-	2	1	1	-	-	1	5
Female	143	-	-	-	-	1	2	-	2	1	-	3
White, both sexes	229	-	-	-	-	3	2	1	1	1	1	6
Male	95	-	-	-	-	2	1	1	-	-	1	4
Female	134	-	-	-	-	1	1	-	1	1	-	2
Black, both sexes	11	-	-	-	-	-	1	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Louisiana												
All races, both sexes	252	-	-	-	-	-	-	-	-	5	2	7
Male	115	-	-	-	-	-	-	-	-	4	2	7
Female	137	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	210	-	-	-	-	-	-	-	-	3	1	2
Male	91	-	-	-	-	-	-	-	-	2	1	2
Female	119	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	40	-	-	-	-	-	-	-	-	2	1	5
Male	23	-	-	-	-	-	-	-	-	2	1	5
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Maine												
All races, both sexes	158	-	-	-	-	-	-	-	-	-	1	-
Male	68	-	-	-	-	-	-	-	-	-	-	-
Female	90	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	157	-	-	-	-	-	-	-	-	-	1	-
Male	67	-	-	-	-	-	-	-	-	-	-	-
Female	90	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	353	-	-	-	-	1	-	-	4	2	6	9
Male	127	-	-	-	-	-	-	-	3	1	2	6
Female	226	-	-	-	-	1	-	-	1	1	4	3
White, both sexes	277	-	-	-	-	1	-	-	1	2	1	7
Male	95	-	-	-	-	-	-	-	-	1	-	5
Female	182	-	-	-	-	1	-	-	1	1	1	2
Black, both sexes	71	-	-	-	-	-	-	-	2	-	5	2
Male	28	-	-	-	-	-	-	-	2	-	2	1
Female	43	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Massachusetts												
All races, both sexes	545	-	-	-	-	-	1	3	1	4	8	6
Male	196	-	-	-	-	-	1	-	1	2	6	6
Female	349	-	-	-	-	-	-	3	-	2	2	-
White, both sexes	525	-	-	-	-	-	1	1	1	4	7	6
Male	191	-	-	-	-	-	1	-	1	2	5	6
Female	334	-	-	-	-	-	-	1	-	2	2	-
Black, both sexes	17	-	-	-	-	-	-	2	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	686	-	-	-	-	-	1	-	3	4	7	11
Male	275	-	-	-	-	-	-	-	2	2	4	5
Female	411	-	-	-	-	-	1	-	1	2	3	6
White, both sexes	633	-	-	-	-	-	1	-	1	1	5	5
Male	251	-	-	-	-	-	-	-	1	-	3	3
Female	382	-	-	-	-	-	1	-	-	1	2	2
Black, both sexes	47	-	-	-	-	-	-	-	2	3	2	5
Male	22	-	-	-	-	-	-	-	1	2	1	1
Female	25	-	-	-	-	-	-	-	1	1	1	4
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Minnesota												
All races, both sexes	409	-	-	-	-	1	-	1	1	-	2	3
Male	161	-	-	-	-	-	-	-	1	-	2	-
Female	248	-	-	-	-	1	-	1	-	-	-	3
White, both sexes	403	-	-	-	-	-	-	1	1	-	2	3
Male	159	-	-	-	-	-	-	-	1	-	2	-
Female	244	-	-	-	-	-	-	1	-	-	-	3
Black, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	116	-	-	-	-	-	-	-	-	3	2	5
Male	54	-	-	-	-	-	-	-	-	1	2	5
Female	62	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	92	-	-	-	-	-	-	-	-	3	1	2
Male	41	-	-	-	-	-	-	-	-	1	1	2
Female	51	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	1	3
Male	12	-	-	-	-	-	-	-	-	-	1	3
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Missouri												
All races, both sexes	451	-	-	-	-	-	-	-	1	3	5	3
Male	167	-	-	-	-	-	-	-	1	2	5	1
Female	284	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	425	-	-	-	-	-	-	-	-	3	5	2
Male	160	-	-	-	-	-	-	-	-	2	5	1
Female	265	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	26	-	-	-	-	-	-	-	1	-	-	1
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	89	-	-	-	1	-	-	-	-	-	-	2
Male	42	-	-	-	1	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	89	-	-	-	1	-	-	-	-	-	-	2
Male	42	-	-	-	1	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Nebraska												
All races, both sexes	168	-	-	-	-	-	-	1	1	1	1	-
Male	61	-	-	-	-	-	-	-	1	1	-	-
Female	107	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	165	-	-	-	-	-	-	1	1	1	-	-
Male	60	-	-	-	-	-	-	-	1	1	-	-
Female	105	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	145	-	-	-	-	-	-	2	2	2	2	4
Male	59	-	-	-	-	-	-	2	1	-	1	1
Female	86	-	-	-	-	-	-	-	1	2	1	3
White, both sexes	135	-	-	-	-	-	-	2	2	2	2	3
Male	55	-	-	-	-	-	-	2	1	-	1	1
Female	80	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
New Hampshire												
All races, both sexes	133	-	-	-	-	-	-	-	1	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	84	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	133	-	-	-	-	-	-	-	1	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	84	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	775	-	-	-	-	-	2	2	4	4	6	7
Male	280	-	-	-	-	-	1	2	2	2	3	4
Female	495	-	-	-	-	-	1	-	2	2	3	3
White, both sexes	720	-	-	-	-	-	1	2	3	2	4	4
Male	258	-	-	-	-	-	1	2	2	1	1	2
Female	462	-	-	-	-	-	-	-	1	1	3	2
Black, both sexes	43	-	-	-	-	-	1	-	1	2	2	3
Male	19	-	-	-	-	-	-	-	-	1	2	2
Female	24	-	-	-	-	-	1	-	1	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
New Mexico												
All races, both sexes	105	-	-	-	-	-	-	1	1	-	4	2
Male	49	-	-	-	-	-	-	-	-	-	4	1
Female	56	-	-	-	-	-	-	1	1	-	-	1
White, both sexes	97	-	-	-	-	-	-	1	1	-	2	1
Male	44	-	-	-	-	-	-	-	-	-	2	-
Female	53	-	-	-	-	-	-	1	1	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,224	-	-	-	-	3	-	1	5	2	2	17
Male	466	-	-	-	-	1	-	-	2	-	1	13
Female	758	-	-	-	-	2	-	1	3	2	1	4
White, both sexes	1,143	-	-	-	-	2	-	1	3	2	1	13
Male	429	-	-	-	-	1	-	-	1	-	1	10
Female	714	-	-	-	-	1	-	1	2	2	-	3
Black, both sexes	70	-	-	-	-	1	-	-	2	-	1	4
Male	33	-	-	-	-	-	-	-	1	-	-	3
Female	37	-	-	-	-	1	-	-	1	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
North Carolina												
All races, both sexes	583	-	-	1	-	1	1	3	4	5	2	7
Male	239	-	-	-	-	1	1	1	1	3	2	5
Female	344	-	-	1	-	-	-	2	3	2	-	2
White, both sexes	510	-	-	1	-	1	1	2	2	2	2	3
Male	210	-	-	-	-	1	1	1	1	2	2	1
Female	300	-	-	1	-	-	-	1	1	-	-	2
Black, both sexes	66	-	-	-	-	-	-	1	2	3	-	4
Male	24	-	-	-	-	-	-	-	-	1	-	4
Female	42	-	-	-	-	-	-	1	2	2	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	71	-	-	-	-	-	-	1	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	69	-	-	-	-	-	-	1	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Ohio												
All races, both sexes	963	-	-	-	-	1	3	1	6	5	7	12
Male	333	-	-	-	-	1	2	-	5	3	3	6
Female	630	-	-	-	-	-	1	1	1	2	4	6
White, both sexes	911	-	-	-	-	-	3	1	6	2	6	8
Male	314	-	-	-	-	-	2	-	5	1	3	6
Female	597	-	-	-	-	-	1	1	1	1	3	2
Black, both sexes	48	-	-	-	-	1	-	-	-	3	1	4
Male	18	-	-	-	-	1	-	-	-	2	-	-
Female	30	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	236	-	-	-	-	1	-	2	-	5	3	6
Male	102	-	-	-	-	-	-	2	-	3	2	2
Female	134	-	-	-	-	1	-	-	-	2	1	4
White, both sexes	212	-	-	-	-	1	-	1	-	3	2	6
Male	90	-	-	-	-	-	-	1	-	2	1	2
Female	122	-	-	-	-	1	-	-	-	1	1	4
Black, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	1	-	1	1	-
Male	6	-	-	-	-	-	-	1	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Oregon												
All races, both sexes	491	-	-	-	-	-	-	-	1	2	1	3
Male	188	-	-	-	-	-	-	-	-	2	1	1
Female	303	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	477	-	-	-	-	-	-	-	1	2	1	2
Male	182	-	-	-	-	-	-	-	-	2	1	-
Female	295	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,515	1	-	-	1	-	1	2	1	4	12	19
Male	565	-	-	-	1	-	-	1	1	4	5	11
Female	950	1	-	-	-	-	1	1	-	-	7	8
White, both sexes	1,446	1	-	-	1	-	1	2	1	3	8	14
Male	533	-	-	-	1	-	-	1	1	3	4	8
Female	913	1	-	-	-	-	1	1	-	-	4	6
Black, both sexes	64	-	-	-	-	-	-	-	-	1	4	5
Male	30	-	-	-	-	-	-	-	-	1	1	3
Female	34	-	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Rhode Island												
All races, both sexes	89	-	-	-	-	-	-	-	-	1	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	87	-	-	-	-	-	-	-	-	1	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	60	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	301	-	-	-	-	-	-	-	-	1	5	4
Male	123	-	-	-	-	-	-	-	-	-	4	1
Female	178	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	254	-	-	-	-	-	-	-	-	-	3	3
Male	105	-	-	-	-	-	-	-	-	-	3	-
Female	149	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	46	-	-	-	-	-	-	-	-	1	2	1
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
South Dakota												
All races, both sexes	67	-	-	-	-	-	-	-	-	-	-	3
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	43	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	398	-	-	-	-	3	1	2	3	4	9	10
Male	179	-	-	-	-	-	-	2	2	2	5	7
Female	219	-	-	-	-	3	1	-	1	2	4	3
White, both sexes	363	-	-	-	-	3	-	2	3	2	9	8
Male	161	-	-	-	-	-	-	2	2	1	5	6
Female	202	-	-	-	-	3	-	-	1	1	4	2
Black, both sexes	35	-	-	-	-	-	1	-	-	2	-	2
Male	18	-	-	-	-	-	-	-	-	1	-	1
Female	17	-	-	-	-	-	1	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Texas												
All races, both sexes	1,065	-	-	1	-	-	7	4	2	2	15	20
Male	412	-	-	1	-	-	5	1	-	1	8	12
Female	653	-	-	-	-	-	2	3	2	1	7	8
White, both sexes	985	-	-	1	-	-	7	4	-	2	13	19
Male	378	-	-	1	-	-	5	1	-	1	6	11
Female	607	-	-	-	-	-	2	3	-	1	7	8
Black, both sexes	70	-	-	-	-	-	-	-	2	-	2	1
Male	27	-	-	-	-	-	-	-	-	-	2	1
Female	43	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	135	-	-	-	-	-	-	1	1	-	-	1
Male	61	-	-	-	-	-	-	-	1	-	-	-
Female	74	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	133	-	-	-	-	-	-	-	1	-	-	1
Male	61	-	-	-	-	-	-	-	1	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Vermont												
All races, both sexes	75	-	-	-	-	-	-	-	-	-	1	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	75	-	-	-	-	-	-	-	-	-	1	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	449	-	-	-	-	1	2	1	5	1	7	5
Male	174	-	-	-	-	1	2	-	4	-	4	4
Female	275	-	-	-	-	-	-	1	1	1	3	1
White, both sexes	404	-	-	-	-	1	2	-	3	1	5	5
Male	150	-	-	-	-	1	2	-	2	-	2	4
Female	254	-	-	-	-	-	-	-	1	1	3	1
Black, both sexes	42	-	-	-	-	-	-	1	2	-	2	-
Male	21	-	-	-	-	-	-	-	2	-	2	-
Female	21	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Washington												
All races, both sexes	679	-	-	-	-	-	2	2	1	1	6	5
Male	265	-	-	-	-	-	1	2	1	-	4	4
Female	414	-	-	-	-	-	1	-	-	1	2	1
White, both sexes	653	-	-	-	-	-	1	2	1	1	6	5
Male	254	-	-	-	-	-	-	2	1	-	4	4
Female	399	-	-	-	-	-	1	-	-	1	2	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	123	-	-	-	-	-	-	1	1	-	1	4
Male	49	-	-	-	-	-	-	1	1	-	1	3
Female	74	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	121	-	-	-	-	-	-	1	1	-	1	4
Male	49	-	-	-	-	-	-	1	1	-	1	3
Female	72	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Wisconsin												
All races, both sexes	548	-	-	-	-	-	1	-	2	-	5	4
Male	205	-	-	-	-	-	1	-	1	-	2	1
Female	343	-	-	-	-	-	-	-	1	-	3	3
White, both sexes	539	-	-	-	-	-	1	-	1	-	5	3
Male	204	-	-	-	-	-	1	-	1	-	2	1
Female	335	-	-	-	-	-	-	-	-	-	3	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	42	-	-	1	-	1	-	-	-	-	-	1
Male	17	-	-	1	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	42	-	-	1	-	1	-	-	-	-	-	1
Male	17	-	-	1	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
United States												
All races, both sexes	384	520	615	887	1,305	2,243	3,798	5,086	4,051	1,472	275	-
Male	220	335	333	468	648	990	1,621	1,883	1,150	306	31	-
Female	164	185	282	419	657	1,253	2,177	3,203	2,901	1,166	244	-
White, both sexes	275	420	535	751	1,161	2,078	3,556	4,862	3,910	1,409	256	-
Male	159	274	283	412	578	924	1,538	1,816	1,110	286	26	-
Female	116	146	252	339	583	1,154	2,018	3,046	2,800	1,123	230	-
Black, both sexes	89	84	57	109	112	127	180	143	96	41	15	-
Male	49	53	33	42	55	52	60	40	21	13	4	-
Female	40	31	24	67	57	75	120	103	75	28	11	-
American Indian 1/, both sexes	5	11	5	8	3	3	8	14	5	-	1	-
Male	3	6	5	4	2	-	4	6	1	-	-	-
Female	2	5	-	4	1	3	4	8	4	-	1	-
Asian or Pacific Islander 2/, both sexes	15	5	18	19	29	35	54	67	40	22	3	-
Male	9	2	12	10	13	14	19	21	18	7	1	-
Female	6	3	6	9	16	21	35	46	22	15	2	-
Alabama												
All races, both sexes	11	2	12	8	21	39	49	65	32	9	3	-
Male	5	1	6	4	14	15	17	20	6	2	-	-
Female	6	1	6	4	7	24	32	45	26	7	3	-
White, both sexes	6	1	7	5	18	30	41	61	30	8	2	-
Male	4	-	3	3	11	14	17	19	5	1	-	-
Female	2	1	4	2	7	16	24	42	25	7	2	-
Black, both sexes	5	1	5	3	3	9	7	4	2	1	1	-
Male	1	1	3	1	3	1	-	1	1	1	-	-
Female	4	-	2	2	-	8	7	3	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Alaska												
All races, both sexes	2	4	-	1	5	2	6	9	5	3	-	-
Male	-	2	-	1	4	1	1	4	2	1	-	-
Female	2	2	-	-	1	1	5	5	3	2	-	-
White, both sexes	1	2	-	1	5	1	4	4	5	3	-	-
Male	-	1	-	1	4	1	1	2	2	1	-	-
Female	1	1	-	-	1	-	3	2	3	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	5	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	1	1	-	-	-	-	-	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Arizona												
All races, both sexes	6	11	11	14	22	30	61	79	56	19	4	-
Male	5	5	7	6	6	16	24	41	21	5	2	-
Female	1	6	4	8	16	14	37	38	35	14	2	-
White, both sexes	6	9	10	13	21	30	60	79	55	19	3	-
Male	5	4	6	5	6	16	23	41	21	5	1	-
Female	1	5	4	8	15	14	37	38	34	14	2	-
Black, both sexes	-	-	1	-	1	-	1	-	-	-	1	-
Male	-	-	1	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Arkansas												
All races, both sexes	2	7	8	11	18	17	37	38	25	11	3	-
Male	-	4	4	4	2	7	17	12	10	2	1	-
Female	2	3	4	7	16	10	20	26	15	9	2	-
White, both sexes	1	6	7	10	17	15	35	37	25	11	3	-
Male	-	3	4	4	2	7	16	12	10	2	1	-
Female	1	3	3	6	15	8	19	25	15	9	2	-
Black, both sexes	1	1	1	1	1	2	2	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	1	2	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	46	39	52	62	129	223	415	580	442	182	27	-
Male	25	25	29	38	72	124	193	238	140	39	5	-
Female	21	14	23	24	57	99	222	342	302	143	22	-
White, both sexes	34	30	40	48	106	199	374	530	417	169	25	-
Male	18	20	21	31	57	114	180	225	129	35	3	-
Female	16	10	19	17	49	85	194	305	288	134	22	-
Black, both sexes	6	7	4	4	9	12	18	14	6	5	1	-
Male	3	3	3	1	7	4	6	2	-	1	1	-
Female	3	4	1	3	2	8	12	12	6	4	-	-
American Indian 1/, both sexes	2	1	1	1	1	-	1	2	-	-	-	-
Male	2	1	1	1	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	7	9	13	12	22	34	19	8	1	-
Male	2	1	4	5	7	6	6	10	11	3	1	-
Female	2	-	3	4	6	6	16	24	8	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Colorado												
All races, both sexes	6	9	13	11	29	30	50	79	43	20	3	-
Male	-	6	8	9	11	11	14	43	11	-	-	-
Female	6	3	5	2	18	19	36	36	32	20	3	-
White, both sexes	5	9	12	11	29	28	49	77	42	20	3	-
Male	-	6	7	9	11	11	14	42	11	-	-	-
Female	5	3	5	2	18	17	35	35	31	20	3	-
Black, both sexes	1	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Connecticut												
All races, both sexes	5	7	6	10	23	27	53	80	66	28	8	-
Male	5	3	2	3	15	9	26	36	19	5	1	-
Female	-	4	4	7	8	18	27	44	47	23	7	-
White, both sexes	2	6	6	8	20	27	52	79	65	27	8	-
Male	2	2	2	3	13	9	25	36	19	5	1	-
Female	-	4	4	5	7	18	27	43	46	22	7	-
Black, both sexes	3	1	-	2	3	-	1	1	1	1	-	-
Male	3	1	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	2	1	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Delaware												
All races, both sexes	2	4	6	8	2	6	10	12	7	4	4	-
Male	2	2	2	4	1	3	2	3	2	1	-	-
Female	-	2	4	4	1	3	8	9	5	3	4	-
White, both sexes	1	3	6	6	2	6	10	12	6	4	4	-
Male	1	1	2	3	1	3	2	3	2	1	-	-
Female	-	2	4	3	1	3	8	9	4	3	4	-
Black, both sexes	1	1	-	2	-	-	-	-	1	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	2	2	2	4	7	4	4	2	-	-
Male	1	-	1	1	1	2	2	-	1	1	-	-
Female	1	-	1	1	1	2	5	4	3	1	-	-
White, both sexes	-	-	1	-	-	3	5	2	2	1	-	-
Male	-	-	-	-	-	2	2	-	1	1	-	-
Female	-	-	1	-	-	1	3	2	1	-	-	-
Black, both sexes	2	-	1	2	2	1	2	2	2	1	-	-
Male	1	-	1	1	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	2	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Florida												
All races, both sexes	17	33	36	69	85	179	264	347	277	122	15	-
Male	16	22	19	37	46	80	123	143	90	32	1	-
Female	1	11	17	32	39	99	141	204	187	90	14	-
White, both sexes	11	27	33	59	77	165	252	336	268	117	13	-
Male	11	18	16	32	42	75	119	140	86	29	1	-
Female	-	9	17	27	35	90	133	196	182	88	12	-
Black, both sexes	6	5	2	10	7	12	11	9	9	4	2	-
Male	5	4	2	5	3	5	3	3	4	2	-	-
Female	1	1	-	5	4	7	8	6	5	2	2	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	2	1	1	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	1	-	-	-	2	-	1	-	-	-	-
Georgia												
All races, both sexes	8	21	11	33	25	48	71	76	59	23	2	-
Male	2	14	9	11	11	17	37	35	22	5	-	-
Female	6	7	2	22	14	31	34	41	37	18	2	-
White, both sexes	3	12	10	25	20	44	63	65	52	20	2	-
Male	1	9	8	10	8	14	33	29	18	4	-	-
Female	2	3	2	15	12	30	30	36	34	16	2	-
Black, both sexes	5	9	1	8	5	4	8	10	7	3	-	-
Male	1	5	1	1	3	3	4	5	4	1	-	-
Female	4	4	-	7	2	1	4	5	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Hawaii												
All races, both sexes	4	-	2	2	8	9	16	22	15	10	2	-
Male	3	-	2	1	4	6	8	11	5	3	-	-
Female	1	-	-	1	4	3	8	11	10	7	2	-
White, both sexes	1	-	1	-	-	3	5	5	1	-	-	-
Male	1	-	1	-	-	3	4	5	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	2	8	6	11	17	14	10	2	-
Male	2	-	1	1	4	3	4	6	5	3	-	-
Female	1	-	-	1	4	3	7	11	9	7	2	-
Idaho												
All races, both sexes	2	1	1	6	9	9	12	38	26	7	-	-
Male	1	1	1	4	4	4	6	12	8	1	-	-
Female	1	-	-	2	5	5	6	26	18	6	-	-
White, both sexes	2	1	1	6	9	9	12	38	26	7	-	-
Male	1	1	1	4	4	4	6	12	8	1	-	-
Female	1	-	-	2	5	5	6	26	18	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Illinois												
All races, both sexes	17	18	18	33	50	94	148	188	162	59	9	-
Male	12	10	15	16	20	37	50	76	43	11	1	-
Female	5	8	3	17	30	57	98	112	119	48	8	-
White, both sexes	13	14	16	27	40	86	139	178	158	58	8	-
Male	10	9	14	14	14	35	50	73	42	11	1	-
Female	3	5	2	13	26	51	89	105	116	47	7	-
Black, both sexes	4	3	-	5	10	6	9	10	4	1	1	-
Male	2	1	-	2	6	1	-	3	1	-	-	-
Female	2	2	-	3	4	5	9	7	3	1	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	1	-	-	-	-	-	-
Indiana												
All races, both sexes	6	15	14	27	31	59	78	108	99	29	8	-
Male	1	8	7	15	17	21	33	30	24	3	1	-
Female	5	7	7	12	14	38	45	78	75	26	7	-
White, both sexes	5	14	11	26	29	58	70	105	98	29	8	-
Male	1	8	4	14	16	21	32	30	23	3	1	-
Female	4	6	7	12	13	37	38	75	75	26	7	-
Black, both sexes	1	1	2	1	2	1	8	2	1	-	-	-
Male	-	-	2	1	1	-	1	-	1	-	-	-
Female	1	1	-	-	1	1	7	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Iowa												
All races, both sexes	1	5	5	6	19	30	38	71	62	24	8	-
Male	1	5	2	6	9	11	15	30	18	4	-	-
Female	-	-	3	-	10	19	23	41	44	20	8	-
White, both sexes	1	5	5	6	19	30	38	70	62	24	8	-
Male	1	5	2	6	9	11	15	30	18	4	-	-
Female	-	-	3	-	10	19	23	40	44	20	8	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	6	2	7	13	16	40	51	60	24	7	-
Male	-	4	2	5	5	6	14	21	17	5	2	-
Female	-	2	-	2	8	10	26	30	43	19	5	-
White, both sexes	-	6	1	7	13	15	40	51	58	22	6	-
Male	-	4	1	5	5	5	14	21	16	5	2	-
Female	-	2	-	2	8	10	26	30	42	17	4	-
Black, both sexes	-	-	1	-	-	1	-	-	1	2	1	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Kentucky												
All races, both sexes	4	8	10	17	10	27	47	50	37	11	3	-
Male	2	7	6	8	4	15	17	17	11	3	-	-
Female	2	1	4	9	6	12	30	33	26	8	3	-
White, both sexes	4	8	7	17	9	27	45	49	34	11	3	-
Male	2	7	4	8	3	15	16	17	11	3	-	-
Female	2	1	3	9	6	12	29	32	23	8	3	-
Black, both sexes	-	-	2	-	1	-	1	1	3	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	9	11	10	22	27	32	51	39	23	13	1	-
Male	6	5	6	8	14	17	21	11	8	5	1	-
Female	3	6	4	14	13	15	30	28	15	8	-	-
White, both sexes	5	7	9	14	24	29	48	36	21	11	-	-
Male	3	3	6	5	13	16	19	11	7	3	-	-
Female	2	4	3	9	11	13	29	25	14	8	-	-
Black, both sexes	4	4	1	7	3	3	2	3	2	2	1	-
Male	3	2	-	2	1	1	2	-	1	2	1	-
Female	1	2	1	5	2	2	-	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Maine												
All races, both sexes	-	5	3	8	9	26	21	38	32	13	2	-
Male	-	5	2	5	4	16	11	17	5	3	-	-
Female	-	-	1	3	5	10	10	21	27	10	2	-
White, both sexes	-	4	3	8	9	26	21	38	32	13	2	-
Male	-	4	2	5	4	16	11	17	5	3	-	-
Female	-	-	1	3	5	10	10	21	27	10	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	10	7	10	18	18	34	75	74	64	21	-	-
Male	4	5	4	6	8	14	35	24	12	3	-	-
Female	6	2	6	12	10	20	40	50	52	18	-	-
White, both sexes	7	4	6	8	9	24	60	67	62	18	-	-
Male	1	3	2	5	5	10	26	22	12	3	-	-
Female	6	1	4	3	4	14	34	45	50	15	-	-
Black, both sexes	2	3	4	9	9	10	13	7	2	3	-	-
Male	2	2	2	1	3	4	7	2	-	-	-	-
Female	-	1	2	8	6	6	6	5	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Massachusetts												
All races, both sexes	7	19	7	15	32	45	89	135	125	41	7	-
Male	5	12	2	5	19	17	33	41	38	7	1	-
Female	2	7	5	10	13	28	56	94	87	34	6	-
White, both sexes	7	19	6	14	30	40	87	132	122	41	7	-
Male	5	12	2	5	19	14	32	41	38	7	1	-
Female	2	7	4	9	11	26	55	91	84	34	6	-
Black, both sexes	-	-	1	1	1	4	1	3	3	-	-	-
Male	-	-	-	-	-	3	1	-	-	-	-	-
Female	-	-	1	1	1	1	-	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Michigan												
All races, both sexes	16	17	19	25	39	79	105	164	140	46	10	-
Male	10	11	13	16	22	34	43	63	38	10	2	-
Female	6	6	6	9	17	45	62	101	102	36	8	-
White, both sexes	8	14	19	21	34	73	101	157	138	45	10	-
Male	5	9	13	14	21	32	41	60	37	10	2	-
Female	3	5	6	7	13	41	60	97	101	35	8	-
Black, both sexes	8	2	-	3	4	6	3	7	1	1	-	-
Male	5	2	-	1	1	2	2	3	1	-	-	-
Female	3	-	-	2	3	4	1	4	-	1	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Minnesota												
All races, both sexes	3	6	9	13	27	43	62	108	87	37	6	-
Male	1	4	6	9	12	22	32	37	28	6	1	-
Female	2	2	3	4	15	21	30	71	59	31	5	-
White, both sexes	2	5	9	13	27	42	61	108	86	37	6	-
Male	1	3	6	9	12	22	31	37	28	6	1	-
Female	1	2	3	4	15	20	30	71	58	31	5	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	8	6	12	10	14	19	20	12	4	1	-
Male	-	4	4	8	3	6	6	7	7	1	-	-
Female	-	4	2	4	7	8	13	13	5	3	1	-
White, both sexes	-	4	5	8	9	12	14	18	12	3	1	-
Male	-	2	4	5	3	5	4	7	7	-	-	-
Female	-	2	1	3	6	7	10	11	5	3	1	-
Black, both sexes	-	3	1	4	1	2	5	2	-	1	-	-
Male	-	1	-	3	-	1	2	-	-	1	-	-
Female	-	2	1	1	1	1	3	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Missouri												
All races, both sexes	11	9	16	17	25	44	90	106	81	35	5	-
Male	7	5	8	8	14	16	40	30	21	7	2	-
Female	4	4	8	9	11	28	50	76	60	28	3	-
White, both sexes	11	8	14	13	21	43	85	102	79	34	5	-
Male	7	4	8	6	13	15	39	30	21	7	2	-
Female	4	4	6	7	8	28	46	72	58	27	3	-
Black, both sexes	-	1	2	4	4	1	5	4	2	1	-	-
Male	-	1	-	2	1	1	1	-	-	-	-	-
Female	-	-	2	2	3	-	4	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	1	-	4	11	10	15	35	9	-	-
Male	1	-	1	-	3	8	8	4	12	3	-	-
Female	-	-	-	-	1	3	2	11	23	6	-	-
White, both sexes	1	-	1	-	4	11	10	15	35	9	-	-
Male	1	-	1	-	3	8	8	4	12	3	-	-
Female	-	-	-	-	1	3	2	11	23	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Nebraska												
All races, both sexes	-	1	1	6	12	17	32	33	43	15	4	-
Male	-	1	-	3	8	6	17	7	12	3	2	-
Female	-	-	1	3	4	11	15	26	31	12	2	-
White, both sexes	-	1	1	6	12	17	31	32	43	15	4	-
Male	-	1	-	3	8	6	16	7	12	3	2	-
Female	-	-	1	3	4	11	15	25	31	12	2	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	4	8	11	22	9	32	22	15	6	1	-
Male	2	4	3	5	10	5	10	9	5	1	-	-
Female	1	-	5	6	12	4	22	13	10	5	1	-
White, both sexes	3	3	8	10	19	9	30	20	15	6	1	-
Male	2	3	3	5	8	5	10	8	5	1	-	-
Female	1	-	5	5	11	4	20	12	10	5	1	-
Black, both sexes	-	1	-	-	3	-	-	1	-	-	-	-
Male	-	1	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
New Hampshire												
All races, both sexes	1	2	4	4	6	14	26	38	28	5	3	-
Male	1	2	2	2	1	5	9	18	8	1	-	-
Female	-	-	2	2	5	9	17	20	20	4	3	-
White, both sexes	1	2	4	4	6	14	26	38	28	5	3	-
Male	1	2	2	2	1	5	9	18	8	1	-	-
Female	-	-	2	2	5	9	17	20	20	4	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	13	12	17	29	50	70	143	176	167	63	10	-
Male	6	8	7	15	25	28	64	58	42	13	-	-
Female	7	4	10	14	25	42	79	118	125	50	10	-
White, both sexes	8	9	15	26	47	67	131	169	162	60	10	-
Male	4	7	5	15	24	27	58	55	41	13	-	-
Female	4	2	10	11	23	40	73	114	121	47	10	-
Black, both sexes	3	1	1	2	2	2	11	6	4	2	-	-
Male	1	-	1	-	1	1	6	3	1	-	-	-
Female	2	1	-	2	1	1	5	3	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	1	1	1	-	1	1	1	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	1	1	-	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
New Mexico												
All races, both sexes	3	3	4	4	8	6	20	27	16	6	-	-
Male	2	3	2	3	3	3	7	12	6	3	-	-
Female	1	-	2	1	5	3	13	15	10	3	-	-
White, both sexes	1	3	4	4	6	6	20	27	16	5	-	-
Male	1	3	2	3	3	3	7	12	6	2	-	-
Female	-	-	2	1	3	3	13	15	10	3	-	-
Black, both sexes	-	-	-	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	16	30	30	41	62	149	216	296	251	77	26	-
Male	11	18	14	24	32	70	99	102	65	14	-	-
Female	5	12	16	17	30	79	117	194	186	63	26	-
White, both sexes	9	24	25	33	56	137	202	287	248	74	26	-
Male	6	15	11	18	29	64	96	101	63	13	-	-
Female	3	9	14	15	27	73	106	186	185	61	26	-
Black, both sexes	6	6	4	6	6	12	9	9	2	2	-	-
Male	4	3	3	5	3	6	2	1	1	1	-	-
Female	2	3	1	1	3	6	7	8	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	4	-	1	1	-	-
Male	1	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	3	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
North Carolina												
All races, both sexes	12	14	18	31	47	54	104	157	89	26	7	-
Male	7	8	9	16	27	22	44	64	23	5	-	-
Female	5	6	9	15	20	32	60	93	66	21	7	-
White, both sexes	9	10	15	27	41	51	93	143	80	22	5	-
Male	5	6	7	13	23	20	41	59	23	4	-	-
Female	4	4	8	14	18	31	52	84	57	18	5	-
Black, both sexes	3	3	3	3	5	3	9	13	8	4	2	-
Male	2	1	2	2	3	2	1	5	-	1	-	-
Female	1	2	1	1	2	1	8	8	8	3	2	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	-	1	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	1	2	2	2	4	7	12	18	12	9	-	-
Male	-	-	2	-	3	4	4	6	2	2	-	-
Female	1	2	-	2	1	3	8	12	10	7	-	-
White, both sexes	-	2	2	2	4	7	12	18	12	9	-	-
Male	-	-	2	-	3	4	4	6	2	2	-	-
Female	-	2	-	2	1	3	8	12	10	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Ohio												
All races, both sexes	17	23	20	36	53	104	176	254	176	59	10	-
Male	8	13	10	17	21	49	71	80	37	7	-	-
Female	9	10	10	19	32	55	105	174	139	52	10	-
White, both sexes	15	19	18	34	47	101	168	247	168	58	10	-
Male	6	10	9	16	19	48	69	77	36	7	-	-
Female	9	9	9	18	28	53	99	170	132	51	10	-
Black, both sexes	2	4	2	2	5	2	7	7	8	-	-	-
Male	2	3	1	1	2	1	1	3	1	-	-	-
Female	-	1	1	1	3	1	6	4	7	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
Oklahoma												
All races, both sexes	9	9	11	10	15	21	48	47	31	13	5	-
Male	6	7	4	3	7	8	27	17	10	4	-	-
Female	3	2	7	7	8	13	21	30	21	9	5	-
White, both sexes	5	7	10	8	14	21	44	45	29	12	4	-
Male	4	5	3	3	6	8	26	17	9	3	-	-
Female	1	2	7	5	8	13	18	28	20	9	4	-
Black, both sexes	4	1	-	1	-	-	4	-	2	1	-	-
Male	2	1	-	-	-	-	1	-	1	1	-	-
Female	2	-	-	1	-	-	3	-	1	-	-	-
American Indian 1/, both sexes	-	1	1	1	1	-	-	2	-	-	1	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Oregon												
All races, both sexes	9	7	5	13	19	55	104	126	98	40	8	-
Male	7	5	4	7	11	18	44	52	24	11	1	-
Female	2	2	1	6	8	37	60	74	74	29	7	-
White, both sexes	8	7	5	13	18	51	100	124	97	40	8	-
Male	6	5	4	7	11	18	42	50	24	11	1	-
Female	2	2	1	6	7	33	58	74	73	29	7	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	1	3	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	23	36	51	55	71	139	284	371	316	107	21	-
Male	9	28	30	29	35	53	123	130	79	22	4	-
Female	14	8	21	26	36	86	161	241	237	85	17	-
White, both sexes	14	29	46	50	69	134	273	366	308	107	19	-
Male	3	22	26	27	35	51	120	128	77	22	4	-
Female	11	7	20	23	34	83	153	238	231	85	15	-
Black, both sexes	8	7	5	5	2	5	10	3	7	-	2	-
Male	5	6	4	2	-	2	2	2	2	-	-	-
Female	3	1	1	3	2	3	8	1	5	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Rhode Island												
All races, both sexes	1	2	4	4	6	6	15	18	22	6	3	-
Male	1	1	3	3	3	2	7	3	4	1	-	-
Female	-	1	1	1	3	4	8	15	18	5	3	-
White, both sexes	1	2	4	4	5	5	15	18	22	6	3	-
Male	1	1	3	3	3	1	7	3	4	1	-	-
Female	-	1	1	1	2	4	8	15	18	5	3	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	11	23	17	23	26	56	78	37	13	4	-
Male	2	9	12	7	10	11	31	22	11	2	1	-
Female	1	2	11	10	13	15	25	56	26	11	3	-
White, both sexes	2	7	20	10	18	22	47	73	34	13	2	-
Male	2	6	12	4	9	8	28	21	10	2	-	-
Female	-	1	8	6	9	14	19	52	24	11	2	-
Black, both sexes	-	4	3	7	5	4	9	5	3	-	2	-
Male	-	3	-	3	1	3	3	1	1	-	1	-
Female	-	1	3	4	4	1	6	4	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
South Dakota												
All races, both sexes	2	2	-	1	4	5	13	17	15	5	-	-
Male	1	2	-	-	3	1	3	7	4	1	-	-
Female	1	-	-	1	1	4	10	10	11	4	-	-
White, both sexes	2	2	-	1	3	5	13	17	15	5	-	-
Male	1	2	-	-	3	1	3	7	4	1	-	-
Female	1	-	-	1	-	4	10	10	11	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	11	12	13	26	36	41	69	85	54	16	3	-
Male	9	10	9	15	22	17	24	29	18	8	-	-
Female	2	2	4	11	14	24	45	56	36	8	3	-
White, both sexes	9	10	9	20	32	37	64	83	53	16	3	-
Male	7	8	7	12	19	16	21	29	18	8	-	-
Female	2	2	2	8	13	21	43	54	35	8	3	-
Black, both sexes	2	2	4	6	4	4	5	2	1	-	-	-
Male	2	2	2	3	3	1	3	-	-	-	-	-
Female	-	-	2	3	1	3	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Texas												
All races, both sexes	27	41	58	63	67	132	179	228	153	57	9	-
Male	14	27	25	32	33	54	68	78	46	7	-	-
Female	13	14	33	31	34	78	111	150	107	50	9	-
White, both sexes	21	33	53	55	59	119	168	219	148	56	8	-
Male	11	24	22	28	28	47	64	76	46	7	-	-
Female	10	9	31	27	31	72	104	143	102	49	8	-
Black, both sexes	6	7	1	6	7	11	11	9	5	1	1	-
Male	3	3	-	2	5	5	4	2	-	-	-	-
Female	3	4	1	4	2	6	7	7	5	1	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	4	2	1	2	-	-	-	-	-	-
Male	-	-	3	2	-	2	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	8	2	2	3	11	15	21	38	23	9	-	-
Male	7	-	1	3	4	6	12	11	13	3	-	-
Female	1	2	1	-	7	9	9	27	10	6	-	-
White, both sexes	8	2	2	3	10	15	21	38	23	9	-	-
Male	7	-	1	3	4	6	12	11	13	3	-	-
Female	1	2	1	-	6	9	9	27	10	6	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Vermont												
All races, both sexes	1	-	-	4	5	7	16	21	15	4	1	-
Male	-	-	-	4	3	3	6	7	4	1	-	-
Female	1	-	-	-	2	4	10	14	11	3	1	-
White, both sexes	1	-	-	4	5	7	16	21	15	4	1	-
Male	-	-	-	4	3	3	6	7	4	1	-	-
Female	1	-	-	-	2	4	10	14	11	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	9	17	21	25	49	79	106	86	24	-	-
Male	5	5	9	10	13	25	34	36	19	3	-	-
Female	6	4	8	11	12	24	45	70	67	21	-	-
White, both sexes	8	6	15	17	22	42	74	101	80	22	-	-
Male	4	3	8	8	11	21	31	32	18	3	-	-
Female	4	3	7	9	11	21	43	69	62	19	-	-
Black, both sexes	3	3	1	4	3	6	5	4	6	2	-	-
Male	1	2	-	2	2	3	3	3	1	-	-	-
Female	2	1	1	2	1	3	2	1	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Washington												
All races, both sexes	9	12	20	17	28	61	120	170	159	51	15	-
Male	4	6	9	10	14	29	47	64	52	17	1	-
Female	5	6	11	7	14	32	73	106	107	34	14	-
White, both sexes	7	10	17	15	28	58	118	163	156	50	15	-
Male	4	5	7	10	14	28	46	61	51	16	1	-
Female	3	5	10	5	14	30	72	102	105	34	14	-
Black, both sexes	2	1	3	-	-	1	-	-	-	1	-	-
Male	-	1	2	-	-	-	-	-	-	1	-	-
Female	2	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	3	2	-	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	1	-	1	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	2	4	1	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	1	3	1	-	-	-
West Virginia												
All races, both sexes	2	4	6	7	11	12	21	25	23	5	-	-
Male	1	2	4	7	5	5	9	7	2	1	-	-
Female	1	2	2	-	6	7	12	18	21	4	-	-
White, both sexes	2	4	6	7	11	12	21	25	21	5	-	-
Male	1	2	4	7	5	5	9	7	2	1	-	-
Female	1	2	2	-	6	7	12	18	19	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Wisconsin												
All races, both sexes	4	10	8	22	28	59	85	131	134	48	7	-
Male	2	6	6	13	10	26	33	54	41	8	1	-
Female	2	4	2	9	18	33	52	77	93	40	6	-
White, both sexes	4	10	7	21	28	57	85	129	134	47	7	-
Male	2	6	5	13	10	26	33	54	41	8	1	-
Female	2	4	2	8	18	31	52	75	93	39	6	-
Black, both sexes	-	-	1	1	-	2	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Wyoming												
All races, both sexes	-	-	3	3	-	8	3	8	12	2	-	-
Male	-	-	-	2	-	5	-	4	4	-	-	-
Female	-	-	3	1	-	3	3	4	8	2	-	-
White, both sexes	-	-	3	3	-	8	3	8	12	2	-	-
Male	-	-	-	2	-	5	-	4	4	-	-	-
Female	-	-	3	1	-	3	3	4	8	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
United States												
All races, both sexes	2,581	-	-	-	2	11	13	20	22	28	39	63
Male	925	-	-	-	1	5	5	7	7	11	15	34
Female	1,656	-	-	-	1	6	8	13	15	17	24	29
White, both sexes	2,369	-	-	-	2	8	12	17	16	23	30	51
Male	846	-	-	-	1	4	5	5	5	9	9	28
Female	1,523	-	-	-	1	4	7	12	11	14	21	23
Black, both sexes	149	-	-	-	-	3	1	1	5	4	8	11
Male	55	-	-	-	-	1	-	1	2	2	5	5
Female	94	-	-	-	-	2	1	-	3	2	3	6
American Indian 1/, both sexes	10	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	1	1	1	1	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	1	1	1	-	-
Alabama												
All races, both sexes	36	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	34	-	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	1	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Arkansas												
All races, both sexes	24	-	-	-	-	1	1	-	1	1	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	19	-	-	-	-	1	1	-	-	1	-	-
White, both sexes	20	-	-	-	-	1	1	-	1	1	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	1	1	-	-	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	247	-	-	-	1	-	1	2	3	4	5	3
Male	101	-	-	-	1	-	-	1	1	2	1	1
Female	146	-	-	-	-	-	1	1	2	2	4	2
White, both sexes	215	-	-	-	1	-	1	1	2	2	4	3
Male	87	-	-	-	1	-	-	-	1	1	1	1
Female	128	-	-	-	-	-	1	1	1	1	3	2
Black, both sexes	9	-	-	-	-	-	-	1	-	1	1	-
Male	5	-	-	-	-	-	-	1	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	1	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Colorado												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	38	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	40	-	-	-	-	-	-	-	-	1	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	35	-	-	-	-	-	-	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Florida												
All races, both sexes	200	-	-	-	-	-	-	-	-	3	5	4
Male	69	-	-	-	-	-	-	-	-	2	2	1
Female	131	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	188	-	-	-	-	-	-	-	-	3	3	4
Male	65	-	-	-	-	-	-	-	-	2	1	1
Female	123	-	-	-	-	-	-	-	-	1	2	3
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Hawaii												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Illinois												
All races, both sexes	120	-	-	-	-	-	-	2	2	-	1	6
Male	42	-	-	-	-	-	-	1	1	-	1	6
Female	78	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	111	-	-	-	-	-	-	2	2	-	-	5
Male	38	-	-	-	-	-	-	1	1	-	-	5
Female	73	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	62	-	-	-	-	-	-	-	2	1	1	1
Male	26	-	-	-	-	-	-	-	2	1	1	1
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	1	1	1	1
Male	24	-	-	-	-	-	-	-	1	1	1	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Iowa												
All races, both sexes	42	-	-	-	1	-	1	-	-	1	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	1	-	1	-	-	1	-	-
White, both sexes	42	-	-	-	1	-	1	-	-	1	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	1	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	24	-	-	-	-	-	-	1	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	23	-	-	-	-	-	-	1	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Kentucky												
All races, both sexes	42	-	-	-	-	2	2	1	1	1	-	2
Male	16	-	-	-	-	2	1	1	-	-	-	1
Female	26	-	-	-	-	-	1	-	1	1	-	1
White, both sexes	36	-	-	-	-	2	1	1	-	1	-	2
Male	14	-	-	-	-	2	1	1	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	5	-	-	-	-	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	25	-	-	-	-	-	-	-	-	3	-	1
Male	11	-	-	-	-	-	-	-	-	2	-	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	3	-	-
Male	8	-	-	-	-	-	-	-	-	2	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Maine												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	5
Male	12	-	-	-	-	-	-	-	-	-	-	3
Female	23	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	5
Male	12	-	-	-	-	-	-	-	-	-	-	3
Female	21	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Massachusetts												
All races, both sexes	62	-	-	-	-	-	-	1	-	2	3	2
Male	26	-	-	-	-	-	-	-	-	1	2	2
Female	36	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	61	-	-	-	-	-	-	1	-	2	3	2
Male	26	-	-	-	-	-	-	-	-	1	2	2
Female	35	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	90	-	-	-	-	-	-	-	-	2	1	4
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	58	-	-	-	-	-	-	-	-	2	1	2
White, both sexes	85	-	-	-	-	-	-	-	-	1	1	2
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	53	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Minnesota												
All races, both sexes	58	-	-	-	-	1	-	-	1	-	-	1
Male	28	-	-	-	-	-	-	-	1	-	-	-
Female	30	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	57	-	-	-	-	-	-	-	1	-	-	1
Male	28	-	-	-	-	-	-	-	1	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Missouri												
All races, both sexes	58	-	-	-	-	-	-	-	1	1	1	-
Male	18	-	-	-	-	-	-	-	1	-	1	-
Female	40	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	1	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	36	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Nebraska												
All races, both sexes	18	-	-	-	-	-	-	1	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	1	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	19	-	-	-	-	-	-	1	1	1	1	1
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	1	1	1
White, both sexes	18	-	-	-	-	-	-	1	1	1	1	1
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
New Hampshire												
All races, both sexes	17	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	95	-	-	-	-	-	-	-	1	-	2	1
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	61	-	-	-	-	-	-	-	1	-	2	-
White, both sexes	88	-	-	-	-	-	-	-	1	-	2	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	1	-	2	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
New Mexico												
All races, both sexes	15	-	-	-	-	-	-	1	1	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	13	-	-	-	-	-	-	1	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	131	-	-	-	-	2	-	1	2	1	1	4
Male	45	-	-	-	-	1	-	-	-	-	1	3
Female	86	-	-	-	-	1	-	1	2	1	-	1
White, both sexes	120	-	-	-	-	1	-	1	2	1	1	3
Male	43	-	-	-	-	1	-	-	-	-	1	3
Female	77	-	-	-	-	-	-	1	2	1	-	-
Black, both sexes	10	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
North Carolina												
All races, both sexes	60	-	-	-	-	-	1	2	1	1	-	1
Male	22	-	-	-	-	-	1	1	-	1	-	1
Female	38	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	52	-	-	-	-	-	1	2	-	-	-	1
Male	19	-	-	-	-	-	1	1	-	-	-	1
Female	33	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Ohio												
All races, both sexes	110	-	-	-	-	1	2	1	1	1	1	1
Male	31	-	-	-	-	1	1	-	-	1	-	-
Female	79	-	-	-	-	-	1	1	1	-	1	1
White, both sexes	103	-	-	-	-	-	2	1	1	1	-	-
Male	30	-	-	-	-	-	1	-	-	1	-	-
Female	73	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	7	-	-	-	-	1	-	-	-	-	1	1
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	30	-	-	-	-	1	-	1	-	-	-	1
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	22	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	26	-	-	-	-	1	-	1	-	-	-	1
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	19	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Oregon												
All races, both sexes	59	-	-	-	-	-	-	-	1	-	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	57	-	-	-	-	-	-	-	1	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	176	-	-	-	-	-	-	1	-	-	3	5
Male	67	-	-	-	-	-	-	-	-	-	1	2
Female	109	-	-	-	-	-	-	1	-	-	2	3
White, both sexes	166	-	-	-	-	-	-	1	-	-	2	3
Male	62	-	-	-	-	-	-	-	-	-	-	2
Female	104	-	-	-	-	-	-	1	-	-	2	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	2
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Rhode Island												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	37	-	-	-	-	-	-	-	-	1	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	37	-	-	-	-	1	-	-	-	1	1	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	1	-	-	-	1	1	-
White, both sexes	32	-	-	-	-	1	-	-	-	1	1	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	1	-	-	-	1	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Texas												
All races, both sexes	147	-	-	-	-	-	1	2	1	-	4	4
Male	48	-	-	-	-	-	-	-	-	-	1	1
Female	99	-	-	-	-	-	1	2	1	-	3	3
White, both sexes	137	-	-	-	-	-	1	2	-	-	3	4
Male	44	-	-	-	-	-	-	-	-	-	-	1
Female	93	-	-	-	-	-	1	2	-	-	3	3
Black, both sexes	8	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Vermont												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	55	-	-	-	-	1	1	-	1	-	1	3
Male	16	-	-	-	-	1	1	-	-	-	-	2
Female	39	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	52	-	-	-	-	1	1	-	1	-	1	3
Male	15	-	-	-	-	1	1	-	-	-	-	2
Female	37	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Washington												
All races, both sexes	77	-	-	-	-	-	1	-	-	1	1	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	1	-	-	1	1	-
White, both sexes	74	-	-	-	-	-	1	-	-	1	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	1	-	-	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Wisconsin												
All races, both sexes	60	-	-	-	-	-	1	-	-	-	-	1
Male	22	-	-	-	-	-	1	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	58	-	-	-	-	-	1	-	-	-	-	1
Male	21	-	-	-	-	-	1	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
United States												
All races, both sexes	70	79	96	146	194	290	461	554	373	101	19	-
Male	31	42	44	72	71	112	166	185	93	21	3	-
Female	39	37	52	74	123	178	295	369	280	80	16	-
White, both sexes	51	66	79	123	176	275	436	534	362	94	14	-
Male	24	35	35	61	62	109	161	183	89	19	2	-
Female	27	31	44	62	114	166	275	351	273	75	12	-
Black, both sexes	15	12	14	14	11	11	16	10	5	5	3	-
Male	5	6	7	7	5	2	4	-	1	2	-	-
Female	10	6	7	7	6	9	12	10	4	3	3	-
American Indian 1/, both sexes	1	-	-	4	1	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	3	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	3	5	6	3	9	10	5	2	2	-
Male	2	1	2	3	4	1	1	2	3	-	1	-
Female	1	-	1	2	2	2	8	8	2	2	1	-
Alabama												
All races, both sexes	2	-	1	1	3	10	8	8	2	-	-	-
Male	1	-	-	1	1	4	2	1	-	-	-	-
Female	1	-	1	-	2	6	6	7	2	-	-	-
White, both sexes	1	-	-	-	2	10	7	8	1	-	-	-
Male	1	-	-	-	-	4	2	1	-	-	-	-
Female	-	-	-	-	2	6	5	7	1	-	-	-
Black, both sexes	1	-	1	1	1	-	1	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Alaska												
All races, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	2	1	-	2	4	3	8	6	5	1	1	-
Male	2	-	-	1	-	2	2	4	3	-	-	-
Female	-	1	-	1	4	1	6	2	2	1	1	-
White, both sexes	2	1	-	2	4	3	8	6	5	1	1	-
Male	2	-	-	1	-	2	2	4	3	-	-	-
Female	-	1	-	1	4	1	6	2	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Arkansas												
All races, both sexes	1	-	2	4	4	2	2	4	1	-	-	-
Male	-	-	-	2	-	1	-	1	-	-	-	-
Female	1	-	2	2	4	1	2	3	1	-	-	-
White, both sexes	-	-	1	3	3	2	2	4	1	-	-	-
Male	-	-	-	2	-	1	-	1	-	-	-	-
Female	-	-	1	1	3	1	2	3	1	-	-	-
Black, both sexes	1	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	9	3	9	19	24	43	54	43	13	3	-
Male	2	7	-	8	9	12	19	20	10	6	1	-
Female	6	2	3	1	10	12	24	34	33	7	2	-
White, both sexes	7	7	3	8	15	24	38	49	38	10	2	-
Male	2	6	-	7	6	12	18	19	7	5	-	-
Female	5	1	3	1	9	12	20	30	31	5	2	-
Black, both sexes	1	2	-	-	1	-	1	-	-	1	-	-
Male	-	1	-	-	1	-	-	-	-	1	-	-
Female	1	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	-	4	5	5	2	1	-
Male	-	-	-	-	2	-	1	1	3	-	1	-
Female	-	-	-	-	1	-	3	4	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Colorado												
All races, both sexes	3	2	1	1	3	5	6	12	4	-	-	-
Male	-	1	-	-	-	2	1	8	3	-	-	-
Female	3	1	1	1	3	3	5	4	1	-	-	-
White, both sexes	3	2	1	1	3	4	6	12	4	-	-	-
Male	-	1	-	-	-	2	1	8	3	-	-	-
Female	3	1	1	1	3	2	5	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	2	3	2	1	3	7	7	4	6	1	-
Male	-	2	1	-	-	1	2	4	1	-	-	-
Female	-	-	2	2	1	2	5	3	3	6	1	-
White, both sexes	-	1	3	1	1	3	6	7	3	5	1	-
Male	-	1	1	-	-	1	1	4	1	-	-	-
Female	-	-	2	1	1	2	5	3	2	5	1	-
Black, both sexes	-	1	-	1	-	-	1	-	1	1	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Delaware												
All races, both sexes	-	2	1	-	2	-	1	3	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	2	1	-	1	-	1	3	-	-	-	-
White, both sexes	-	2	1	-	2	-	1	3	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	2	1	-	1	-	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	2	2	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Florida												
All races, both sexes	3	7	4	10	13	26	40	42	36	6	1	-
Male	2	1	1	5	6	9	15	15	9	1	-	-
Female	1	6	3	5	7	17	25	27	27	5	1	-
White, both sexes	2	7	4	9	13	22	40	40	35	6	-	-
Male	2	1	1	4	6	8	15	15	8	1	-	-
Female	-	6	3	5	7	14	25	25	27	5	-	-
Black, both sexes	1	-	-	1	-	4	-	2	1	-	1	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	1	-	-	-	-	3	-	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	2	2	6	2	9	9	2	4	1	-	-
Male	1	1	2	1	-	1	6	-	3	-	-	-
Female	1	1	-	5	2	8	3	2	1	1	-	-
White, both sexes	2	1	1	5	2	9	9	1	4	-	-	-
Male	1	1	1	1	-	1	6	-	3	-	-	-
Female	1	-	-	4	2	8	3	1	1	-	-	-
Black, both sexes	-	1	1	1	-	-	-	1	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Hawaii												
All races, both sexes	-	-	-	1	2	2	4	4	-	-	1	-
Male	-	-	-	1	2	1	-	2	-	-	-	-
Female	-	-	-	-	-	1	4	2	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	2	4	3	-	-	1	-
Male	-	-	-	1	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	4	2	-	-	1	-
Idaho												
All races, both sexes	1	-	-	1	1	-	1	3	4	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	1	-	-	1	1	-	-	3	3	-	-	-
White, both sexes	1	-	-	1	1	-	1	3	4	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	1	-	-	1	1	-	-	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Illinois												
All races, both sexes	2	1	2	8	7	12	17	27	24	7	2	-
Male	2	-	2	3	5	2	4	11	2	2	-	-
Female	-	1	-	5	2	10	13	16	22	5	2	-
White, both sexes	1	1	2	5	7	10	17	27	24	7	1	-
Male	1	-	2	2	5	2	4	11	2	2	-	-
Female	-	1	-	3	2	8	13	16	22	5	1	-
Black, both sexes	1	-	-	2	-	2	-	-	-	-	1	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	2	2	3	8	10	12	11	3	4	-	-
Male	-	1	2	1	5	4	4	2	1	1	-	-
Female	2	1	-	2	3	6	8	9	2	3	-	-
White, both sexes	1	2	1	3	8	10	11	11	3	4	-	-
Male	-	1	1	1	5	4	4	2	1	1	-	-
Female	1	1	-	2	3	6	7	9	2	3	-	-
Black, both sexes	1	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Iowa												
All races, both sexes	-	2	2	2	6	7	3	12	3	1	-	-
Male	-	2	-	2	2	1	3	6	1	-	-	-
Female	-	-	2	-	4	6	-	6	2	1	-	-
White, both sexes	-	2	2	2	6	7	3	12	3	1	-	-
Male	-	2	-	2	2	1	3	6	1	-	-	-
Female	-	-	2	-	4	6	-	6	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	1	1	3	3	6	6	1	1	-
Male	-	-	-	-	-	2	3	1	1	-	1	-
Female	-	-	-	1	1	1	-	5	5	1	-	-
White, both sexes	-	-	-	1	1	3	3	6	6	-	1	-
Male	-	-	-	-	-	2	3	1	1	-	1	-
Female	-	-	-	1	1	1	-	5	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Kentucky												
All races, both sexes	-	2	3	3	2	3	8	6	4	2	-	-
Male	-	2	2	1	1	-	2	2	1	-	-	-
Female	-	-	1	2	1	3	6	4	3	2	-	-
White, both sexes	-	2	-	3	2	3	7	6	4	2	-	-
Male	-	2	-	1	1	-	2	2	1	-	-	-
Female	-	-	-	2	1	3	5	4	3	2	-	-
Black, both sexes	-	-	2	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	3	1	3	5	6	1	1	-	-
Male	1	-	-	2	1	1	1	-	1	1	-	-
Female	-	-	-	1	-	2	4	6	-	-	-	-
White, both sexes	1	-	-	2	1	3	5	6	1	-	-	-
Male	1	-	-	1	1	1	1	-	1	-	-	-
Female	-	-	-	1	-	2	4	6	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Maine												
All races, both sexes	-	2	-	2	3	3	2	4	2	2	-	-
Male	-	2	-	1	-	2	1	1	-	-	-	-
Female	-	-	-	1	3	1	1	3	2	2	-	-
White, both sexes	-	2	-	2	3	3	2	4	2	2	-	-
Male	-	2	-	1	-	2	1	1	-	-	-	-
Female	-	-	-	1	3	1	1	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	1	3	2	-	4	6	8	3	-	-	-
Male	1	-	-	-	-	1	3	2	2	-	-	-
Female	2	1	3	2	-	3	3	6	1	-	-	-
White, both sexes	3	1	2	1	-	4	6	8	3	-	-	-
Male	1	-	-	-	-	1	3	2	2	-	-	-
Female	2	1	2	1	-	3	3	6	1	-	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Massachusetts												
All races, both sexes	1	4	1	2	4	2	9	12	16	3	-	-
Male	1	2	1	-	2	1	4	4	6	-	-	-
Female	-	2	-	2	2	1	5	8	10	3	-	-
White, both sexes	1	4	1	2	4	2	9	11	16	3	-	-
Male	1	2	1	-	2	1	4	4	6	-	-	-
Female	-	2	-	2	2	1	5	7	10	3	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	4	5	2	8	10	18	17	17	1	-	-
Male	1	2	3	-	5	5	7	3	4	-	-	-
Female	-	2	2	2	3	5	11	14	13	1	-	-
White, both sexes	1	4	5	2	7	9	18	17	17	1	-	-
Male	1	2	3	-	5	5	7	3	4	-	-	-
Female	-	2	2	2	2	4	11	14	13	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Minnesota												
All races, both sexes	1	3	2	2	4	8	12	14	8	1	-	-
Male	1	2	2	2	1	6	5	4	4	-	-	-
Female	-	1	-	-	3	2	7	10	4	1	-	-
White, both sexes	1	3	2	2	4	8	12	14	8	1	-	-
Male	1	2	2	2	1	6	5	4	4	-	-	-
Female	-	1	-	-	3	2	7	10	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	2	3	-	2	3	2	-	-	-
Male	-	-	-	2	-	-	-	3	-	-	-	-
Female	-	-	1	-	3	-	2	-	2	-	-	-
White, both sexes	-	-	1	2	3	-	2	3	2	-	-	-
Male	-	-	-	2	-	-	-	3	-	-	-	-
Female	-	-	1	-	3	-	2	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Missouri												
All races, both sexes	1	3	1	2	3	2	12	12	16	3	-	-
Male	1	2	-	2	1	1	5	2	2	-	-	-
Female	-	1	1	-	2	1	7	10	14	3	-	-
White, both sexes	1	2	-	1	1	2	11	12	16	3	-	-
Male	1	1	-	1	1	1	5	2	2	-	-	-
Female	-	1	-	-	-	1	6	10	14	3	-	-
Black, both sexes	-	1	1	1	2	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	1	-	3	3	1	-	-
Male	-	-	-	-	-	1	-	-	2	1	-	-
Female	-	-	-	-	1	-	-	3	1	-	-	-
White, both sexes	-	-	-	-	1	1	-	3	3	1	-	-
Male	-	-	-	-	-	1	-	-	2	1	-	-
Female	-	-	-	-	1	-	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Nebraska												
All races, both sexes	-	-	-	1	-	2	5	4	4	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	4	4	4	-	-	-
White, both sexes	-	-	-	1	-	2	5	3	4	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	4	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	2	3	2	1	3	1	2	-	-	-
Male	-	-	1	-	-	-	2	-	2	-	-	-
Female	-	-	1	3	2	1	1	1	-	-	-	-
White, both sexes	-	-	2	2	2	1	3	1	2	-	-	-
Male	-	-	1	-	-	-	2	-	2	-	-	-
Female	-	-	1	2	2	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
New Hampshire												
All races, both sexes	-	-	1	-	2	2	1	6	2	1	1	-
Male	-	-	1	-	-	1	-	5	1	-	-	-
Female	-	-	-	-	2	1	1	1	1	1	1	-
White, both sexes	-	-	1	-	2	2	1	6	2	1	1	-
Male	-	-	1	-	-	1	-	5	1	-	-	-
Female	-	-	-	-	2	1	1	1	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	3	3	8	11	14	16	12	18	3	1	-
Male	1	2	1	4	4	4	9	6	2	-	-	-
Female	1	1	2	4	7	10	7	6	16	3	1	-
White, both sexes	1	2	3	7	11	13	14	12	18	3	1	-
Male	-	1	1	4	4	4	8	6	2	-	-	-
Female	1	1	2	3	7	9	6	6	16	3	1	-
Black, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
New Mexico												
All races, both sexes	1	-	-	-	1	1	1	5	-	3	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	1	-	-	-	-	-	1	5	-	2	-	-
White, both sexes	-	-	-	-	1	1	1	5	-	3	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	5	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	2	4	10	6	18	22	28	19	7	-	-
Male	1	1	1	6	2	9	7	9	3	1	-	-
Female	3	1	3	4	4	9	15	19	16	6	-	-
White, both sexes	3	2	4	8	4	18	19	27	19	7	-	-
Male	1	1	1	5	1	9	7	9	3	1	-	-
Female	2	1	3	3	3	9	12	18	16	6	-	-
Black, both sexes	1	-	-	1	2	-	3	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
North Carolina												
All races, both sexes	1	4	2	6	7	5	11	8	8	2	-	-
Male	1	1	1	1	4	2	4	3	1	-	-	-
Female	-	3	1	5	3	3	7	5	7	2	-	-
White, both sexes	-	2	2	5	6	5	11	8	7	2	-	-
Male	-	-	1	1	4	2	4	3	1	-	-	-
Female	-	2	1	4	2	3	7	5	6	2	-	-
Black, both sexes	1	2	-	1	1	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	3	1	2	3	2	1	-	-
Male	-	-	-	-	2	1	-	1	-	-	-	-
Female	1	-	-	-	1	-	2	2	2	1	-	-
White, both sexes	-	-	-	-	3	1	2	3	2	1	-	-
Male	-	-	-	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	2	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Ohio												
All races, both sexes	1	1	3	5	11	15	22	27	12	5	-	-
Male	-	1	-	3	3	6	7	6	2	-	-	-
Female	1	-	3	2	8	9	15	21	10	5	-	-
White, both sexes	1	1	3	5	10	15	21	25	12	5	-	-
Male	-	1	-	3	3	6	7	6	2	-	-	-
Female	1	-	3	2	7	9	14	19	10	5	-	-
Black, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	-	2	2	2	3	7	4	2	1	-	-
Male	3	-	-	-	-	2	1	-	1	-	-	-
Female	1	-	2	2	2	1	6	4	1	1	-	-
White, both sexes	2	-	2	1	2	3	6	4	2	1	-	-
Male	2	-	-	-	-	2	1	-	1	-	-	-
Female	-	-	2	1	2	1	5	4	1	1	-	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Oregon												
All races, both sexes	2	2	-	-	3	5	11	18	9	5	1	-
Male	2	1	-	-	-	3	5	8	1	2	-	-
Female	-	1	-	-	3	2	6	10	8	3	1	-
White, both sexes	2	2	-	-	3	4	11	18	9	5	1	-
Male	2	1	-	-	-	3	5	8	1	2	-	-
Female	-	1	-	-	3	1	6	10	8	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	5	11	8	12	19	32	41	22	7	5	-
Male	1	5	8	6	4	7	8	17	5	2	1	-
Female	4	-	3	2	8	12	24	24	17	5	4	-
White, both sexes	3	4	9	8	12	18	32	41	22	7	4	-
Male	-	4	6	6	4	7	8	17	5	2	1	-
Female	3	-	3	2	8	11	24	24	17	5	3	-
Black, both sexes	1	1	2	-	-	1	-	-	-	-	1	-
Male	-	1	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Rhode Island												
All races, both sexes	-	1	2	2	1	1	3	3	-	1	-	-
Male	-	-	2	1	1	-	1	-	-	1	-	-
Female	-	1	-	1	-	1	2	3	-	-	-	-
White, both sexes	-	1	2	2	1	1	3	3	-	1	-	-
Male	-	-	2	1	1	-	1	-	-	1	-	-
Female	-	1	-	1	-	1	2	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	1	3	2	9	9	6	3	1	-	-
Male	-	-	1	2	-	3	5	2	1	-	-	-
Female	-	1	-	1	2	6	4	4	2	1	-	-
White, both sexes	-	-	1	1	2	7	7	5	3	1	-	-
Male	-	-	1	-	-	2	3	2	1	-	-	-
Female	-	-	-	1	2	5	4	3	2	1	-	-
Black, both sexes	-	1	-	2	-	2	2	1	-	-	-	-
Male	-	-	-	2	-	1	2	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
South Dakota												
All races, both sexes	1	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	1	3	3	4	3	8	8	3	-	-	-
Male	1	1	1	1	2	2	2	4	-	-	-	-
Female	-	-	2	2	2	1	6	4	3	-	-	-
White, both sexes	-	1	1	3	3	3	8	7	3	-	-	-
Male	-	1	1	1	1	2	2	4	-	-	-	-
Female	-	-	-	2	2	1	6	3	3	-	-	-
Black, both sexes	1	-	2	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Texas												
All races, both sexes	7	5	11	12	11	16	28	29	13	3	-	-
Male	3	2	4	7	3	3	7	11	6	-	-	-
Female	4	3	7	5	8	13	21	18	7	3	-	-
White, both sexes	5	3	10	11	10	16	27	29	13	3	-	-
Male	2	2	4	6	2	3	7	11	6	-	-	-
Female	3	1	6	5	8	13	20	18	7	3	-	-
Black, both sexes	2	2	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	2	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	1	-	-	2	2	1	4	1	1	-	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	2	2	-	4	-	1	-	-
White, both sexes	1	1	-	-	2	2	1	4	1	1	-	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	2	2	-	4	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Vermont												
All races, both sexes	-	-	-	-	-	-	2	5	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	5	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	5	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	3	3	-	5	9	15	11	-	-	-
Male	-	-	2	1	-	2	4	3	-	-	-	-
Female	2	-	1	2	-	3	5	12	11	-	-	-
White, both sexes	1	-	2	3	-	5	9	15	10	-	-	-
Male	-	-	1	1	-	2	4	3	-	-	-	-
Female	1	-	1	2	-	3	5	12	10	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Washington												
All races, both sexes	3	3	3	5	2	4	16	21	13	3	1	-
Male	1	1	1	2	-	1	5	7	4	1	-	-
Female	2	2	2	3	2	3	11	14	9	2	1	-
White, both sexes	3	2	3	4	2	4	16	20	13	3	1	-
Male	1	-	1	2	-	1	5	7	4	1	-	-
Female	2	2	2	2	2	3	11	13	9	2	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	2	-	2	4	3	2	-	-	-
Male	-	-	1	2	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	2	3	3	1	-	-	-
White, both sexes	-	-	1	2	-	2	4	3	1	-	-	-
Male	-	-	1	2	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	2	3	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Wisconsin												
All races, both sexes	-	1	4	2	5	7	7	15	15	2	-	-
Male	-	-	2	1	2	2	4	5	5	-	-	-
Female	-	1	2	1	3	5	3	10	10	2	-	-
White, both sexes	-	1	3	2	5	7	7	14	15	2	-	-
Male	-	-	1	1	2	2	4	5	5	-	-	-
Female	-	1	2	1	3	5	3	9	10	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
United States												
All races, both sexes	13,752	-	-	2	1	5	8	12	27	30	67	93
Male	5,468	-	-	1	1	1	4	9	20	21	46	60
Female	8,284	-	-	1	-	4	4	3	7	9	21	33
White, both sexes	12,942	-	-	2	1	5	7	8	19	18	59	71
Male	5,122	-	-	1	1	1	3	7	16	13	43	45
Female	7,820	-	-	1	-	4	4	1	3	5	16	26
Black, both sexes	589	-	-	-	-	-	-	3	5	11	8	18
Male	243	-	-	-	-	-	-	1	2	7	3	12
Female	346	-	-	-	-	-	-	2	3	4	5	6
American Indian 1/, both sexes	38	-	-	-	-	-	-	1	-	1	-	3
Male	23	-	-	-	-	-	-	1	-	1	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	183	-	-	-	-	-	1	-	3	-	-	1
Male	80	-	-	-	-	-	1	-	2	-	-	-
Female	103	-	-	-	-	-	-	-	1	-	-	1
Alabama												
All races, both sexes	169	-	-	-	-	-	-	-	1	-	1	-
Male	67	-	-	-	-	-	-	-	-	-	1	-
Female	102	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	144	-	-	-	-	-	-	-	-	-	1	-
Male	59	-	-	-	-	-	-	-	-	-	1	-
Female	85	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Alaska												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	190	-	-	-	-	-	-	1	4	-	-	-
Male	92	-	-	-	-	-	-	1	3	-	-	-
Female	98	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	186	-	-	-	-	-	-	1	3	-	-	-
Male	90	-	-	-	-	-	-	1	2	-	-	-
Female	96	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Arkansas												
All races, both sexes	111	-	-	-	-	-	1	-	-	-	-	2
Male	46	-	-	-	-	-	-	-	-	-	-	1
Female	65	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	108	-	-	-	-	-	1	-	-	-	-	2
Male	45	-	-	-	-	-	-	-	-	-	-	1
Female	63	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,587	-	-	-	-	-	-	3	3	3	2	6
Male	679	-	-	-	-	-	-	2	2	3	2	3
Female	908	-	-	-	-	-	-	1	1	-	-	3
White, both sexes	1,442	-	-	-	-	-	-	2	2	3	2	6
Male	617	-	-	-	-	-	-	2	2	3	2	3
Female	825	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	58	-	-	-	-	-	-	1	1	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	80	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Colorado												
All races, both sexes	182	-	-	-	-	-	-	-	1	-	1	3
Male	76	-	-	-	-	-	-	-	1	-	-	3
Female	106	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	181	-	-	-	-	-	-	-	1	-	1	3
Male	76	-	-	-	-	-	-	-	1	-	-	3
Female	105	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	212	-	-	-	-	-	-	-	-	-	-	1
Male	88	-	-	-	-	-	-	-	-	-	-	1
Female	124	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	209	-	-	-	-	-	-	-	-	-	-	1
Male	86	-	-	-	-	-	-	-	-	-	-	1
Female	123	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Delaware												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Florida												
All races, both sexes	918	-	-	-	-	-	-	-	2	1	8	5
Male	409	-	-	-	-	-	-	-	2	1	3	3
Female	509	-	-	-	-	-	-	-	-	-	5	2
White, both sexes	869	-	-	-	-	-	-	-	1	-	6	3
Male	387	-	-	-	-	-	-	-	1	-	3	3
Female	482	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	44	-	-	-	-	-	-	-	1	1	2	2
Male	20	-	-	-	-	-	-	-	1	1	-	-
Female	24	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	259	-	-	-	-	-	-	1	-	-	3	-
Male	116	-	-	-	-	-	-	1	-	-	2	-
Female	143	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	228	-	-	-	-	-	-	-	-	-	3	-
Male	97	-	-	-	-	-	-	-	-	-	2	-
Female	131	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	30	-	-	-	-	-	-	1	-	-	-	-
Male	18	-	-	-	-	-	-	1	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Hawaii												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	1	-
Male	31	-	-	-	-	-	-	-	-	-	1	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	-	-	-	1	-
Male	31	-	-	-	-	-	-	-	-	-	1	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Illinois												
All races, both sexes	519	-	-	-	-	-	-	-	-	3	4	3
Male	205	-	-	-	-	-	-	-	-	3	2	3
Female	314	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	485	-	-	-	-	-	-	-	-	1	1	3
Male	193	-	-	-	-	-	-	-	-	1	1	3
Female	292	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	2	3	-
Male	11	-	-	-	-	-	-	-	-	2	1	-
Female	20	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	293	-	-	-	-	-	-	-	2	1	3	3
Male	95	-	-	-	-	-	-	-	-	1	2	3
Female	198	-	-	-	-	-	-	-	2	-	1	-
White, both sexes	280	-	-	-	-	-	-	-	2	1	3	2
Male	91	-	-	-	-	-	-	-	-	1	2	2
Female	189	-	-	-	-	-	-	-	2	-	1	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Iowa												
All races, both sexes	208	-	-	-	-	-	-	-	-	-	-	-
Male	73	-	-	-	-	-	-	-	-	-	-	-
Female	135	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	207	-	-	-	-	-	-	-	-	-	-	-
Male	73	-	-	-	-	-	-	-	-	-	-	-
Female	134	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	156	-	-	-	-	-	-	-	-	1	2	1
Male	59	-	-	-	-	-	-	-	-	1	1	1
Female	97	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	150	-	-	-	-	-	-	-	-	1	2	-
Male	56	-	-	-	-	-	-	-	-	1	1	-
Female	94	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Kentucky												
All races, both sexes	124	-	-	-	-	1	1	-	-	-	1	3
Male	52	-	-	-	-	-	-	-	-	-	1	2
Female	72	-	-	-	-	1	1	-	-	-	-	1
White, both sexes	119	-	-	-	-	1	1	-	-	-	1	1
Male	51	-	-	-	-	-	-	-	-	-	1	1
Female	68	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Louisiana												
All races, both sexes	157	-	-	-	-	-	-	-	-	1	-	4
Male	77	-	-	-	-	-	-	-	-	1	-	4
Female	80	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	139	-	-	-	-	-	-	-	-	-	-	1
Male	64	-	-	-	-	-	-	-	-	-	-	1
Female	75	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	1	-	3
Male	12	-	-	-	-	-	-	-	-	1	-	3
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Maine												
All races, both sexes	106	-	-	-	-	-	-	-	-	-	-	-
Male	48	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	106	-	-	-	-	-	-	-	-	-	-	-
Male	48	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	234	-	-	-	-	-	-	-	1	-	3	3
Male	83	-	-	-	-	-	-	-	1	-	1	2
Female	151	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	195	-	-	-	-	-	-	-	-	-	1	2
Male	65	-	-	-	-	-	-	-	-	-	-	2
Female	130	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	35	-	-	-	-	-	-	-	-	-	2	1
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Massachusetts												
All races, both sexes	383	-	-	-	-	-	-	1	-	-	-	2
Male	128	-	-	-	-	-	-	-	-	-	-	2
Female	255	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	372	-	-	-	-	-	-	-	-	-	-	2
Male	126	-	-	-	-	-	-	-	-	-	-	2
Female	246	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	410	-	-	-	-	-	1	-	1	-	3	4
Male	164	-	-	-	-	-	-	-	1	-	3	3
Female	246	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	387	-	-	-	-	-	1	-	1	-	3	2
Male	154	-	-	-	-	-	-	-	1	-	3	1
Female	233	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Minnesota												
All races, both sexes	284	-	-	-	-	-	-	1	-	-	1	1
Male	109	-	-	-	-	-	-	-	-	-	1	-
Female	175	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	280	-	-	-	-	-	-	1	-	-	1	1
Male	107	-	-	-	-	-	-	-	-	-	1	-
Female	173	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	73	-	-	-	-	-	-	-	-	3	1	1
Male	32	-	-	-	-	-	-	-	-	1	1	1
Female	41	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	57	-	-	-	-	-	-	-	-	3	1	-
Male	23	-	-	-	-	-	-	-	-	1	1	-
Female	34	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Missouri												
All races, both sexes	282	-	-	-	-	-	-	-	-	1	4	2
Male	107	-	-	-	-	-	-	-	-	1	4	1
Female	175	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	271	-	-	-	-	-	-	-	-	1	4	2
Male	104	-	-	-	-	-	-	-	-	1	4	1
Female	167	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	55	-	-	-	1	-	-	-	-	-	-	-
Male	28	-	-	-	1	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	1	-	-	-	-	-	-	-
Male	28	-	-	-	1	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Nebraska												
All races, both sexes	104	-	-	-	-	-	-	-	1	-	-	-
Male	40	-	-	-	-	-	-	-	1	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	103	-	-	-	-	-	-	-	1	-	-	-
Male	39	-	-	-	-	-	-	-	1	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	78	-	-	-	-	-	-	1	-	-	1	-
Male	32	-	-	-	-	-	-	1	-	-	1	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	1	-	-	1	-
Male	29	-	-	-	-	-	-	1	-	-	1	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
New Hampshire												
All races, both sexes	99	-	-	-	-	-	-	-	-	-	-	-
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	99	-	-	-	-	-	-	-	-	-	-	-
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	502	-	-	-	-	-	1	1	-	2	-	4
Male	182	-	-	-	-	-	1	1	-	2	-	2
Female	320	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	474	-	-	-	-	-	1	1	-	1	-	2
Male	169	-	-	-	-	-	1	1	-	1	-	1
Female	305	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	23	-	-	-	-	-	-	-	-	1	-	2
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
New Mexico												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	760	-	-	-	-	1	-	-	2	-	-	5
Male	290	-	-	-	-	-	-	-	2	-	-	4
Female	470	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	734	-	-	-	-	1	-	-	1	-	-	5
Male	276	-	-	-	-	-	-	-	1	-	-	4
Female	458	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	19	-	-	-	-	-	-	-	1	-	-	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
North Carolina												
All races, both sexes	388	-	-	1	-	1	-	-	1	1	1	1
Male	160	-	-	-	-	1	-	-	-	-	1	1
Female	228	-	-	1	-	-	-	-	1	1	-	-
White, both sexes	345	-	-	1	-	1	-	-	-	-	1	-
Male	144	-	-	-	-	1	-	-	-	-	1	-
Female	201	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	41	-	-	-	-	-	-	-	1	1	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Ohio												
All races, both sexes	611	-	-	-	-	-	-	-	2	3	3	4
Male	211	-	-	-	-	-	-	-	2	1	2	1
Female	400	-	-	-	-	-	-	-	-	2	1	3
White, both sexes	585	-	-	-	-	-	-	-	2	1	3	2
Male	203	-	-	-	-	-	-	-	2	-	2	1
Female	382	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	25	-	-	-	-	-	-	-	-	2	-	2
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	137	-	-	-	-	-	-	1	-	4	2	2
Male	61	-	-	-	-	-	-	1	-	2	1	1
Female	76	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	126	-	-	-	-	-	-	-	-	2	2	2
Male	56	-	-	-	-	-	-	-	-	1	1	1
Female	70	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	-	1	-	-
Male	2	-	-	-	-	-	-	1	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Oregon												
All races, both sexes	350	-	-	-	-	-	-	-	-	1	1	-
Male	129	-	-	-	-	-	-	-	-	1	1	-
Female	221	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	342	-	-	-	-	-	-	-	-	1	1	-
Male	126	-	-	-	-	-	-	-	-	1	1	-
Female	216	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	976	-	-	-	-	-	1	-	-	2	4	9
Male	343	-	-	-	-	-	-	-	-	2	3	5
Female	633	-	-	-	-	-	1	-	-	-	1	4
White, both sexes	947	-	-	-	-	-	1	-	-	1	4	8
Male	329	-	-	-	-	-	-	-	-	1	3	4
Female	618	-	-	-	-	-	1	-	-	-	1	4
Black, both sexes	26	-	-	-	-	-	-	-	-	1	-	1
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Rhode Island												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	178	-	-	-	-	-	-	-	-	-	2	2
Male	77	-	-	-	-	-	-	-	-	-	2	-
Female	101	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	157	-	-	-	-	-	-	-	-	-	2	2
Male	70	-	-	-	-	-	-	-	-	-	2	-
Female	87	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
South Dakota												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	248	-	-	-	-	2	-	-	1	1	5	4
Male	104	-	-	-	-	-	-	-	1	-	2	3
Female	144	-	-	-	-	2	-	-	-	1	3	1
White, both sexes	233	-	-	-	-	2	-	-	1	-	5	3
Male	97	-	-	-	-	-	-	-	1	-	2	2
Female	136	-	-	-	-	2	-	-	-	-	3	1
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Texas												
All races, both sexes	655	-	-	1	-	-	2	-	-	1	5	9
Male	255	-	-	1	-	-	2	-	-	1	4	5
Female	400	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	614	-	-	1	-	-	2	-	-	1	4	8
Male	236	-	-	1	-	-	2	-	-	1	3	4
Female	378	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	36	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	-	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Vermont												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	286	-	-	-	-	-	-	-	1	-	1	-
Male	114	-	-	-	-	-	-	-	1	-	-	-
Female	172	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	261	-	-	-	-	-	-	-	1	-	1	-
Male	104	-	-	-	-	-	-	-	1	-	-	-
Female	157	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Washington												
All races, both sexes	476	-	-	-	-	-	1	1	1	-	3	2
Male	191	-	-	-	-	-	1	1	1	-	3	1
Female	285	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	460	-	-	-	-	-	-	1	1	-	3	2
Male	182	-	-	-	-	-	-	1	1	-	3	1
Female	278	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	80	-	-	-	-	-	-	1	1	-	-	3
Male	35	-	-	-	-	-	-	1	1	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	79	-	-	-	-	-	-	1	1	-	-	3
Male	35	-	-	-	-	-	-	1	1	-	-	2
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Wisconsin												
All races, both sexes	362	-	-	-	-	-	-	-	2	-	1	2
Male	148	-	-	-	-	-	-	-	1	-	1	1
Female	214	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	360	-	-	-	-	-	-	-	1	-	1	2
Male	148	-	-	-	-	-	-	-	1	-	1	1
Female	212	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
United States												
All races, both sexes	140	206	287	439	717	1,331	2,470	3,570	3,016	1,130	201	-
Male	93	142	159	246	400	621	1,113	1,388	881	241	21	-
Female	47	64	128	193	317	710	1,357	2,182	2,135	889	180	-
White, both sexes	111	181	260	394	649	1,238	2,323	3,414	2,909	1,079	194	-
Male	72	125	140	227	364	579	1,056	1,334	852	223	20	-
Female	39	56	120	167	285	659	1,267	2,080	2,057	856	174	-
Black, both sexes	23	18	19	36	51	65	116	99	77	34	6	-
Male	17	13	13	15	28	31	44	29	16	11	1	-
Female	6	5	6	21	23	34	72	70	61	23	5	-
American Indian 1/, both sexes	1	5	1	2	1	1	7	12	3	-	-	-
Male	1	3	1	1	1	-	4	6	1	-	-	-
Female	-	2	-	1	-	1	3	6	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	2	7	7	16	27	24	45	27	17	1	-
Male	3	1	5	3	7	11	9	19	12	7	-	-
Female	2	1	2	4	9	16	15	26	15	10	1	-
Alabama												
All races, both sexes	4	2	9	1	15	21	34	44	26	9	2	-
Male	3	1	6	1	11	8	13	15	6	2	-	-
Female	1	1	3	-	4	13	21	29	20	7	2	-
White, both sexes	3	1	5	1	14	15	30	40	25	8	1	-
Male	3	-	3	1	10	8	13	14	5	1	-	-
Female	-	1	2	-	4	7	17	26	20	7	1	-
Black, both sexes	1	1	4	-	1	6	4	4	1	1	1	-
Male	-	1	3	-	1	-	-	1	1	1	-	-
Female	1	-	1	-	-	6	4	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Alaska												
All races, both sexes	-	1	-	1	4	2	3	7	5	1	-	-
Male	-	-	-	1	4	1	1	3	2	-	-	-
Female	-	1	-	-	-	1	2	4	3	1	-	-
White, both sexes	-	-	-	1	4	1	3	3	5	1	-	-
Male	-	-	-	1	4	1	1	1	2	-	-	-
Female	-	-	-	-	-	-	2	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	4	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Arizona												
All races, both sexes	1	2	8	7	13	18	30	57	38	10	1	-
Male	1	1	4	2	6	10	17	29	14	3	1	-
Female	-	1	4	5	7	8	13	28	24	7	-	-
White, both sexes	1	2	7	7	12	18	30	57	37	10	1	-
Male	1	1	3	2	6	10	17	29	14	3	1	-
Female	-	1	4	5	6	8	13	28	23	7	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Arkansas												
All races, both sexes	1	2	2	2	9	11	25	22	22	9	3	-
Male	-	1	2	1	2	4	16	8	9	1	1	-
Female	1	1	-	1	7	7	9	14	13	8	2	-
White, both sexes	1	2	2	2	9	10	24	21	22	9	3	-
Male	-	1	2	1	2	4	15	8	9	1	1	-
Female	1	1	-	1	7	6	9	13	13	8	2	-
Black, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	24	14	28	35	85	158	290	432	346	139	19	-
Male	13	9	15	22	50	91	140	181	114	29	3	-
Female	11	5	13	13	35	67	150	251	232	110	16	-
White, both sexes	18	12	19	30	68	140	265	397	328	131	19	-
Male	9	7	9	19	39	84	132	170	107	26	3	-
Female	9	5	10	11	29	56	133	227	221	105	16	-
Black, both sexes	2	-	4	2	8	8	13	11	5	3	-	-
Male	2	-	3	1	6	2	5	1	-	-	-	-
Female	-	-	1	1	2	6	8	10	5	3	-	-
American Indian 1/, both sexes	1	1	1	-	1	-	1	2	-	-	-	-
Male	1	1	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	4	3	8	10	11	22	13	5	-	-
Male	1	1	2	2	4	5	2	9	7	3	-	-
Female	2	-	2	1	4	5	9	13	6	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Colorado												
All races, both sexes	-	4	8	7	20	17	25	47	30	17	2	-
Male	-	4	4	6	9	7	11	26	5	-	-	-
Female	-	-	4	1	11	10	14	21	25	17	2	-
White, both sexes	-	4	8	7	20	17	24	47	30	17	2	-
Male	-	4	4	6	9	7	11	26	5	-	-	-
Female	-	-	4	1	11	10	13	21	25	17	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	2	2	3	3	11	18	33	61	55	18	5	-
Male	2	1	1	1	8	7	19	25	18	4	1	-
Female	-	1	2	2	3	11	14	36	37	14	4	-
White, both sexes	1	2	3	3	10	18	33	60	55	18	5	-
Male	1	1	1	1	7	7	19	25	18	4	1	-
Female	-	1	2	2	3	11	14	35	37	14	4	-
Black, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Delaware												
All races, both sexes	2	1	3	6	-	3	6	8	6	4	4	-
Male	2	1	1	3	-	1	1	3	2	1	-	-
Female	-	-	2	3	-	2	5	5	4	3	4	-
White, both sexes	1	1	3	5	-	3	6	8	5	4	4	-
Male	1	1	1	3	-	1	1	3	2	1	-	-
Female	-	-	2	2	-	2	5	5	3	3	4	-
Black, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	-	4	4	4	2	-	-
Male	-	-	-	-	-	-	1	-	1	1	-	-
Female	-	-	1	-	-	-	3	4	3	1	-	-
White, both sexes	-	-	1	-	-	-	2	2	2	1	-	-
Male	-	-	-	-	-	-	1	-	1	1	-	-
Female	-	-	1	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	2	2	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Florida												
All races, both sexes	7	7	18	36	45	96	164	233	193	91	12	-
Male	7	6	13	20	26	46	86	103	65	27	1	-
Female	-	1	5	16	19	50	78	130	128	64	11	-
White, both sexes	4	6	16	33	41	88	159	228	187	86	11	-
Male	4	5	11	19	24	42	85	102	63	24	1	-
Female	-	1	5	14	17	46	74	126	124	62	10	-
Black, both sexes	3	1	2	3	3	6	5	4	6	4	1	-
Male	3	1	2	1	1	4	1	1	2	2	-	-
Female	-	-	-	2	2	2	4	3	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Georgia												
All races, both sexes	-	11	4	14	18	28	45	66	46	21	2	-
Male	-	8	4	5	9	13	21	31	17	5	-	-
Female	-	3	-	9	9	15	24	35	29	16	2	-
White, both sexes	-	8	4	12	14	24	42	60	40	19	2	-
Male	-	6	4	4	7	10	20	27	13	4	-	-
Female	-	2	-	8	7	14	22	33	27	15	2	-
Black, both sexes	-	3	-	2	4	4	3	5	6	2	-	-
Male	-	2	-	1	2	3	1	3	4	1	-	-
Female	-	1	-	1	2	1	2	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Hawaii												
All races, both sexes	1	-	-	1	4	7	9	16	10	9	1	-
Male	1	-	-	-	2	5	7	7	3	3	-	-
Female	-	-	-	1	2	2	2	9	7	6	1	-
White, both sexes	1	-	-	-	-	3	4	2	1	-	-	-
Male	1	-	-	-	-	3	3	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	4	4	5	14	9	9	1	-
Male	-	-	-	-	2	2	4	5	3	3	-	-
Female	-	-	-	1	2	2	1	9	6	6	1	-
Idaho												
All races, both sexes	-	1	1	2	4	7	7	27	16	6	-	-
Male	-	1	1	2	3	3	3	10	6	1	-	-
Female	-	-	-	-	1	4	4	17	10	5	-	-
White, both sexes	-	1	1	2	4	7	7	27	16	6	-	-
Male	-	1	1	2	3	3	3	10	6	1	-	-
Female	-	-	-	-	1	4	4	17	10	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Illinois												
All races, both sexes	6	10	10	14	27	57	103	125	110	41	6	-
Male	6	7	9	8	10	27	34	53	34	8	1	-
Female	-	3	1	6	17	30	69	72	76	33	5	-
White, both sexes	6	7	10	13	23	55	97	116	107	40	6	-
Male	6	6	9	8	7	26	34	50	33	8	1	-
Female	-	1	1	5	16	29	63	66	74	32	5	-
Black, both sexes	-	2	-	1	4	-	6	9	3	1	-	-
Male	-	1	-	-	3	-	-	3	1	-	-	-
Female	-	1	-	1	1	-	6	6	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Indiana												
All races, both sexes	-	7	6	12	13	40	43	67	70	19	7	-
Male	-	4	-	7	6	14	21	20	16	1	-	-
Female	-	3	6	5	7	26	22	47	54	18	7	-
White, both sexes	-	7	6	11	12	39	37	65	69	19	7	-
Male	-	4	-	6	6	14	20	20	15	1	-	-
Female	-	3	6	5	6	25	17	45	54	18	7	-
Black, both sexes	-	-	-	1	1	1	6	2	1	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	1	5	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Iowa												
All races, both sexes	1	2	3	1	10	19	34	56	54	20	8	-
Male	1	2	2	1	5	8	12	23	17	2	-	-
Female	-	-	1	-	5	11	22	33	37	18	8	-
White, both sexes	1	2	3	1	10	19	34	55	54	20	8	-
Male	1	2	2	1	5	8	12	23	17	2	-	-
Female	-	-	1	-	5	11	22	32	37	18	8	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	3	-	5	6	8	27	35	44	19	5	-
Male	-	2	-	4	3	4	8	16	13	5	1	-
Female	-	1	-	1	3	4	19	19	31	14	4	-
White, both sexes	-	3	-	5	6	7	27	35	42	18	4	-
Male	-	2	-	4	3	3	8	16	12	5	1	-
Female	-	1	-	1	3	4	19	19	30	13	3	-
Black, both sexes	-	-	-	-	-	1	-	-	1	1	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Kentucky												
All races, both sexes	1	1	2	9	3	14	24	31	26	5	2	-
Male	1	1	1	5	1	10	8	12	7	3	-	-
Female	-	-	1	4	2	4	16	19	19	2	2	-
White, both sexes	1	1	2	9	3	14	24	30	24	5	2	-
Male	1	1	1	5	1	10	8	12	7	3	-	-
Female	-	-	1	4	2	4	16	18	17	2	2	-
Black, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	6	5	8	18	19	33	30	18	11	-	-
Male	4	2	4	3	11	12	17	10	5	4	-	-
Female	-	4	1	5	7	7	16	20	13	7	-	-
White, both sexes	2	6	5	5	17	17	31	28	17	10	-	-
Male	2	2	4	1	11	11	15	10	4	3	-	-
Female	-	4	1	4	6	6	16	18	13	7	-	-
Black, both sexes	2	-	-	2	1	2	2	2	1	1	-	-
Male	2	-	-	1	-	1	2	-	1	1	-	-
Female	-	-	-	1	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Maine												
All races, both sexes	-	1	-	4	3	18	14	28	26	10	2	-
Male	-	1	-	3	2	11	9	15	5	2	-	-
Female	-	-	-	1	1	7	5	13	21	8	2	-
White, both sexes	-	1	-	4	3	18	14	28	26	10	2	-
Male	-	1	-	3	2	11	9	15	5	2	-	-
Female	-	-	-	1	1	7	5	13	21	8	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	4	3	9	11	15	56	53	54	18	-	-
Male	1	3	2	4	7	7	26	18	9	2	-	-
Female	3	1	1	5	4	8	30	35	45	16	-	-
White, both sexes	3	3	2	4	7	11	46	48	53	15	-	-
Male	-	3	1	3	5	5	19	16	9	2	-	-
Female	3	-	1	1	2	6	27	32	44	13	-	-
Black, both sexes	-	1	1	4	4	4	9	5	1	3	-	-
Male	-	-	1	1	2	2	6	2	-	-	-	-
Female	-	1	-	3	2	2	3	3	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Massachusetts												
All races, both sexes	3	8	4	5	19	30	66	104	98	37	6	-
Male	2	6	1	3	14	9	25	31	27	7	1	-
Female	1	2	3	2	5	21	41	73	71	30	5	-
White, both sexes	3	8	4	5	18	28	64	102	95	37	6	-
Male	2	6	1	3	14	8	24	31	27	7	1	-
Female	1	2	3	2	4	20	40	71	68	30	5	-
Black, both sexes	-	-	-	-	-	2	1	2	3	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Michigan												
All races, both sexes	4	10	5	12	17	39	61	117	92	37	7	-
Male	2	6	2	7	9	19	24	52	25	10	1	-
Female	2	4	3	5	8	20	37	65	67	27	6	-
White, both sexes	2	8	5	10	15	35	59	111	92	36	7	-
Male	1	5	2	6	9	17	24	49	25	10	1	-
Female	1	3	3	4	6	18	35	62	67	26	6	-
Black, both sexes	2	1	-	2	2	4	1	6	-	1	-	-
Male	1	1	-	1	-	2	-	3	-	-	-	-
Female	1	-	-	1	2	2	1	3	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Minnesota												
All races, both sexes	-	3	6	9	19	26	35	79	67	31	6	-
Male	-	2	3	6	10	12	16	31	21	6	1	-
Female	-	1	3	3	9	14	19	48	46	25	5	-
White, both sexes	-	2	6	9	19	25	34	79	66	31	6	-
Male	-	1	3	6	10	12	15	31	21	6	1	-
Female	-	1	3	3	9	13	19	48	45	25	5	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	4	3	5	5	10	16	12	8	4	1	-
Male	-	3	2	3	3	4	5	2	6	1	-	-
Female	-	1	1	2	2	6	11	10	2	3	1	-
White, both sexes	-	2	2	2	5	9	11	10	8	3	1	-
Male	-	1	2	1	3	3	3	2	6	-	-	-
Female	-	1	-	1	2	6	8	8	2	3	1	-
Black, both sexes	-	1	1	3	-	1	5	2	-	1	-	-
Male	-	1	-	2	-	1	2	-	-	1	-	-
Female	-	-	1	1	-	-	3	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Missouri												
All races, both sexes	3	3	6	10	11	30	60	72	55	22	3	-
Male	2	2	5	3	8	12	26	23	15	4	1	-
Female	1	1	1	7	3	18	34	49	40	18	2	-
White, both sexes	3	3	6	9	10	29	58	69	53	21	3	-
Male	2	2	5	3	7	11	25	23	15	4	1	-
Female	1	1	1	6	3	18	33	46	38	17	2	-
Black, both sexes	-	-	-	1	1	1	2	3	2	1	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	1	-	2	6	7	7	23	7	-	-
Male	1	-	1	-	2	4	5	2	10	2	-	-
Female	-	-	-	-	-	2	2	5	13	5	-	-
White, both sexes	1	-	1	-	2	6	7	7	23	7	-	-
Male	1	-	1	-	2	4	5	2	10	2	-	-
Female	-	-	-	-	-	2	2	5	13	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Nebraska												
All races, both sexes	-	-	-	3	7	7	17	22	32	11	4	-
Male	-	-	-	2	4	2	13	5	9	2	2	-
Female	-	-	-	1	3	5	4	17	23	9	2	-
White, both sexes	-	-	-	3	7	7	16	22	32	11	4	-
Male	-	-	-	2	4	2	12	5	9	2	2	-
Female	-	-	-	1	3	5	4	17	23	9	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	1	5	13	4	20	18	7	6	1	-
Male	-	1	-	3	8	2	5	8	2	1	-	-
Female	-	-	1	2	5	2	15	10	5	5	1	-
White, both sexes	-	1	1	5	11	4	18	16	7	6	1	-
Male	-	1	-	3	6	2	5	7	2	1	-	-
Female	-	-	1	2	5	2	13	9	5	5	1	-
Black, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
New Hampshire												
All races, both sexes	1	2	2	4	3	9	19	29	25	4	1	-
Male	1	2	1	2	1	4	8	13	7	1	-	-
Female	-	-	1	2	2	5	11	16	18	3	1	-
White, both sexes	1	2	2	4	3	9	19	29	25	4	1	-
Male	1	2	1	2	1	4	8	13	7	1	-	-
Female	-	-	1	2	2	5	11	16	18	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	3	11	13	25	37	94	126	125	52	6	-
Male	1	3	5	6	16	16	43	44	33	9	-	-
Female	1	-	6	7	9	21	51	82	92	43	6	-
White, both sexes	2	3	10	12	24	35	86	121	121	49	6	-
Male	1	3	4	6	15	15	38	42	32	9	-	-
Female	1	-	6	6	9	20	48	79	89	40	6	-
Black, both sexes	-	-	-	1	1	1	8	4	3	2	-	-
Male	-	-	-	-	1	1	5	2	1	-	-	-
Female	-	-	-	1	-	-	3	2	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	1	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
New Mexico												
All races, both sexes	1	2	1	3	6	2	15	14	11	3	-	-
Male	1	2	-	2	2	-	6	8	4	2	-	-
Female	-	-	1	1	4	2	9	6	7	1	-	-
White, both sexes	1	2	1	3	4	2	15	14	11	2	-	-
Male	1	2	-	2	2	-	6	8	4	1	-	-
Female	-	-	1	1	2	2	9	6	7	1	-	-
Black, both sexes	-	-	-	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	9	9	15	25	76	140	206	189	61	18	-
Male	4	5	6	8	17	33	73	74	51	13	-	-
Female	-	4	3	7	8	43	67	132	138	48	18	-
White, both sexes	3	9	9	12	24	72	134	202	186	58	18	-
Male	3	5	6	6	16	31	70	73	49	12	-	-
Female	-	4	3	6	8	41	64	129	137	46	18	-
Black, both sexes	-	-	-	2	1	4	3	4	2	2	-	-
Male	-	-	-	2	1	2	2	1	1	1	-	-
Female	-	-	-	-	-	2	1	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	2	-	1	1	-	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
North Carolina												
All races, both sexes	3	6	8	16	24	34	72	124	71	19	5	-
Male	2	5	6	10	14	16	29	52	20	3	-	-
Female	1	1	2	6	10	18	43	72	51	16	5	-
White, both sexes	2	6	7	14	22	31	64	114	63	15	4	-
Male	1	5	5	8	12	14	27	48	20	2	-	-
Female	1	1	2	6	10	17	37	66	43	13	4	-
Black, both sexes	1	-	1	1	2	3	7	10	8	4	1	-
Male	1	-	1	1	2	2	1	4	-	1	-	-
Female	-	-	-	-	-	1	6	6	8	3	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	1	1	4	6	12	7	4	-	-
Male	-	-	1	-	1	2	3	4	-	1	-	-
Female	-	1	-	1	-	2	3	8	7	3	-	-
White, both sexes	-	1	1	1	1	4	6	12	7	4	-	-
Male	-	-	1	-	1	2	3	4	-	1	-	-
Female	-	1	-	1	-	2	3	8	7	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Ohio												
All races, both sexes	11	10	11	22	27	58	99	175	136	42	8	-
Male	6	6	5	11	12	30	40	60	30	5	-	-
Female	5	4	6	11	15	28	59	115	106	37	8	-
White, both sexes	9	10	10	21	24	57	95	172	129	42	8	-
Male	4	6	5	10	11	30	39	58	30	5	-	-
Female	5	4	5	11	13	27	56	114	99	37	8	-
Black, both sexes	2	-	1	1	3	-	4	3	7	-	-	-
Male	2	-	-	1	1	-	1	2	-	-	-	-
Female	-	-	1	-	2	-	3	1	7	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	3	7	6	9	12	33	28	19	8	2	-
Male	1	1	3	3	4	3	22	11	6	2	-	-
Female	-	2	4	3	5	9	11	17	13	6	2	-
White, both sexes	1	3	7	5	9	12	30	27	17	7	2	-
Male	1	1	3	3	4	3	21	11	5	1	-	-
Female	-	2	4	2	5	9	9	16	12	6	2	-
Black, both sexes	-	-	-	1	-	-	3	-	2	1	-	-
Male	-	-	-	-	-	-	1	-	1	1	-	-
Female	-	-	-	1	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Oregon												
All races, both sexes	4	4	4	11	10	34	72	94	79	30	6	-
Male	2	3	3	5	6	9	29	38	22	9	1	-
Female	2	1	1	6	4	25	43	56	57	21	5	-
White, both sexes	4	4	4	11	9	32	69	92	79	30	6	-
Male	2	3	3	5	6	9	28	36	22	9	1	-
Female	2	1	1	6	3	23	41	56	57	21	5	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10	14	23	25	40	79	177	262	238	80	12	-
Male	5	9	12	13	19	30	79	89	60	15	2	-
Female	5	5	11	12	21	49	98	173	178	65	10	-
White, both sexes	6	13	22	24	40	75	171	259	231	80	12	-
Male	3	8	11	12	19	28	76	88	59	15	2	-
Female	3	5	11	12	21	47	95	171	172	65	10	-
Black, both sexes	4	1	1	1	-	4	5	2	6	-	-	-
Male	2	1	1	1	-	2	2	1	1	-	-	-
Female	2	-	-	-	-	2	3	1	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Rhode Island												
All races, both sexes	-	-	1	1	3	2	6	13	15	4	3	-
Male	-	-	1	1	2	1	2	2	2	-	-	-
Female	-	-	-	-	1	1	4	11	13	4	3	-
White, both sexes	-	-	1	1	2	2	6	13	15	4	3	-
Male	-	-	1	1	2	1	2	2	2	-	-	-
Female	-	-	-	-	-	1	4	11	13	4	3	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	5	9	5	9	11	40	58	24	9	4	-
Male	-	4	5	2	7	5	23	17	9	2	1	-
Female	-	1	4	3	2	6	17	41	15	7	3	-
White, both sexes	-	4	8	2	9	10	33	55	21	9	2	-
Male	-	3	5	1	7	4	22	16	8	2	-	-
Female	-	1	3	1	2	6	11	39	13	7	2	-
Black, both sexes	-	1	1	3	-	1	7	3	3	-	2	-
Male	-	1	-	1	-	1	1	1	1	-	1	-
Female	-	-	1	2	-	-	6	2	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
South Dakota												
All races, both sexes	1	-	-	1	1	-	8	13	9	5	-	-
Male	1	-	-	-	1	-	3	5	4	1	-	-
Female	-	-	-	1	-	-	5	8	5	4	-	-
White, both sexes	1	-	-	1	1	-	8	13	9	5	-	-
Male	1	-	-	-	1	-	3	5	4	1	-	-
Female	-	-	-	1	-	-	5	8	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	5	3	10	17	28	48	57	45	15	2	-
Male	4	3	2	7	11	13	17	18	15	8	-	-
Female	1	2	1	3	6	15	31	39	30	7	2	-
White, both sexes	4	4	3	10	15	26	43	56	44	15	2	-
Male	3	2	2	7	10	13	14	18	15	8	-	-
Female	1	2	1	3	5	13	29	38	29	7	2	-
Black, both sexes	1	1	-	-	2	2	5	1	1	-	-	-
Male	1	1	-	-	1	-	3	-	-	-	-	-
Female	-	-	-	-	1	2	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Texas												
All races, both sexes	12	17	27	31	39	83	117	146	111	48	6	-
Male	7	14	10	16	20	38	47	50	34	6	-	-
Female	5	3	17	15	19	45	70	96	77	42	6	-
White, both sexes	11	14	25	28	35	75	110	139	108	47	6	-
Male	6	12	8	16	18	33	44	48	34	6	-	-
Female	5	2	17	12	17	42	66	91	74	41	6	-
Black, both sexes	1	3	-	3	3	6	7	7	3	1	-	-
Male	1	2	-	-	2	3	3	2	-	-	-	-
Female	-	1	-	3	1	3	4	5	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	2	-	-	-	-	-	-
Male	-	-	2	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	4	-	1	1	2	7	11	23	19	6	-	-
Male	3	-	1	1	-	3	6	9	12	2	-	-
Female	1	-	-	-	2	4	5	14	7	4	-	-
White, both sexes	4	-	1	1	1	7	11	23	19	6	-	-
Male	3	-	1	1	-	3	6	9	12	2	-	-
Female	1	-	-	-	1	4	5	14	7	4	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Vermont												
All races, both sexes	-	-	-	4	5	7	11	14	12	4	1	-
Male	-	-	-	4	3	3	2	7	2	1	-	-
Female	-	-	-	-	2	4	9	7	10	3	1	-
White, both sexes	-	-	-	4	5	7	11	14	12	4	1	-
Male	-	-	-	4	3	3	2	7	2	1	-	-
Female	-	-	-	-	2	4	9	7	10	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	6	10	12	18	27	53	73	61	20	-	-
Male	3	3	4	6	10	17	25	28	14	3	-	-
Female	1	3	6	6	8	10	28	45	47	17	-	-
White, both sexes	3	4	9	11	16	24	49	69	56	18	-	-
Male	3	2	4	6	9	15	23	25	13	3	-	-
Female	-	2	5	5	7	9	26	44	43	15	-	-
Black, both sexes	1	2	1	1	2	3	4	3	5	2	-	-
Male	-	1	-	-	1	2	2	2	1	-	-	-
Female	1	1	1	1	1	1	2	1	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Washington												
All races, both sexes	3	2	11	10	20	42	92	112	122	42	12	-
Male	1	1	5	6	10	22	37	47	40	14	1	-
Female	2	1	6	4	10	20	55	65	82	28	11	-
White, both sexes	2	2	10	9	20	40	91	107	119	41	12	-
Male	1	1	4	6	10	21	36	44	39	13	1	-
Female	1	1	6	3	10	19	55	63	80	28	11	-
Black, both sexes	1	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	3	2	-	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	1	2	1	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
West Virginia												
All races, both sexes	2	2	3	5	6	7	11	17	17	5	-	-
Male	1	2	2	5	4	4	5	6	1	1	-	-
Female	1	-	1	-	2	3	6	11	16	4	-	-
White, both sexes	2	2	3	5	6	7	11	17	16	5	-	-
Male	1	2	2	5	4	4	5	6	1	1	-	-
Female	1	-	1	-	2	3	6	11	15	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Wisconsin												
All races, both sexes	3	5	4	15	15	38	54	90	93	34	6	-
Male	1	4	4	10	7	16	24	42	30	6	1	-
Female	2	1	-	5	8	22	30	48	63	28	5	-
White, both sexes	3	5	4	15	15	38	54	89	93	34	6	-
Male	1	4	4	10	7	16	24	42	30	6	1	-
Female	2	1	-	5	8	22	30	47	63	28	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	2	-	6	1	4	7	1	-	-
Male	-	-	-	2	-	3	-	2	2	-	-	-
Female	-	-	1	-	-	3	1	2	5	1	-	-
White, both sexes	-	-	1	2	-	6	1	4	7	1	-	-
Male	-	-	-	2	-	3	-	2	2	-	-	-
Female	-	-	1	-	-	3	1	2	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
United States												
All races, both sexes	5,053	1	-	2	1	1	11	18	34	35	84	120
Male	2,039	-	-	1	1	-	8	12	22	23	50	82
Female	3,014	1	-	1	-	1	3	6	12	12	34	38
White, both sexes	4,468	1	-	2	1	1	9	11	23	26	57	86
Male	1,781	-	-	1	1	-	8	8	16	17	36	60
Female	2,687	1	-	1	-	1	1	3	7	9	21	26
Black, both sexes	474	-	-	-	-	-	2	7	10	8	23	31
Male	214	-	-	-	-	-	-	4	6	5	13	21
Female	260	-	-	-	-	-	2	3	4	3	10	10
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	86	-	-	-	-	-	-	-	1	1	2	2
Male	30	-	-	-	-	-	-	-	-	1	-	-
Female	56	-	-	-	-	-	-	-	1	-	2	2
Alabama												
All races, both sexes	57	-	-	-	-	-	-	-	2	-	3	3
Male	18	-	-	-	-	-	-	-	1	-	-	3
Female	39	-	-	-	-	-	-	-	1	-	3	-
White, both sexes	41	-	-	-	-	-	-	-	2	-	2	-
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	29	-	-	-	-	-	-	-	1	-	2	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	3
Male	6	-	-	-	-	-	-	-	-	-	-	3
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Alaska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	104	-	-	-	-	-	1	-	-	1	4	3
Male	43	-	-	-	-	-	1	-	-	1	3	2
Female	61	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	97	-	-	-	-	-	1	-	-	1	3	2
Male	38	-	-	-	-	-	1	-	-	1	2	2
Female	59	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Arkansas												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	421	-	-	1	-	-	1	2	1	2	5	10
Male	185	-	-	-	-	-	1	2	-	2	4	9
Female	236	-	-	1	-	-	-	-	1	-	1	1
White, both sexes	361	-	-	1	-	-	1	1	-	1	4	9
Male	162	-	-	-	-	-	1	1	-	1	4	9
Female	199	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	25	-	-	-	-	-	-	1	-	-	-	-
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	1	1	1	1
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	21	-	-	-	-	-	-	-	1	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Colorado												
All races, both sexes	80	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	74	-	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	71	-	-	-	-	-	-	1	2	-	1	1
Male	30	-	-	-	-	-	-	1	2	-	1	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	64	-	-	-	-	-	-	-	1	-	1	1
Male	25	-	-	-	-	-	-	-	1	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	1	1	-	-	-
Male	5	-	-	-	-	-	-	1	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Delaware												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	10	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Florida												
All races, both sexes	371	-	-	-	-	-	-	1	2	3	2	9
Male	156	-	-	-	-	-	-	-	1	2	1	7
Female	215	-	-	-	-	-	-	1	1	1	1	2
White, both sexes	335	-	-	-	-	-	-	1	2	3	2	6
Male	138	-	-	-	-	-	-	-	1	2	1	6
Female	197	-	-	-	-	-	-	1	1	1	1	-
Black, both sexes	32	-	-	-	-	-	-	-	-	-	-	3
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	94	-	-	-	-	-	-	-	-	1	3	7
Male	43	-	-	-	-	-	-	-	-	1	2	5
Female	51	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	62	-	-	-	-	-	-	-	-	1	-	4
Male	29	-	-	-	-	-	-	-	-	1	-	3
Female	33	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	31	-	-	-	-	-	-	-	-	-	2	3
Male	14	-	-	-	-	-	-	-	-	-	2	2
Female	17	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Hawaii												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	2	1
Male	12	-	-	-	-	-	-	-	-	-	2	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	32	-	-	-	-	-	-	-	-	-	2	1
Male	12	-	-	-	-	-	-	-	-	-	2	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Illinois												
All races, both sexes	189	-	-	-	-	-	-	1	2	3	3	2
Male	67	-	-	-	-	-	-	1	1	1	2	1
Female	122	-	-	-	-	-	-	-	1	2	1	1
White, both sexes	162	-	-	-	-	-	-	-	2	2	2	1
Male	58	-	-	-	-	-	-	-	1	1	2	-
Female	104	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	25	-	-	-	-	-	-	1	-	1	1	1
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	139	-	-	-	-	-	-	1	-	1	3	1
Male	55	-	-	-	-	-	-	1	-	1	2	1
Female	84	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	132	-	-	-	-	-	-	1	-	1	3	-
Male	51	-	-	-	-	-	-	1	-	1	2	-
Female	81	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Iowa												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Kentucky												
All races, both sexes	77	-	-	-	-	-	-	-	1	-	-	3
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	74	-	-	-	-	-	-	-	1	-	-	3
Male	30	-	-	-	-	-	-	-	-	-	-	2
Female	44	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	70	-	-	-	-	-	-	-	-	1	2	2
Male	27	-	-	-	-	-	-	-	-	1	2	2
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	1	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Maine												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	84	-	-	-	-	1	-	-	3	2	3	1
Male	32	-	-	-	-	-	-	-	2	1	1	1
Female	52	-	-	-	-	1	-	-	1	1	2	-
White, both sexes	49	-	-	-	-	1	-	-	1	2	-	-
Male	18	-	-	-	-	-	-	-	-	1	-	-
Female	31	-	-	-	-	1	-	-	1	1	-	-
Black, both sexes	34	-	-	-	-	-	-	-	2	-	3	1
Male	13	-	-	-	-	-	-	-	2	-	1	1
Female	21	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Massachusetts												
All races, both sexes	100	-	-	-	-	-	1	1	1	2	5	2
Male	42	-	-	-	-	-	1	-	1	1	4	2
Female	58	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	92	-	-	-	-	-	1	-	1	2	4	2
Male	39	-	-	-	-	-	1	-	1	1	3	2
Female	53	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	7	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	186	-	-	-	-	-	-	-	2	2	3	3
Male	79	-	-	-	-	-	-	-	1	2	1	-
Female	107	-	-	-	-	-	-	-	1	-	2	3
White, both sexes	161	-	-	-	-	-	-	-	-	-	1	1
Male	65	-	-	-	-	-	-	-	-	-	-	-
Female	96	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	23	-	-	-	-	-	-	-	2	2	2	2
Male	13	-	-	-	-	-	-	-	1	2	1	-
Female	10	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Minnesota												
All races, both sexes	67	-	-	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	43	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	66	-	-	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	42	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	4
Male	16	-	-	-	-	-	-	-	-	-	-	4
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Missouri												
All races, both sexes	111	-	-	-	-	-	-	-	-	1	-	1
Male	42	-	-	-	-	-	-	-	-	1	-	-
Female	69	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	103	-	-	-	-	-	-	-	-	1	-	-
Male	41	-	-	-	-	-	-	-	-	1	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Nebraska												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	48	-	-	-	-	-	-	-	1	1	-	3
Male	21	-	-	-	-	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	45	-	-	-	-	-	-	-	1	1	-	2
Male	20	-	-	-	-	-	-	-	1	-	-	1
Female	25	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
New Hampshire												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	178	-	-	-	-	-	1	1	3	2	4	2
Male	64	-	-	-	-	-	-	1	2	-	3	1
Female	114	-	-	-	-	-	1	-	1	2	1	1
White, both sexes	158	-	-	-	-	-	-	1	2	1	2	2
Male	59	-	-	-	-	-	-	1	2	-	1	1
Female	99	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	16	-	-	-	-	-	1	-	1	1	2	-
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	11	-	-	-	-	-	1	-	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
New Mexico												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	3	1
Male	17	-	-	-	-	-	-	-	-	-	3	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	2	1
Male	15	-	-	-	-	-	-	-	-	-	2	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	333	-	-	-	-	-	-	-	1	1	1	8
Male	131	-	-	-	-	-	-	-	-	-	-	6
Female	202	-	-	-	-	-	-	-	1	1	1	2
White, both sexes	289	-	-	-	-	-	-	-	-	1	-	5
Male	110	-	-	-	-	-	-	-	-	-	-	3
Female	179	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	41	-	-	-	-	-	-	-	1	-	1	3
Male	21	-	-	-	-	-	-	-	-	-	-	3
Female	20	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
North Carolina												
All races, both sexes	135	-	-	-	-	-	-	1	2	3	1	5
Male	57	-	-	-	-	-	-	-	1	2	1	3
Female	78	-	-	-	-	-	-	1	1	1	-	2
White, both sexes	113	-	-	-	-	-	-	-	2	2	1	2
Male	47	-	-	-	-	-	-	-	1	2	1	-
Female	66	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	18	-	-	-	-	-	-	1	-	1	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	3
Female	11	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	20	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Ohio												
All races, both sexes	242	-	-	-	-	-	1	-	3	1	3	7
Male	91	-	-	-	-	-	1	-	3	1	1	5
Female	151	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	223	-	-	-	-	-	1	-	3	-	3	6
Male	81	-	-	-	-	-	1	-	3	-	1	5
Female	142	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	16	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	69	-	-	-	-	-	-	-	-	1	1	3
Male	33	-	-	-	-	-	-	-	-	1	1	1
Female	36	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	60	-	-	-	-	-	-	-	-	1	-	3
Male	27	-	-	-	-	-	-	-	-	1	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Oregon												
All races, both sexes	82	-	-	-	-	-	-	-	-	1	-	1
Male	36	-	-	-	-	-	-	-	-	1	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	78	-	-	-	-	-	-	-	-	1	-	1
Male	34	-	-	-	-	-	-	-	-	1	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	363	1	-	-	1	-	-	1	1	2	5	5
Male	155	-	-	-	1	-	-	1	1	2	1	4
Female	208	1	-	-	-	-	-	-	-	-	4	1
White, both sexes	333	1	-	-	1	-	-	1	1	2	2	3
Male	142	-	-	-	1	-	-	1	1	2	1	2
Female	191	1	-	-	-	-	-	-	-	-	1	1
Black, both sexes	29	-	-	-	-	-	-	-	-	-	3	2
Male	13	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Rhode Island												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	2	2
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	55	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	69	-	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
South Dakota												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	113	-	-	-	-	-	1	2	2	2	3	6
Male	61	-	-	-	-	-	-	2	1	2	3	4
Female	52	-	-	-	-	-	1	-	1	-	-	2
White, both sexes	98	-	-	-	-	-	-	2	2	1	3	5
Male	52	-	-	-	-	-	-	2	1	1	3	4
Female	46	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	15	-	-	-	-	-	1	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Texas												
All races, both sexes	263	-	-	-	-	-	4	2	1	1	6	7
Male	109	-	-	-	-	-	3	1	-	-	3	6
Female	154	-	-	-	-	-	1	1	1	1	3	1
White, both sexes	234	-	-	-	-	-	4	2	-	1	6	7
Male	98	-	-	-	-	-	3	1	-	-	3	6
Female	136	-	-	-	-	-	1	1	-	1	3	1
Black, both sexes	26	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	47	-	-	-	-	-	-	-	1	-	-	1
Male	21	-	-	-	-	-	-	-	1	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	47	-	-	-	-	-	-	-	1	-	-	1
Male	21	-	-	-	-	-	-	-	1	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Vermont												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	108	-	-	-	-	-	1	1	3	1	5	2
Male	44	-	-	-	-	-	1	-	3	-	4	2
Female	64	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	91	-	-	-	-	-	1	-	1	1	3	2
Male	31	-	-	-	-	-	1	-	1	-	2	2
Female	60	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	16	-	-	-	-	-	-	1	2	-	2	-
Male	12	-	-	-	-	-	-	-	2	-	2	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Washington												
All races, both sexes	126	-	-	-	-	-	-	1	-	-	2	3
Male	51	-	-	-	-	-	-	1	-	-	1	3
Female	75	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	119	-	-	-	-	-	-	1	-	-	2	3
Male	50	-	-	-	-	-	-	1	-	-	1	3
Female	69	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Wisconsin												
All races, both sexes	126	-	-	-	-	-	-	-	-	-	4	1
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	91	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	121	-	-	-	-	-	-	-	-	-	4	-
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	86	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	17	-	-	1	-	-	-	-	-	-	-	1
Male	6	-	-	1	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	1	-	-	-	-	-	-	-	1
Male	6	-	-	1	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
United States												
All races, both sexes	174	235	232	302	394	622	867	962	662	241	55	-
Male	96	151	130	150	177	257	342	310	176	44	7	-
Female	78	84	102	152	217	365	525	652	486	197	48	-
White, both sexes	113	173	196	234	336	565	797	914	639	236	48	-
Male	63	114	108	124	152	236	321	299	169	44	4	-
Female	50	59	88	110	184	329	476	615	470	192	44	-
Black, both sexes	51	54	24	59	50	51	48	34	14	2	6	-
Male	27	34	13	20	22	19	12	11	4	-	3	-
Female	24	20	11	39	28	32	36	23	10	2	3	-
American Indian 1/, both sexes	3	6	4	2	1	1	1	2	1	-	1	-
Male	2	3	4	2	1	-	-	-	-	-	-	-
Female	1	3	-	-	-	1	1	2	1	-	1	-
Asian or Pacific Islander 2/, both sexes	7	2	8	7	7	5	21	12	8	3	-	-
Male	4	-	5	4	2	2	9	-	3	-	-	-
Female	3	2	3	3	5	3	12	12	5	3	-	-
Alabama												
All races, both sexes	5	-	2	6	3	8	7	13	4	-	1	-
Male	1	-	-	2	2	3	2	4	-	-	-	-
Female	4	-	2	4	1	5	5	9	4	-	1	-
White, both sexes	2	-	2	4	2	5	4	13	4	-	1	-
Male	-	-	-	2	1	2	2	4	-	-	-	-
Female	2	-	2	2	1	3	2	9	4	-	1	-
Black, both sexes	3	-	-	2	1	3	2	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	2	-	-	2	-	2	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Alaska												
All races, both sexes	2	3	-	-	1	-	2	2	-	1	-	-
Male	-	2	-	-	-	-	-	1	-	-	-	-
Female	2	1	-	-	1	-	2	1	-	1	-	-
White, both sexes	1	2	-	-	1	-	1	1	-	1	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	1	-	1	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	3	8	3	5	5	9	23	16	13	8	2	-
Male	2	4	3	3	-	4	5	8	4	2	1	-
Female	1	4	-	2	5	5	18	8	9	6	1	-
White, both sexes	3	6	3	4	5	9	22	16	13	8	1	-
Male	2	3	3	2	-	4	4	8	4	2	-	-
Female	1	3	-	2	5	5	18	8	9	6	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Arkansas												
All races, both sexes	-	5	4	5	5	4	10	12	2	2	-	-
Male	-	3	2	1	-	2	1	3	1	1	-	-
Female	-	2	2	4	5	2	9	9	1	1	-	-
White, both sexes	-	4	4	5	5	3	9	12	2	2	-	-
Male	-	2	2	1	-	2	1	3	1	1	-	-
Female	-	2	2	4	5	1	8	9	1	1	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	14	16	21	18	25	41	82	94	53	30	5	-
Male	10	9	14	8	13	21	34	37	16	4	1	-
Female	4	7	7	10	12	20	48	57	37	26	4	-
White, both sexes	9	11	18	10	23	35	71	84	51	28	4	-
Male	7	7	12	5	12	18	30	36	15	4	-	-
Female	2	4	6	5	11	17	41	48	36	24	4	-
Black, both sexes	3	5	-	2	-	4	4	3	1	1	1	-
Male	1	2	-	-	-	2	1	1	-	-	1	-
Female	2	3	-	2	-	2	3	2	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	6	2	2	7	7	1	1	-	-
Male	1	-	2	3	1	1	3	-	1	-	-	-
Female	-	-	1	3	1	1	4	7	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Colorado												
All races, both sexes	3	3	4	3	6	8	19	20	9	3	1	-
Male	-	1	4	3	2	2	2	9	3	-	-	-
Female	3	2	-	-	4	6	17	11	6	3	1	-
White, both sexes	2	3	3	3	6	7	19	18	8	3	1	-
Male	-	1	3	3	2	2	2	8	3	-	-	-
Female	2	2	-	-	4	5	17	10	5	3	1	-
Black, both sexes	1	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Connecticut												
All races, both sexes	3	3	-	5	11	6	13	12	7	4	2	-
Male	3	-	-	2	7	1	5	7	-	1	-	-
Female	-	3	-	3	4	5	8	5	7	3	2	-
White, both sexes	1	3	-	4	9	6	13	12	7	4	2	-
Male	1	-	-	2	6	1	5	7	-	1	-	-
Female	-	3	-	2	3	5	8	5	7	3	2	-
Black, both sexes	2	-	-	1	2	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Delaware												
All races, both sexes	-	1	2	2	-	3	3	1	1	-	-	-
Male	-	1	1	1	-	2	1	-	-	-	-	-
Female	-	-	1	1	-	1	2	1	1	-	-	-
White, both sexes	-	-	2	1	-	3	3	1	1	-	-	-
Male	-	-	1	-	-	2	1	-	-	-	-	-
Female	-	-	1	1	-	1	2	1	1	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	2	2	2	1	-	-	-	-	-
Male	1	-	-	1	1	1	1	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	2	2	1	-	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Florida												
All races, both sexes	7	19	14	23	27	57	60	72	48	25	2	-
Male	7	15	5	12	14	25	22	25	16	4	-	-
Female	-	4	9	11	13	32	38	47	32	21	2	-
White, both sexes	5	14	13	17	23	55	53	68	46	25	2	-
Male	5	12	4	9	12	25	19	23	15	4	-	-
Female	-	2	9	8	11	30	34	45	31	21	2	-
Black, both sexes	2	4	-	6	4	2	6	3	2	-	-	-
Male	2	3	-	3	2	-	2	2	1	-	-	-
Female	-	1	-	3	2	2	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	6	8	5	13	5	11	17	8	9	1	-	-
Male	1	5	3	5	2	3	10	4	2	-	-	-
Female	5	3	2	8	3	8	7	4	7	1	-	-
White, both sexes	1	3	5	8	4	11	12	4	8	1	-	-
Male	-	2	3	5	1	3	7	2	2	-	-	-
Female	1	1	2	3	3	8	5	2	6	1	-	-
Black, both sexes	5	5	-	5	1	-	5	4	1	-	-	-
Male	1	3	-	-	1	-	3	2	-	-	-	-
Female	4	2	-	5	-	-	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Hawaii												
All races, both sexes	3	-	2	-	2	-	3	2	5	1	-	-
Male	2	-	2	-	-	-	1	2	2	-	-	-
Female	1	-	-	-	2	-	2	-	3	1	-	-
White, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	2	-	2	-	5	1	-	-
Male	2	-	1	-	-	-	-	-	2	-	-	-
Female	1	-	-	-	2	-	2	-	3	1	-	-
Idaho												
All races, both sexes	1	-	-	3	4	2	4	8	6	1	-	-
Male	1	-	-	2	1	1	2	2	1	-	-	-
Female	-	-	-	1	3	1	2	6	5	1	-	-
White, both sexes	1	-	-	3	4	2	4	8	6	1	-	-
Male	1	-	-	2	1	1	2	2	1	-	-	-
Female	-	-	-	1	3	1	2	6	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Illinois												
All races, both sexes	9	7	6	11	16	25	28	36	28	11	1	-
Male	4	3	4	5	5	8	12	12	7	1	-	-
Female	5	4	2	6	11	17	16	24	21	10	1	-
White, both sexes	6	6	4	9	10	21	25	35	27	11	1	-
Male	3	3	3	4	2	7	12	12	7	1	-	-
Female	3	3	1	5	8	14	13	23	20	10	1	-
Black, both sexes	3	1	-	2	6	4	3	1	1	-	-	-
Male	1	-	-	1	3	1	-	-	-	-	-	-
Female	2	1	-	1	3	3	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	4	6	6	12	10	9	23	30	26	6	1	-
Male	1	3	5	7	6	3	8	8	7	1	1	-
Female	3	3	1	5	4	6	15	22	19	5	-	-
White, both sexes	4	5	4	12	9	9	22	29	26	6	1	-
Male	1	3	3	7	5	3	8	8	7	1	1	-
Female	3	2	1	5	4	6	14	21	19	5	-	-
Black, both sexes	-	1	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Iowa												
All races, both sexes	-	1	-	3	3	4	1	3	5	3	-	-
Male	-	1	-	3	2	2	-	1	-	2	-	-
Female	-	-	-	-	1	2	1	2	5	1	-	-
White, both sexes	-	1	-	3	3	4	1	3	5	3	-	-
Male	-	1	-	3	2	2	-	1	-	2	-	-
Female	-	-	-	-	1	2	1	2	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	3	2	1	6	5	10	10	10	4	1	-
Male	-	2	2	1	2	-	3	4	3	-	-	-
Female	-	1	-	-	4	5	7	6	7	4	1	-
White, both sexes	-	3	1	1	6	5	10	10	10	4	1	-
Male	-	2	1	1	2	-	3	4	3	-	-	-
Female	-	1	-	-	4	5	7	6	7	4	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Kentucky												
All races, both sexes	3	5	5	5	5	10	15	13	7	4	1	-
Male	1	4	3	2	2	5	7	3	3	-	-	-
Female	2	1	2	3	3	5	8	10	4	4	1	-
White, both sexes	3	5	5	5	4	10	14	13	6	4	1	-
Male	1	4	3	2	1	5	6	3	3	-	-	-
Female	2	1	2	3	3	5	8	10	3	4	1	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	5	5	11	8	10	13	3	4	1	1	-
Male	1	3	2	3	2	4	3	1	2	-	1	-
Female	3	2	3	8	6	6	10	2	2	1	-	-
White, both sexes	2	1	4	7	6	9	12	2	3	1	-	-
Male	-	1	2	3	1	4	3	1	2	-	-	-
Female	2	-	2	4	5	5	9	1	1	1	-	-
Black, both sexes	2	4	1	4	2	1	-	1	1	-	1	-
Male	1	2	-	-	1	-	-	-	-	-	1	-
Female	1	2	1	4	1	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Maine												
All races, both sexes	-	2	3	2	3	5	5	6	4	1	-	-
Male	-	2	2	1	2	3	1	1	-	1	-	-
Female	-	-	1	1	1	2	4	5	4	-	-	-
White, both sexes	-	1	3	2	3	5	5	6	4	1	-	-
Male	-	1	2	1	2	3	1	1	-	1	-	-
Female	-	-	1	1	1	2	4	5	4	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	2	4	7	7	15	13	13	7	3	-	-
Male	2	2	2	2	1	6	6	4	1	1	-	-
Female	1	-	2	5	6	9	7	9	6	2	-	-
White, both sexes	1	-	2	3	2	9	8	11	6	3	-	-
Male	-	-	1	2	-	4	4	4	1	1	-	-
Female	1	-	1	1	2	5	4	7	5	2	-	-
Black, both sexes	2	2	2	4	5	6	4	2	1	-	-	-
Male	2	2	1	-	1	2	1	-	-	-	-	-
Female	-	-	1	4	4	4	3	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Massachusetts												
All races, both sexes	3	7	2	8	9	13	14	19	11	1	1	-
Male	2	4	-	2	3	7	4	6	5	-	-	-
Female	1	3	2	6	6	6	10	13	6	1	1	-
White, both sexes	3	7	1	7	8	10	14	19	11	1	1	-
Male	2	4	-	2	3	5	4	6	5	-	-	-
Female	1	3	1	5	5	5	10	13	6	1	1	-
Black, both sexes	-	-	1	1	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	11	3	9	11	14	30	26	30	31	8	3	-
Male	7	3	8	9	8	10	12	8	9	-	1	-
Female	4	-	1	2	6	20	14	22	22	8	2	-
White, both sexes	5	2	9	9	12	29	24	29	29	8	3	-
Male	3	2	8	8	7	10	10	8	8	-	1	-
Female	2	-	1	1	5	19	14	21	21	8	2	-
Black, both sexes	6	1	-	1	2	1	2	1	1	-	-	-
Male	4	1	-	-	1	-	2	-	1	-	-	-
Female	2	-	-	1	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Minnesota												
All races, both sexes	2	-	1	2	4	9	15	15	12	5	-	-
Male	-	-	1	1	1	4	11	2	3	-	-	-
Female	2	-	-	1	3	5	4	13	9	5	-	-
White, both sexes	1	-	1	2	4	9	15	15	12	5	-	-
Male	-	-	1	1	1	4	11	2	3	-	-	-
Female	1	-	-	1	3	5	4	13	9	5	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	4	2	5	2	4	1	5	2	-	-	-
Male	-	1	2	3	-	2	1	2	1	-	-	-
Female	-	3	-	2	2	2	-	3	1	-	-	-
White, both sexes	-	2	2	4	1	3	1	5	2	-	-	-
Male	-	1	2	2	-	2	1	2	1	-	-	-
Female	-	1	-	2	1	1	-	3	1	-	-	-
Black, both sexes	-	2	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	2	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Missouri												
All races, both sexes	7	3	9	5	11	12	18	22	10	10	2	-
Male	4	1	3	3	5	3	9	5	4	3	1	-
Female	3	2	6	2	6	9	9	17	6	7	1	-
White, both sexes	7	3	8	3	10	12	16	21	10	10	2	-
Male	4	1	3	2	5	3	9	5	4	3	1	-
Female	3	2	5	1	5	9	7	16	6	7	1	-
Black, both sexes	-	-	1	2	1	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	4	3	5	9	1	-	-
Male	-	-	-	-	1	3	3	2	-	-	-	-
Female	-	-	-	-	-	1	-	3	9	1	-	-
White, both sexes	-	-	-	-	1	4	3	5	9	1	-	-
Male	-	-	-	-	1	3	3	2	-	-	-	-
Female	-	-	-	-	-	1	-	3	9	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Nebraska												
All races, both sexes	-	1	1	2	5	8	10	7	7	4	-	-
Male	-	1	-	1	4	3	3	2	3	1	-	-
Female	-	-	1	1	1	5	7	5	4	3	-	-
White, both sexes	-	1	1	2	5	8	10	7	7	4	-	-
Male	-	1	-	1	4	3	3	2	3	1	-	-
Female	-	-	1	1	1	5	7	5	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	3	5	3	7	4	9	3	6	-	-	-
Male	2	3	2	2	2	3	3	1	1	-	-	-
Female	1	-	3	1	5	1	6	2	5	-	-	-
White, both sexes	3	2	5	3	6	4	9	3	6	-	-	-
Male	2	2	2	2	2	3	3	1	1	-	-	-
Female	1	-	3	1	4	1	6	2	5	-	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
New Hampshire												
All races, both sexes	-	-	1	-	1	3	6	3	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	1	3	5	3	1	-	1	-
White, both sexes	-	-	1	-	1	3	6	3	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	1	3	5	3	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	6	3	8	14	19	33	38	24	8	3	-
Male	4	3	1	5	5	8	12	8	7	4	-	-
Female	5	3	2	3	9	11	21	30	17	4	3	-
White, both sexes	5	4	2	7	12	19	31	36	23	8	3	-
Male	3	3	-	5	5	8	12	7	7	4	-	-
Female	2	1	2	2	7	11	19	29	16	4	3	-
Black, both sexes	3	1	1	1	1	-	1	2	1	-	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	2	1	-	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
New Mexico												
All races, both sexes	1	1	3	1	1	3	4	8	5	-	-	-
Male	1	1	2	1	-	2	1	4	2	-	-	-
Female	-	-	1	-	1	1	3	4	3	-	-	-
White, both sexes	-	1	3	1	1	3	4	8	5	-	-	-
Male	-	1	2	1	-	2	1	4	2	-	-	-
Female	-	-	1	-	1	1	3	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	8	19	17	16	31	55	54	62	43	9	8	-
Male	6	12	7	10	13	28	19	19	11	-	-	-
Female	2	7	10	6	18	27	35	43	32	9	8	-
White, both sexes	3	13	12	13	28	47	49	58	43	9	8	-
Male	2	9	4	7	12	24	19	19	11	-	-	-
Female	1	4	8	6	16	23	30	39	32	9	8	-
Black, both sexes	5	6	4	3	3	8	3	4	-	-	-	-
Male	4	3	3	3	1	4	-	-	-	-	-	-
Female	1	3	1	-	2	4	3	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
North Carolina												
All races, both sexes	8	4	8	9	16	15	21	25	10	5	2	-
Male	4	2	2	5	9	4	11	9	2	2	-	-
Female	4	2	6	4	7	11	10	16	8	3	2	-
White, both sexes	7	2	6	8	13	15	18	21	10	5	1	-
Male	4	1	1	4	7	4	10	8	2	2	-	-
Female	3	1	5	4	6	11	8	13	8	3	1	-
Black, both sexes	1	1	2	1	2	-	2	3	-	-	1	-
Male	-	-	1	1	1	-	-	1	-	-	-	-
Female	1	1	1	-	1	-	2	2	-	-	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	1	-	2	4	3	3	4	-	-
Male	-	-	1	-	-	1	1	1	2	1	-	-
Female	-	1	-	1	-	1	3	2	1	3	-	-
White, both sexes	-	1	1	1	-	2	4	3	3	4	-	-
Male	-	-	1	-	-	1	1	1	2	1	-	-
Female	-	1	-	1	-	1	3	2	1	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Ohio												
All races, both sexes	5	12	6	9	15	31	55	52	28	12	2	-
Male	2	6	5	3	6	13	24	14	5	2	-	-
Female	3	6	1	6	9	18	31	38	23	10	2	-
White, both sexes	5	8	5	8	13	29	52	50	27	11	2	-
Male	2	3	4	3	5	12	23	13	4	2	-	-
Female	3	5	1	5	8	17	29	37	23	9	2	-
Black, both sexes	-	4	1	1	1	2	2	2	1	-	-	-
Male	-	3	1	-	1	1	-	1	1	-	-	-
Female	-	1	-	1	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	4	6	2	2	4	6	8	15	10	4	3	-
Male	2	6	1	-	3	3	4	6	3	2	-	-
Female	2	-	1	2	1	3	4	9	7	2	3	-
White, both sexes	2	4	1	2	3	6	8	14	10	4	2	-
Male	1	4	-	-	2	3	4	6	3	2	-	-
Female	1	-	1	2	1	3	4	8	7	2	2	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	1	-	-	1	-	-	1	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Oregon												
All races, both sexes	3	1	1	2	6	16	21	14	10	5	1	-
Male	3	1	1	2	5	6	10	6	1	-	-	-
Female	-	-	-	-	1	10	11	8	9	5	1	-
White, both sexes	2	1	1	2	6	15	20	14	9	5	1	-
Male	2	1	1	2	5	6	9	6	1	-	-	-
Female	-	-	-	-	1	9	11	8	8	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	8	17	17	22	19	41	75	68	56	20	4	-
Male	3	14	10	10	12	16	36	24	14	5	1	-
Female	5	3	7	12	7	25	39	44	42	15	3	-
White, both sexes	5	12	15	18	17	41	70	66	55	20	3	-
Male	-	10	9	9	12	16	36	23	13	5	1	-
Female	5	2	6	9	5	25	34	43	42	15	2	-
Black, both sexes	3	5	2	4	2	-	5	1	1	-	1	-
Male	3	4	1	1	-	-	-	1	1	-	-	-
Female	-	1	1	3	2	-	5	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Rhode Island												
All races, both sexes	1	1	1	1	2	3	6	2	7	1	-	-
Male	1	1	-	1	-	1	4	1	2	-	-	-
Female	-	-	1	-	2	2	2	1	5	1	-	-
White, both sexes	1	1	1	1	2	2	6	2	7	1	-	-
Male	1	1	-	1	-	-	4	1	2	-	-	-
Female	-	-	1	-	2	2	2	1	5	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	5	13	9	12	6	7	14	10	3	-	-
Male	2	5	6	3	3	3	3	3	1	-	-	-
Female	1	-	7	6	9	3	4	11	9	3	-	-
White, both sexes	2	3	11	7	7	5	7	13	10	3	-	-
Male	2	3	6	3	2	2	3	3	1	-	-	-
Female	-	-	5	4	5	3	4	10	9	3	-	-
Black, both sexes	-	2	2	2	5	1	-	1	-	-	-	-
Male	-	2	-	-	1	1	-	-	-	-	-	-
Female	-	-	2	2	4	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
South Dakota												
All races, both sexes	-	2	-	-	1	5	5	3	6	-	-	-
Male	-	2	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	4	5	2	6	-	-	-
White, both sexes	-	2	-	-	1	5	5	3	6	-	-	-
Male	-	2	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	4	5	2	6	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	6	7	13	15	10	13	20	6	1	1	-
Male	4	6	6	7	9	2	5	7	3	-	-	-
Female	1	-	1	6	6	8	8	13	3	1	1	-
White, both sexes	5	5	5	7	14	8	13	20	6	1	1	-
Male	4	5	4	4	8	1	5	7	3	-	-	-
Female	1	-	1	3	6	7	8	13	3	1	1	-
Black, both sexes	-	1	2	6	1	2	-	-	-	-	-	-
Male	-	1	2	3	1	1	-	-	-	-	-	-
Female	-	-	-	3	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Texas												
All races, both sexes	8	19	20	20	17	33	34	53	29	6	3	-
Male	4	11	11	9	10	13	14	17	6	1	-	-
Female	4	8	9	11	7	20	20	36	23	5	3	-
White, both sexes	5	16	18	16	14	28	31	51	27	6	2	-
Male	3	10	10	6	8	11	13	17	6	1	-	-
Female	2	6	8	10	6	17	18	34	21	5	2	-
Black, both sexes	3	2	1	3	3	5	3	2	2	-	1	-
Male	1	1	-	2	2	2	1	-	-	-	-	-
Female	2	1	1	1	1	3	2	2	2	-	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	1	1	2	7	6	9	11	3	2	-	-
Male	3	-	-	2	4	3	5	2	-	1	-	-
Female	-	1	1	-	3	3	4	9	3	1	-	-
White, both sexes	3	1	1	2	7	6	9	11	3	2	-	-
Male	3	-	-	2	4	3	5	2	-	1	-	-
Female	-	1	1	-	3	3	4	9	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Vermont												
All races, both sexes	1	-	-	-	-	-	3	2	2	-	-	-
Male	-	-	-	-	-	-	2	-	2	-	-	-
Female	1	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	1	-	-	-	-	-	3	2	2	-	-	-
Male	-	-	-	-	-	-	2	-	2	-	-	-
Female	1	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	3	4	6	7	17	17	18	14	4	-	-
Male	2	2	3	3	3	6	5	5	5	-	-	-
Female	3	1	1	3	4	11	12	13	9	4	-	-
White, both sexes	4	2	4	3	6	13	16	17	14	4	-	-
Male	1	1	3	1	2	4	4	4	5	-	-	-
Female	3	1	1	2	4	9	12	13	9	4	-	-
Black, both sexes	1	1	-	3	1	3	1	1	-	-	-	-
Male	1	1	-	2	1	1	1	1	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Washington												
All races, both sexes	3	7	6	2	6	15	12	37	24	6	2	-
Male	2	4	3	2	4	6	5	10	8	2	-	-
Female	1	3	3	-	2	9	7	27	16	4	2	-
White, both sexes	2	6	4	2	6	14	11	36	24	6	2	-
Male	2	4	2	2	4	6	5	10	8	2	-	-
Female	-	2	2	-	2	8	6	26	16	4	2	-
Black, both sexes	1	-	2	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
West Virginia												
All races, both sexes	-	2	2	-	5	3	6	5	4	-	-	-
Male	-	-	1	-	1	1	3	1	-	-	-	-
Female	-	2	1	-	4	2	3	4	4	-	-	-
White, both sexes	-	2	2	-	5	3	6	5	4	-	-	-
Male	-	-	1	-	1	1	3	1	-	-	-	-
Female	-	2	1	-	4	2	3	4	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Wisconsin												
All races, both sexes	1	4	-	5	8	14	24	26	26	12	1	-
Male	1	2	-	2	1	8	5	7	6	2	-	-
Female	-	2	-	3	7	6	19	19	20	10	1	-
White, both sexes	1	4	-	4	8	12	24	26	26	11	1	-
Male	1	2	-	2	1	8	5	7	6	2	-	-
Female	-	2	-	2	7	4	19	19	20	9	1	-
Black, both sexes	-	-	-	1	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	2	1	-	1	2	3	5	1	-	-
Male	-	-	-	-	-	1	-	1	2	-	-	-
Female	-	-	2	1	-	-	2	2	3	1	-	-
White, both sexes	-	-	2	1	-	1	2	3	5	1	-	-
Male	-	-	-	-	-	1	-	1	2	-	-	-
Female	-	-	2	1	-	-	2	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
United States												
All races, both sexes	24,703	115	41	18	41	92	178	264	368	503	820	1,169
Male	14,399	65	22	12	30	71	123	200	268	357	570	822
Female	10,304	50	19	6	11	21	55	64	100	146	250	347
White, both sexes	19,865	79	24	10	21	66	115	161	212	309	550	794
Male	11,500	43	13	7	16	51	82	126	162	222	385	556
Female	8,365	36	11	3	5	15	33	35	50	87	165	238
Black, both sexes	4,255	29	14	7	19	25	51	93	143	180	243	343
Male	2,521	17	8	4	14	19	32	67	95	122	165	239
Female	1,734	12	6	3	5	6	19	26	48	58	78	104
American Indian 1/, both sexes	113	2	-	-	-	-	4	2	4	7	11	9
Male	78	2	-	-	-	-	3	1	3	7	8	8
Female	35	-	-	-	-	-	1	1	1	-	3	1
Asian or Pacific Islander 2/, both sexes	470	5	3	1	1	1	8	8	9	7	16	23
Male	300	3	1	1	-	1	6	6	8	6	12	19
Female	170	2	2	-	1	-	2	2	1	1	4	4
Alabama												
All races, both sexes	446	-	-	2	-	2	3	8	8	11	9	15
Male	250	-	-	-	-	2	2	5	7	3	3	8
Female	196	-	-	2	-	-	1	3	1	8	6	7
White, both sexes	314	-	-	1	-	1	1	6	4	6	6	9
Male	176	-	-	-	-	1	1	5	3	1	1	4
Female	138	-	-	1	-	-	-	1	1	5	5	5
Black, both sexes	132	-	-	1	-	1	2	2	4	5	3	6
Male	74	-	-	-	-	1	1	-	4	2	2	4
Female	58	-	-	1	-	-	1	2	-	3	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Alaska												
All races, both sexes	41	-	-	-	-	-	1	1	-	1	2	3
Male	28	-	-	-	-	-	1	-	-	1	2	1
Female	13	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	2
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	1	1	-	1	-	1
Male	5	-	-	-	-	-	1	-	-	1	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	301	6	2	-	1	1	6	2	3	11	6	21
Male	200	4	1	-	1	-	5	2	3	10	6	17
Female	101	2	1	-	-	1	1	-	-	1	-	4
White, both sexes	269	4	2	-	1	-	4	2	3	7	4	19
Male	175	2	1	-	1	-	3	2	3	6	4	16
Female	94	2	1	-	-	-	1	-	-	1	-	3
Black, both sexes	19	1	-	-	-	1	-	-	-	2	1	2
Male	16	1	-	-	-	-	-	-	-	2	1	1
Female	3	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	13	1	-	-	-	-	2	-	-	2	1	-
Male	9	1	-	-	-	-	2	-	-	2	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Arkansas												
All races, both sexes	239	5	2	-	1	1	6	4	5	3	8	13
Male	141	3	1	-	1	1	4	2	5	2	4	8
Female	98	2	1	-	-	-	2	2	-	1	4	5
White, both sexes	182	3	1	-	1	1	3	3	3	2	2	4
Male	108	2	1	-	1	1	2	2	3	2	1	3
Female	74	1	-	-	-	-	1	1	-	-	1	1
Black, both sexes	57	2	1	-	-	-	3	1	2	1	6	9
Male	33	1	-	-	-	-	2	-	2	-	3	5
Female	24	1	1	-	-	-	1	1	-	1	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,970	13	4	2	5	8	28	38	58	72	153	196
Male	1,849	6	3	2	5	8	18	32	47	47	119	142
Female	1,121	7	1	-	-	-	10	6	11	25	34	54
White, both sexes	2,380	8	3	2	4	7	22	28	43	55	118	153
Male	1,495	4	2	2	4	7	14	23	37	35	91	113
Female	885	4	1	-	-	-	8	5	6	20	27	40
Black, both sexes	369	2	-	-	1	1	3	5	11	15	27	30
Male	217	1	-	-	1	1	2	5	6	11	21	18
Female	152	1	-	-	-	-	1	-	5	4	6	12
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	1	5
Male	14	-	-	-	-	-	-	-	-	-	-	5
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	205	3	1	-	-	-	3	5	4	2	7	8
Male	123	1	1	-	-	-	2	4	4	1	7	6
Female	82	2	-	-	-	-	1	1	-	1	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Colorado												
All races, both sexes	234	1	-	-	1	1	5	2	6	7	12	12
Male	135	-	-	-	-	-	4	1	4	5	7	9
Female	99	1	-	-	1	1	1	1	2	2	5	3
White, both sexes	218	1	-	-	1	1	5	2	5	6	12	11
Male	122	-	-	-	-	-	4	1	3	4	7	8
Female	96	1	-	-	1	1	1	1	2	2	5	3
Black, both sexes	10	-	-	-	-	-	-	-	1	1	-	1
Male	7	-	-	-	-	-	-	-	1	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	304	-	-	-	-	-	3	4	-	11	9	13
Male	185	-	-	-	-	-	2	4	-	10	2	10
Female	119	-	-	-	-	-	1	-	-	1	7	3
White, both sexes	278	-	-	-	-	-	3	3	-	8	6	10
Male	167	-	-	-	-	-	2	3	-	7	2	8
Female	111	-	-	-	-	-	1	-	-	1	4	2
Black, both sexes	25	-	-	-	-	-	-	1	-	3	3	3
Male	17	-	-	-	-	-	-	1	-	3	-	2
Female	8	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Delaware												
All races, both sexes	99	2	-	-	-	-	1	-	2	2	2	3
Male	56	1	-	-	-	-	1	-	1	2	2	2
Female	43	1	-	-	-	-	-	-	1	-	-	1
White, both sexes	78	-	-	-	-	-	1	-	-	2	-	3
Male	45	-	-	-	-	-	1	-	-	2	-	2
Female	33	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	19	1	-	-	-	-	-	-	2	-	2	-
Male	10	-	-	-	-	-	-	-	1	-	2	-
Female	9	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	32	1	-	-	-	1	1	2	-	1	2	-
Male	20	-	-	-	-	1	-	1	-	1	2	-
Female	12	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	28	1	-	-	-	1	-	2	-	1	2	-
Male	18	-	-	-	-	1	-	1	-	1	2	-
Female	10	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Florida												
All races, both sexes	2,263	7	5	1	2	3	15	17	25	28	46	88
Male	1,311	3	3	1	-	1	12	11	17	21	32	55
Female	952	4	2	-	2	2	3	6	8	7	14	33
White, both sexes	1,910	5	2	-	-	3	9	9	15	11	27	62
Male	1,110	2	1	-	-	1	8	7	9	9	22	37
Female	800	3	1	-	-	2	1	2	6	2	5	25
Black, both sexes	345	2	2	1	2	-	6	8	10	17	19	26
Male	194	1	2	1	-	-	4	4	8	12	10	18
Female	151	1	-	-	2	-	2	4	2	5	9	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	824	6	2	-	3	3	4	12	12	22	24	30
Male	487	5	1	-	1	2	1	9	7	14	16	24
Female	337	1	1	-	2	1	3	3	5	8	8	6
White, both sexes	511	4	1	-	1	2	1	6	3	6	9	15
Male	301	3	1	-	-	1	1	5	2	6	7	12
Female	210	1	-	-	1	1	-	1	1	-	2	3
Black, both sexes	307	2	1	-	2	1	3	6	9	16	15	15
Male	182	2	-	-	1	1	-	4	5	8	9	12
Female	125	-	1	-	1	-	3	2	4	8	6	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Hawaii												
All races, both sexes	164	-	1	1	-	-	2	2	1	4	5	6
Male	109	-	-	1	-	-	2	1	1	3	2	4
Female	55	-	1	-	-	-	-	1	-	1	3	2
White, both sexes	38	-	-	-	-	-	-	-	-	1	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	124	-	1	1	-	-	2	2	1	3	5	6
Male	88	-	-	1	-	-	2	1	1	3	2	4
Female	36	-	1	-	-	-	-	1	-	-	3	2
Idaho												
All races, both sexes	50	-	-	-	-	-	-	1	-	-	-	3
Male	30	-	-	-	-	-	-	1	-	-	-	3
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	1	-	-	-	3
Male	29	-	-	-	-	-	-	1	-	-	-	3
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Illinois												
All races, both sexes	1,031	2	1	-	1	13	7	16	17	27	37	42
Male	605	2	-	-	1	12	5	12	11	18	24	28
Female	426	-	1	-	-	1	2	4	6	9	13	14
White, both sexes	781	1	1	-	-	7	4	11	7	12	22	19
Male	450	1	-	-	-	6	3	9	5	8	14	14
Female	331	-	1	-	-	1	1	2	2	4	8	5
Black, both sexes	240	1	-	-	1	6	3	5	9	15	15	23
Male	149	1	-	-	1	6	2	3	6	10	10	14
Female	91	-	-	-	-	-	1	2	3	5	5	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	581	7	-	1	-	4	5	3	9	8	18	27
Male	327	3	-	-	-	4	3	2	4	6	12	17
Female	254	4	-	1	-	-	2	1	5	2	6	10
White, both sexes	516	2	-	1	-	4	5	3	4	8	14	23
Male	289	1	-	-	-	4	3	2	2	6	10	13
Female	227	1	-	1	-	-	2	1	2	2	4	10
Black, both sexes	65	5	-	-	-	-	-	-	5	-	4	4
Male	38	2	-	-	-	-	-	-	2	-	2	4
Female	27	3	-	-	-	-	-	-	3	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Iowa												
All races, both sexes	187	-	-	-	-	1	1	1	-	1	8	7
Male	100	-	-	-	-	1	1	1	-	-	5	4
Female	87	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	180	-	-	-	-	1	1	1	-	1	7	7
Male	95	-	-	-	-	1	1	1	-	-	4	4
Female	85	-	-	-	-	-	-	-	-	1	3	3
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	201	-	-	-	-	1	-	2	1	2	9	5
Male	97	-	-	-	-	-	-	1	1	-	4	3
Female	104	-	-	-	-	1	-	1	-	2	5	2
White, both sexes	176	-	-	-	-	1	-	2	-	1	7	5
Male	84	-	-	-	-	-	-	1	-	-	3	3
Female	92	-	-	-	-	1	-	1	-	1	4	2
Black, both sexes	22	-	-	-	-	-	-	-	-	1	2	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Kentucky												
All races, both sexes	319	-	-	-	-	-	-	3	3	4	5	11
Male	186	-	-	-	-	-	-	2	3	3	5	6
Female	133	-	-	-	-	-	-	1	-	1	-	5
White, both sexes	295	-	-	-	-	-	-	2	3	3	4	8
Male	172	-	-	-	-	-	-	1	3	3	4	4
Female	123	-	-	-	-	-	-	1	-	-	-	4
Black, both sexes	24	-	-	-	-	-	-	1	-	1	1	3
Male	14	-	-	-	-	-	-	1	-	-	1	2
Female	10	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	480	5	-	-	-	1	5	8	11	6	15	15
Male	258	4	-	-	-	-	3	5	6	5	8	10
Female	222	1	-	-	-	1	2	3	5	1	7	5
White, both sexes	331	2	-	-	-	-	3	4	6	4	8	7
Male	183	2	-	-	-	-	2	3	3	3	5	5
Female	148	-	-	-	-	-	1	1	3	1	3	2
Black, both sexes	148	3	-	-	-	1	2	4	5	2	7	8
Male	74	2	-	-	-	-	1	2	3	2	3	5
Female	74	1	-	-	-	1	1	2	2	-	4	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Maine												
All races, both sexes	122	-	-	-	1	2	-	1	2	2	3	8
Male	77	-	-	-	1	2	-	1	2	1	2	7
Female	45	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	120	-	-	-	1	2	-	1	2	2	3	8
Male	75	-	-	-	1	2	-	1	2	1	2	7
Female	45	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	523	1	-	-	3	-	1	4	6	14	14	16
Male	285	1	-	-	2	-	1	4	5	9	10	10
Female	238	-	-	-	1	-	-	-	1	5	4	6
White, both sexes	357	1	-	-	1	-	-	2	2	5	5	10
Male	197	1	-	-	1	-	-	2	1	5	5	6
Female	160	-	-	-	-	-	-	-	1	-	-	4
Black, both sexes	163	-	-	-	2	-	1	2	4	9	9	6
Male	86	-	-	-	1	-	1	2	4	4	5	4
Female	77	-	-	-	1	-	-	-	-	5	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Massachusetts												
All races, both sexes	417	1	-	-	1	1	2	3	4	5	5	16
Male	239	1	-	-	1	1	2	3	3	3	4	13
Female	178	-	-	-	-	-	-	-	1	2	1	3
White, both sexes	394	1	-	-	-	1	2	3	3	3	4	14
Male	227	1	-	-	-	1	2	3	3	2	3	11
Female	167	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	20	-	-	-	1	-	-	-	1	2	1	1
Male	11	-	-	-	1	-	-	-	-	1	1	1
Female	9	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,046	2	2	2	1	2	11	13	18	41	50	70
Male	586	2	2	1	-	2	9	10	15	29	39	41
Female	460	-	-	1	1	-	2	3	3	12	11	29
White, both sexes	822	1	1	2	1	1	4	7	8	23	30	41
Male	456	1	1	1	-	1	4	5	7	18	26	27
Female	366	-	-	1	1	-	-	2	1	5	4	14
Black, both sexes	217	1	1	-	-	1	6	6	10	18	19	29
Male	125	1	1	-	-	1	5	5	8	11	12	14
Female	92	-	-	-	-	-	1	1	2	7	7	15
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Minnesota												
All races, both sexes	270	-	1	-	-	1	1	2	4	3	7	12
Male	150	-	-	-	-	1	1	1	3	1	2	10
Female	120	-	1	-	-	-	-	1	1	2	5	2
White, both sexes	258	-	1	-	-	1	1	2	3	3	6	7
Male	143	-	-	-	-	1	1	1	3	1	1	6
Female	115	-	1	-	-	-	-	1	-	2	5	1
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	4
Male	5	-	-	-	-	-	-	-	-	-	-	3
Female	3	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	226	2	-	1	3	1	3	5	8	1	7	9
Male	116	1	-	-	3	1	1	3	4	1	5	6
Female	110	1	-	1	-	-	2	2	4	-	2	3
White, both sexes	124	2	-	-	1	-	1	1	1	-	-	-
Male	65	1	-	-	1	-	1	1	1	-	-	-
Female	59	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	101	-	-	1	2	1	2	4	7	1	7	9
Male	50	-	-	-	2	1	-	2	3	1	5	6
Female	51	-	-	1	-	-	2	2	4	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Missouri												
All races, both sexes	539	-	-	-	2	2	4	3	6	9	9	20
Male	294	-	-	-	1	1	2	3	2	6	6	17
Female	245	-	-	-	1	1	2	-	4	3	3	3
White, both sexes	448	-	-	-	2	2	1	3	2	5	6	12
Male	242	-	-	-	1	1	-	3	1	4	4	10
Female	206	-	-	-	1	1	1	-	1	1	2	2
Black, both sexes	88	-	-	-	-	-	3	-	4	4	3	8
Male	49	-	-	-	-	-	2	-	1	2	2	7
Female	39	-	-	-	-	-	1	-	3	2	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	56	-	-	-	-	-	-	1	-	1	2	-
Male	39	-	-	-	-	-	-	1	-	1	2	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	2	-
Male	36	-	-	-	-	-	-	-	-	-	2	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Nebraska												
All races, both sexes	114	-	-	-	-	-	-	1	1	2	1	3
Male	69	-	-	-	-	-	-	1	-	2	1	2
Female	45	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	100	-	-	-	-	-	-	-	1	2	-	2
Male	60	-	-	-	-	-	-	-	-	2	-	1
Female	40	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	13	-	-	-	-	-	-	1	-	-	1	1
Male	9	-	-	-	-	-	-	1	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	238	1	1	-	-	-	1	3	8	7	14	11
Male	166	1	1	-	-	-	1	2	7	5	8	8
Female	72	-	-	-	-	-	-	1	1	2	6	3
White, both sexes	198	1	-	-	-	-	-	2	6	6	11	10
Male	142	1	-	-	-	-	-	1	5	4	7	7
Female	56	-	-	-	-	-	-	1	1	2	4	3
Black, both sexes	26	-	1	-	-	-	-	-	-	1	2	1
Male	17	-	1	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	1	-	-	-
Male	4	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	1	-	1	-	1	-
Male	3	-	-	-	-	-	1	-	1	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
New Hampshire												
All races, both sexes	76	-	1	-	-	1	2	-	-	2	3	2
Male	44	-	1	-	-	-	2	-	-	1	1	1
Female	32	-	-	-	-	1	-	-	-	1	2	1
White, both sexes	76	-	1	-	-	1	2	-	-	2	3	2
Male	44	-	1	-	-	-	2	-	-	1	1	1
Female	32	-	-	-	-	1	-	-	-	1	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	841	1	2	-	1	1	1	10	16	12	19	32
Male	509	1	1	-	1	1	-	10	15	10	17	28
Female	332	-	1	-	-	-	1	-	1	2	2	4
White, both sexes	682	1	1	-	1	1	1	4	9	8	13	14
Male	398	1	-	-	1	1	-	4	9	6	11	13
Female	284	-	1	-	-	-	1	-	-	2	2	1
Black, both sexes	150	-	1	-	-	-	-	6	6	3	6	18
Male	105	-	1	-	-	-	-	6	5	3	6	15
Female	45	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
New Mexico												
All races, both sexes	119	1	-	-	-	-	2	1	4	3	5	6
Male	66	1	-	-	-	-	1	1	3	3	3	5
Female	53	-	-	-	-	-	1	-	1	-	2	1
White, both sexes	106	1	-	-	-	-	2	1	3	2	3	5
Male	60	1	-	-	-	-	1	1	3	2	1	4
Female	46	-	-	-	-	-	1	-	-	-	2	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	1	2	-
Male	3	-	-	-	-	-	-	-	-	1	2	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,163	3	2	1	3	6	2	14	15	23	45	70
Male	694	3	2	1	2	5	1	12	10	17	29	49
Female	469	-	-	-	1	1	1	2	5	6	16	21
White, both sexes	1,002	3	1	-	2	4	2	8	11	17	31	50
Male	593	3	1	-	1	3	1	7	6	12	20	32
Female	409	-	-	-	1	1	1	1	5	5	11	18
Black, both sexes	142	-	1	1	1	2	-	6	4	6	12	18
Male	89	-	1	1	1	2	-	5	4	5	8	15
Female	53	-	-	-	-	-	-	1	-	1	4	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	2	2
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
North Carolina												
All races, both sexes	889	1	3	2	5	5	10	10	6	18	34	33
Male	532	-	1	2	4	4	6	6	5	15	25	25
Female	357	1	2	-	1	1	4	4	1	3	9	8
White, both sexes	596	1	3	1	-	3	4	5	2	7	20	20
Male	355	-	1	1	-	3	2	3	2	6	15	15
Female	241	1	2	-	-	-	2	2	-	1	5	5
Black, both sexes	281	-	-	1	5	2	5	5	4	11	13	13
Male	170	-	-	1	4	1	4	3	3	9	9	10
Female	111	-	-	-	1	1	1	2	1	2	4	3
American Indian 1/, both sexes	6	-	-	-	-	-	1	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	40	-	-	-	1	-	1	-	1	-	-	-
Male	22	-	-	-	1	-	-	-	1	-	-	-
Female	18	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	38	-	-	-	1	-	1	-	-	-	-	-
Male	21	-	-	-	1	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Ohio												
All races, both sexes	1,046	9	3	1	-	5	4	12	12	18	30	51
Male	593	3	1	1	-	5	2	8	10	13	19	35
Female	453	6	2	-	-	-	2	4	2	5	11	16
White, both sexes	867	7	2	1	-	4	4	7	9	11	24	35
Male	481	2	1	1	-	4	2	4	8	8	15	26
Female	386	5	1	-	-	-	2	3	1	3	9	9
Black, both sexes	179	2	1	-	-	1	-	5	3	7	6	16
Male	112	1	-	-	-	1	-	4	2	5	4	9
Female	67	1	1	-	-	-	-	1	1	2	2	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	278	2	-	-	-	-	5	7	9	3	12	22
Male	164	2	-	-	-	-	3	7	8	2	8	17
Female	114	-	-	-	-	-	2	-	1	1	4	5
White, both sexes	235	1	-	-	-	-	4	5	5	2	11	15
Male	132	1	-	-	-	-	2	5	4	2	7	11
Female	103	-	-	-	-	-	2	-	1	-	4	4
Black, both sexes	31	-	-	-	-	-	1	2	3	1	-	6
Male	21	-	-	-	-	-	1	2	3	-	-	5
Female	10	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	12	1	-	-	-	-	-	-	1	-	1	1
Male	11	1	-	-	-	-	-	-	1	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Oregon												
All races, both sexes	209	1	-	-	-	-	2	3	4	5	6	9
Male	120	-	-	-	-	-	-	1	4	4	4	6
Female	89	1	-	-	-	-	2	2	-	1	2	3
White, both sexes	197	1	-	-	-	-	2	3	2	5	5	8
Male	113	-	-	-	-	-	-	1	2	4	4	5
Female	84	1	-	-	-	-	2	2	-	1	1	3
Black, both sexes	6	-	-	-	-	-	-	-	2	-	-	-
Male	5	-	-	-	-	-	-	-	2	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,476	6	1	-	2	4	3	13	12	21	41	59
Male	833	3	-	-	2	2	3	9	9	15	31	45
Female	643	3	1	-	-	2	-	4	3	6	10	14
White, both sexes	1,293	6	-	-	1	3	2	5	6	13	32	43
Male	718	3	-	-	1	2	2	4	4	9	22	31
Female	575	3	-	-	-	1	-	1	2	4	10	12
Black, both sexes	180	-	1	-	1	1	1	8	6	8	9	15
Male	113	-	-	-	1	-	1	5	5	6	9	13
Female	67	-	1	-	-	1	-	3	1	2	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Rhode Island												
All races, both sexes	45	-	-	-	-	1	-	2	-	1	1	3
Male	26	-	-	-	-	1	-	2	-	1	1	2
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	1	-	1	-	1	1	3
Male	25	-	-	-	-	1	-	1	-	1	1	2
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	422	4	1	-	-	2	2	4	6	5	11	20
Male	240	3	-	-	-	1	1	4	1	4	9	13
Female	182	1	1	-	-	1	1	-	5	1	2	7
White, both sexes	261	1	-	-	-	1	-	2	3	2	6	11
Male	155	1	-	-	-	1	-	2	1	1	4	8
Female	106	-	-	-	-	-	-	-	2	1	2	3
Black, both sexes	161	3	1	-	-	1	2	2	3	3	5	9
Male	85	2	-	-	-	-	1	2	-	3	5	5
Female	76	1	1	-	-	1	1	-	3	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
South Dakota												
All races, both sexes	41	1	-	-	-	-	-	-	-	1	-	1
Male	24	1	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	40	-	-	-	-	-	-	-	-	1	-	1
Male	23	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	488	4	1	-	-	3	5	1	7	9	25	16
Male	284	2	-	-	-	2	3	-	4	5	15	10
Female	204	2	1	-	-	1	2	1	3	4	10	6
White, both sexes	348	4	1	-	-	1	1	-	3	5	11	9
Male	192	2	-	-	-	-	1	-	1	3	4	7
Female	156	2	1	-	-	1	-	-	2	2	7	2
Black, both sexes	135	-	-	-	-	1	4	1	4	4	14	7
Male	89	-	-	-	-	1	2	-	3	2	11	3
Female	46	-	-	-	-	-	2	1	1	2	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Texas												
All races, both sexes	1,377	12	3	3	3	8	11	17	34	35	54	90
Male	800	6	2	2	3	5	9	14	25	25	40	60
Female	577	6	1	1	-	3	2	3	9	10	14	30
White, both sexes	1,107	11	1	1	2	7	7	10	21	24	38	59
Male	630	5	1	1	2	4	6	7	16	17	29	36
Female	477	6	-	-	-	3	1	3	5	7	9	23
Black, both sexes	256	-	2	2	1	1	4	7	13	11	16	28
Male	161	-	1	1	1	1	3	7	9	8	11	21
Female	95	-	1	1	-	-	1	-	4	3	5	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	1	-	-	-	-	-	-	-	-	-	3
Male	8	1	-	-	-	-	-	-	-	-	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	78	1	-	-	-	-	1	-	2	-	1	4
Male	47	-	-	-	-	-	1	-	2	-	1	4
Female	31	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	76	1	-	-	-	-	1	-	2	-	-	4
Male	45	-	-	-	-	-	1	-	2	-	-	4
Female	31	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Vermont												
All races, both sexes	46	-	-	-	-	1	-	-	-	2	1	1
Male	26	-	-	-	-	-	-	-	-	2	-	-
Female	20	-	-	-	-	1	-	-	-	-	1	1
White, both sexes	45	-	-	-	-	1	-	-	-	2	1	1
Male	25	-	-	-	-	-	-	-	-	2	-	-
Female	20	-	-	-	-	1	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	692	3	3	1	-	3	5	2	8	14	21	37
Male	387	1	2	1	-	2	5	2	5	11	15	30
Female	305	2	1	-	-	1	-	-	3	3	6	7
White, both sexes	482	2	2	1	-	2	4	1	2	8	12	16
Male	267	1	1	1	-	2	4	1	2	7	8	12
Female	215	1	1	-	-	-	-	-	-	1	4	4
Black, both sexes	198	1	1	-	-	1	-	1	6	6	9	20
Male	110	-	1	-	-	-	-	1	3	4	7	17
Female	88	1	-	-	-	1	-	-	3	2	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	1	-	-	-	-	1
Male	10	-	-	-	-	-	1	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Washington												
All races, both sexes	400	2	-	-	-	2	-	4	5	13	10	21
Male	235	1	-	-	-	2	-	3	2	10	6	15
Female	165	1	-	-	-	-	-	1	3	3	4	6
White, both sexes	352	1	-	-	-	2	-	3	4	7	7	18
Male	205	-	-	-	-	2	-	3	2	4	5	12
Female	147	1	-	-	-	-	-	-	2	3	2	6
Black, both sexes	29	1	-	-	-	-	-	1	1	4	1	2
Male	20	1	-	-	-	-	-	-	-	4	-	2
Female	9	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	145	-	-	-	-	-	3	-	2	1	3	3
Male	93	-	-	-	-	-	2	-	2	-	2	2
Female	52	-	-	-	-	-	1	-	-	1	1	1
White, both sexes	138	-	-	-	-	-	3	-	2	1	3	3
Male	89	-	-	-	-	-	2	-	2	-	2	2
Female	49	-	-	-	-	-	1	-	-	1	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Wisconsin												
All races, both sexes	492	2	-	-	1	1	3	2	6	11	15	14
Male	252	1	-	-	-	1	3	-	3	9	9	10
Female	240	1	-	-	1	-	-	2	3	2	6	4
White, both sexes	448	2	-	-	-	-	3	2	3	9	14	13
Male	232	1	-	-	-	-	3	-	1	7	9	10
Female	216	1	-	-	-	-	-	2	2	2	5	3
Black, both sexes	32	-	-	-	-	1	-	-	3	1	1	1
Male	16	-	-	-	-	1	-	-	2	1	-	-
Female	16	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	1	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	23	-	-	-	-	-	-	-	1	-	1	-
Male	13	-	-	-	-	-	-	-	1	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	1	-	1	-
Male	13	-	-	-	-	-	-	-	1	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
United States												
All races, both sexes	1,325	1,568	1,498	1,690	2,140	2,854	3,557	3,370	2,182	789	119	2
Male	916	1,093	1,004	1,112	1,359	1,701	1,978	1,603	838	229	24	2
Female	409	475	494	578	781	1,153	1,579	1,767	1,344	560	95	-
White, both sexes	889	1,097	1,093	1,303	1,713	2,429	3,133	3,054	1,991	720	101	1
Male	620	769	752	872	1,098	1,486	1,774	1,459	777	208	21	1
Female	269	328	341	431	615	943	1,359	1,595	1,214	512	80	-
Black, both sexes	409	422	364	335	368	357	367	257	153	59	16	1
Male	273	287	228	205	221	176	174	106	47	18	3	1
Female	136	135	136	130	147	181	193	151	106	41	13	-
American Indian 1/, both sexes	7	12	14	11	10	10	6	3	1	-	-	-
Male	5	8	8	6	6	6	4	3	-	-	-	-
Female	2	4	6	5	4	4	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	20	37	27	41	49	58	51	56	37	10	2	-
Male	18	29	16	29	34	33	26	35	14	3	-	-
Female	2	8	11	12	15	25	25	21	23	7	2	-
Alabama												
All races, both sexes	27	23	18	32	42	59	71	68	38	6	4	-
Male	21	16	11	20	30	38	43	27	13	-	1	-
Female	6	7	7	12	12	21	28	41	25	6	3	-
White, both sexes	12	15	10	16	31	47	52	58	32	4	3	-
Male	11	11	5	12	24	30	31	24	11	-	1	-
Female	1	4	5	4	7	17	21	34	21	4	2	-
Black, both sexes	15	8	8	16	11	12	19	10	6	2	1	-
Male	10	5	6	8	6	8	12	3	2	-	-	-
Female	5	3	2	8	5	4	7	7	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Alaska												
All races, both sexes	3	4	2	5	9	2	2	3	2	1	-	-
Male	3	3	2	4	6	2	1	1	1	-	-	-
Female	-	1	-	1	3	-	1	2	1	1	-	-
White, both sexes	3	3	-	2	8	2	2	2	2	1	-	-
Male	3	2	-	1	5	2	1	1	1	-	-	-
Female	-	1	-	1	3	-	1	1	1	1	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	20	22	28	22	27	25	43	32	16	7	-	-
Male	15	15	16	18	21	15	21	15	11	4	-	-
Female	5	7	12	4	6	10	22	17	5	3	-	-
White, both sexes	16	17	25	20	26	24	40	32	16	7	-	-
Male	12	11	14	16	21	15	18	15	11	4	-	-
Female	4	6	11	4	5	9	22	17	5	3	-	-
Black, both sexes	2	3	1	2	-	1	3	-	-	-	-	-
Male	2	3	1	2	-	-	3	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	2	2	2	-	1	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Arkansas												
All races, both sexes	8	9	14	18	25	26	27	30	29	4	1	-
Male	7	6	8	11	18	19	14	14	11	2	-	-
Female	1	3	6	7	7	7	13	16	18	2	1	-
White, both sexes	6	6	12	18	16	22	23	28	25	2	1	-
Male	5	3	7	11	12	17	12	13	9	1	-	-
Female	1	3	5	7	4	5	11	15	16	1	1	-
Black, both sexes	2	3	2	-	9	4	4	2	4	2	-	-
Male	2	3	1	-	6	2	2	1	2	1	-	-
Female	-	-	1	-	3	2	2	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	176	204	179	186	241	293	399	392	217	87	19	-
Male	124	151	128	129	159	176	227	202	93	29	2	-
Female	52	53	51	57	82	117	172	190	124	58	17	-
White, both sexes	123	151	135	133	197	238	336	344	187	78	15	-
Male	85	116	103	97	129	150	195	177	83	26	2	-
Female	38	35	32	36	68	88	141	167	104	52	13	-
Black, both sexes	44	32	33	32	29	29	28	24	14	6	3	-
Male	31	18	17	18	18	12	17	12	6	2	-	-
Female	13	14	16	14	11	17	11	12	8	4	3	-
American Indian 1/, both sexes	-	1	2	2	2	1	1	-	1	-	-	-
Male	-	1	2	2	2	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	9	20	9	19	13	25	34	24	15	3	1	-
Male	8	16	6	12	10	13	14	13	4	1	-	-
Female	1	4	3	7	3	12	20	11	11	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Colorado												
All races, both sexes	18	15	10	12	23	22	34	24	19	9	1	-
Male	13	12	6	11	17	16	14	7	8	1	-	-
Female	5	3	4	1	6	6	20	17	11	8	1	-
White, both sexes	17	12	9	11	20	20	33	23	19	9	1	-
Male	12	9	5	10	15	14	14	7	8	1	-	-
Female	5	3	4	1	5	6	19	16	11	8	1	-
Black, both sexes	-	2	1	-	1	1	1	1	-	-	-	-
Male	-	2	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	-	-	-	-	-	-	-
Male	-	1	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	10	13	22	21	21	34	51	46	32	12	2	-
Male	9	9	19	14	16	24	29	21	13	3	-	-
Female	1	4	3	7	5	10	22	25	19	9	2	-
White, both sexes	9	10	19	17	20	33	51	43	32	12	2	-
Male	8	7	17	10	15	24	29	19	13	3	-	-
Female	1	3	2	7	5	9	22	24	19	9	2	-
Black, both sexes	1	3	3	4	1	1	-	2	-	-	-	-
Male	1	2	2	4	1	-	-	1	-	-	-	-
Female	-	1	1	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Delaware												
All races, both sexes	8	9	6	4	7	11	13	14	10	4	1	-
Male	6	4	3	3	3	8	7	6	5	1	1	-
Female	2	5	3	1	4	3	6	8	5	3	-	-
White, both sexes	7	7	3	3	5	9	12	11	10	4	1	-
Male	5	3	1	2	3	6	7	6	5	1	1	-
Female	2	4	2	1	2	3	5	5	5	3	-	-
Black, both sexes	1	2	3	1	2	2	1	2	-	-	-	-
Male	1	1	2	1	-	2	-	-	-	-	-	-
Female	-	1	1	-	2	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	1	1	2	4	4	5	2	1	-	-	-
Male	4	1	1	1	2	3	1	2	-	-	-	-
Female	-	-	-	1	2	1	4	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	1	1	2	4	3	5	1	-	-	-	-
Male	4	1	1	1	2	2	1	1	-	-	-	-
Female	-	-	-	1	2	1	4	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Florida												
All races, both sexes	107	117	109	156	186	291	352	345	240	109	14	-
Male	68	83	80	103	115	177	199	178	111	37	4	-
Female	39	34	29	53	71	114	153	167	129	72	10	-
White, both sexes	60	92	84	132	157	259	316	320	230	105	12	-
Male	38	67	62	87	98	161	182	171	109	35	4	-
Female	22	25	22	45	59	98	134	149	121	70	8	-
Black, both sexes	46	24	25	22	28	32	36	23	10	4	2	-
Male	29	15	18	14	16	16	17	5	2	2	-	-
Female	17	9	7	8	12	16	19	18	8	2	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	-	-	2	-	-	-	-
Male	1	1	-	1	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	52	75	64	60	86	96	104	107	44	17	1	-
Male	37	50	35	43	56	53	61	45	17	10	-	-
Female	15	25	29	17	30	43	43	62	27	7	1	-
White, both sexes	21	33	30	41	57	68	75	85	38	14	1	-
Male	15	25	18	29	36	40	44	32	17	7	-	-
Female	6	8	12	12	21	28	31	53	21	7	1	-
Black, both sexes	31	42	34	19	28	26	29	20	5	3	-	-
Male	22	25	17	14	19	12	17	11	-	3	-	-
Female	9	17	17	5	9	14	12	9	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	2	1	-	-	-
Male	-	-	-	-	1	1	-	2	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Hawaii												
All races, both sexes	11	8	14	6	21	22	12	24	15	8	1	-
Male	10	6	9	4	16	17	8	14	9	2	-	-
Female	1	2	5	2	5	5	4	10	6	6	1	-
White, both sexes	3	4	1	2	6	5	3	7	5	1	-	-
Male	3	3	1	-	5	2	2	1	3	-	-	-
Female	-	1	-	2	1	3	1	6	2	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	4	12	4	15	17	9	17	10	7	-	-
Male	7	3	7	4	11	15	6	13	6	2	-	-
Female	1	1	5	-	4	2	3	4	4	5	-	-
Idaho												
All races, both sexes	1	4	4	3	8	7	7	7	4	1	-	-
Male	1	2	2	3	4	6	4	3	1	-	-	-
Female	-	2	2	-	4	1	3	4	3	1	-	-
White, both sexes	1	4	4	3	8	6	7	7	4	1	-	-
Male	1	2	2	3	4	5	4	3	1	-	-	-
Female	-	2	2	-	4	1	3	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Illinois												
All races, both sexes	58	65	53	73	106	108	129	130	113	28	5	-
Male	45	48	37	47	67	62	80	60	34	10	2	-
Female	13	17	16	26	39	46	49	70	79	18	3	-
White, both sexes	34	38	40	57	75	82	114	120	107	26	4	-
Male	29	29	28	37	47	48	72	55	33	10	2	-
Female	5	9	12	20	28	34	42	65	74	16	2	-
Black, both sexes	24	27	13	15	28	25	14	8	6	2	-	-
Male	16	19	9	9	18	13	7	4	1	-	-	-
Female	8	8	4	6	10	12	7	4	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	3	1	1	2	-	-	1	-
Male	-	-	-	1	2	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	1	-
Indiana												
All races, both sexes	25	35	39	50	50	69	94	75	48	14	-	-
Male	18	18	27	32	31	40	56	32	18	4	-	-
Female	7	17	12	18	19	29	38	43	30	10	-	-
White, both sexes	19	26	33	42	47	62	88	74	47	14	-	-
Male	15	13	23	26	28	36	53	32	18	4	-	-
Female	4	13	10	16	19	26	35	42	29	10	-	-
Black, both sexes	6	9	6	8	3	7	6	1	1	-	-	-
Male	3	5	4	6	3	4	3	-	-	-	-	-
Female	3	4	2	2	-	3	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Iowa												
All races, both sexes	6	15	11	7	12	22	30	27	28	10	-	-
Male	3	12	8	5	6	15	18	9	10	2	-	-
Female	3	3	3	2	6	7	12	18	18	8	-	-
White, both sexes	6	14	11	6	10	22	28	27	28	10	-	-
Male	3	11	8	4	5	15	17	9	10	2	-	-
Female	3	3	3	2	5	7	11	18	18	8	-	-
Black, both sexes	-	1	-	1	2	-	2	-	-	-	-	-
Male	-	1	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	8	9	9	10	22	23	28	32	19	19	2	-
Male	6	8	5	3	14	13	13	17	6	3	-	-
Female	2	1	4	7	8	10	15	15	13	16	2	-
White, both sexes	6	6	8	8	19	19	25	30	19	18	2	-
Male	4	5	5	3	12	11	12	16	6	3	-	-
Female	2	1	3	5	7	8	13	14	13	15	2	-
Black, both sexes	2	3	1	2	3	3	3	1	-	1	-	-
Male	2	3	-	-	2	1	1	-	-	-	-	-
Female	-	-	1	2	1	2	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Kentucky												
All races, both sexes	17	23	19	45	23	37	55	49	16	9	-	-
Male	12	15	13	31	12	23	28	23	7	3	-	-
Female	5	8	6	14	11	14	27	26	9	6	-	-
White, both sexes	13	18	16	44	23	36	53	47	16	9	-	-
Male	8	12	12	31	12	22	28	22	7	3	-	-
Female	5	6	4	13	11	14	25	25	9	6	-	-
Black, both sexes	4	5	3	1	-	1	2	2	-	-	-	-
Male	4	3	1	-	-	1	-	1	-	-	-	-
Female	-	2	2	1	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	19	34	38	34	45	55	70	60	45	14	-	-
Male	13	26	23	21	22	36	36	22	16	2	-	-
Female	6	8	15	13	23	19	34	38	29	12	-	-
White, both sexes	10	14	20	24	26	42	59	50	41	11	-	-
Male	7	10	13	18	14	30	31	19	16	2	-	-
Female	3	4	7	6	12	12	28	31	25	9	-	-
Black, both sexes	9	19	18	10	19	13	11	10	4	3	-	-
Male	6	15	10	3	8	6	5	3	-	-	-	-
Female	3	4	8	7	11	7	6	7	4	3	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Maine												
All races, both sexes	11	6	3	3	10	18	22	13	9	6	2	-
Male	7	6	2	2	9	12	13	7	1	1	1	-
Female	4	-	1	1	1	6	9	6	8	5	1	-
White, both sexes	10	5	3	3	10	18	22	13	9	6	2	-
Male	6	5	2	2	9	12	13	7	1	1	1	-
Female	4	-	1	1	1	6	9	6	8	5	1	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	24	42	32	37	49	52	59	78	71	19	1	-
Male	13	29	25	21	32	28	25	37	27	5	1	-
Female	11	13	7	16	17	24	34	41	44	14	-	-
White, both sexes	8	24	17	21	30	39	46	67	61	18	-	-
Male	5	16	14	12	22	24	21	34	23	5	-	-
Female	3	8	3	9	8	15	25	33	38	13	-	-
Black, both sexes	16	18	14	15	19	13	12	11	10	1	1	-
Male	8	13	10	8	10	4	4	3	4	-	1	-
Female	8	5	4	7	9	9	8	8	6	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Massachusetts												
All races, both sexes	21	14	21	30	21	53	75	78	47	16	3	-
Male	17	12	11	24	11	30	44	39	15	5	-	-
Female	4	2	10	6	10	23	31	39	32	11	3	-
White, both sexes	18	13	20	29	21	51	75	75	44	14	3	-
Male	15	11	10	23	11	30	44	38	14	5	-	-
Female	3	2	10	6	10	21	31	37	30	9	3	-
Black, both sexes	3	1	1	1	-	-	-	3	3	2	-	-
Male	2	1	1	1	-	-	-	1	1	-	-	-
Female	1	-	-	-	-	-	-	2	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	54	68	54	56	84	114	133	136	91	34	10	-
Male	36	46	33	30	48	65	76	55	34	12	1	-
Female	18	22	21	26	36	49	57	81	57	22	9	-
White, both sexes	31	50	42	47	66	96	116	127	88	31	9	-
Male	19	35	26	26	39	56	66	54	32	11	1	-
Female	12	15	16	21	27	40	50	73	56	20	8	-
Black, both sexes	22	17	12	9	17	17	17	9	2	3	1	-
Male	17	10	7	4	8	8	10	1	1	1	-	-
Female	5	7	5	5	9	9	7	8	1	2	1	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Minnesota												
All races, both sexes	9	13	23	15	20	32	35	45	31	14	2	-
Male	5	9	17	7	12	21	21	19	15	5	-	-
Female	4	4	6	8	8	11	14	26	16	9	2	-
White, both sexes	8	13	22	14	20	32	35	44	30	14	2	-
Male	4	9	17	7	12	21	21	18	15	5	-	-
Female	4	4	5	7	8	11	14	26	15	9	2	-
Black, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Mississippi												
All races, both sexes	14	6	13	21	18	24	35	31	20	3	1	-
Male	8	4	6	14	12	10	16	12	8	1	-	-
Female	6	2	7	7	6	14	19	19	12	2	1	-
White, both sexes	7	1	9	14	10	16	26	23	9	2	1	-
Male	5	-	5	8	7	7	13	10	4	1	-	-
Female	2	1	4	6	3	9	13	13	5	1	1	-
Black, both sexes	7	5	4	7	8	7	9	8	11	1	-	-
Male	3	4	1	6	5	2	3	2	4	-	-	-
Female	4	1	3	1	3	5	6	6	7	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Missouri												
All races, both sexes	30	35	29	36	52	68	88	75	49	21	1	-
Male	19	15	15	22	36	41	49	32	22	5	-	-
Female	11	20	14	14	16	27	39	43	27	16	1	-
White, both sexes	20	23	23	28	47	57	80	70	46	20	1	-
Male	13	10	12	16	33	32	45	31	21	5	-	-
Female	7	13	11	12	14	25	35	39	25	15	1	-
Black, both sexes	10	12	4	8	4	11	8	5	3	1	-	-
Male	6	5	1	6	2	9	4	1	1	-	-	-
Female	4	7	3	2	2	2	4	4	2	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	2	5	3	5	10	12	6	3	-	-	-
Male	6	1	4	1	3	7	6	4	3	-	-	-
Female	-	1	1	2	2	3	6	2	-	-	-	-
White, both sexes	6	2	5	3	4	10	12	5	3	-	-	-
Male	6	1	4	1	3	7	6	3	3	-	-	-
Female	-	1	1	2	1	3	6	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Nebraska												
All races, both sexes	5	6	7	8	14	19	17	17	9	4	-	-
Male	4	5	6	3	10	11	7	9	7	1	-	-
Female	1	1	1	5	4	8	10	8	2	3	-	-
White, both sexes	5	4	4	7	14	17	15	16	9	4	-	-
Male	4	4	3	3	10	11	6	8	7	1	-	-
Female	1	-	1	4	4	6	9	8	2	3	-	-
Black, both sexes	-	1	3	1	-	2	2	1	-	-	-	-
Male	-	1	3	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	10	23	21	30	28	35	23	11	2	-	-
Male	7	8	17	11	17	25	28	15	4	1	-	-
Female	2	2	6	10	13	3	7	8	7	1	-	-
White, both sexes	7	6	20	15	22	28	32	21	9	2	-	-
Male	5	5	15	10	14	25	25	13	4	1	-	-
Female	2	1	5	5	8	3	7	8	5	1	-	-
Black, both sexes	2	3	3	3	4	-	3	1	2	-	-	-
Male	2	3	2	1	2	-	3	1	-	-	-	-
Female	-	-	1	2	2	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	3	3	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	3	3	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
New Hampshire												
All races, both sexes	4	6	6	1	3	14	6	10	12	2	1	-
Male	2	5	5	1	3	11	1	6	4	-	-	-
Female	2	1	1	-	-	3	5	4	8	2	1	-
White, both sexes	4	6	6	1	3	14	6	10	12	2	1	-
Male	2	5	5	1	3	11	1	6	4	-	-	-
Female	2	1	1	-	-	3	5	4	8	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	40	60	47	49	71	99	130	138	78	31	3	-
Male	34	43	34	31	51	56	69	73	24	9	1	-
Female	6	17	13	18	20	43	61	65	54	22	2	-
White, both sexes	28	41	31	43	54	87	117	124	72	29	3	-
Male	23	31	23	28	39	49	65	65	20	8	1	-
Female	5	10	8	15	15	38	52	59	52	21	2	-
Black, both sexes	12	18	15	5	17	11	12	13	5	2	-	-
Male	11	11	10	2	12	7	3	8	4	1	-	-
Female	1	7	5	3	5	4	9	5	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	1	1	1	1	-	-	-
Male	-	1	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
New Mexico												
All races, both sexes	8	15	8	6	4	18	11	15	10	1	1	-
Male	4	9	5	1	3	11	5	8	3	-	-	-
Female	4	6	3	5	1	7	6	7	7	1	1	-
White, both sexes	7	12	7	4	4	18	10	15	10	1	1	-
Male	4	7	5	1	3	11	5	8	3	-	-	-
Female	3	5	2	3	1	7	5	7	7	1	1	-
Black, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	2	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	56	66	74	62	79	127	181	166	119	42	7	-
Male	33	50	47	50	51	83	108	82	46	10	3	-
Female	23	16	27	12	28	44	73	84	73	32	4	-
White, both sexes	38	52	59	50	71	119	170	159	111	38	6	-
Male	23	42	40	40	47	80	101	79	43	9	3	-
Female	15	10	19	10	24	39	69	80	68	29	3	-
Black, both sexes	17	13	13	9	6	6	10	7	5	4	1	-
Male	9	7	6	8	3	3	6	3	1	1	-	-
Female	8	6	7	1	3	3	4	4	4	3	1	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	2	2	1	-	3	-	-	-
Male	-	1	-	2	1	-	1	-	2	-	-	-
Female	-	-	1	-	1	2	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
North Carolina												
All races, both sexes	45	76	62	67	77	99	141	107	67	18	3	-
Male	30	54	39	46	61	58	76	48	21	5	1	-
Female	15	22	23	21	16	41	65	59	46	13	2	-
White, both sexes	20	45	31	34	56	77	108	85	59	14	1	-
Male	16	29	20	24	42	48	66	39	19	4	-	-
Female	4	16	11	10	14	29	42	46	40	10	1	-
Black, both sexes	25	30	29	29	19	22	32	22	8	4	2	-
Male	14	25	19	19	17	10	9	9	2	1	1	-
Female	11	5	10	10	2	12	23	13	6	3	1	-
American Indian 1/, both sexes	-	-	1	2	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	2	-	-	-	-	-	-	-
Male	-	-	-	2	2	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	1	1	2	6	4	7	9	2	3	1	-
Male	1	1	1	-	5	1	6	5	-	-	-	-
Female	-	-	-	2	1	3	1	4	2	3	1	-
White, both sexes	1	1	1	1	6	4	7	9	2	3	1	-
Male	1	1	1	-	5	1	6	5	-	-	-	-
Female	-	-	-	1	1	3	1	4	2	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Ohio												
All races, both sexes	70	70	58	81	98	111	131	154	98	27	3	-
Male	45	47	34	49	64	60	81	79	31	6	-	-
Female	25	23	24	32	34	51	50	75	67	21	3	-
White, both sexes	51	55	44	64	81	94	120	139	89	24	2	-
Male	32	36	27	39	51	49	71	71	29	5	-	-
Female	19	19	17	25	30	45	49	68	60	19	2	-
Black, both sexes	19	15	14	17	17	17	11	15	9	3	1	-
Male	13	11	7	10	13	11	10	8	2	1	-	-
Female	6	4	7	7	4	6	1	7	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	16	7	15	30	33	43	31	24	8	1	-
Male	6	11	3	12	19	18	20	15	9	3	1	-
Female	4	5	4	3	11	15	23	16	15	5	-	-
White, both sexes	7	11	7	13	24	30	39	31	22	7	1	-
Male	5	7	3	11	14	15	19	15	8	2	1	-
Female	2	4	4	2	10	15	20	16	14	5	-	-
Black, both sexes	3	2	-	2	4	2	2	-	2	1	-	-
Male	1	1	-	1	3	2	-	-	1	1	-	-
Female	2	1	-	1	1	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	3	-	-	2	1	2	-	-	-	-	-
Male	-	3	-	-	2	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Oregon												
All races, both sexes	15	11	11	11	21	25	29	26	18	10	2	-
Male	8	9	8	8	12	14	17	15	5	5	-	-
Female	7	2	3	3	9	11	12	11	13	5	2	-
White, both sexes	14	9	9	11	20	25	28	25	18	10	2	-
Male	7	8	7	8	12	14	16	15	5	5	-	-
Female	7	1	2	3	8	11	12	10	13	5	2	-
Black, both sexes	1	1	1	-	1	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	58	69	73	93	126	197	264	230	149	48	7	-
Male	40	44	54	68	76	110	147	106	57	10	2	-
Female	18	25	19	25	50	87	117	124	92	38	5	-
White, both sexes	48	56	61	74	103	176	249	217	145	46	7	-
Male	34	34	44	55	64	101	141	100	55	10	2	-
Female	14	22	17	19	39	75	108	117	90	36	5	-
Black, both sexes	10	13	12	17	23	21	15	13	4	2	-	-
Male	6	10	10	12	12	9	6	6	2	-	-	-
Female	4	3	2	5	11	12	9	7	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Rhode Island												
All races, both sexes	2	1	3	3	5	4	4	5	6	4	-	-
Male	2	-	2	3	2	3	1	2	4	-	-	-
Female	-	1	1	-	3	1	3	3	2	4	-	-
White, both sexes	2	1	3	3	5	4	4	5	6	3	-	-
Male	2	-	2	3	2	3	1	2	4	-	-	-
Female	-	1	1	-	3	1	3	3	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	31	25	35	44	43	51	51	42	37	7	-	1
Male	21	18	26	24	23	30	28	19	12	2	-	1
Female	10	7	9	20	20	21	23	23	25	5	-	-
White, both sexes	15	12	16	29	27	36	35	38	24	3	-	-
Male	11	8	13	19	15	21	21	19	9	1	-	-
Female	4	4	3	10	12	15	14	19	15	2	-	-
Black, both sexes	16	13	19	15	16	15	16	4	13	4	-	1
Male	10	10	13	5	8	9	7	-	3	1	-	1
Female	6	3	6	10	8	6	9	4	10	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
South Dakota												
All races, both sexes	1	1	1	4	2	4	8	7	8	1	1	-
Male	1	1	1	1	1	4	7	2	3	1	-	-
Female	-	-	-	3	1	-	1	5	5	-	1	-
White, both sexes	1	1	1	4	2	4	8	7	8	1	1	-
Male	1	1	1	1	1	4	7	2	3	1	-	-
Female	-	-	-	3	1	-	1	5	5	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	20	34	41	42	45	47	81	59	27	18	2	1
Male	13	24	27	29	27	29	41	36	10	5	1	1
Female	7	10	14	13	18	18	40	23	17	13	1	-
White, both sexes	13	22	21	26	37	37	68	48	24	16	-	1
Male	8	16	14	18	21	23	34	26	9	4	-	1
Female	5	6	7	8	16	14	34	22	15	12	-	-
Black, both sexes	7	12	20	15	7	10	12	10	3	2	2	-
Male	5	8	13	11	6	6	6	9	1	1	1	-
Female	2	4	7	4	1	4	6	1	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Texas												
All races, both sexes	97	103	103	87	114	167	162	145	89	34	6	-
Male	67	72	70	56	60	85	92	62	30	14	1	-
Female	30	31	33	31	54	82	70	83	59	20	5	-
White, both sexes	73	75	81	67	95	145	145	135	75	30	5	-
Male	50	52	56	42	49	79	83	59	23	12	1	-
Female	23	23	25	25	46	66	62	76	52	18	4	-
Black, both sexes	22	27	22	19	18	19	17	10	12	4	1	-
Male	15	20	14	13	11	5	9	3	6	2	-	-
Female	7	7	8	6	7	14	8	7	6	2	1	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	2	-	-	2	-	-	-
Male	1	-	-	1	-	1	-	-	1	-	-	-
Female	-	1	-	-	1	1	-	-	1	-	-	-
Utah												
All races, both sexes	3	5	7	7	9	8	14	7	8	1	-	-
Male	3	2	5	2	5	5	9	4	4	-	-	-
Female	-	3	2	5	4	3	5	3	4	1	-	-
White, both sexes	2	5	7	7	9	8	14	7	8	1	-	-
Male	2	2	5	2	5	5	9	4	4	-	-	-
Female	-	3	2	5	4	3	5	3	4	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Vermont												
All races, both sexes	1	1	4	3	7	7	7	5	5	1	-	-
Male	1	1	2	2	5	3	4	4	2	-	-	-
Female	-	-	2	1	2	4	3	1	3	1	-	-
White, both sexes	1	-	4	3	7	7	7	5	5	1	-	-
Male	1	-	2	2	5	3	4	4	2	-	-	-
Female	-	-	2	1	2	4	3	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	36	43	42	52	56	80	100	108	52	23	3	-
Male	22	27	28	33	31	50	45	50	21	5	1	-
Female	14	16	14	19	25	30	55	58	31	18	2	-
White, both sexes	18	21	22	29	41	61	77	93	45	22	3	-
Male	10	13	14	18	26	40	37	43	21	5	1	-
Female	8	8	8	11	15	21	40	50	24	17	2	-
Black, both sexes	17	20	19	22	15	17	21	14	7	1	-	-
Male	11	12	13	14	5	10	6	6	-	-	-	-
Female	6	8	6	8	10	7	15	8	7	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	-	2	2	1	-	-	-	-
Male	1	2	1	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Washington												
All races, both sexes	35	33	34	35	27	51	55	39	26	7	1	-
Male	25	22	21	21	17	29	34	15	11	1	-	-
Female	10	11	13	14	10	22	21	24	15	6	1	-
White, both sexes	34	26	28	32	22	46	52	37	25	7	1	-
Male	24	17	17	18	14	28	33	14	11	1	-	-
Female	10	9	11	14	8	18	19	23	14	6	1	-
Black, both sexes	-	5	3	3	3	2	3	-	-	-	-	-
Male	-	4	3	3	2	-	1	-	-	-	-	-
Female	-	1	-	-	1	2	2	-	-	-	-	-
American Indian 1/, both sexes	1	2	2	-	1	1	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	2	-	2	1	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
West Virginia												
All races, both sexes	6	12	8	18	17	22	20	16	14	-	-	-
Male	4	11	6	14	12	14	11	7	6	-	-	-
Female	2	1	2	4	5	8	9	9	8	-	-	-
White, both sexes	6	10	7	18	17	20	20	14	14	-	-	-
Male	4	9	5	14	12	13	11	7	6	-	-	-
Female	2	1	2	4	5	7	9	7	8	-	-	-
Black, both sexes	-	2	1	-	-	2	-	2	-	-	-	-
Male	-	2	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Wisconsin												
All races, both sexes	24	31	22	31	36	59	73	76	54	25	6	-
Male	16	21	17	22	25	32	30	34	15	4	-	-
Female	8	10	5	9	11	27	43	42	39	21	6	-
White, both sexes	19	23	21	29	31	53	71	75	49	25	6	-
Male	13	15	16	22	22	32	30	33	14	4	-	-
Female	6	8	5	7	9	21	41	42	35	21	6	-
Black, both sexes	5	6	1	2	2	3	1	1	4	-	-	-
Male	3	4	1	-	2	-	-	1	1	-	-	-
Female	2	2	-	2	-	3	1	-	3	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	2	-	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	1	-	-	-
Wyoming												
All races, both sexes	1	2	1	1	3	3	2	6	2	-	-	-
Male	1	1	-	1	1	2	1	4	-	-	-	-
Female	-	1	1	-	2	1	1	2	2	-	-	-
White, both sexes	1	2	1	1	3	3	2	6	2	-	-	-
Male	1	1	-	1	1	2	1	4	-	-	-	-
Female	-	1	1	-	2	1	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
United States												
All races, both sexes	37,739	36	19	13	36	73	128	152	201	282	467	911
Male	16,671	15	10	8	21	49	81	100	117	180	287	560
Female	21,068	21	9	5	15	24	47	52	84	102	180	351
White, both sexes	32,170	21	11	10	29	47	88	99	128	181	322	684
Male	13,995	10	6	7	16	32	61	65	75	122	201	429
Female	18,175	11	5	3	13	15	27	34	53	59	121	255
Black, both sexes	4,795	13	6	2	6	21	35	46	66	98	134	203
Male	2,279	5	4	-	5	13	18	29	37	56	77	120
Female	2,516	8	2	2	1	8	17	17	29	42	57	83
American Indian 1/, both sexes	148	1	-	-	-	2	1	-	1	-	5	13
Male	74	-	-	-	-	1	-	-	-	-	4	5
Female	74	1	-	-	-	1	1	-	1	-	1	8
Asian or Pacific Islander 2/, both sexes	626	1	2	1	1	3	4	7	6	3	6	11
Male	323	-	-	1	-	3	2	6	5	2	5	6
Female	303	1	2	-	1	-	2	1	1	1	1	5
Alabama												
All races, both sexes	2,616	6	1	1	2	1	7	17	29	52	59	100
Male	1,245	3	1	-	-	1	5	14	18	35	40	57
Female	1,371	3	-	1	2	-	2	3	11	17	19	43
White, both sexes	1,916	5	-	-	2	1	6	9	20	26	39	61
Male	909	3	-	-	-	1	5	8	11	20	28	33
Female	1,007	2	-	-	2	-	1	1	9	6	11	28
Black, both sexes	688	1	1	1	-	-	1	7	8	26	20	39
Male	330	-	1	-	-	-	-	5	6	15	12	24
Female	358	1	-	1	-	-	1	2	2	11	8	15
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	1	1	-	-	-
Male	5	-	-	-	-	-	-	1	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Alaska												
All races, both sexes	44	1	-	-	-	-	-	1	-	-	-	3
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	25	1	-	-	-	-	-	1	-	-	-	2
White, both sexes	36	-	-	-	-	-	-	1	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	1	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	361	-	-	-	1	2	2	1	-	3	1	7
Male	160	-	-	-	-	2	1	1	-	3	1	3
Female	201	-	-	-	1	-	1	-	-	-	-	4
White, both sexes	340	-	-	-	1	1	2	1	-	3	1	5
Male	148	-	-	-	-	1	1	1	-	3	1	2
Female	192	-	-	-	1	-	1	-	-	-	-	3
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Arkansas												
All races, both sexes	388	1	-	1	1	-	1	2	1	5	7	18
Male	188	1	-	1	1	-	1	2	1	2	4	10
Female	200	-	-	-	-	-	-	-	-	3	3	8
White, both sexes	324	-	-	1	1	-	-	2	1	1	3	14
Male	154	-	-	1	1	-	-	2	1	1	1	7
Female	170	-	-	-	-	-	-	-	-	-	2	7
Black, both sexes	63	1	-	-	-	-	1	-	-	4	4	4
Male	34	1	-	-	-	-	1	-	-	1	3	3
Female	29	-	-	-	-	-	-	-	-	3	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,410	1	1	1	4	2	2	6	5	9	3	13
Male	583	-	1	1	2	1	2	4	3	7	3	6
Female	827	1	-	-	2	1	-	2	2	2	-	7
White, both sexes	1,262	-	1	1	4	2	2	6	4	8	3	11
Male	520	-	1	1	2	1	2	4	3	6	3	5
Female	742	-	-	-	2	1	-	2	1	2	-	6
Black, both sexes	52	-	-	-	-	-	-	-	1	1	-	1
Male	24	-	-	-	-	-	-	-	-	1	-	1
Female	28	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	88	1	-	-	-	-	-	-	-	-	-	1
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	51	1	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Colorado												
All races, both sexes	286	-	-	-	1	2	3	-	-	-	4	5
Male	104	-	-	-	1	-	2	-	-	-	4	4
Female	182	-	-	-	-	2	1	-	-	-	-	1
White, both sexes	282	-	-	-	1	1	3	-	-	-	3	5
Male	102	-	-	-	1	-	2	-	-	-	3	4
Female	180	-	-	-	-	1	1	-	-	-	-	1
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	904	1	-	-	-	4	1	2	2	1	8	18
Male	389	1	-	-	-	2	1	2	1	-	6	9
Female	515	-	-	-	-	2	-	-	1	1	2	9
White, both sexes	856	1	-	-	-	4	1	1	1	1	5	16
Male	368	1	-	-	-	2	1	1	1	-	4	7
Female	488	-	-	-	-	2	-	-	-	1	1	9
Black, both sexes	42	-	-	-	-	-	-	1	1	-	3	2
Male	19	-	-	-	-	-	-	1	-	-	2	2
Female	23	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Delaware												
All races, both sexes	90	-	-	-	-	-	-	1	2	-	1	2
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	1	2	-	1	1
White, both sexes	75	-	-	-	-	-	-	1	1	-	-	2
Male	35	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	1	1	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	48	-	1	-	-	-	-	-	-	1	1	1
Male	20	-	1	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	38	-	1	-	-	-	-	-	-	-	1	1
Male	15	-	1	-	-	-	-	-	-	-	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Florida												
All races, both sexes	1,515	1	1	2	3	3	5	3	2	8	12	30
Male	683	1	-	2	3	2	2	2	2	5	9	20
Female	832	-	1	-	-	1	3	1	-	3	3	10
White, both sexes	1,364	-	-	2	3	3	3	2	1	7	9	19
Male	607	-	-	2	3	2	2	2	1	5	7	13
Female	757	-	-	-	-	1	1	-	-	2	2	6
Black, both sexes	147	1	1	-	-	-	2	1	1	1	3	11
Male	75	1	-	-	-	-	-	-	1	-	2	7
Female	72	-	1	-	-	-	2	1	-	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,150	3	1	1	1	5	5	6	13	18	26	56
Male	948	1	-	-	-	3	-	4	9	15	16	30
Female	1,202	2	1	1	1	2	5	2	4	3	10	26
White, both sexes	1,598	2	1	1	1	1	1	2	8	12	16	35
Male	700	1	-	-	-	1	-	1	4	10	10	21
Female	898	1	1	1	1	-	1	1	4	2	6	14
Black, both sexes	537	1	-	-	-	3	3	4	3	6	10	21
Male	239	-	-	-	-	2	-	3	3	5	6	9
Female	298	1	-	-	-	1	3	1	-	1	4	12
American Indian 1/, both sexes	4	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	1	-	2	-	-	-
Male	6	-	-	-	-	-	-	-	2	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Hawaii												
All races, both sexes	340	-	-	-	-	1	1	1	2	-	2	6
Male	173	-	-	-	-	1	-	1	1	-	1	3
Female	167	-	-	-	-	-	1	-	1	-	1	3
White, both sexes	60	-	-	-	-	-	-	-	1	-	-	1
Male	28	-	-	-	-	-	-	-	1	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	276	-	-	-	-	1	-	1	1	-	2	4
Male	144	-	-	-	-	1	-	1	-	-	1	3
Female	132	-	-	-	-	-	-	-	1	-	1	1
Idaho												
All races, both sexes	148	-	1	1	-	-	-	-	2	-	1	3
Male	65	-	-	-	-	-	-	-	2	-	-	1
Female	83	-	1	1	-	-	-	-	-	-	1	2
White, both sexes	146	-	1	1	-	-	-	-	2	-	1	3
Male	63	-	-	-	-	-	-	-	2	-	-	1
Female	83	-	1	1	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Illinois												
All races, both sexes	1,572	-	1	-	2	6	12	15	16	17	22	42
Male	638	-	1	-	1	4	9	11	6	10	13	24
Female	934	-	-	-	1	2	3	4	10	7	9	18
White, both sexes	1,354	-	1	-	1	3	9	9	9	11	15	32
Male	549	-	1	-	1	2	6	7	5	6	9	19
Female	805	-	-	-	-	1	3	2	4	5	6	13
Black, both sexes	193	-	-	-	-	2	3	4	7	5	7	10
Male	79	-	-	-	-	1	3	3	1	4	4	5
Female	114	-	-	-	-	1	-	1	6	1	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	1	1	-	2	-	1	-	-
Male	10	-	-	-	-	1	-	1	-	-	-	-
Female	15	-	-	-	1	-	-	1	-	1	-	-
Indiana												
All races, both sexes	981	4	1	-	2	-	3	6	5	7	20	25
Male	449	2	1	-	-	-	1	3	4	4	13	16
Female	532	2	-	-	2	-	2	3	1	3	7	9
White, both sexes	907	1	1	-	2	-	2	5	4	6	19	24
Male	411	1	1	-	-	-	1	3	3	3	12	16
Female	496	-	-	-	2	-	1	2	1	3	7	8
Black, both sexes	73	3	-	-	-	-	1	1	1	1	1	1
Male	38	1	-	-	-	-	-	-	1	1	1	-
Female	35	2	-	-	-	-	1	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Iowa												
All races, both sexes	309	-	-	-	-	-	1	-	1	5	-	4
Male	132	-	-	-	-	-	-	-	-	4	-	3
Female	177	-	-	-	-	-	1	-	1	1	-	1
White, both sexes	306	-	-	-	-	-	1	-	1	4	-	4
Male	130	-	-	-	-	-	-	-	-	3	-	3
Female	176	-	-	-	-	-	1	-	1	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	375	1	1	-	-	3	2	2	1	4	2	3
Male	148	-	1	-	-	2	2	2	1	2	2	2
Female	227	1	-	-	-	1	-	-	-	2	-	1
White, both sexes	353	1	-	-	-	2	2	1	-	4	2	3
Male	137	-	-	-	-	1	2	1	-	2	2	2
Female	216	1	-	-	-	1	-	-	-	2	-	1
Black, both sexes	19	-	1	-	-	-	-	1	1	-	-	-
Male	9	-	1	-	-	-	-	1	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Kentucky												
All races, both sexes	914	-	1	-	1	-	3	3	9	6	19	36
Male	365	-	-	-	1	-	1	3	3	4	11	23
Female	549	-	1	-	-	-	2	-	6	2	8	13
White, both sexes	836	-	1	-	1	-	2	1	5	5	15	32
Male	336	-	-	-	1	-	-	1	2	3	9	21
Female	500	-	1	-	-	-	2	-	3	2	6	11
Black, both sexes	77	-	-	-	-	-	1	2	4	1	3	4
Male	29	-	-	-	-	-	1	2	1	1	2	2
Female	48	-	-	-	-	-	-	-	3	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	667	-	1	-	1	1	6	7	5	5	10	16
Male	299	-	1	-	1	1	3	3	1	4	7	9
Female	368	-	-	-	-	-	3	4	4	1	3	7
White, both sexes	458	-	1	-	-	1	3	3	1	4	7	10
Male	206	-	1	-	-	1	2	2	-	4	5	6
Female	252	-	-	-	-	-	1	1	1	-	2	4
Black, both sexes	208	-	-	-	1	-	3	4	4	1	3	6
Male	93	-	-	-	1	-	1	1	1	-	2	3
Female	115	-	-	-	-	-	2	3	3	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Maine												
All races, both sexes	197	-	-	-	-	-	-	-	-	1	1	3
Male	105	-	-	-	-	-	-	-	-	1	1	3
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	195	-	-	-	-	-	-	-	-	1	-	3
Male	104	-	-	-	-	-	-	-	-	1	-	3
Female	91	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	602	1	1	1	-	2	5	3	8	4	6	14
Male	270	-	-	-	-	1	3	3	7	3	4	7
Female	332	1	1	1	-	1	2	-	1	1	2	7
White, both sexes	430	-	-	1	-	1	3	2	6	2	3	7
Male	188	-	-	-	-	1	3	2	5	1	2	3
Female	242	-	-	1	-	-	-	-	1	1	1	4
Black, both sexes	159	1	-	-	-	1	2	1	2	2	3	7
Male	76	-	-	-	-	-	-	1	2	2	2	4
Female	83	1	-	-	-	1	2	-	-	-	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Massachusetts												
All races, both sexes	1,018	-	-	-	-	1	2	4	-	1	8	15
Male	438	-	-	-	-	1	1	3	-	1	5	11
Female	580	-	-	-	-	-	1	1	-	-	3	4
White, both sexes	974	-	-	-	-	-	2	2	-	-	5	13
Male	413	-	-	-	-	-	1	1	-	-	2	10
Female	561	-	-	-	-	-	1	1	-	-	3	3
Black, both sexes	36	-	-	-	-	1	-	2	-	1	3	2
Male	22	-	-	-	-	1	-	2	-	1	3	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	877	1	-	1	-	2	3	4	4	5	11	14
Male	352	-	-	1	-	1	2	4	3	4	8	11
Female	525	1	-	-	-	1	1	-	1	1	3	3
White, both sexes	779	1	-	1	-	-	3	1	3	3	9	12
Male	311	-	-	1	-	-	2	1	3	3	7	9
Female	468	1	-	-	-	-	1	-	-	-	2	3
Black, both sexes	92	-	-	-	-	2	-	3	1	2	2	2
Male	36	-	-	-	-	1	-	3	-	1	1	2
Female	56	-	-	-	-	1	-	-	1	1	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Minnesota												
All races, both sexes	582	1	-	-	-	-	2	-	3	2	7	11
Male	246	-	-	-	-	-	-	-	2	1	4	8
Female	336	1	-	-	-	-	2	-	1	1	3	3
White, both sexes	571	1	-	-	-	-	2	-	2	2	7	10
Male	238	-	-	-	-	-	-	-	1	1	4	7
Female	333	1	-	-	-	-	2	-	1	1	3	3
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	709	-	-	-	1	-	2	5	2	6	15	22
Male	350	-	-	-	1	-	2	1	1	6	10	13
Female	359	-	-	-	-	-	-	4	1	-	5	9
White, both sexes	456	-	-	-	-	-	-	3	1	2	7	12
Male	212	-	-	-	-	-	-	1	-	2	5	5
Female	244	-	-	-	-	-	-	2	1	-	2	7
Black, both sexes	252	-	-	-	1	-	2	2	1	4	8	10
Male	137	-	-	-	1	-	2	-	1	4	5	8
Female	115	-	-	-	-	-	-	2	-	-	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Missouri												
All races, both sexes	879	-	2	-	-	3	4	2	4	5	20	30
Male	385	-	1	-	-	2	1	1	2	3	12	19
Female	494	-	1	-	-	1	3	1	2	2	8	11
White, both sexes	812	-	1	-	-	3	2	1	3	4	18	23
Male	359	-	-	-	-	2	-	1	1	3	12	16
Female	453	-	1	-	-	1	2	-	2	1	6	7
Black, both sexes	63	-	1	-	-	-	1	1	1	1	2	7
Male	26	-	1	-	-	-	1	-	1	-	-	3
Female	37	-	-	-	-	-	-	1	-	1	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	141	-	-	-	-	-	1	1	1	3	-	3
Male	66	-	-	-	-	-	1	1	1	1	-	2
Female	75	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	130	-	-	-	-	-	1	1	1	3	-	1
Male	61	-	-	-	-	-	1	1	1	1	-	1
Female	69	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Nebraska												
All races, both sexes	374	-	-	-	1	-	-	-	1	1	4	7
Male	163	-	-	-	1	-	-	-	1	1	2	4
Female	211	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	366	-	-	-	1	-	-	-	1	1	4	7
Male	158	-	-	-	1	-	-	-	1	1	2	4
Female	208	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	745	-	-	1	-	-	5	8	10	7	24	32
Male	426	-	-	1	-	-	3	4	7	3	12	24
Female	319	-	-	-	-	-	2	4	3	4	12	8
White, both sexes	611	-	-	1	-	-	3	5	5	2	18	24
Male	352	-	-	1	-	-	2	2	3	1	10	17
Female	259	-	-	-	-	-	1	3	2	1	8	7
Black, both sexes	100	-	-	-	-	-	1	3	4	5	6	7
Male	49	-	-	-	-	-	1	2	3	2	2	6
Female	51	-	-	-	-	-	-	1	1	3	4	1
American Indian 1/, both sexes	8	-	-	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	1	-	-	-
Male	21	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Hampshire												
All races, both sexes	125	-	-	-	1	2	-	-	1	2	4	1
Male	54	-	-	-	1	-	-	-	1	1	1	1
Female	71	-	-	-	-	2	-	-	-	1	3	-
White, both sexes	124	-	-	-	1	2	-	-	1	2	3	1
Male	53	-	-	-	1	-	-	-	1	1	-	1
Female	71	-	-	-	-	2	-	-	-	1	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,095	1	-	-	-	-	3	4	3	6	12	17
Male	477	1	-	-	-	-	2	4	2	4	6	9
Female	618	-	-	-	-	-	1	-	1	2	6	8
White, both sexes	964	-	-	-	-	-	2	3	3	3	10	14
Male	417	-	-	-	-	-	2	3	2	2	6	8
Female	547	-	-	-	-	-	-	-	1	1	4	6
Black, both sexes	106	1	-	-	-	-	1	1	-	2	2	2
Male	46	1	-	-	-	-	-	1	-	1	-	1
Female	60	-	-	-	-	-	1	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Mexico												
All races, both sexes	153	-	-	-	-	1	-	-	-	-	-	1
Male	61	-	-	-	-	1	-	-	-	-	-	-
Female	92	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	142	-	-	-	-	1	-	-	-	-	-	-
Male	55	-	-	-	-	1	-	-	-	-	-	-
Female	87	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,714	1	1	-	3	6	8	7	10	13	20	38
Male	742	1	-	-	2	6	8	3	6	10	11	20
Female	972	-	1	-	1	-	-	4	4	3	9	18
White, both sexes	1,573	1	1	-	2	5	4	6	8	11	12	31
Male	666	1	-	-	2	5	4	2	5	9	6	17
Female	907	-	1	-	-	-	-	4	3	2	6	14
Black, both sexes	116	-	-	-	1	1	3	1	2	2	6	5
Male	60	-	-	-	-	1	3	1	1	1	3	3
Female	56	-	-	-	1	-	-	-	1	1	3	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	1	-	-	-	2	1
Male	15	-	-	-	-	-	1	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
North Carolina												
All races, both sexes	1,349	1	1	-	-	1	4	4	7	10	13	24
Male	594	-	-	-	-	1	2	2	4	5	8	14
Female	755	1	1	-	-	-	2	2	3	5	5	10
White, both sexes	1,063	1	-	-	-	-	3	3	3	4	6	22
Male	464	-	-	-	-	-	1	2	2	3	4	13
Female	599	1	-	-	-	-	2	1	1	1	2	9
Black, both sexes	269	-	-	-	-	1	1	1	4	6	6	1
Male	120	-	-	-	-	1	1	-	2	2	3	-
Female	149	-	-	-	-	-	-	1	2	4	3	1
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	1	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	76	-	-	-	-	-	-	-	-	-	1	1
Male	36	-	-	-	-	-	-	-	-	-	1	-
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	72	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Ohio												
All races, both sexes	1,385	-	-	-	-	1	3	2	5	3	11	35
Male	549	-	-	-	-	1	2	1	1	2	7	26
Female	836	-	-	-	-	-	1	1	4	1	4	9
White, both sexes	1,262	-	-	-	-	1	2	1	4	3	9	29
Male	493	-	-	-	-	1	2	-	1	2	6	23
Female	769	-	-	-	-	-	-	1	3	1	3	6
Black, both sexes	118	-	-	-	-	-	1	1	1	-	2	6
Male	53	-	-	-	-	-	-	1	-	-	1	3
Female	65	-	-	-	-	-	1	-	1	-	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	393	-	-	-	1	3	-	2	1	1	6	6
Male	163	-	-	-	1	2	-	1	-	1	3	4
Female	230	-	-	-	-	1	-	1	1	-	3	2
White, both sexes	341	-	-	-	-	2	-	2	-	-	5	3
Male	136	-	-	-	-	1	-	1	-	-	2	3
Female	205	-	-	-	-	1	-	1	-	-	3	-
Black, both sexes	24	-	-	-	1	-	-	-	-	1	-	1
Male	15	-	-	-	1	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	-	-	-	-	1	-	-	1	-	1	2
Male	12	-	-	-	-	1	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	1	-	-	2
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Oregon												
All races, both sexes	438	-	-	-	2	1	2	2	1	2	2	7
Male	172	-	-	-	1	1	1	2	1	2	1	4
Female	266	-	-	-	1	-	1	-	-	-	1	3
White, both sexes	427	-	-	-	2	1	2	-	1	2	2	7
Male	167	-	-	-	1	1	1	-	1	2	1	4
Female	260	-	-	-	1	-	1	-	-	-	1	3
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,714	1	2	-	4	1	9	8	9	13	23	55
Male	1,214	-	2	-	2	-	6	5	3	7	15	38
Female	1,500	1	-	-	2	1	3	3	6	6	8	17
White, both sexes	2,452	-	2	-	3	1	5	8	7	7	14	45
Male	1,083	-	2	-	1	-	3	5	2	4	8	33
Female	1,369	-	-	-	2	1	2	3	5	3	6	12
Black, both sexes	252	1	-	-	1	-	4	-	2	6	9	10
Male	127	-	-	-	1	-	3	-	1	3	7	5
Female	125	1	-	-	-	-	1	-	1	3	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Rhode Island												
All races, both sexes	91	-	-	-	-	-	1	-	-	-	-	2
Male	30	-	-	-	-	-	1	-	-	-	-	1
Female	61	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	91	-	-	-	-	-	1	-	-	-	-	2
Male	30	-	-	-	-	-	1	-	-	-	-	1
Female	61	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	825	1	-	-	-	1	1	3	5	8	10	28
Male	394	-	-	-	-	-	1	1	4	4	5	20
Female	431	1	-	-	-	1	-	2	1	4	5	8
White, both sexes	529	1	-	-	-	-	-	1	2	4	1	15
Male	245	-	-	-	-	-	-	-	1	3	1	11
Female	284	1	-	-	-	-	-	1	1	1	-	4
Black, both sexes	290	-	-	-	-	1	1	2	3	4	9	13
Male	147	-	-	-	-	-	1	1	3	1	4	9
Female	143	-	-	-	-	1	-	1	-	3	5	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
South Dakota												
All races, both sexes	47	-	-	-	-	-	-	-	1	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	26	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	1	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	26	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	832	3	-	-	1	2	1	5	8	12	14	31
Male	376	2	-	-	-	2	1	4	6	7	8	21
Female	456	1	-	-	1	-	-	1	2	5	6	10
White, both sexes	708	2	-	-	1	1	1	4	3	9	12	28
Male	310	1	-	-	-	1	1	3	2	5	7	19
Female	398	1	-	-	1	-	-	1	1	4	5	9
Black, both sexes	121	1	-	-	-	1	-	-	4	3	2	3
Male	63	1	-	-	-	1	-	-	3	2	1	2
Female	58	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	1	-	-	-
Male	2	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Texas												
All races, both sexes	1,888	3	-	1	1	5	7	5	13	14	28	42
Male	845	2	-	1	-	3	7	1	6	9	13	21
Female	1,043	1	-	-	1	2	-	4	7	5	15	21
White, both sexes	1,594	2	-	1	1	3	7	3	9	4	20	31
Male	693	2	-	1	-	1	7	1	5	3	10	16
Female	901	-	-	-	1	2	-	2	4	1	10	15
Black, both sexes	258	1	-	-	-	2	-	2	4	10	8	9
Male	134	-	-	-	-	2	-	-	1	6	3	4
Female	124	1	-	-	-	-	-	2	3	4	5	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	193	-	-	-	1	1	1	2	2	3	1	6
Male	78	-	-	-	1	1	-	1	2	-	1	3
Female	115	-	-	-	-	-	1	1	-	3	-	3
White, both sexes	185	-	-	-	1	1	1	2	2	3	1	5
Male	73	-	-	-	1	1	-	1	2	-	1	2
Female	112	-	-	-	-	-	1	1	-	3	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Vermont												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,473	2	-	2	1	7	2	5	5	12	22	43
Male	680	-	-	1	1	4	1	3	4	6	14	30
Female	793	2	-	1	-	3	1	2	1	6	8	13
White, both sexes	1,162	2	-	-	-	4	1	5	1	10	14	28
Male	535	-	-	-	-	3	1	3	1	5	9	20
Female	627	2	-	-	-	1	-	2	-	5	5	8
Black, both sexes	297	-	-	1	1	3	1	-	4	2	8	15
Male	136	-	-	-	1	1	-	-	3	1	5	10
Female	161	-	-	1	-	2	1	-	1	1	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	1	-	-	-	-	-	-	-	-
Male	9	-	-	1	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Washington												
All races, both sexes	367	-	1	-	-	-	4	2	-	4	-	12
Male	160	-	-	-	-	-	3	2	-	2	-	6
Female	207	-	1	-	-	-	1	-	-	2	-	6
White, both sexes	347	-	-	-	-	-	3	1	-	3	-	10
Male	152	-	-	-	-	-	3	1	-	2	-	5
Female	195	-	-	-	-	-	-	-	-	1	-	5
Black, both sexes	9	-	1	-	-	-	1	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	1	-	-	-	1	-	-	1	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	515	1	-	-	-	1	-	1	1	2	2	8
Male	232	-	-	-	-	1	-	1	-	-	1	6
Female	283	1	-	-	-	-	-	-	1	2	1	2
White, both sexes	488	-	-	-	-	1	-	1	1	2	1	8
Male	222	-	-	-	-	1	-	1	-	-	-	6
Female	266	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	26	1	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	16	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Wisconsin												
All races, both sexes	681	-	-	-	-	2	2	-	1	1	3	9
Male	297	-	-	-	-	2	2	-	1	1	2	6
Female	384	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	651	-	-	-	-	1	1	-	1	1	2	9
Male	281	-	-	-	-	1	1	-	1	1	1	6
Female	370	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	24	-	-	-	-	1	-	-	-	-	-	-
Male	12	-	-	-	-	1	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	87	-	-	-	-	-	2	-	-	-	-	5
Male	32	-	-	-	-	-	1	-	-	-	-	4
Female	55	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	86	-	-	-	-	-	2	-	-	-	-	5
Male	31	-	-	-	-	-	1	-	-	-	-	4
Female	55	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
United States												
All races, both sexes	1,222	1,671	2,004	2,262	3,001	4,519	6,229	6,700	5,161	2,157	493	2
Male	806	1,064	1,257	1,343	1,603	2,194	2,626	2,368	1,452	459	69	2
Female	416	607	747	919	1,398	2,325	3,603	4,332	3,709	1,698	424	-
White, both sexes	902	1,280	1,573	1,787	2,435	3,880	5,469	6,101	4,739	1,973	411	-
Male	612	824	994	1,069	1,321	1,887	2,322	2,162	1,310	416	54	-
Female	290	456	579	718	1,114	1,993	3,147	3,939	3,429	1,557	357	-
Black, both sexes	292	351	388	402	493	545	631	499	344	152	66	2
Male	174	210	237	223	249	257	245	162	110	34	12	2
Female	118	141	151	179	244	288	386	337	234	118	54	-
American Indian 1/, both sexes	7	10	11	16	17	15	20	13	9	4	3	-
Male	7	9	7	12	7	7	10	2	3	-	-	-
Female	-	1	4	4	10	8	10	11	6	4	3	-
Asian or Pacific Islander 2/, both sexes	21	30	32	57	56	79	109	87	69	28	13	-
Male	13	21	19	39	26	43	49	42	29	9	3	-
Female	8	9	13	18	30	36	60	45	40	19	10	-
Alabama												
All races, both sexes	138	177	198	193	222	320	369	348	242	114	20	-
Male	91	114	129	119	124	152	137	114	69	18	4	-
Female	47	63	69	74	98	168	232	234	173	96	16	-
White, both sexes	98	115	136	141	154	253	284	276	190	91	9	-
Male	66	71	89	91	93	123	108	92	51	15	1	-
Female	32	44	47	50	61	130	176	184	139	76	8	-
Black, both sexes	40	62	59	52	66	66	83	70	52	23	11	-
Male	25	43	38	28	31	29	27	22	18	3	3	-
Female	15	19	21	24	35	37	56	48	34	20	8	-
American Indian 1/, both sexes	-	-	1	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	-	1	1	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Alaska												
All races, both sexes	1	2	2	2	3	5	7	9	7	1	-	-
Male	1	2	2	1	2	4	1	5	-	-	-	-
Female	-	-	-	1	1	1	6	4	7	1	-	-
White, both sexes	-	2	2	2	3	4	5	8	7	1	-	-
Male	-	2	2	1	2	4	1	4	-	-	-	-
Female	-	-	-	1	1	-	4	4	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	15	14	16	22	30	50	62	64	48	19	4	-
Male	9	7	10	19	16	19	26	23	16	4	-	-
Female	6	7	6	3	14	31	36	41	32	15	4	-
White, both sexes	10	14	15	22	30	47	59	63	45	18	3	-
Male	5	7	9	19	16	17	24	23	15	4	-	-
Female	5	7	6	3	14	30	35	40	30	14	3	-
Black, both sexes	2	-	-	-	-	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	1	-	-	3	1	1	2	1	1	-
Male	2	-	1	-	-	2	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	1	1	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Arkansas												
All races, both sexes	12	11	26	34	38	48	63	61	41	13	4	-
Male	8	9	18	21	18	28	24	24	9	6	-	-
Female	4	2	8	13	20	20	39	37	32	7	4	-
White, both sexes	9	8	17	28	32	44	61	54	36	9	3	-
Male	6	8	11	17	15	26	23	21	9	4	-	-
Female	3	-	6	11	17	18	38	33	27	5	3	-
Black, both sexes	3	3	9	6	6	4	2	6	5	4	1	-
Male	2	1	7	4	3	2	1	3	-	2	-	-
Female	1	2	2	2	3	2	1	3	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
California												
All races, both sexes	22	43	42	48	74	145	245	322	270	121	31	-
Male	18	22	25	27	35	67	111	118	88	35	7	-
Female	4	21	17	21	39	78	134	204	182	86	24	-
White, both sexes	15	36	35	37	63	124	218	305	249	111	27	-
Male	12	18	22	20	30	57	101	115	80	32	5	-
Female	3	18	13	17	33	67	117	190	169	79	22	-
Black, both sexes	3	5	2	4	6	6	6	4	9	3	1	-
Male	3	3	-	1	3	4	2	-	2	3	1	-
Female	-	2	2	3	3	2	4	4	7	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	2	2	1	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	2	2	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	4	2	5	7	3	13	19	12	12	6	3	-
Male	3	1	3	6	2	6	6	3	6	-	1	-
Female	1	1	2	1	1	7	13	9	6	6	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Colorado												
All races, both sexes	2	5	12	18	19	31	54	54	56	18	2	-
Male	2	2	3	8	7	19	25	13	10	4	-	-
Female	-	3	9	10	12	12	29	41	46	14	2	-
White, both sexes	2	5	12	18	17	31	54	54	56	18	2	-
Male	2	2	3	8	6	19	25	13	10	4	-	-
Female	-	3	9	10	11	12	29	41	46	14	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	20	40	35	45	52	92	158	184	144	73	24	-
Male	18	25	23	29	28	44	65	72	51	10	2	-
Female	2	15	12	16	24	48	93	112	93	63	22	-
White, both sexes	20	38	32	40	49	84	147	180	142	72	22	-
Male	18	23	20	26	28	41	62	71	51	10	1	-
Female	2	15	12	14	21	43	85	109	91	62	21	-
Black, both sexes	-	1	3	4	1	7	11	4	2	1	1	-
Male	-	1	3	2	-	3	3	1	-	-	1	-
Female	-	-	-	2	1	4	8	3	2	1	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	1	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Delaware												
All races, both sexes	3	6	6	3	10	11	16	7	13	8	1	-
Male	-	4	3	1	9	4	5	3	8	2	-	-
Female	3	2	3	2	1	7	11	4	5	6	1	-
White, both sexes	1	6	3	3	6	10	15	6	12	8	1	-
Male	-	4	2	1	6	4	5	2	8	2	-	-
Female	1	2	1	2	-	6	10	4	4	6	1	-
Black, both sexes	2	-	3	-	4	1	1	1	1	-	-	-
Male	-	-	1	-	3	-	-	1	-	-	-	-
Female	2	-	2	-	1	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	1	-	4	5	1	10	6	7	6	-	-
Male	1	1	-	2	4	-	3	-	2	4	-	-
Female	3	-	-	2	1	1	7	6	5	2	-	-
White, both sexes	1	1	-	-	-	-	1	-	3	3	-	-
Male	1	1	-	-	-	-	-	-	1	2	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-
Black, both sexes	3	-	-	4	5	1	9	6	4	3	-	-
Male	-	-	-	2	4	-	3	-	1	2	-	-
Female	3	-	-	2	1	1	6	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Florida												
All races, both sexes	32	58	60	59	113	181	266	320	241	95	19	1
Male	18	34	39	35	59	96	119	117	87	25	5	1
Female	14	24	21	24	54	85	147	203	154	70	14	-
White, both sexes	25	42	49	42	98	162	256	302	229	91	19	-
Male	15	25	34	25	52	85	115	109	82	23	5	-
Female	10	17	15	17	46	77	141	193	147	68	14	-
Black, both sexes	7	14	11	17	14	18	10	18	12	4	-	1
Male	3	8	5	10	7	11	4	8	5	2	-	1
Female	4	6	6	7	7	7	6	10	7	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
Georgia												
All races, both sexes	101	108	153	149	170	259	329	360	246	121	19	-
Male	61	74	91	89	95	117	124	131	63	24	1	-
Female	40	34	62	60	75	142	205	229	183	97	18	-
White, both sexes	66	71	111	104	124	196	248	284	207	98	9	-
Male	46	48	61	65	70	93	92	107	51	19	-	-
Female	20	23	50	39	54	103	156	177	156	79	9	-
Black, both sexes	34	36	42	44	46	60	79	74	38	23	10	-
Male	15	25	30	23	25	22	31	22	12	5	1	-
Female	19	11	12	21	21	38	48	52	26	18	9	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	2	2	1	1	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Hawaii												
All races, both sexes	4	12	15	28	29	37	69	54	53	18	8	-
Male	1	10	10	16	12	21	36	28	23	7	2	-
Female	3	2	5	12	17	16	33	26	30	11	6	-
White, both sexes	2	-	2	4	3	9	10	10	12	4	2	-
Male	1	-	2	1	2	2	7	6	5	-	1	-
Female	1	-	-	3	1	7	3	4	7	4	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	12	13	24	25	28	58	44	41	14	6	-
Male	-	10	8	15	9	19	29	22	18	7	1	-
Female	2	2	5	9	16	9	29	22	23	7	5	-
Idaho												
All races, both sexes	4	6	10	5	5	17	29	31	24	5	4	-
Male	1	6	5	4	2	8	12	16	7	1	-	-
Female	3	-	5	1	3	9	17	15	17	4	4	-
White, both sexes	4	5	10	5	4	17	29	31	24	5	4	-
Male	1	5	5	4	1	8	12	16	7	1	-	-
Female	3	-	5	1	3	9	17	15	17	4	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Illinois												
All races, both sexes	42	65	80	100	108	172	270	274	207	92	29	-
Male	23	39	49	65	44	67	114	85	53	16	4	-
Female	19	26	31	35	64	105	156	189	154	76	25	-
White, both sexes	31	55	68	79	79	148	229	264	196	88	27	-
Male	19	36	42	52	36	57	97	83	51	16	4	-
Female	12	19	26	27	43	91	132	181	145	72	23	-
Black, both sexes	11	9	11	17	26	20	40	8	8	3	2	-
Male	4	3	7	10	7	8	17	1	1	-	-	-
Female	7	6	4	7	19	12	23	7	7	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	4	3	4	1	2	3	1	-	-
Male	-	-	-	3	1	2	-	1	1	-	-	-
Female	-	1	1	1	2	2	1	1	2	1	-	-
Indiana												
All races, both sexes	47	64	59	83	87	105	142	138	119	47	17	-
Male	34	44	45	58	46	55	57	38	18	7	3	-
Female	13	20	14	25	41	50	85	100	101	40	14	-
White, both sexes	38	58	54	73	77	98	134	136	115	46	14	-
Male	27	38	42	52	40	54	54	38	16	7	3	-
Female	11	20	12	21	37	44	80	98	99	39	11	-
Black, both sexes	8	6	5	10	10	7	8	2	4	1	3	-
Male	7	6	3	6	6	1	3	-	2	-	-	-
Female	1	-	2	4	4	6	5	2	2	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Iowa												
All races, both sexes	10	12	14	20	20	33	46	53	52	30	8	-
Male	8	9	7	13	11	20	21	19	12	4	1	-
Female	2	3	7	7	9	13	25	34	40	26	7	-
White, both sexes	9	12	14	20	20	33	45	53	52	30	8	-
Male	7	9	7	13	11	20	21	19	12	4	1	-
Female	2	3	7	7	9	13	24	34	40	26	7	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	12	11	26	14	21	42	67	63	62	37	1	-
Male	7	10	17	7	7	17	24	18	20	6	1	-
Female	5	1	9	7	14	25	43	45	42	31	-	-
White, both sexes	9	11	23	13	18	40	66	59	61	37	1	-
Male	5	10	16	7	7	15	23	17	20	6	1	-
Female	4	1	7	6	11	25	43	42	41	31	-	-
Black, both sexes	2	-	3	1	3	2	1	3	1	-	-	-
Male	1	-	1	-	-	2	1	1	-	-	-	-
Female	1	-	2	1	3	-	-	2	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Kentucky												
All races, both sexes	26	43	42	71	74	104	150	147	113	52	14	-
Male	17	26	19	36	41	45	51	58	20	6	-	-
Female	9	17	23	35	33	59	99	89	93	46	14	-
White, both sexes	23	38	34	64	72	95	142	138	109	46	13	-
Male	16	24	16	33	41	39	49	56	19	6	-	-
Female	7	14	18	31	31	56	93	82	90	40	13	-
Black, both sexes	3	5	8	7	2	9	8	9	4	6	1	-
Male	1	2	3	3	-	6	2	2	1	-	-	-
Female	2	3	5	4	2	3	6	7	3	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	32	33	46	40	73	91	95	95	72	34	4	-
Male	19	16	30	20	35	45	42	37	15	10	-	-
Female	13	17	16	20	38	46	53	58	57	24	4	-
White, both sexes	22	16	29	26	46	63	73	73	53	23	4	-
Male	14	8	17	13	21	31	34	29	11	7	-	-
Female	8	8	12	13	25	32	39	44	42	16	4	-
Black, both sexes	10	17	17	14	26	28	22	22	19	11	-	-
Male	5	8	13	7	14	14	8	8	4	3	-	-
Female	5	9	4	7	12	14	14	14	15	8	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Maine												
All races, both sexes	6	12	10	18	23	19	28	47	19	6	4	-
Male	6	8	8	13	14	12	13	19	5	2	-	-
Female	-	4	2	5	9	7	15	28	14	4	4	-
White, both sexes	6	12	10	18	23	19	28	46	19	6	4	-
Male	6	8	8	13	14	12	13	19	5	2	-	-
Female	-	4	2	5	9	7	15	27	14	4	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	12	21	28	42	50	79	96	112	81	30	6	-
Male	7	12	18	24	27	36	44	42	25	7	-	-
Female	5	9	10	18	23	43	52	70	56	23	6	-
White, both sexes	6	10	17	23	31	57	68	94	67	28	4	-
Male	3	5	11	13	16	29	32	35	20	7	-	-
Female	3	5	6	10	15	28	36	59	47	21	4	-
Black, both sexes	6	9	11	16	17	21	26	17	14	1	2	-
Male	4	6	7	8	10	7	11	7	5	-	-	-
Female	2	3	4	8	7	14	15	10	9	1	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	3	2	1	1	1	-	1	-	-
Male	-	1	-	3	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Massachusetts												
All races, both sexes	17	37	56	47	66	114	171	205	179	71	24	-
Male	8	30	34	28	39	58	80	72	48	16	3	-
Female	9	7	22	19	27	56	91	133	131	55	21	-
White, both sexes	12	35	51	43	64	110	167	199	178	71	22	-
Male	4	28	32	27	38	56	79	68	48	16	3	-
Female	8	7	19	16	26	54	88	131	130	55	19	-
Black, both sexes	4	1	5	3	2	3	2	5	1	-	1	-
Male	3	1	2	1	1	2	1	3	-	-	-	-
Female	1	-	3	2	1	1	1	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	1	1	1	-	-	1	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	1	-
Michigan												
All races, both sexes	16	26	43	39	60	79	161	198	142	48	20	-
Male	9	15	27	21	32	40	60	66	34	10	4	-
Female	7	11	16	18	28	39	101	132	108	38	16	-
White, both sexes	9	20	31	33	49	67	150	187	135	47	18	-
Male	5	12	22	19	28	35	56	63	32	10	3	-
Female	4	8	9	14	21	32	94	124	103	37	15	-
Black, both sexes	6	5	11	6	11	12	10	10	7	1	1	-
Male	3	2	4	2	4	5	3	3	2	-	-	-
Female	3	3	7	4	7	7	7	7	5	1	1	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	1	-	-	1	-
Male	1	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Minnesota												
All races, both sexes	16	18	24	28	37	71	93	112	101	48	8	-
Male	13	11	16	15	18	40	35	45	26	10	2	-
Female	3	7	8	13	19	31	58	67	75	38	6	-
White, both sexes	15	15	23	26	37	71	91	112	101	48	8	-
Male	12	8	15	14	18	40	35	45	26	10	2	-
Female	3	7	8	12	19	31	56	67	75	38	6	-
Black, both sexes	-	3	-	-	-	-	-	-	-	-	-	-
Male	-	3	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	2	-	-	-	-	-
Male	1	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Mississippi												
All races, both sexes	33	46	35	55	72	95	100	101	75	39	5	-
Male	20	34	21	28	40	46	48	36	30	12	1	-
Female	13	12	14	27	32	49	52	65	45	27	4	-
White, both sexes	17	25	19	37	44	64	71	74	51	26	3	-
Male	10	19	11	20	26	27	36	26	15	8	1	-
Female	7	6	8	17	18	37	35	48	36	18	2	-
Black, both sexes	16	20	16	18	28	31	29	27	24	13	2	-
Male	10	14	10	8	14	19	12	10	15	4	-	-
Female	6	6	6	10	14	12	17	17	9	9	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Missouri												
All races, both sexes	38	47	51	48	56	105	122	160	111	54	17	-
Male	20	32	29	31	30	49	59	57	30	4	3	-
Female	18	15	22	17	26	56	63	103	81	50	14	-
White, both sexes	33	43	47	40	55	92	113	156	110	52	16	-
Male	18	29	26	29	29	45	54	57	30	4	3	-
Female	15	14	21	11	26	47	59	99	80	48	13	-
Black, both sexes	5	4	4	8	1	11	8	4	1	2	1	-
Male	2	3	3	2	1	4	5	-	-	-	-	-
Female	3	1	1	6	-	7	3	4	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	5	4	5	4	11	14	30	27	17	10	5	-
Male	4	3	2	3	7	4	19	13	2	3	-	-
Female	1	1	3	1	4	10	11	14	15	7	5	-
White, both sexes	4	3	3	4	10	14	29	26	16	10	4	-
Male	3	2	2	3	6	4	18	13	2	3	-	-
Female	1	1	1	1	4	10	11	13	14	7	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	2	-	1	-	1	1	1	-	1	-
Male	1	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Nebraska												
All races, both sexes	7	9	16	19	34	38	58	72	56	40	11	-
Male	6	6	8	14	17	23	32	22	14	11	1	-
Female	1	3	8	5	17	15	26	50	42	29	10	-
White, both sexes	7	9	13	17	34	35	58	72	56	40	11	-
Male	6	6	7	12	17	21	32	22	14	11	1	-
Female	1	3	6	5	17	14	26	50	42	29	10	-
Black, both sexes	-	-	2	1	-	3	-	-	-	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	2	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	47	64	85	73	76	87	99	75	37	13	2	-
Male	32	45	61	45	41	49	51	33	13	2	-	-
Female	15	19	24	28	35	38	48	42	24	11	2	-
White, both sexes	32	51	73	64	62	73	87	67	33	9	2	-
Male	22	37	56	40	34	40	46	28	12	1	-	-
Female	10	14	17	24	28	33	41	39	21	8	2	-
Black, both sexes	11	11	10	5	11	8	8	6	3	1	-	-
Male	6	6	4	1	4	4	3	4	1	-	-	-
Female	5	5	6	4	7	4	5	2	2	1	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	1	1	1	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	4	1	2	3	2	6	4	1	-	2	-	-
Male	4	1	1	3	2	5	2	1	-	1	-	-
Female	-	-	1	-	-	1	2	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Hampshire												
All races, both sexes	3	6	2	5	13	14	23	23	16	8	1	-
Male	2	5	2	4	7	9	6	7	6	1	-	-
Female	1	1	-	1	6	5	17	16	10	7	1	-
White, both sexes	3	6	2	5	13	14	23	23	16	8	1	-
Male	2	5	2	4	7	9	6	7	6	1	-	-
Female	1	1	-	1	6	5	17	16	10	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	21	33	41	63	89	148	184	233	165	63	9	-
Male	13	22	24	38	44	83	72	81	57	14	1	-
Female	8	11	17	25	45	65	112	152	108	49	8	-
White, both sexes	15	27	34	46	76	132	161	217	152	62	7	-
Male	10	19	19	28	37	76	61	76	53	14	1	-
Female	5	8	15	18	39	56	100	141	99	48	6	-
Black, both sexes	6	6	5	13	11	12	19	13	10	1	1	-
Male	3	3	3	8	5	4	10	3	3	-	-	-
Female	3	3	2	5	6	8	9	10	7	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	4	2	4	4	3	3	-	1	-
Male	-	-	2	2	2	3	1	2	1	-	-	-
Female	-	-	-	2	-	1	3	1	2	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Mexico												
All races, both sexes	1	3	10	6	10	21	28	31	24	15	2	-
Male	1	2	5	3	8	7	14	12	6	2	-	-
Female	-	1	5	3	2	14	14	19	18	13	2	-
White, both sexes	1	3	9	4	9	19	28	29	23	15	1	-
Male	1	2	4	2	7	6	14	11	5	2	-	-
Female	-	1	5	2	2	13	14	18	18	13	1	-
Black, both sexes	-	-	-	-	-	2	-	-	1	-	1	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	1	1	1	-	-	1	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
New York												
All races, both sexes	34	44	53	69	124	209	300	359	290	110	15	-
Male	27	21	35	39	68	110	138	136	76	25	-	-
Female	7	23	18	30	56	99	162	223	214	85	15	-
White, both sexes	27	34	44	58	115	194	278	341	282	105	14	-
Male	22	19	29	30	64	102	124	128	72	25	-	-
Female	5	15	15	28	51	92	154	213	210	80	14	-
Black, both sexes	7	8	7	9	8	13	19	14	5	4	1	-
Male	5	1	5	7	3	6	11	5	4	-	-	-
Female	2	7	2	2	5	7	8	9	1	4	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	2	1	2	3	4	2	1	-	-
Male	-	1	-	2	1	2	3	3	-	-	-	-
Female	-	1	1	-	-	-	-	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
North Carolina												
All races, both sexes	35	52	86	86	120	174	228	240	182	65	16	-
Male	25	28	54	57	61	89	100	83	50	7	4	-
Female	10	24	32	29	59	85	128	157	132	58	12	-
White, both sexes	24	30	58	60	86	141	187	204	163	57	11	-
Male	16	16	36	40	44	71	89	73	46	6	2	-
Female	8	14	22	20	42	70	98	131	117	51	9	-
Black, both sexes	11	22	27	24	31	30	38	35	19	7	5	-
Male	9	12	17	15	16	16	9	10	4	1	2	-
Female	2	10	10	9	15	14	29	25	15	6	3	-
American Indian 1/, both sexes	-	-	1	2	1	2	2	1	-	1	-	-
Male	-	-	1	2	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	1	1	5	6	11	12	15	10	6	2	-
Male	3	1	-	4	4	4	8	5	3	3	-	-
Female	2	-	1	1	2	7	4	10	7	3	2	-
White, both sexes	4	1	1	4	6	11	12	15	10	6	2	-
Male	2	1	-	4	4	4	8	5	3	3	-	-
Female	2	-	1	-	2	7	4	10	7	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Ohio												
All races, both sexes	38	53	63	79	101	190	231	264	214	83	9	-
Male	26	28	38	39	48	83	76	93	56	20	2	-
Female	12	25	25	40	53	107	155	171	158	63	7	-
White, both sexes	28	42	54	70	90	172	218	248	203	80	8	-
Male	20	22	33	34	42	74	73	86	52	20	2	-
Female	8	20	21	36	48	98	145	162	151	60	6	-
Black, both sexes	9	10	8	7	11	18	13	16	11	3	1	-
Male	6	5	4	4	6	9	3	7	4	-	-	-
Female	3	5	4	3	5	9	10	9	7	3	1	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	13	13	17	20	41	45	81	71	43	21	8	-
Male	10	10	9	10	22	24	32	20	9	4	1	-
Female	3	3	8	10	19	21	49	51	34	17	7	-
White, both sexes	13	10	13	13	32	42	71	67	39	21	8	-
Male	10	7	6	5	21	21	27	19	8	4	1	-
Female	3	3	7	8	11	21	44	48	31	17	7	-
Black, both sexes	-	3	3	3	4	1	4	3	-	-	-	-
Male	-	3	3	1	1	1	2	1	-	-	-	-
Female	-	-	-	2	3	-	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	4	5	2	6	1	3	-	-	-
Male	-	-	-	4	-	2	3	-	1	-	-	-
Female	-	-	1	-	5	-	3	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Oregon												
All races, both sexes	11	13	21	22	26	44	75	84	95	22	6	-
Male	5	7	13	14	18	21	21	28	25	7	-	-
Female	6	6	8	8	8	23	54	56	70	15	6	-
White, both sexes	11	11	20	21	26	43	73	83	94	22	6	-
Male	5	6	12	14	18	21	21	28	24	7	-	-
Female	6	5	8	7	8	22	52	55	70	15	6	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	-	-	-	-
Pennsylvania												
All races, both sexes	81	109	148	171	220	365	495	478	356	133	32	1
Male	56	70	102	100	113	169	227	166	99	28	5	1
Female	25	39	46	71	107	196	268	312	257	105	27	-
White, both sexes	66	91	128	144	194	331	453	458	342	125	28	-
Male	46	61	90	84	98	156	210	157	92	26	5	-
Female	20	30	38	60	96	175	243	301	250	99	23	-
Black, both sexes	14	16	19	26	26	32	41	18	14	8	4	1
Male	10	8	12	15	15	13	17	7	7	2	-	1
Female	4	8	7	11	11	19	24	11	7	6	4	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	-	-	2	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	1	1	1	-	-	2	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Rhode Island												
All races, both sexes	2	-	1	5	5	13	14	23	15	7	3	-
Male	2	-	-	3	4	4	4	5	4	1	1	-
Female	-	-	1	2	1	9	10	18	11	6	2	-
White, both sexes	2	-	1	5	5	13	14	23	15	7	3	-
Male	2	-	-	3	4	4	4	5	4	1	1	-
Female	-	-	1	2	1	9	10	18	11	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	43	47	63	60	87	99	134	124	77	28	6	-
Male	29	36	36	34	54	51	61	34	16	7	1	-
Female	14	11	27	26	33	48	73	90	61	21	5	-
White, both sexes	18	21	30	39	55	69	96	92	58	23	4	-
Male	13	17	19	23	34	36	41	28	11	6	1	-
Female	5	4	11	16	21	33	55	64	47	17	3	-
Black, both sexes	25	26	33	21	31	28	36	32	18	5	2	-
Male	16	19	17	11	20	14	19	6	5	1	-	-
Female	9	7	16	10	11	14	17	26	13	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
South Dakota												
All races, both sexes	1	2	-	3	-	1	6	11	11	6	3	-
Male	-	2	-	3	-	-	3	4	5	1	1	-
Female	1	-	-	-	-	1	3	7	6	5	2	-
White, both sexes	1	1	-	3	-	1	6	11	11	6	3	-
Male	-	1	-	3	-	-	3	4	5	1	1	-
Female	1	-	-	-	-	1	3	7	6	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	41	56	54	54	70	80	146	134	84	31	5	-
Male	27	36	32	32	32	30	51	44	32	8	1	-
Female	14	20	22	22	38	50	95	90	52	23	4	-
White, both sexes	34	52	38	43	57	64	134	122	73	26	4	-
Male	23	34	20	22	24	27	47	40	26	7	1	-
Female	11	18	18	21	33	37	87	82	47	19	3	-
Black, both sexes	7	4	16	11	12	16	12	12	11	5	1	-
Male	4	2	12	10	7	3	4	4	6	1	-	-
Female	3	2	4	1	5	13	8	8	5	4	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Texas												
All races, both sexes	53	93	90	113	159	246	320	312	258	105	20	-
Male	37	51	62	62	96	119	123	128	78	23	3	-
Female	16	42	28	51	63	127	197	184	180	82	17	-
White, both sexes	33	74	72	91	125	208	279	279	240	97	15	-
Male	23	42	49	46	74	97	108	113	75	19	1	-
Female	10	32	23	45	51	111	171	166	165	78	14	-
Black, both sexes	17	16	16	21	30	31	37	28	15	6	5	-
Male	11	7	12	16	18	20	14	13	2	3	2	-
Female	6	9	4	5	12	11	23	15	13	3	3	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	1	3	7	4	5	3	2	-	-
Male	2	2	-	-	3	2	1	2	1	1	-	-
Female	-	1	1	1	-	5	3	3	2	1	-	-
Utah												
All races, both sexes	2	9	11	8	22	25	28	32	27	9	3	-
Male	1	5	10	4	9	11	10	10	8	1	-	-
Female	1	4	1	4	13	14	18	22	19	8	3	-
White, both sexes	2	9	11	8	20	23	28	31	26	9	2	-
Male	1	5	10	4	8	9	10	10	7	1	-	-
Female	1	4	1	4	12	14	18	21	19	8	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Vermont												
All races, both sexes	1	4	2	-	3	4	10	12	13	6	1	-
Male	1	2	-	-	1	1	4	3	2	1	-	-
Female	-	2	2	-	2	3	6	9	11	5	1	-
White, both sexes	1	4	2	-	3	4	10	12	13	6	1	-
Male	1	2	-	-	1	1	4	3	2	1	-	-
Female	-	2	2	-	2	3	6	9	11	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	69	70	80	94	146	184	210	247	189	64	19	-
Male	51	45	50	54	82	102	96	72	48	13	3	-
Female	18	25	30	40	64	82	114	175	141	51	16	-
White, both sexes	54	48	60	65	107	143	174	216	163	55	12	-
Male	40	33	39	37	63	78	84	63	42	13	1	-
Female	14	15	21	28	44	65	90	153	121	42	11	-
Black, both sexes	15	22	19	25	37	41	35	27	25	9	7	-
Male	11	12	10	14	18	24	11	8	5	-	2	-
Female	4	10	9	11	19	17	24	19	20	9	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	4	2	-	1	4	1	-	-	-
Male	-	-	1	3	1	-	1	1	1	-	-	-
Female	-	-	-	1	1	-	-	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Washington												
All races, both sexes	10	10	17	20	28	42	59	74	59	20	5	-
Male	4	6	11	11	17	20	25	29	18	6	-	-
Female	6	4	6	9	11	22	34	45	41	14	5	-
White, both sexes	9	8	17	19	25	40	57	71	59	20	5	-
Male	4	5	11	11	15	19	25	27	18	6	-	-
Female	5	3	6	8	10	21	32	44	41	14	5	-
Black, both sexes	-	1	-	-	-	-	2	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	2	-	1	-	-	-	-
Male	-	1	-	-	1	1	-	1	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	17	24	23	27	45	69	103	88	62	35	6	-
Male	14	16	12	16	27	28	52	30	21	7	-	-
Female	3	8	11	11	18	41	51	58	41	28	6	-
White, both sexes	14	23	23	26	42	67	97	87	59	31	5	-
Male	11	15	12	16	26	28	49	30	21	6	-	-
Female	3	8	11	10	16	39	48	57	38	25	5	-
Black, both sexes	3	1	-	1	3	2	6	1	3	3	1	-
Male	3	1	-	-	1	-	3	-	-	1	-	-
Female	-	-	-	1	2	2	3	1	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Wisconsin												
All races, both sexes	14	28	25	37	40	84	112	137	122	53	11	-
Male	8	19	12	22	25	40	57	53	31	14	2	-
Female	6	9	13	15	15	44	55	84	91	39	9	-
White, both sexes	13	24	22	33	37	83	107	134	120	52	11	-
Male	8	18	12	19	24	40	52	51	31	13	2	-
Female	5	6	10	14	13	43	55	83	89	39	9	-
Black, both sexes	1	4	2	4	3	-	4	2	2	1	-	-
Male	-	1	-	3	1	-	4	1	-	1	-	-
Female	1	3	2	1	2	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	6	1	4	8	6	25	9	14	4	-	-
Male	2	2	1	2	3	4	8	1	4	-	-	-
Female	1	4	-	2	5	2	17	8	10	4	-	-
White, both sexes	3	5	1	4	8	6	25	9	14	4	-	-
Male	2	1	1	2	3	4	8	1	4	-	-	-
Female	1	4	-	2	5	2	17	8	10	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
United States												
All races, both sexes	60,337	19	9	4	8	13	24	26	60	107	225	463
Male	23,918	11	5	1	4	8	14	16	40	62	137	293
Female	36,419	8	4	3	4	5	10	10	20	45	88	170
White, both sexes	53,980	8	8	3	5	7	11	12	26	55	137	273
Male	21,236	4	4	-	3	3	7	9	17	29	80	180
Female	32,744	4	4	3	2	4	4	3	9	26	57	93
Black, both sexes	5,564	11	1	1	3	6	11	11	33	46	82	180
Male	2,335	7	1	1	1	5	7	5	22	27	51	105
Female	3,229	4	-	-	2	1	4	6	11	19	31	75
American Indian 1/, both sexes	218	-	-	-	-	-	2	1	-	4	5	7
Male	101	-	-	-	-	-	-	-	-	4	5	5
Female	117	-	-	-	-	-	2	1	-	-	-	2
Asian or Pacific Islander 2/, both sexes	575	-	-	-	-	-	-	2	1	2	1	3
Male	246	-	-	-	-	-	-	2	1	2	1	3
Female	329	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	2,046	1	-	-	-	2	2	-	5	7	20	41
Male	822	-	-	-	-	2	2	-	5	5	13	29
Female	1,224	1	-	-	-	-	-	-	-	2	7	12
White, both sexes	1,619	-	-	-	-	1	2	-	3	4	12	22
Male	640	-	-	-	-	1	2	-	3	2	8	17
Female	979	-	-	-	-	-	-	-	-	2	4	5
Black, both sexes	422	1	-	-	-	1	-	-	2	2	8	19
Male	180	-	-	-	-	1	-	-	2	2	5	12
Female	242	1	-	-	-	-	-	-	-	-	3	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Alaska												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	745	-	1	-	-	1	-	1	-	1	-	2
Male	330	-	-	-	-	-	-	1	-	-	-	2
Female	415	-	1	-	-	1	-	-	-	1	-	-
White, both sexes	719	-	1	-	-	1	-	1	-	1	-	2
Male	317	-	-	-	-	-	-	1	-	-	-	2
Female	402	-	1	-	-	1	-	-	-	1	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Arkansas												
All races, both sexes	710	-	-	-	1	1	-	-	1	-	4	10
Male	297	-	-	-	-	-	-	-	1	-	2	6
Female	413	-	-	-	1	1	-	-	-	-	2	4
White, both sexes	636	-	-	-	-	1	-	-	-	-	2	5
Male	264	-	-	-	-	-	-	-	-	-	1	3
Female	372	-	-	-	-	1	-	-	-	-	1	2
Black, both sexes	71	-	-	-	1	-	-	-	1	-	2	5
Male	31	-	-	-	-	-	-	-	1	-	1	3
Female	40	-	-	-	1	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,781	-	-	-	1	1	1	1	4	7	14	34
Male	1,926	-	-	-	-	1	1	1	3	4	8	22
Female	2,855	-	-	-	1	-	-	-	1	3	6	12
White, both sexes	4,233	-	-	-	1	-	1	-	2	3	10	20
Male	1,695	-	-	-	-	-	1	-	1	1	5	13
Female	2,538	-	-	-	1	-	-	-	1	2	5	7
Black, both sexes	297	-	-	-	-	1	-	-	1	2	3	12
Male	128	-	-	-	-	1	-	-	1	1	2	7
Female	169	-	-	-	-	-	-	-	-	1	1	5
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	1	-	2
Male	11	-	-	-	-	-	-	-	-	1	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	226	-	-	-	-	-	-	1	1	1	1	-
Male	92	-	-	-	-	-	-	1	1	1	1	-
Female	134	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Colorado												
All races, both sexes	712	-	-	-	1	-	-	-	-	1	2	2
Male	278	-	-	-	1	-	-	-	-	-	2	1
Female	434	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	682	-	-	-	1	-	-	-	-	1	2	1
Male	268	-	-	-	1	-	-	-	-	-	2	-
Female	414	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	736	-	-	-	-	-	-	-	-	-	1	4
Male	288	-	-	-	-	-	-	-	-	-	1	1
Female	448	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	705	-	-	-	-	-	-	-	-	-	1	3
Male	278	-	-	-	-	-	-	-	-	-	1	1
Female	427	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Delaware												
All races, both sexes	127	-	-	-	-	-	-	-	-	1	-	2
Male	54	-	-	-	-	-	-	-	-	1	-	2
Female	73	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	111	-	-	-	-	-	-	-	-	-	-	2
Male	49	-	-	-	-	-	-	-	-	-	-	2
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	41	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Florida												
All races, both sexes	1,900	1	-	-	-	-	-	1	1	1	6	10
Male	793	1	-	-	-	-	-	1	-	-	2	6
Female	1,107	-	-	-	-	-	-	-	1	1	4	4
White, both sexes	1,759	1	-	-	-	-	-	-	-	1	3	2
Male	733	1	-	-	-	-	-	-	-	-	1	2
Female	1,026	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	134	-	-	-	-	-	-	1	1	-	3	8
Male	59	-	-	-	-	-	-	1	-	-	1	4
Female	75	-	-	-	-	-	-	-	1	-	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,168	1	1	-	-	-	2	4	4	12	19	29
Male	887	1	1	-	-	-	2	2	2	6	11	19
Female	1,281	-	-	-	-	-	-	2	2	6	8	10
White, both sexes	1,652	-	1	-	-	-	-	-	-	3	12	12
Male	678	-	1	-	-	-	-	-	-	1	9	10
Female	974	-	-	-	-	-	-	-	-	2	3	2
Black, both sexes	498	1	-	-	-	-	2	4	4	9	7	17
Male	206	1	-	-	-	-	2	2	2	5	2	9
Female	292	-	-	-	-	-	-	2	2	4	5	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Hawaii												
All races, both sexes	146	-	-	-	-	-	-	1	-	1	1	3
Male	75	-	-	-	-	-	-	1	-	1	1	2
Female	71	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	101	-	-	-	-	-	-	1	-	1	-	2
Male	55	-	-	-	-	-	-	1	-	1	-	2
Female	46	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	270	-	-	-	-	-	-	-	-	-	1	1
Male	103	-	-	-	-	-	-	-	-	-	-	-
Female	167	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	266	-	-	-	-	-	-	-	-	-	1	-
Male	103	-	-	-	-	-	-	-	-	-	-	-
Female	163	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Illinois												
All races, both sexes	3,103	3	-	1	-	-	1	1	-	8	4	16
Male	1,207	1	-	-	-	-	1	1	-	5	3	12
Female	1,896	2	-	1	-	-	-	-	-	3	1	4
White, both sexes	2,794	1	-	1	-	-	1	-	-	5	2	8
Male	1,074	-	-	-	-	-	1	-	-	3	1	8
Female	1,720	1	-	1	-	-	-	-	-	2	1	-
Black, both sexes	293	2	-	-	-	-	-	1	-	3	2	8
Male	125	1	-	-	-	-	-	1	-	2	2	4
Female	168	1	-	-	-	-	-	-	-	1	-	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,941	-	-	-	-	2	1	-	1	2	10	21
Male	757	-	-	-	-	2	-	-	1	1	5	14
Female	1,184	-	-	-	-	-	1	-	-	1	5	7
White, both sexes	1,816	-	-	-	-	1	-	-	-	1	8	14
Male	701	-	-	-	-	1	-	-	-	1	3	10
Female	1,115	-	-	-	-	-	-	-	-	-	5	4
Black, both sexes	125	-	-	-	-	1	1	-	1	1	2	7
Male	56	-	-	-	-	1	-	-	1	-	2	4
Female	69	-	-	-	-	-	1	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Iowa												
All races, both sexes	285	-	-	-	-	-	-	-	-	-	1	-
Male	103	-	-	-	-	-	-	-	-	-	1	-
Female	182	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	277	-	-	-	-	-	-	-	-	-	1	-
Male	102	-	-	-	-	-	-	-	-	-	1	-
Female	175	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	994	-	-	-	-	-	-	-	-	-	3	3
Male	377	-	-	-	-	-	-	-	-	-	3	3
Female	617	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	943	-	-	-	-	-	-	-	-	-	1	2
Male	359	-	-	-	-	-	-	-	-	-	1	2
Female	584	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	43	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Kentucky												
All races, both sexes	1,381	-	-	-	-	1	-	1	1	4	8	13
Male	533	-	-	-	-	1	-	1	1	1	6	10
Female	848	-	-	-	-	-	-	-	-	3	2	3
White, both sexes	1,313	-	-	-	-	-	-	1	1	2	8	11
Male	504	-	-	-	-	-	-	1	1	1	6	9
Female	809	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	64	-	-	-	-	1	-	-	-	2	-	2
Male	29	-	-	-	-	1	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,291	-	1	-	1	-	1	-	3	4	9	15
Male	537	-	1	-	-	-	-	-	1	3	7	10
Female	754	-	-	-	1	-	1	-	2	1	2	5
White, both sexes	984	-	-	-	1	-	1	-	1	1	2	6
Male	405	-	-	-	-	-	-	-	-	1	2	4
Female	579	-	-	-	1	-	1	-	1	-	-	2
Black, both sexes	304	-	1	-	-	-	-	-	2	3	7	9
Male	131	-	1	-	-	-	-	-	1	2	5	6
Female	173	-	-	-	-	-	-	-	1	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Maine												
All races, both sexes	312	-	-	-	-	1	-	-	-	2	-	2
Male	129	-	-	-	-	-	-	-	-	1	-	1
Female	183	-	-	-	-	1	-	-	-	1	-	1
White, both sexes	310	-	-	-	-	1	-	-	-	1	-	2
Male	128	-	-	-	-	-	-	-	-	-	-	1
Female	182	-	-	-	-	1	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	679	-	-	-	-	-	-	1	3	2	2	7
Male	258	-	-	-	-	-	-	1	1	1	2	5
Female	421	-	-	-	-	-	-	-	2	1	-	2
White, both sexes	556	-	-	-	-	-	-	1	-	-	1	2
Male	208	-	-	-	-	-	-	1	-	-	1	1
Female	348	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	111	-	-	-	-	-	-	-	3	2	1	5
Male	46	-	-	-	-	-	-	-	1	1	1	4
Female	65	-	-	-	-	-	-	-	2	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Massachusetts												
All races, both sexes	1,798	-	1	-	-	-	-	1	1	-	3	5
Male	665	-	1	-	-	-	-	1	1	-	2	5
Female	1,133	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1,745	-	1	-	-	-	-	1	1	-	3	4
Male	645	-	1	-	-	-	-	1	1	-	2	4
Female	1,100	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,588	1	-	-	-	-	1	1	6	1	8	12
Male	1,005	-	-	-	-	-	1	-	5	1	5	8
Female	1,583	1	-	-	-	-	-	1	1	-	3	4
White, both sexes	2,338	-	-	-	-	-	-	-	1	1	4	5
Male	891	-	-	-	-	-	-	-	1	1	1	2
Female	1,447	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	233	1	-	-	-	-	1	1	5	-	4	6
Male	102	-	-	-	-	-	1	-	4	-	4	5
Female	131	1	-	-	-	-	-	1	1	-	-	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Minnesota												
All races, both sexes	1,177	-	-	1	-	-	1	-	1	-	2	3
Male	488	-	-	-	-	-	-	-	1	-	1	2
Female	689	-	-	1	-	-	1	-	-	-	1	1
White, both sexes	1,145	-	-	1	-	-	1	-	1	-	1	2
Male	470	-	-	-	-	-	-	-	1	-	1	1
Female	675	-	-	1	-	-	1	-	-	-	-	1
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,438	-	-	-	-	-	3	1	2	4	6	18
Male	546	-	-	-	-	-	1	-	1	3	2	7
Female	892	-	-	-	-	-	2	1	1	1	4	11
White, both sexes	1,045	-	-	-	-	-	1	-	1	2	3	7
Male	401	-	-	-	-	-	-	-	-	2	1	3
Female	644	-	-	-	-	-	1	-	1	-	2	4
Black, both sexes	388	-	-	-	-	-	2	1	1	2	3	11
Male	142	-	-	-	-	-	1	-	1	1	1	4
Female	246	-	-	-	-	-	1	1	-	1	2	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Missouri												
All races, both sexes	1,498	-	-	-	-	-	-	-	1	3	2	12
Male	544	-	-	-	-	-	-	-	1	3	1	7
Female	954	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	1,406	-	-	-	-	-	-	-	1	3	1	11
Male	508	-	-	-	-	-	-	-	1	3	-	7
Female	898	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	91	-	-	-	-	-	-	-	-	-	1	1
Male	36	-	-	-	-	-	-	-	-	-	1	-
Female	55	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	335	-	-	-	-	-	-	-	1	-	-	1
Male	139	-	-	-	-	-	-	-	1	-	-	-
Female	196	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	327	-	-	-	-	-	-	-	1	-	-	-
Male	135	-	-	-	-	-	-	-	1	-	-	-
Female	192	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Nebraska												
All races, both sexes	574	-	-	-	-	-	1	-	-	2	-	6
Male	217	-	-	-	-	-	1	-	-	-	-	4
Female	357	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	556	-	-	-	-	-	1	-	-	2	-	5
Male	209	-	-	-	-	-	1	-	-	-	-	3
Female	347	-	-	-	-	-	-	-	-	2	-	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	530	1	-	-	-	-	-	-	2	1	1	5
Male	245	-	-	-	-	-	-	-	1	1	-	3
Female	285	1	-	-	-	-	-	-	1	-	1	2
White, both sexes	487	1	-	-	-	-	-	-	2	1	-	3
Male	226	-	-	-	-	-	-	-	1	1	-	2
Female	261	1	-	-	-	-	-	-	1	-	-	1
Black, both sexes	26	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
New Hampshire												
All races, both sexes	271	-	-	-	-	-	-	-	-	-	-	-
Male	100	-	-	-	-	-	-	-	-	-	-	-
Female	171	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	271	-	-	-	-	-	-	-	-	-	-	-
Male	100	-	-	-	-	-	-	-	-	-	-	-
Female	171	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,523	3	-	-	-	-	1	-	-	3	9	10
Male	603	2	-	-	-	-	-	-	-	3	6	6
Female	920	1	-	-	-	-	1	-	-	-	3	4
White, both sexes	1,374	1	-	-	-	-	-	-	-	3	4	7
Male	541	-	-	-	-	-	-	-	-	3	4	3
Female	833	1	-	-	-	-	-	-	-	-	-	4
Black, both sexes	138	2	-	-	-	-	1	-	-	-	5	3
Male	58	2	-	-	-	-	-	-	-	-	2	3
Female	80	-	-	-	-	-	1	-	-	-	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
New Mexico												
All races, both sexes	418	-	-	-	-	-	-	-	1	-	1	5
Male	162	-	-	-	-	-	-	-	-	-	1	3
Female	256	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	407	-	-	-	-	-	-	-	1	-	-	5
Male	156	-	-	-	-	-	-	-	-	-	-	3
Female	251	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,784	1	-	-	-	-	-	1	2	5	6	11
Male	1,098	-	-	-	-	-	-	-	1	3	1	8
Female	1,686	1	-	-	-	-	-	1	1	2	5	3
White, both sexes	2,559	-	-	-	-	-	-	1	1	3	6	6
Male	1,016	-	-	-	-	-	-	-	-	3	1	6
Female	1,543	-	-	-	-	-	-	1	1	-	5	-
Black, both sexes	197	1	-	-	-	-	-	-	1	2	-	5
Male	70	-	-	-	-	-	-	-	1	-	-	2
Female	127	1	-	-	-	-	-	-	-	2	-	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
North Carolina												
All races, both sexes	1,601	2	1	-	-	-	1	2	2	2	9	10
Male	620	2	-	-	-	-	-	1	2	2	6	8
Female	981	-	1	-	-	-	1	1	-	-	3	2
White, both sexes	1,310	1	1	-	-	-	-	1	-	1	3	4
Male	494	1	-	-	-	-	-	1	-	1	2	4
Female	816	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	281	1	-	-	-	-	1	1	2	1	6	6
Male	118	1	-	-	-	-	-	-	2	1	4	4
Female	163	-	-	-	-	-	1	1	-	-	2	2
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	109	-	-	-	-	-	-	-	-	-	1	-
Male	32	-	-	-	-	-	-	-	-	-	1	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	101	-	-	-	-	-	-	-	-	-	1	-
Male	29	-	-	-	-	-	-	-	-	-	1	-
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Ohio												
All races, both sexes	1,891	1	1	-	-	-	-	1	1	4	6	10
Male	737	1	1	-	-	-	-	1	1	3	4	6
Female	1,154	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	1,733	1	1	-	-	-	-	1	1	2	5	5
Male	674	1	1	-	-	-	-	1	1	1	4	3
Female	1,059	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	153	-	-	-	-	-	-	-	-	2	1	5
Male	60	-	-	-	-	-	-	-	-	2	-	3
Female	93	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,330	-	-	-	-	-	-	2	1	3	6	8
Male	537	-	-	-	-	-	-	1	-	3	5	5
Female	793	-	-	-	-	-	-	1	1	-	1	3
White, both sexes	1,208	-	-	-	-	-	-	1	-	1	4	6
Male	485	-	-	-	-	-	-	1	-	1	3	4
Female	723	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	65	-	-	-	-	-	-	-	1	1	-	1
Male	24	-	-	-	-	-	-	-	-	1	-	-
Female	41	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	54	-	-	-	-	-	-	1	-	1	2	1
Male	28	-	-	-	-	-	-	-	-	1	2	1
Female	26	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Oregon												
All races, both sexes	783	-	-	-	-	-	-	-	-	1	1	3
Male	343	-	-	-	-	-	-	-	-	-	1	3
Female	440	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	762	-	-	-	-	-	-	-	-	1	1	2
Male	335	-	-	-	-	-	-	-	-	-	1	2
Female	427	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,681	1	-	1	-	2	1	1	2	4	6	19
Male	1,371	1	-	1	-	-	1	-	1	2	5	12
Female	2,310	-	-	-	-	2	-	1	1	2	1	7
White, both sexes	3,434	1	-	-	-	1	-	1	-	2	4	12
Male	1,264	1	-	-	-	-	-	-	-	-	3	7
Female	2,170	-	-	-	-	1	-	1	-	2	1	5
Black, both sexes	238	-	-	1	-	1	1	-	2	2	2	7
Male	102	-	-	1	-	-	1	-	1	2	2	5
Female	136	-	-	-	-	1	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Rhode Island												
All races, both sexes	166	-	-	-	-	-	-	-	-	-	-	1
Male	64	-	-	-	-	-	-	-	-	-	-	-
Female	102	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	161	-	-	-	-	-	-	-	-	-	-	1
Male	61	-	-	-	-	-	-	-	-	-	-	-
Female	100	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	979	1	-	-	-	-	1	-	3	2	7	13
Male	435	-	-	-	-	-	1	-	1	2	4	9
Female	544	1	-	-	-	-	-	-	2	-	3	4
White, both sexes	740	1	-	-	-	-	1	-	1	-	3	8
Male	332	-	-	-	-	-	1	-	-	-	1	6
Female	408	1	-	-	-	-	-	-	1	-	2	2
Black, both sexes	235	-	-	-	-	-	-	-	2	2	4	5
Male	103	-	-	-	-	-	-	-	1	2	3	3
Female	132	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
South Dakota												
All races, both sexes	64	-	-	-	-	-	2	-	-	-	2	-
Male	24	-	-	-	-	-	-	-	-	-	2	-
Female	40	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	2	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	742	-	-	-	1	-	-	-	2	4	6	12
Male	277	-	-	-	1	-	-	-	2	1	4	6
Female	465	-	-	-	-	-	-	-	-	3	2	6
White, both sexes	626	-	-	-	-	-	-	-	2	2	3	8
Male	233	-	-	-	-	-	-	-	2	1	2	4
Female	393	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	114	-	-	-	1	-	-	-	-	2	3	4
Male	44	-	-	-	1	-	-	-	-	-	2	2
Female	70	-	-	-	-	-	-	-	-	2	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Texas												
All races, both sexes	4,371	1	3	-	1	2	3	3	8	5	25	48
Male	1,784	1	1	-	1	2	2	2	5	2	10	29
Female	2,587	-	2	-	-	-	1	1	3	3	15	19
White, both sexes	3,877	-	3	-	1	1	2	1	5	3	17	34
Male	1,579	-	1	-	1	1	1	1	3	1	5	20
Female	2,298	-	2	-	-	-	1	-	2	2	12	14
Black, both sexes	466	1	-	-	-	1	1	2	3	2	8	14
Male	194	1	-	-	-	1	1	1	2	1	5	9
Female	272	-	-	-	-	-	-	1	1	1	3	5
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	617	-	-	-	-	-	-	1	-	1	1	2
Male	226	-	-	-	-	-	-	-	-	-	1	1
Female	391	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	608	-	-	-	-	-	-	1	-	1	1	2
Male	221	-	-	-	-	-	-	-	-	-	1	1
Female	387	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Vermont												
All races, both sexes	69	-	-	1	1	-	-	-	-	-	-	-
Male	26	-	-	-	1	-	-	-	-	-	-	-
Female	43	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	69	-	-	1	1	-	-	-	-	-	-	-
Male	26	-	-	-	1	-	-	-	-	-	-	-
Female	43	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,717	1	-	-	1	-	1	1	1	4	8	15
Male	690	1	-	-	-	-	1	1	1	3	6	9
Female	1,027	-	-	-	1	-	-	-	-	1	2	6
White, both sexes	1,388	-	-	-	-	-	-	1	-	1	1	6
Male	526	-	-	-	-	-	-	1	-	1	1	4
Female	862	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	311	1	-	-	1	-	1	-	1	3	7	9
Male	155	1	-	-	-	-	1	-	1	2	5	5
Female	156	-	-	-	1	-	-	-	-	1	2	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Washington												
All races, both sexes	534	-	-	-	-	-	-	-	-	2	-	3
Male	225	-	-	-	-	-	-	-	-	1	-	1
Female	309	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	513	-	-	-	-	-	-	-	-	-	-	3
Male	215	-	-	-	-	-	-	-	-	-	-	1
Female	298	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	2	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	638	-	-	-	-	-	-	-	-	3	1	6
Male	221	-	-	-	-	-	-	-	-	-	-	4
Female	417	-	-	-	-	-	-	-	-	3	1	2
White, both sexes	623	-	-	-	-	-	-	-	-	3	1	5
Male	215	-	-	-	-	-	-	-	-	-	-	3
Female	408	-	-	-	-	-	-	-	-	3	1	2
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Wisconsin												
All races, both sexes	1,523	-	-	-	-	-	-	-	-	-	2	7
Male	628	-	-	-	-	-	-	-	-	-	2	2
Female	895	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	1,484	-	-	-	-	-	-	-	-	-	2	6
Male	612	-	-	-	-	-	-	-	-	-	2	2
Female	872	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	146	-	-	-	-	-	-	-	-	-	1	1
Male	67	-	-	-	-	-	-	-	-	-	-	-
Female	79	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	144	-	-	-	-	-	-	-	-	-	1	1
Male	66	-	-	-	-	-	-	-	-	-	-	-
Female	78	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
United States												
All races, both sexes	762	1,202	1,599	2,275	3,685	6,547	10,762	13,590	12,139	5,491	1,324	3
Male	487	724	924	1,286	1,916	3,223	4,652	5,067	3,630	1,197	219	2
Female	275	478	675	989	1,769	3,324	6,110	8,523	8,509	4,294	1,105	1
White, both sexes	484	864	1,259	1,824	3,112	5,788	9,778	12,635	11,346	5,142	1,200	3
Male	310	546	738	1,058	1,616	2,898	4,266	4,755	3,404	1,111	196	2
Female	174	318	521	766	1,496	2,890	5,512	7,880	7,942	4,031	1,004	1
Black, both sexes	254	311	316	413	517	654	845	806	651	305	107	-
Male	162	160	177	209	267	282	327	257	177	69	16	-
Female	92	151	139	204	250	372	518	549	474	236	91	-
American Indian 1/, both sexes	8	8	14	20	20	26	39	29	25	7	3	-
Male	3	7	5	13	11	11	13	14	8	-	2	-
Female	5	1	9	7	9	15	26	15	17	7	1	-
Asian or Pacific Islander 2/, both sexes	16	19	10	18	36	79	100	120	117	37	14	-
Male	12	11	4	6	22	32	46	41	41	17	5	-
Female	4	8	6	12	14	47	54	79	76	20	9	-
Alabama												
All races, both sexes	56	75	79	120	162	241	351	380	345	123	36	-
Male	36	45	47	67	81	101	140	118	101	28	2	-
Female	20	30	32	53	81	140	211	262	244	95	34	-
White, both sexes	27	55	53	86	116	199	288	324	295	107	25	-
Male	16	36	33	52	58	81	116	103	88	23	1	-
Female	11	19	20	34	58	118	172	221	207	84	24	-
Black, both sexes	29	20	26	34	46	41	63	55	48	16	11	-
Male	20	9	14	15	23	20	24	14	13	5	1	-
Female	9	11	12	19	23	21	39	41	35	11	10	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Alaska												
All races, both sexes	2	1	2	-	6	7	10	14	11	5	2	-
Male	2	1	-	-	3	2	5	5	4	-	1	-
Female	-	-	2	-	3	5	5	9	7	5	1	-
White, both sexes	1	1	1	-	4	3	9	11	9	4	2	-
Male	1	1	-	-	1	1	5	4	4	-	1	-
Female	-	-	1	-	3	2	4	7	5	4	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	1	4	1	2	2	1	-	-
Male	1	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	1	-	-	3	1	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	10	12	22	33	50	88	143	176	143	50	12	-
Male	4	8	16	19	25	48	67	74	49	13	4	-
Female	6	4	6	14	25	40	76	102	94	37	8	-
White, both sexes	9	12	21	29	48	83	138	171	142	49	11	-
Male	3	8	15	17	23	46	64	73	49	13	3	-
Female	6	4	6	12	25	37	74	98	93	36	8	-
Black, both sexes	1	-	-	1	-	1	2	3	-	1	-	-
Male	1	-	-	1	-	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	1	3	2	4	3	2	1	-	1	-
Male	-	-	1	1	2	1	1	-	-	-	1	-
Female	-	-	-	2	-	3	2	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Arkansas												
All races, both sexes	9	22	20	28	46	92	120	169	120	56	11	-
Male	7	14	13	18	25	49	56	56	37	12	1	-
Female	2	8	7	10	21	43	64	113	83	44	10	-
White, both sexes	5	16	15	22	41	90	114	161	108	47	9	-
Male	4	9	10	14	22	48	55	55	33	9	1	-
Female	1	7	5	8	19	42	59	106	75	38	8	-
Black, both sexes	4	5	5	5	5	2	5	8	12	9	2	-
Male	3	4	3	4	3	1	-	1	4	3	-	-
Female	1	1	2	1	2	1	5	7	8	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
California												
All races, both sexes	72	121	113	167	290	444	887	1,076	1,002	439	106	1
Male	52	69	60	88	154	201	401	407	339	96	19	-
Female	20	52	53	79	136	243	486	669	663	343	87	1
White, both sexes	49	95	95	131	248	387	793	978	913	409	97	1
Male	36	56	50	73	136	181	361	375	306	83	17	-
Female	13	39	45	58	112	206	432	603	607	326	80	1
Black, both sexes	19	19	15	31	29	30	48	43	33	10	1	-
Male	13	10	9	13	12	12	17	13	14	3	-	-
Female	6	9	6	18	17	18	31	30	19	7	1	-
American Indian 1/, both sexes	-	-	1	1	1	6	6	5	1	-	1	-
Male	-	-	-	1	-	1	3	2	1	-	-	-
Female	-	-	1	-	1	5	3	3	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	7	2	4	12	21	40	50	55	20	7	-
Male	3	3	1	1	6	7	20	17	18	10	2	-
Female	1	4	1	3	6	14	20	33	37	10	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Colorado												
All races, both sexes	4	7	15	25	32	68	116	178	156	85	20	-
Male	3	5	10	12	14	35	53	66	47	24	5	-
Female	1	2	5	13	18	33	63	112	109	61	15	-
White, both sexes	4	6	15	23	30	65	111	170	151	82	20	-
Male	3	4	10	11	13	34	51	64	46	24	5	-
Female	1	2	5	12	17	31	60	106	105	58	15	-
Black, both sexes	-	1	-	2	2	-	3	7	2	1	-	-
Male	-	1	-	1	1	-	1	2	1	-	-	-
Female	-	-	-	1	1	-	2	5	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	2	1	2	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	2	1	1	2	-	-	-
Connecticut												
All races, both sexes	2	5	9	26	36	85	136	175	159	76	22	-
Male	2	3	5	11	20	52	59	63	57	11	3	-
Female	-	2	4	15	16	33	77	112	102	65	19	-
White, both sexes	1	5	9	22	31	84	131	171	154	72	21	-
Male	1	3	5	11	17	51	57	62	57	10	2	-
Female	-	2	4	11	14	33	74	109	97	62	19	-
Black, both sexes	-	-	-	4	5	1	5	4	4	3	1	-
Male	-	-	-	-	3	1	2	1	-	1	1	-
Female	-	-	-	4	2	-	3	3	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Delaware												
All races, both sexes	-	5	8	5	12	13	15	32	23	10	1	-
Male	-	3	7	2	5	5	9	9	9	1	1	-
Female	-	2	1	3	7	8	6	23	14	9	-	-
White, both sexes	-	3	8	3	11	11	14	29	21	8	1	-
Male	-	3	7	1	5	5	8	8	8	1	1	-
Female	-	-	1	2	6	6	6	21	13	7	-	-
Black, both sexes	-	2	-	2	1	2	1	3	2	1	-	-
Male	-	-	-	1	-	-	1	1	1	-	-	-
Female	-	2	-	1	1	2	-	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	1	1	6	4	4	11	9	9	3	3	-
Male	1	1	1	3	2	2	6	2	3	-	1	-
Female	1	-	-	3	2	2	5	7	6	3	2	-
White, both sexes	-	1	-	-	-	2	2	4	3	1	1	-
Male	-	1	-	-	-	2	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	3	3	1	1	-
Black, both sexes	2	-	1	6	4	2	9	5	6	2	2	-
Male	1	-	1	3	2	-	5	1	3	-	1	-
Female	1	-	-	3	2	2	4	4	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Florida												
All races, both sexes	20	24	48	63	107	183	331	440	433	189	42	-
Male	15	13	29	36	61	88	158	167	152	54	10	-
Female	5	11	19	27	46	95	173	273	281	135	32	-
White, both sexes	14	16	36	58	97	166	315	423	405	181	41	-
Male	11	11	20	33	53	80	150	166	141	54	10	-
Female	3	5	16	25	44	86	165	257	264	127	31	-
Black, both sexes	6	7	12	5	10	16	16	14	26	8	1	-
Male	4	2	9	3	8	8	8	-	11	-	-	-
Female	2	5	3	2	2	8	8	14	15	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	3	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	2	2	-	-	-
Georgia												
All races, both sexes	60	76	88	124	164	249	384	441	344	132	34	-
Male	30	57	57	63	83	113	168	137	105	25	5	-
Female	30	19	31	61	81	136	216	304	239	107	29	-
White, both sexes	30	46	58	82	109	193	321	365	283	111	26	-
Male	15	37	38	43	54	89	155	117	83	21	5	-
Female	15	9	20	39	55	104	166	248	200	90	21	-
Black, both sexes	29	30	29	42	52	53	58	73	60	20	8	-
Male	15	20	19	20	26	24	13	20	22	4	-	-
Female	14	10	10	22	26	29	45	53	38	16	8	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	2	3	5	3	1	1	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	1	-	1	-	-	3	5	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Hawaii												
All races, both sexes	5	1	4	7	7	20	20	29	31	13	3	-
Male	4	1	2	5	6	9	11	11	14	5	2	-
Female	1	-	2	2	1	11	9	18	17	8	1	-
White, both sexes	-	-	-	3	1	7	7	12	7	6	-	-
Male	-	-	-	3	1	4	3	4	3	1	-	-
Female	-	-	-	-	-	3	4	8	4	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	4	4	6	13	13	17	24	7	3	-
Male	4	1	2	2	5	5	8	7	11	4	2	-
Female	1	-	2	2	1	8	5	10	13	3	1	-
Idaho												
All races, both sexes	1	-	3	7	14	24	44	68	69	27	11	-
Male	-	-	3	6	7	9	21	25	22	7	3	-
Female	1	-	-	1	7	15	23	43	47	20	8	-
White, both sexes	1	-	3	7	14	24	43	67	69	26	11	-
Male	-	-	3	6	7	9	21	25	22	7	3	-
Female	1	-	-	1	7	15	22	42	47	19	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Illinois												
All races, both sexes	36	61	89	107	168	350	574	652	659	292	81	-
Male	26	32	48	52	94	172	249	254	175	66	16	-
Female	10	29	41	55	74	178	325	398	484	226	65	-
White, both sexes	23	39	70	90	133	313	511	607	634	278	78	-
Male	15	22	38	48	77	152	222	238	171	63	15	-
Female	8	17	32	42	56	161	289	369	463	215	63	-
Black, both sexes	13	21	19	17	34	36	56	40	24	14	3	-
Male	11	10	10	4	16	20	23	13	4	3	1	-
Female	2	11	9	13	18	16	33	27	20	11	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	7	4	1	-	-	-
Male	-	-	-	-	1	-	4	2	-	-	-	-
Female	-	1	-	-	-	1	3	2	1	-	-	-
Indiana												
All races, both sexes	17	34	47	89	124	195	345	425	416	170	42	-
Male	6	21	34	51	62	112	133	155	122	35	3	-
Female	11	13	13	38	62	83	212	270	294	135	39	-
White, both sexes	14	27	39	78	114	182	328	409	398	161	42	-
Male	5	16	27	45	58	107	126	150	118	31	3	-
Female	9	11	12	33	56	75	202	259	280	130	39	-
Black, both sexes	3	7	8	11	10	13	17	16	18	9	-	-
Male	1	5	7	6	4	5	7	5	4	4	-	-
Female	2	2	1	5	6	8	10	11	14	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Iowa												
All races, both sexes	-	3	6	5	15	30	46	54	59	51	15	-
Male	-	-	3	4	4	15	20	20	20	15	1	-
Female	-	3	3	1	11	15	26	34	39	36	14	-
White, both sexes	-	2	6	5	14	28	45	52	59	51	14	-
Male	-	-	3	4	3	15	20	20	20	15	1	-
Female	-	2	3	1	11	13	25	32	39	36	13	-
Black, both sexes	-	1	-	-	1	1	1	2	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Kansas												
All races, both sexes	4	13	15	24	56	81	156	224	252	128	35	-
Male	4	7	8	18	25	35	70	94	67	39	4	-
Female	-	6	7	6	31	46	86	130	185	89	31	-
White, both sexes	2	13	12	19	52	75	150	213	246	124	34	-
Male	2	7	8	15	24	35	68	89	67	37	4	-
Female	-	6	4	4	28	40	82	124	179	87	30	-
Black, both sexes	2	-	3	3	2	6	4	11	5	4	1	-
Male	2	-	-	2	1	-	1	5	-	2	-	-
Female	-	-	3	1	1	6	3	6	5	2	1	-
American Indian 1/, both sexes	-	-	-	2	1	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Kentucky												
All races, both sexes	17	41	48	76	105	185	235	295	234	97	20	-
Male	15	25	23	47	51	82	100	91	54	21	4	-
Female	2	16	25	29	54	103	135	204	180	76	16	-
White, both sexes	16	36	46	70	100	174	225	280	230	94	19	-
Male	14	21	23	41	48	78	95	89	53	21	3	-
Female	2	15	23	29	52	96	130	191	177	73	16	-
Black, both sexes	1	4	2	6	5	10	10	13	4	3	1	-
Male	1	4	-	6	3	4	5	2	1	-	1	-
Female	-	-	2	-	2	6	5	11	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	2	-	-	-	-
Louisiana												
All races, both sexes	23	42	53	65	90	169	225	262	220	91	17	-
Male	11	24	26	36	46	85	96	93	81	13	4	-
Female	12	18	27	29	44	84	129	169	139	78	13	-
White, both sexes	8	29	32	40	62	124	182	217	190	73	15	-
Male	4	18	17	24	34	65	75	78	69	10	4	-
Female	4	11	15	16	28	59	107	139	121	63	11	-
Black, both sexes	15	13	21	25	26	45	43	45	30	17	2	-
Male	7	6	9	12	11	20	21	15	12	3	-	-
Female	8	7	12	13	15	25	22	30	18	14	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Maine												
All races, both sexes	1	4	7	7	15	27	68	80	60	31	7	-
Male	1	2	6	4	10	19	28	34	17	4	2	-
Female	-	2	1	3	5	8	40	46	43	27	5	-
White, both sexes	1	4	7	7	15	27	67	80	60	31	7	-
Male	1	2	6	4	10	19	28	34	17	4	2	-
Female	-	2	1	3	5	8	39	46	43	27	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	12	19	21	44	65	127	160	132	62	17	1
Male	2	5	9	10	26	27	55	63	36	11	3	1
Female	2	7	10	11	18	38	72	97	96	51	14	-
White, both sexes	1	6	14	15	33	51	98	140	121	57	15	1
Male	-	2	7	6	20	19	49	56	32	10	3	1
Female	1	4	7	9	13	32	49	84	89	47	12	-
Black, both sexes	3	6	5	5	11	13	22	19	10	4	2	-
Male	2	3	2	4	6	7	5	7	3	-	-	-
Female	1	3	3	1	5	6	17	12	7	4	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	7	1	1	1	-	-
Male	-	-	-	-	-	1	1	-	1	1	-	-
Female	-	-	-	-	-	-	6	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Massachusetts												
All races, both sexes	12	11	27	31	79	180	339	441	414	201	52	-
Male	9	9	14	17	40	108	145	158	113	35	7	-
Female	3	2	13	14	39	72	194	283	301	166	45	-
White, both sexes	11	11	27	29	74	166	330	433	405	198	51	-
Male	9	9	14	16	38	101	141	154	112	35	7	-
Female	2	2	13	13	36	65	189	279	293	163	44	-
Black, both sexes	-	-	-	2	4	11	7	8	8	2	1	-
Male	-	-	-	1	1	6	4	4	1	-	-	-
Female	-	-	-	1	3	5	3	4	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	3	2	-	1	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	2	2	-	1	1	-	-
Michigan												
All races, both sexes	23	41	53	88	144	283	465	617	549	249	46	-
Male	18	28	28	45	69	146	207	230	151	57	6	-
Female	5	13	25	43	75	137	258	387	398	192	40	-
White, both sexes	15	25	43	74	121	246	426	585	518	233	41	-
Male	12	18	23	39	59	131	186	216	144	53	5	-
Female	3	7	20	35	62	115	240	369	374	180	36	-
Black, both sexes	8	15	10	13	20	34	36	30	29	16	4	-
Male	6	9	5	5	9	12	19	13	5	4	1	-
Female	2	6	5	8	11	22	17	17	24	12	3	-
American Indian 1/, both sexes	-	1	-	1	2	1	2	1	-	-	-	-
Male	-	1	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	1	2	-	1	-
Male	-	-	-	-	1	2	1	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Minnesota												
All races, both sexes	7	4	14	12	45	109	194	297	301	143	43	-
Male	6	3	9	10	33	58	99	130	90	37	9	-
Female	1	1	5	2	12	51	95	167	211	106	34	-
White, both sexes	4	4	14	11	37	104	192	291	297	143	42	-
Male	4	3	9	9	28	54	98	127	89	37	9	-
Female	-	1	5	2	9	50	94	164	208	106	33	-
Black, both sexes	2	-	-	1	5	2	-	3	2	-	1	-
Male	2	-	-	1	3	2	-	1	-	-	-	-
Female	-	-	-	-	2	-	-	2	2	-	1	-
American Indian 1/, both sexes	1	-	-	-	2	1	1	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	3	2	-	-	-
Male	-	-	-	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	1	1	-	1	1	-	-	-
Mississippi												
All races, both sexes	27	50	36	83	88	183	234	292	258	121	32	-
Male	17	31	21	55	47	88	89	92	66	19	7	-
Female	10	19	15	28	41	95	145	200	192	102	25	-
White, both sexes	10	23	19	57	67	134	182	232	202	87	18	-
Male	7	15	11	38	38	70	73	71	55	13	4	-
Female	3	8	8	19	29	64	109	161	147	74	14	-
Black, both sexes	17	26	17	26	21	48	51	58	56	34	14	-
Male	10	15	10	17	9	17	15	21	11	6	3	-
Female	7	11	7	9	12	31	36	37	45	28	11	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Missouri												
All races, both sexes	17	21	28	53	96	152	275	341	322	133	42	-
Male	9	13	13	24	58	65	109	115	86	32	8	-
Female	8	8	15	29	38	87	166	226	236	101	34	-
White, both sexes	14	16	27	49	81	136	260	329	311	128	39	-
Male	8	10	12	23	46	61	107	110	82	30	8	-
Female	6	6	15	26	35	75	153	219	229	98	31	-
Black, both sexes	3	5	1	4	15	16	14	12	11	5	3	-
Male	1	3	1	1	12	4	2	5	4	2	-	-
Female	2	2	-	3	3	12	12	7	7	3	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	6	12	6	12	26	51	91	88	39	1	-
Male	1	3	10	5	4	14	23	42	27	9	-	-
Female	-	3	2	1	8	12	28	49	61	30	1	-
White, both sexes	1	6	11	6	10	25	49	91	87	39	1	-
Male	1	3	9	5	3	13	22	42	27	9	-	-
Female	-	3	2	1	7	12	27	49	60	30	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	1	2	-	1	-	-	-
Male	-	-	1	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Nebraska												
All races, both sexes	12	10	13	18	27	58	94	107	136	68	22	-
Male	8	6	7	10	12	29	44	33	41	18	4	-
Female	4	4	6	8	15	29	50	74	95	50	18	-
White, both sexes	11	9	12	16	26	55	92	105	134	68	20	-
Male	7	6	6	9	11	28	43	32	41	18	4	-
Female	4	3	6	7	15	27	49	73	93	50	16	-
Black, both sexes	1	1	1	1	-	2	2	2	2	-	2	-
Male	1	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	2	-	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	16	24	39	50	57	117	118	67	20	7	-
Male	1	9	10	27	30	34	52	49	21	5	2	-
Female	4	7	14	12	20	23	65	69	46	15	5	-
White, both sexes	5	16	19	33	47	49	110	108	67	19	7	-
Male	1	9	8	24	29	28	51	44	21	5	2	-
Female	4	7	11	9	18	21	59	64	46	14	5	-
Black, both sexes	-	-	4	3	3	4	3	5	-	1	-	-
Male	-	-	2	3	1	3	-	3	-	-	-	-
Female	-	-	2	-	2	1	3	2	-	1	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	4	2	4	-	-	-	-
Male	-	-	-	-	-	3	1	2	-	-	-	-
Female	-	-	-	2	-	1	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
New Hampshire												
All races, both sexes	2	-	3	3	11	25	54	68	68	28	9	-
Male	2	-	2	2	8	12	24	25	19	6	-	-
Female	-	-	1	1	3	13	30	43	49	22	9	-
White, both sexes	2	-	3	3	11	25	54	68	68	28	9	-
Male	2	-	2	2	8	12	24	25	19	6	-	-
Female	-	-	1	1	3	13	30	43	49	22	9	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	10	22	35	39	83	173	266	362	320	151	36	-
Male	6	10	22	21	45	88	121	142	96	30	5	-
Female	4	12	13	18	38	85	145	220	224	121	31	-
White, both sexes	6	14	24	28	67	155	242	340	306	142	35	-
Male	5	7	16	14	34	81	114	137	92	27	4	-
Female	1	7	8	14	33	74	128	203	214	115	31	-
Black, both sexes	4	8	11	10	14	17	24	19	10	9	1	-
Male	1	3	6	7	10	6	7	5	2	3	1	-
Female	3	5	5	3	4	11	17	14	8	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	-	2	4	-	-	-
Male	-	-	-	-	1	1	-	-	2	-	-	-
Female	-	-	-	1	1	-	-	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
New Mexico												
All races, both sexes	1	5	7	23	21	46	77	93	84	45	9	-
Male	-	4	5	12	12	25	32	29	27	10	2	-
Female	1	1	2	11	9	21	45	64	57	35	7	-
White, both sexes	1	5	7	23	20	44	76	91	81	45	8	-
Male	-	4	5	12	11	24	31	29	26	10	1	-
Female	1	1	2	11	9	20	45	62	55	35	7	-
Black, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	17	49	62	86	145	318	475	667	578	293	68	-
Male	12	28	35	49	70	174	192	269	175	70	11	-
Female	5	21	27	37	75	144	283	398	403	223	57	-
White, both sexes	11	37	50	65	125	288	439	630	552	283	62	-
Male	7	23	31	40	62	158	177	260	169	68	11	-
Female	4	14	19	25	63	130	262	370	383	215	51	-
Black, both sexes	5	10	12	19	17	26	31	34	21	8	5	-
Male	4	4	4	7	7	15	13	8	4	1	-	-
Female	1	6	8	12	10	11	18	26	17	7	5	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	2	2	4	4	3	5	2	1	-
Male	1	-	-	2	1	1	2	1	2	1	-	-
Female	-	1	-	-	1	3	2	2	3	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
North Carolina												
All races, both sexes	26	32	56	75	115	208	298	333	286	113	30	-
Male	17	16	29	41	57	96	139	95	84	23	2	-
Female	9	16	27	34	58	112	159	238	202	90	28	-
White, both sexes	12	19	32	54	92	176	248	290	255	95	26	-
Male	9	13	16	26	46	81	113	81	78	20	2	-
Female	3	6	16	28	46	95	135	209	177	75	24	-
Black, both sexes	13	13	22	20	23	30	49	42	29	18	4	-
Male	7	3	12	14	11	13	25	14	4	3	-	-
Female	6	10	10	6	12	17	24	28	25	15	4	-
American Indian 1/, both sexes	1	-	2	1	-	1	-	-	2	-	-	-
Male	1	-	1	1	-	1	-	-	2	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	2	2	1	6	5	10	28	30	22	2	-
Male	-	1	-	1	4	3	5	12	4	1	-	-
Female	-	1	2	-	2	2	5	16	26	21	2	-
White, both sexes	-	1	2	1	4	3	8	27	30	22	2	-
Male	-	-	-	1	3	2	5	12	4	1	-	-
Female	-	1	2	-	1	1	3	15	26	21	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	2	2	2	1	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Ohio												
All races, both sexes	20	34	46	54	129	224	374	455	347	152	32	-
Male	9	21	25	25	63	109	150	180	104	29	5	-
Female	11	13	21	29	66	115	224	275	243	123	27	-
White, both sexes	9	27	36	46	116	202	349	431	334	141	26	-
Male	3	17	21	23	54	101	143	169	100	26	5	-
Female	6	10	15	23	62	101	206	262	234	115	21	-
Black, both sexes	11	7	9	8	13	21	25	23	12	10	6	-
Male	6	4	4	2	9	7	7	10	3	3	-	-
Female	5	3	5	6	4	14	18	13	9	7	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	1	1	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	23	33	52	67	104	157	244	274	236	95	25	-
Male	17	18	31	39	54	84	90	97	73	16	4	-
Female	6	15	21	28	50	73	154	177	163	79	21	-
White, both sexes	18	27	44	55	89	144	229	260	216	90	24	-
Male	14	15	27	31	48	78	84	91	69	15	4	-
Female	4	12	17	24	41	66	145	169	147	75	20	-
Black, both sexes	3	4	2	6	12	11	7	5	7	4	1	-
Male	3	1	2	3	4	4	3	1	1	1	-	-
Female	-	3	-	3	8	7	4	4	6	3	1	-
American Indian 1/, both sexes	2	2	5	6	3	2	8	9	11	1	-	-
Male	-	2	2	5	2	2	3	5	3	-	-	-
Female	2	-	3	1	1	-	5	4	8	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Oregon												
All races, both sexes	14	9	13	14	35	75	121	194	183	100	20	-
Male	5	6	11	9	20	38	52	96	71	25	6	-
Female	9	3	2	5	15	37	69	98	112	75	14	-
White, both sexes	13	9	12	13	35	73	117	187	180	100	19	-
Male	5	6	11	9	20	37	51	93	70	25	5	-
Female	8	3	1	4	15	36	66	94	110	75	14	-
Black, both sexes	-	-	-	-	-	-	3	3	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	1	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	3	1	-	1	-
Male	-	-	-	-	-	-	-	2	-	-	1	-
Female	-	-	-	1	-	1	1	1	1	-	-	-
Pennsylvania												
All races, both sexes	26	53	78	121	222	396	655	921	778	322	72	-
Male	15	30	49	70	114	185	263	333	223	56	10	-
Female	11	23	29	51	108	211	392	588	555	266	62	-
White, both sexes	20	41	64	103	201	357	617	884	746	312	68	-
Male	13	23	40	61	102	171	250	319	211	54	9	-
Female	7	18	24	42	99	186	367	565	535	258	59	-
Black, both sexes	5	12	14	17	21	36	37	35	31	10	4	-
Male	1	7	9	9	12	13	12	13	11	2	1	-
Female	4	5	5	8	9	23	25	22	20	8	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	3	1	2	1	-	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	1	-	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Rhode Island												
All races, both sexes	-	2	2	10	10	20	31	32	38	18	2	-
Male	-	2	2	6	6	8	14	12	7	7	-	-
Female	-	-	-	4	4	12	17	20	31	11	2	-
White, both sexes	-	2	2	9	10	19	30	30	38	18	2	-
Male	-	2	2	5	6	7	14	11	7	7	-	-
Female	-	-	-	4	4	12	16	19	31	11	2	-
Black, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	24	40	45	64	71	112	172	202	156	57	9	-
Male	18	22	29	37	34	58	77	75	53	14	1	-
Female	6	18	16	27	37	54	95	127	103	43	8	-
White, both sexes	13	24	26	41	58	90	132	168	121	47	6	-
Male	8	14	16	26	28	50	61	68	40	13	-	-
Female	5	10	10	15	30	40	71	100	81	34	6	-
Black, both sexes	11	16	19	22	13	22	38	33	35	10	3	-
Male	10	8	13	11	6	8	16	7	13	1	1	-
Female	1	8	6	11	7	14	22	26	22	9	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
South Dakota												
All races, both sexes	-	-	1	1	1	5	9	11	17	9	6	-
Male	-	-	-	1	-	4	6	3	6	1	1	-
Female	-	-	1	-	1	1	3	8	11	8	5	-
White, both sexes	-	-	1	1	1	5	8	11	17	9	6	-
Male	-	-	-	1	-	4	5	3	6	1	1	-
Female	-	-	1	-	1	1	3	8	11	8	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19	16	39	34	53	89	143	129	128	59	8	-
Male	10	11	22	19	22	40	47	53	29	10	-	-
Female	9	5	17	15	31	49	96	76	99	49	8	-
White, both sexes	9	8	37	23	42	77	125	118	113	54	5	-
Male	4	7	21	14	19	34	43	48	24	10	-	-
Female	5	1	16	9	23	43	82	70	89	44	5	-
Black, both sexes	10	8	2	11	11	12	18	11	13	5	3	-
Male	6	4	1	5	3	6	4	5	5	-	-	-
Female	4	4	1	6	8	6	14	6	8	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Texas												
All races, both sexes	81	97	146	191	299	501	780	956	753	391	77	-
Male	51	57	83	114	164	238	344	370	209	85	14	-
Female	30	40	63	77	135	263	436	586	544	306	63	-
White, both sexes	59	72	113	167	259	443	705	880	688	358	66	-
Male	37	47	64	103	141	216	315	341	195	75	11	-
Female	22	25	49	64	118	227	390	539	493	283	55	-
Black, both sexes	21	22	32	22	39	55	71	68	61	32	11	-
Male	13	8	19	9	23	22	29	26	12	9	3	-
Female	8	14	13	13	16	33	42	42	49	23	8	-
American Indian 1/, both sexes	-	2	1	1	-	-	-	2	1	-	-	-
Male	-	1	-	1	-	-	-	2	1	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	3	4	6	3	1	-	-
Male	1	1	-	1	-	-	-	1	1	1	-	-
Female	-	-	-	-	1	3	4	5	2	-	-	-
Utah												
All races, both sexes	5	6	6	18	35	51	126	172	134	52	6	1
Male	2	3	4	8	21	22	61	57	29	15	1	1
Female	3	3	2	10	14	29	65	115	105	37	5	-
White, both sexes	4	5	6	18	34	49	125	170	134	52	5	1
Male	1	2	4	8	20	21	60	57	29	15	1	1
Female	3	3	2	10	14	28	65	113	105	37	4	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	2	-	2	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Vermont												
All races, both sexes	-	-	1	1	1	7	9	18	14	14	2	-
Male	-	-	-	1	-	3	5	5	7	3	1	-
Female	-	-	1	-	1	4	4	13	7	11	1	-
White, both sexes	-	-	1	1	1	7	9	18	14	14	2	-
Male	-	-	-	1	-	3	5	5	7	3	1	-
Female	-	-	1	-	1	4	4	13	7	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	23	52	43	70	125	185	301	359	326	160	41	-
Male	20	39	23	44	69	91	138	120	95	26	3	-
Female	3	13	20	26	56	94	163	239	231	134	38	-
White, both sexes	12	28	30	42	89	156	246	313	288	142	33	-
Male	11	21	19	26	45	75	107	103	88	22	2	-
Female	1	7	11	16	44	81	139	210	200	120	31	-
Black, both sexes	11	23	12	27	34	26	52	44	36	16	7	-
Male	9	17	3	17	23	14	28	17	7	4	1	-
Female	2	6	9	10	11	12	24	27	29	12	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	2	3	3	1	2	1	1	-
Male	-	1	1	-	1	2	3	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	2	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Washington												
All races, both sexes	6	3	9	9	25	44	93	125	138	58	19	-
Male	3	1	4	8	12	19	38	60	53	21	4	-
Female	3	2	5	1	13	25	55	65	85	37	15	-
White, both sexes	4	2	9	8	23	41	89	121	136	58	19	-
Male	2	1	4	7	11	17	36	58	53	21	4	-
Female	2	1	5	1	12	24	53	63	83	37	15	-
Black, both sexes	-	1	-	1	1	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	2	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	3	2	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	2	2	-	-	-
West Virginia												
All races, both sexes	4	7	19	16	45	81	123	145	119	53	16	-
Male	1	3	10	8	15	33	52	55	32	6	2	-
Female	3	4	9	8	30	48	71	90	87	47	14	-
White, both sexes	3	7	18	16	44	78	121	142	119	50	16	-
Male	1	3	9	8	15	32	52	54	32	4	2	-
Female	2	4	9	8	29	46	69	88	87	46	14	-
Black, both sexes	1	-	1	-	1	3	2	3	-	3	-	-
Male	-	-	1	-	-	1	-	1	-	2	-	-
Female	1	-	-	-	1	2	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Wisconsin												
All races, both sexes	12	19	29	35	57	145	260	366	362	182	47	-
Male	9	13	18	22	27	83	112	168	116	49	7	-
Female	3	6	11	13	30	62	148	198	246	133	40	-
White, both sexes	11	13	28	34	52	141	248	361	359	182	47	-
Male	8	10	17	22	26	80	108	166	115	49	7	-
Female	3	3	11	12	26	61	140	195	244	133	40	-
Black, both sexes	1	3	1	1	4	4	8	3	3	-	-	-
Male	1	-	1	-	1	3	3	-	1	-	-	-
Female	-	3	-	1	3	1	5	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	-	-	-	1	2	-	-	-	-
Male	-	3	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	-	2	2	8	4	12	28	37	32	13	6	-
Male	-	1	1	4	2	7	17	18	12	2	3	-
Female	-	1	1	4	2	5	11	19	20	11	3	-
White, both sexes	-	1	2	8	3	12	28	37	32	13	6	-
Male	-	1	1	4	1	7	17	18	12	2	3	-
Female	-	-	1	4	2	5	11	19	20	11	3	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
United States												
All races, both sexes	55,923	5	5	2	3	7	17	20	45	84	172	376
Male	21,908	3	3	-	1	6	10	11	33	48	101	236
Female	34,015	2	2	2	2	1	7	9	12	36	71	140
White, both sexes	50,271	2	5	2	3	2	9	8	18	42	99	220
Male	19,583	2	3	-	1	1	6	6	13	23	55	143
Female	30,688	-	2	2	2	1	3	2	5	19	44	77
Black, both sexes	4,939	3	-	-	-	5	7	9	26	37	67	148
Male	2,010	1	-	-	-	5	4	3	19	20	40	86
Female	2,929	2	-	-	-	-	3	6	7	17	27	62
American Indian 1/, both sexes	198	-	-	-	-	-	1	1	-	3	5	6
Male	92	-	-	-	-	-	-	-	-	3	5	5
Female	106	-	-	-	-	-	1	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	515	-	-	-	-	-	-	2	1	2	1	2
Male	223	-	-	-	-	-	-	2	1	2	1	2
Female	292	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1,809	-	-	-	-	1	2	-	3	4	11	23
Male	707	-	-	-	-	1	2	-	3	2	8	17
Female	1,102	-	-	-	-	-	-	-	-	2	3	6
White, both sexes	1,491	-	-	-	-	-	2	-	2	3	6	14
Male	586	-	-	-	-	-	2	-	2	1	4	11
Female	905	-	-	-	-	-	-	-	-	2	2	3
Black, both sexes	315	-	-	-	-	1	-	-	1	1	5	9
Male	120	-	-	-	-	1	-	-	1	1	4	6
Female	195	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Alaska												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	699	-	-	-	-	1	-	-	-	1	-	2
Male	308	-	-	-	-	-	-	-	-	-	-	2
Female	391	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	677	-	-	-	-	1	-	-	-	1	-	2
Male	297	-	-	-	-	-	-	-	-	-	-	2
Female	380	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Arkansas												
All races, both sexes	663	-	-	-	-	-	-	-	1	-	3	10
Male	276	-	-	-	-	-	-	-	1	-	1	6
Female	387	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	594	-	-	-	-	-	-	-	-	-	1	5
Male	246	-	-	-	-	-	-	-	-	-	-	3
Female	348	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	66	-	-	-	-	-	-	-	1	-	2	5
Male	28	-	-	-	-	-	-	-	1	-	1	3
Female	38	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,679	-	-	-	1	1	1	1	4	6	14	34
Male	1,885	-	-	-	-	1	1	1	3	4	8	22
Female	2,794	-	-	-	1	-	-	-	1	2	6	12
White, both sexes	4,141	-	-	-	1	-	1	-	2	2	10	20
Male	1,658	-	-	-	-	-	1	-	1	1	5	13
Female	2,483	-	-	-	1	-	-	-	1	1	5	7
Black, both sexes	292	-	-	-	-	1	-	-	1	2	3	12
Male	124	-	-	-	-	1	-	-	1	1	2	7
Female	168	-	-	-	-	-	-	-	-	1	1	5
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	1	-	2
Male	11	-	-	-	-	-	-	-	-	1	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	221	-	-	-	-	-	-	1	1	1	1	-
Male	92	-	-	-	-	-	-	1	1	1	1	-
Female	129	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Colorado												
All races, both sexes	664	-	-	-	1	-	-	-	-	1	1	2
Male	251	-	-	-	1	-	-	-	-	-	1	1
Female	413	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	634	-	-	-	1	-	-	-	-	1	1	1
Male	241	-	-	-	1	-	-	-	-	-	1	-
Female	393	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	638	-	-	-	-	-	-	-	-	-	1	4
Male	250	-	-	-	-	-	-	-	-	-	1	1
Female	388	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	612	-	-	-	-	-	-	-	-	-	1	3
Male	243	-	-	-	-	-	-	-	-	-	1	1
Female	369	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Delaware												
All races, both sexes	124	-	-	-	-	-	-	-	-	1	-	2
Male	53	-	-	-	-	-	-	-	-	1	-	2
Female	71	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	111	-	-	-	-	-	-	-	-	-	-	2
Male	49	-	-	-	-	-	-	-	-	-	-	2
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	2
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	37	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Florida												
All races, both sexes	1,750	-	-	-	-	-	-	-	-	1	5	8
Male	727	-	-	-	-	-	-	-	-	-	2	5
Female	1,023	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	1,627	-	-	-	-	-	-	-	-	1	2	2
Male	678	-	-	-	-	-	-	-	-	-	1	2
Female	949	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	117	-	-	-	-	-	-	-	-	-	3	6
Male	48	-	-	-	-	-	-	-	-	-	1	3
Female	69	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,969	-	1	-	-	-	2	4	3	12	17	25
Male	800	-	1	-	-	-	2	2	2	6	9	17
Female	1,169	-	-	-	-	-	-	2	1	6	8	8
White, both sexes	1,507	-	1	-	-	-	-	-	-	3	11	10
Male	615	-	1	-	-	-	-	-	-	1	8	9
Female	892	-	-	-	-	-	-	-	-	2	3	1
Black, both sexes	445	-	-	-	-	-	2	4	3	9	6	15
Male	182	-	-	-	-	-	2	2	2	5	1	8
Female	263	-	-	-	-	-	-	2	1	4	5	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Hawaii												
All races, both sexes	121	-	-	-	-	-	-	1	-	1	1	2
Male	67	-	-	-	-	-	-	1	-	1	1	1
Female	54	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	39	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	82	-	-	-	-	-	-	1	-	1	-	1
Male	49	-	-	-	-	-	-	1	-	1	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	259	-	-	-	-	-	-	-	-	-	1	-
Male	98	-	-	-	-	-	-	-	-	-	-	-
Female	161	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	256	-	-	-	-	-	-	-	-	-	1	-
Male	98	-	-	-	-	-	-	-	-	-	-	-
Female	158	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Illinois												
All races, both sexes	2,907	-	-	-	-	-	1	-	-	6	4	12
Male	1,123	-	-	-	-	-	1	-	-	4	3	10
Female	1,784	-	-	-	-	-	-	-	-	2	1	2
White, both sexes	2,624	-	-	-	-	-	1	-	-	5	2	6
Male	1,001	-	-	-	-	-	1	-	-	3	1	6
Female	1,623	-	-	-	-	-	-	-	-	2	1	-
Black, both sexes	270	-	-	-	-	-	-	-	-	1	2	6
Male	116	-	-	-	-	-	-	-	-	1	2	4
Female	154	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,823	-	-	-	-	1	1	-	1	2	6	19
Male	693	-	-	-	-	1	-	-	1	1	2	12
Female	1,130	-	-	-	-	-	1	-	-	1	4	7
White, both sexes	1,704	-	-	-	-	-	-	-	-	1	5	13
Male	641	-	-	-	-	-	-	-	-	1	1	9
Female	1,063	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	119	-	-	-	-	1	1	-	1	1	1	6
Male	52	-	-	-	-	1	-	-	1	-	1	3
Female	67	-	-	-	-	-	1	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Iowa												
All races, both sexes	251	-	-	-	-	-	-	-	-	-	1	-
Male	88	-	-	-	-	-	-	-	-	-	1	-
Female	163	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	245	-	-	-	-	-	-	-	-	-	1	-
Male	87	-	-	-	-	-	-	-	-	-	1	-
Female	158	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	946	-	-	-	-	-	-	-	-	-	3	3
Male	350	-	-	-	-	-	-	-	-	-	3	3
Female	596	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	901	-	-	-	-	-	-	-	-	-	1	2
Male	335	-	-	-	-	-	-	-	-	-	1	2
Female	566	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	38	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Kentucky												
All races, both sexes	1,306	-	-	-	-	1	-	1	-	3	6	11
Male	502	-	-	-	-	1	-	1	-	-	4	8
Female	804	-	-	-	-	-	-	-	-	3	2	3
White, both sexes	1,240	-	-	-	-	-	-	1	-	1	6	9
Male	473	-	-	-	-	-	-	1	-	-	4	7
Female	767	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	64	-	-	-	-	1	-	-	-	2	-	2
Male	29	-	-	-	-	1	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,233	-	-	-	1	-	1	-	3	3	8	13
Male	505	-	-	-	-	-	-	-	1	2	6	9
Female	728	-	-	-	1	-	1	-	2	1	2	4
White, both sexes	939	-	-	-	1	-	1	-	1	1	2	4
Male	381	-	-	-	-	-	-	-	-	1	2	3
Female	558	-	-	-	1	-	1	-	1	-	-	1
Black, both sexes	291	-	-	-	-	-	-	-	2	2	6	9
Male	123	-	-	-	-	-	-	-	1	1	4	6
Female	168	-	-	-	-	-	-	-	1	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Maine												
All races, both sexes	288	-	-	-	-	-	-	-	-	2	-	2
Male	116	-	-	-	-	-	-	-	-	1	-	1
Female	172	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	286	-	-	-	-	-	-	-	-	1	-	2
Male	115	-	-	-	-	-	-	-	-	-	-	1
Female	171	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	655	-	-	-	-	-	-	1	2	1	2	5
Male	247	-	-	-	-	-	-	1	1	-	2	4
Female	408	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	534	-	-	-	-	-	-	1	-	-	1	-
Male	198	-	-	-	-	-	-	1	-	-	1	-
Female	336	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	109	-	-	-	-	-	-	-	2	1	1	5
Male	45	-	-	-	-	-	-	-	1	-	1	4
Female	64	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Massachusetts												
All races, both sexes	1,664	-	-	-	-	-	-	-	-	-	3	4
Male	611	-	-	-	-	-	-	-	-	-	2	4
Female	1,053	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1,615	-	-	-	-	-	-	-	-	-	3	4
Male	592	-	-	-	-	-	-	-	-	-	2	4
Female	1,023	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,435	1	-	-	-	-	-	1	6	-	7	11
Male	930	-	-	-	-	-	-	-	5	-	4	7
Female	1,505	1	-	-	-	-	-	1	1	-	3	4
White, both sexes	2,208	-	-	-	-	-	-	-	1	-	4	5
Male	831	-	-	-	-	-	-	-	1	-	1	2
Female	1,377	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	212	1	-	-	-	-	-	1	5	-	3	5
Male	89	-	-	-	-	-	-	-	4	-	3	4
Female	123	1	-	-	-	-	-	1	1	-	-	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Minnesota												
All races, both sexes	1,077	-	-	1	-	-	1	-	1	-	1	2
Male	440	-	-	-	-	-	-	-	1	-	-	2
Female	637	-	-	1	-	-	1	-	-	-	1	-
White, both sexes	1,049	-	-	1	-	-	1	-	1	-	-	1
Male	424	-	-	-	-	-	-	-	1	-	-	1
Female	625	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,312	-	-	-	-	-	2	1	2	4	3	17
Male	485	-	-	-	-	-	1	-	1	3	2	6
Female	827	-	-	-	-	-	1	1	1	1	1	11
White, both sexes	959	-	-	-	-	-	-	-	1	2	1	6
Male	361	-	-	-	-	-	-	-	-	2	1	2
Female	598	-	-	-	-	-	-	-	1	-	-	4
Black, both sexes	349	-	-	-	-	-	2	1	1	2	2	11
Male	121	-	-	-	-	-	1	-	1	1	1	4
Female	228	-	-	-	-	-	1	1	-	1	1	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Missouri												
All races, both sexes	1,424	-	-	-	-	-	-	-	1	2	1	9
Male	524	-	-	-	-	-	-	-	1	2	1	5
Female	900	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	1,340	-	-	-	-	-	-	-	1	2	-	8
Male	490	-	-	-	-	-	-	-	1	2	-	5
Female	850	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	83	-	-	-	-	-	-	-	-	-	1	1
Male	34	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	316	-	-	-	-	-	-	-	1	-	-	1
Male	134	-	-	-	-	-	-	-	1	-	-	-
Female	182	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	310	-	-	-	-	-	-	-	1	-	-	-
Male	130	-	-	-	-	-	-	-	1	-	-	-
Female	180	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Nebraska												
All races, both sexes	498	-	-	-	-	-	1	-	-	-	-	2
Male	168	-	-	-	-	-	1	-	-	-	-	1
Female	330	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	480	-	-	-	-	-	1	-	-	-	-	1
Male	160	-	-	-	-	-	1	-	-	-	-	-
Female	320	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	496	-	-	-	-	-	-	-	1	1	1	4
Male	228	-	-	-	-	-	-	-	-	1	-	3
Female	268	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	457	-	-	-	-	-	-	-	1	1	-	3
Male	209	-	-	-	-	-	-	-	-	1	-	2
Female	248	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
New Hampshire												
All races, both sexes	253	-	-	-	-	-	-	-	-	-	-	-
Male	93	-	-	-	-	-	-	-	-	-	-	-
Female	160	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	253	-	-	-	-	-	-	-	-	-	-	-
Male	93	-	-	-	-	-	-	-	-	-	-	-
Female	160	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,380	-	-	-	-	-	-	-	-	3	6	5
Male	539	-	-	-	-	-	-	-	-	3	3	3
Female	841	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	1,247	-	-	-	-	-	-	-	-	3	1	3
Male	485	-	-	-	-	-	-	-	-	3	1	1
Female	762	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	122	-	-	-	-	-	-	-	-	-	5	2
Male	50	-	-	-	-	-	-	-	-	-	2	2
Female	72	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
New Mexico												
All races, both sexes	394	-	-	-	-	-	-	-	-	-	1	5
Male	149	-	-	-	-	-	-	-	-	-	1	3
Female	245	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	383	-	-	-	-	-	-	-	-	-	-	5
Male	143	-	-	-	-	-	-	-	-	-	-	3
Female	240	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,555	1	-	-	-	-	-	-	1	4	4	8
Male	995	-	-	-	-	-	-	-	1	2	-	6
Female	1,560	1	-	-	-	-	-	-	-	2	4	2
White, both sexes	2,350	-	-	-	-	-	-	-	-	2	4	4
Male	921	-	-	-	-	-	-	-	-	2	-	4
Female	1,429	-	-	-	-	-	-	-	-	-	4	-
Black, both sexes	180	1	-	-	-	-	-	-	1	2	-	4
Male	65	-	-	-	-	-	-	-	1	-	-	2
Female	115	1	-	-	-	-	-	-	-	2	-	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
North Carolina												
All races, both sexes	1,454	-	-	-	-	-	1	2	1	2	4	7
Male	553	-	-	-	-	-	-	1	1	2	3	6
Female	901	-	-	-	-	-	1	1	-	-	1	1
White, both sexes	1,204	-	-	-	-	-	-	1	-	1	-	2
Male	444	-	-	-	-	-	-	1	-	1	-	2
Female	760	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	243	-	-	-	-	-	1	1	1	1	4	5
Male	103	-	-	-	-	-	-	-	1	1	3	4
Female	140	-	-	-	-	-	1	1	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	97	-	-	-	-	-	-	-	-	-	1	-
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	89	-	-	-	-	-	-	-	-	-	1	-
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	65	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Ohio												
All races, both sexes	1,772	1	1	-	-	-	-	1	1	4	3	10
Male	694	1	1	-	-	-	-	1	1	3	1	6
Female	1,078	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	1,626	1	1	-	-	-	-	1	1	2	2	5
Male	633	1	1	-	-	-	-	1	1	1	1	3
Female	993	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	143	-	-	-	-	-	-	-	-	2	1	5
Male	59	-	-	-	-	-	-	-	-	2	-	3
Female	84	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,248	-	-	-	-	-	-	2	1	3	6	8
Male	506	-	-	-	-	-	-	1	-	3	5	5
Female	742	-	-	-	-	-	-	1	1	-	1	3
White, both sexes	1,138	-	-	-	-	-	-	1	-	1	4	6
Male	460	-	-	-	-	-	-	1	-	1	3	4
Female	678	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	57	-	-	-	-	-	-	-	1	1	-	1
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	38	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	50	-	-	-	-	-	-	1	-	1	2	1
Male	27	-	-	-	-	-	-	-	-	1	2	1
Female	23	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Oregon												
All races, both sexes	745	-	-	-	-	-	-	-	-	1	1	1
Male	326	-	-	-	-	-	-	-	-	-	1	1
Female	419	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	724	-	-	-	-	-	-	-	-	1	1	-
Male	318	-	-	-	-	-	-	-	-	-	1	-
Female	406	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,411	1	-	-	-	-	-	1	2	2	5	16
Male	1,247	1	-	-	-	-	-	-	1	1	4	9
Female	2,164	-	-	-	-	-	-	1	1	1	1	7
White, both sexes	3,199	1	-	-	-	-	-	1	-	1	3	11
Male	1,162	1	-	-	-	-	-	-	-	-	2	6
Female	2,037	-	-	-	-	-	-	1	-	1	1	5
Black, both sexes	205	-	-	-	-	-	-	-	2	1	2	5
Male	82	-	-	-	-	-	-	-	1	1	2	3
Female	123	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Rhode Island												
All races, both sexes	158	-	-	-	-	-	-	-	-	-	-	1
Male	62	-	-	-	-	-	-	-	-	-	-	-
Female	96	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	153	-	-	-	-	-	-	-	-	-	-	1
Male	59	-	-	-	-	-	-	-	-	-	-	-
Female	94	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	869	-	-	-	-	-	-	-	3	2	6	8
Male	374	-	-	-	-	-	-	-	1	2	3	4
Female	495	-	-	-	-	-	-	-	2	-	3	4
White, both sexes	668	-	-	-	-	-	-	-	1	-	2	6
Male	293	-	-	-	-	-	-	-	-	-	-	4
Female	375	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	197	-	-	-	-	-	-	-	2	2	4	2
Male	81	-	-	-	-	-	-	-	1	2	3	-
Female	116	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
South Dakota												
All races, both sexes	57	-	-	-	-	-	1	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	36	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	1	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	686	-	-	-	-	-	-	-	2	4	5	10
Male	252	-	-	-	-	-	-	-	2	1	3	5
Female	434	-	-	-	-	-	-	-	-	3	2	5
White, both sexes	575	-	-	-	-	-	-	-	2	2	2	6
Male	212	-	-	-	-	-	-	-	2	1	1	3
Female	363	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	109	-	-	-	-	-	-	-	-	2	3	4
Male	40	-	-	-	-	-	-	-	-	-	2	2
Female	69	-	-	-	-	-	-	-	-	2	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Texas												
All races, both sexes	3,988	-	3	-	-	2	3	2	5	4	19	42
Male	1,599	-	1	-	-	2	2	1	5	1	7	26
Female	2,389	-	2	-	-	-	1	1	-	3	12	16
White, both sexes	3,551	-	3	-	-	1	2	-	3	2	13	29
Male	1,425	-	1	-	-	1	1	-	3	-	4	18
Female	2,126	-	2	-	-	-	1	-	-	2	9	11
Black, both sexes	415	-	-	-	-	1	1	2	2	2	6	13
Male	167	-	-	-	-	1	1	1	2	1	3	8
Female	248	-	-	-	-	-	-	1	-	1	3	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	579	-	-	-	-	-	-	1	-	-	1	2
Male	216	-	-	-	-	-	-	-	-	-	1	1
Female	363	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	572	-	-	-	-	-	-	1	-	-	1	2
Male	212	-	-	-	-	-	-	-	-	-	1	1
Female	360	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Vermont												
All races, both sexes	67	-	-	1	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	67	-	-	1	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,453	1	-	-	-	-	-	1	-	1	5	11
Male	560	1	-	-	-	-	-	1	-	1	3	7
Female	893	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	1,207	-	-	-	-	-	-	1	-	1	1	5
Male	442	-	-	-	-	-	-	1	-	1	1	3
Female	765	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	231	1	-	-	-	-	-	-	-	-	4	6
Male	110	1	-	-	-	-	-	-	-	-	2	4
Female	121	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Washington												
All races, both sexes	494	-	-	-	-	-	-	-	-	2	-	1
Male	204	-	-	-	-	-	-	-	-	1	-	-
Female	290	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	476	-	-	-	-	-	-	-	-	-	-	1
Male	196	-	-	-	-	-	-	-	-	-	-	-
Female	280	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	2	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	516	-	-	-	-	-	-	-	-	1	1	5
Male	182	-	-	-	-	-	-	-	-	-	-	4
Female	334	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	503	-	-	-	-	-	-	-	-	1	1	4
Male	176	-	-	-	-	-	-	-	-	-	-	3
Female	327	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Wisconsin												
All races, both sexes	1,496	-	-	-	-	-	-	-	-	-	2	6
Male	615	-	-	-	-	-	-	-	-	-	2	1
Female	881	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	1,460	-	-	-	-	-	-	-	-	-	2	5
Male	601	-	-	-	-	-	-	-	-	-	2	1
Female	859	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	139	-	-	-	-	-	-	-	-	-	-	1
Male	64	-	-	-	-	-	-	-	-	-	-	-
Female	75	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	137	-	-	-	-	-	-	-	-	-	-	1
Male	63	-	-	-	-	-	-	-	-	-	-	-
Female	74	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
United States												
All races, both sexes	639	1,029	1,412	2,034	3,339	6,025	10,021	12,786	11,487	5,188	1,224	3
Male	396	619	806	1,143	1,721	2,940	4,287	4,768	3,444	1,128	202	2
Female	243	410	606	891	1,618	3,085	5,734	8,018	8,043	4,060	1,022	1
White, both sexes	404	744	1,121	1,639	2,832	5,338	9,126	11,915	10,753	4,875	1,111	3
Male	251	467	651	944	1,463	2,657	3,944	4,483	3,234	1,054	180	2
Female	153	277	470	695	1,369	2,681	5,182	7,432	7,519	3,821	931	1
Black, both sexes	216	264	272	362	453	589	769	741	601	273	97	-
Male	133	138	148	183	226	244	291	233	162	59	15	-
Female	83	126	124	179	227	345	478	508	439	214	82	-
American Indian 1/, both sexes	7	7	12	20	19	23	35	26	23	7	3	-
Male	3	6	4	13	10	9	12	12	8	-	2	-
Female	4	1	8	7	9	14	23	14	15	7	1	-
Asian or Pacific Islander 2/, both sexes	12	14	7	13	35	75	91	104	110	33	13	-
Male	9	8	3	3	22	30	40	40	40	15	5	-
Female	3	6	4	10	13	45	51	64	70	18	8	-
Alabama												
All races, both sexes	38	57	61	101	135	222	321	357	324	115	34	-
Male	23	35	37	61	66	91	126	113	95	25	2	-
Female	15	22	24	40	69	131	195	244	229	90	32	-
White, both sexes	19	42	44	77	105	185	272	311	282	103	24	-
Male	12	29	30	49	54	75	108	101	85	22	1	-
Female	7	13	14	28	51	110	164	210	197	81	23	-
Black, both sexes	19	15	17	24	30	36	49	45	41	12	10	-
Male	11	6	7	12	12	16	18	11	10	3	1	-
Female	8	9	10	12	18	20	31	34	31	9	9	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Alaska												
All races, both sexes	2	1	2	-	6	6	9	13	11	5	1	-
Male	2	1	-	-	3	2	4	5	4	-	-	-
Female	-	-	2	-	3	4	5	8	7	5	1	-
White, both sexes	1	1	1	-	4	3	8	10	9	4	1	-
Male	1	1	-	-	1	1	4	4	4	-	-	-
Female	-	-	1	-	3	2	4	6	5	4	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	1	3	1	2	2	1	-	-
Male	1	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	1	-	-	2	1	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	9	9	20	31	47	78	137	170	134	48	12	-
Male	4	6	14	18	23	42	63	71	48	13	4	-
Female	5	3	6	13	24	36	74	99	86	35	8	-
White, both sexes	8	9	19	27	45	75	133	166	133	47	11	-
Male	3	6	13	16	21	42	60	70	48	13	3	-
Female	5	3	6	11	24	33	73	96	85	34	8	-
Black, both sexes	1	-	-	1	-	-	2	3	-	1	-	-
Male	1	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	1	3	2	3	2	1	1	-	1	-
Male	-	-	1	1	2	-	1	-	-	-	1	-
Female	-	-	-	2	-	3	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Arkansas												
All races, both sexes	9	21	17	22	43	85	113	161	115	52	11	-
Male	7	14	10	13	24	46	53	53	36	11	1	-
Female	2	7	7	9	19	39	60	108	79	41	10	-
White, both sexes	5	15	13	18	38	83	107	153	104	43	9	-
Male	4	9	8	11	21	45	52	52	32	8	1	-
Female	1	6	5	7	17	38	55	101	72	35	8	-
Black, both sexes	4	5	4	3	5	2	5	8	11	9	2	-
Male	3	4	2	2	3	1	-	1	4	3	-	-
Female	1	1	2	1	2	1	5	7	7	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
California												
All races, both sexes	71	116	106	165	277	431	871	1,054	990	431	104	1
Male	51	68	55	87	146	194	394	402	337	93	18	-
Female	20	48	51	78	131	237	477	652	653	338	86	1
White, both sexes	49	91	89	129	236	374	779	959	902	401	95	1
Male	36	55	46	72	129	174	354	371	304	80	16	-
Female	13	36	43	57	107	200	425	588	598	321	79	1
Black, both sexes	18	19	14	31	28	30	47	42	33	10	1	-
Male	12	10	8	13	11	12	17	12	14	3	-	-
Female	6	9	6	18	17	18	30	30	19	7	1	-
American Indian 1/, both sexes	-	-	1	1	1	6	6	5	1	-	1	-
Male	-	-	-	1	-	1	3	2	1	-	-	-
Female	-	-	1	-	1	5	3	3	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	6	2	4	12	21	39	48	54	20	7	-
Male	3	3	1	1	6	7	20	17	18	10	2	-
Female	1	3	1	3	6	14	19	31	36	10	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Colorado												
All races, both sexes	4	7	12	21	28	60	106	169	150	82	20	-
Male	3	5	7	11	11	30	46	61	46	23	5	-
Female	1	2	5	10	17	30	60	108	104	59	15	-
White, both sexes	4	6	12	19	26	57	101	161	145	79	20	-
Male	3	4	7	10	10	29	44	59	45	23	5	-
Female	1	2	5	9	16	28	57	102	100	56	15	-
Black, both sexes	-	1	-	2	2	-	3	7	2	1	-	-
Male	-	1	-	1	1	-	1	2	1	-	-	-
Female	-	-	-	1	1	-	2	5	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	2	1	2	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	2	1	1	2	-	-	-
Connecticut												
All races, both sexes	-	3	7	21	34	75	111	156	141	69	16	-
Male	-	2	4	8	19	47	46	58	51	11	2	-
Female	-	1	3	13	15	28	65	98	90	58	14	-
White, both sexes	-	3	7	18	29	75	106	152	136	66	16	-
Male	-	2	4	8	16	47	44	57	51	10	2	-
Female	-	1	3	10	13	28	62	95	85	56	14	-
Black, both sexes	-	-	-	3	5	-	5	4	4	3	-	-
Male	-	-	-	-	3	-	2	1	-	1	-	-
Female	-	-	-	3	2	-	3	3	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Delaware												
All races, both sexes	-	4	8	5	12	13	14	32	22	10	1	-
Male	-	3	7	2	5	5	8	9	9	1	1	-
Female	-	1	1	3	7	8	6	23	13	9	-	-
White, both sexes	-	3	8	3	11	11	14	29	21	8	1	-
Male	-	3	7	1	5	5	8	8	8	1	1	-
Female	-	-	1	2	6	6	6	21	13	7	-	-
Black, both sexes	-	1	-	2	1	2	-	3	1	1	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	1	-	1	1	2	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	1	1	5	3	3	9	9	9	3	3	-
Male	1	1	1	3	1	1	4	2	3	-	1	-
Female	1	-	-	2	2	2	5	7	6	3	2	-
White, both sexes	-	1	-	-	-	1	2	4	3	1	1	-
Male	-	1	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	3	3	1	1	-
Black, both sexes	2	-	1	5	3	2	7	5	6	2	2	-
Male	1	-	1	3	1	-	3	1	3	-	1	-
Female	1	-	-	2	2	2	4	4	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Florida												
All races, both sexes	18	17	41	57	99	166	304	409	405	180	40	-
Male	14	10	24	33	55	78	145	156	144	51	10	-
Female	4	7	17	24	44	88	159	253	261	129	30	-
White, both sexes	13	10	33	53	90	150	291	392	379	172	39	-
Male	10	8	18	30	48	71	140	155	134	51	10	-
Female	3	2	15	23	42	79	151	237	245	121	29	-
Black, both sexes	5	6	8	4	9	15	13	14	25	8	1	-
Male	4	2	6	3	7	7	5	-	10	-	-	-
Female	1	4	2	1	2	8	8	14	15	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	3	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	2	1	-	-	-
Georgia												
All races, both sexes	53	67	81	108	150	219	351	416	314	118	28	-
Male	26	52	52	55	75	102	151	131	92	21	4	-
Female	27	15	29	53	75	117	200	285	222	97	24	-
White, both sexes	27	41	54	75	100	171	291	347	256	99	21	-
Male	13	33	36	39	49	82	139	113	71	17	4	-
Female	14	8	18	36	51	89	152	234	185	82	17	-
Black, both sexes	25	26	26	33	47	45	55	67	57	18	7	-
Male	13	19	16	16	23	20	12	18	21	4	-	-
Female	12	7	10	17	24	25	43	49	36	14	7	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	2	3	5	2	1	1	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	1	-	1	-	-	3	5	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Hawaii												
All races, both sexes	5	1	2	4	7	19	16	22	30	8	2	-
Male	4	1	1	4	6	9	8	11	14	3	2	-
Female	1	-	1	-	1	10	8	11	16	5	-	-
White, both sexes	-	-	-	2	1	7	6	10	7	4	-	-
Male	-	-	-	2	1	4	2	4	3	1	-	-
Female	-	-	-	-	-	3	4	6	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	2	2	6	12	10	12	23	4	2	-
Male	4	1	1	2	5	5	6	7	11	2	2	-
Female	1	-	1	-	1	7	4	5	12	2	-	-
Idaho												
All races, both sexes	1	-	3	6	14	22	43	66	66	26	11	-
Male	-	-	3	5	7	9	20	24	21	6	3	-
Female	1	-	-	1	7	13	23	42	45	20	8	-
White, both sexes	1	-	3	6	14	22	42	65	66	25	11	-
Male	-	-	3	5	7	9	20	24	21	6	3	-
Female	1	-	-	1	7	13	22	41	45	19	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Illinois												
All races, both sexes	28	55	82	96	159	330	530	617	631	281	75	-
Male	19	28	45	49	88	163	228	240	168	62	15	-
Female	9	27	37	47	71	167	302	377	463	219	60	-
White, both sexes	15	37	64	80	125	297	470	574	607	269	72	-
Male	8	20	35	45	72	144	204	224	164	60	14	-
Female	7	17	29	35	53	153	266	350	443	209	58	-
Black, both sexes	13	17	18	16	33	32	55	38	24	12	3	-
Male	11	8	10	4	15	19	22	13	4	2	1	-
Female	2	9	8	12	18	13	33	25	20	10	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	5	4	-	-	-	-
Male	-	-	-	-	1	-	2	2	-	-	-	-
Female	-	1	-	-	-	1	3	2	-	-	-	-
Indiana												
All races, both sexes	16	32	42	77	108	184	321	407	406	160	40	-
Male	6	19	30	42	56	105	118	146	118	33	3	-
Female	10	13	12	35	52	79	203	261	288	127	37	-
White, both sexes	13	25	34	67	98	171	305	391	388	153	40	-
Male	5	14	23	37	52	100	111	141	114	30	3	-
Female	8	11	11	30	46	71	194	250	274	123	37	-
Black, both sexes	3	7	8	10	10	13	16	16	18	7	-	-
Male	1	5	7	5	4	5	7	5	4	3	-	-
Female	2	2	1	5	6	8	9	11	14	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Iowa												
All races, both sexes	-	3	5	5	13	28	39	47	55	45	10	-
Male	-	-	2	4	4	13	17	16	18	12	1	-
Female	-	3	3	1	9	15	22	31	37	33	9	-
White, both sexes	-	2	5	5	12	26	38	46	55	45	10	-
Male	-	-	2	4	3	13	17	16	18	12	1	-
Female	-	2	3	1	9	13	21	30	37	33	9	-
Black, both sexes	-	1	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Kansas												
All races, both sexes	2	11	13	23	52	76	144	217	244	124	34	-
Male	2	5	6	17	23	33	63	88	64	39	4	-
Female	-	6	7	6	29	43	81	129	180	85	30	-
White, both sexes	1	11	10	18	49	70	139	208	238	121	33	-
Male	1	5	6	14	22	33	61	85	64	37	4	-
Female	-	6	4	4	27	37	78	123	174	84	29	-
Black, both sexes	1	-	3	3	1	6	4	9	5	3	1	-
Male	1	-	-	2	1	-	1	3	-	2	-	-
Female	-	-	3	1	-	6	3	6	5	1	1	-
American Indian 1/, both sexes	-	-	-	2	1	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Kentucky												
All races, both sexes	16	39	42	74	101	173	223	284	221	92	19	-
Male	14	25	21	46	50	76	94	88	50	20	4	-
Female	2	14	21	28	51	97	129	196	171	72	15	-
White, both sexes	15	35	40	68	96	162	213	270	217	89	18	-
Male	13	21	21	40	47	72	89	86	49	20	3	-
Female	2	14	19	28	49	90	124	184	168	69	15	-
Black, both sexes	1	4	2	6	5	10	10	13	4	3	1	-
Male	1	4	-	6	3	4	5	2	1	-	1	-
Female	-	-	2	-	2	6	5	11	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Louisiana												
All races, both sexes	22	37	49	62	86	163	219	253	210	87	16	-
Male	11	19	23	35	43	81	92	90	76	13	4	-
Female	11	18	26	27	43	82	127	163	134	74	12	-
White, both sexes	8	25	29	37	59	119	177	209	181	70	15	-
Male	4	14	15	23	31	62	72	76	64	10	4	-
Female	4	11	14	14	28	57	105	133	117	60	11	-
Black, both sexes	14	12	20	25	25	44	42	44	29	16	1	-
Male	7	5	8	12	11	19	20	14	12	3	-	-
Female	7	7	12	13	14	25	22	30	17	13	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Maine												
All races, both sexes	-	2	4	5	14	26	62	78	57	29	7	-
Male	-	1	3	2	10	18	24	34	16	4	2	-
Female	-	1	1	3	4	8	38	44	41	25	5	-
White, both sexes	-	2	4	5	14	26	61	78	57	29	7	-
Male	-	1	3	2	10	18	24	34	16	4	2	-
Female	-	1	1	3	4	8	37	44	41	25	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	12	18	20	41	62	122	157	129	61	17	1
Male	2	5	9	9	23	26	53	61	36	11	3	1
Female	2	7	9	11	18	36	69	96	93	50	14	-
White, both sexes	1	6	13	14	30	48	93	137	118	56	15	1
Male	-	2	7	5	17	18	47	54	32	10	3	1
Female	1	4	6	9	13	30	46	83	86	46	12	-
Black, both sexes	3	6	5	5	11	13	22	19	10	4	2	-
Male	2	3	2	4	6	7	5	7	3	-	-	-
Female	1	3	3	1	5	6	17	12	7	4	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	7	1	1	1	-	-
Male	-	-	-	-	-	1	1	-	1	1	-	-
Female	-	-	-	-	-	-	6	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Massachusetts												
All races, both sexes	9	10	26	27	68	164	311	413	388	192	49	-
Male	7	8	13	14	36	103	132	145	108	33	6	-
Female	2	2	13	13	32	61	179	268	280	159	43	-
White, both sexes	9	10	26	25	63	150	302	405	381	189	48	-
Male	7	8	13	13	34	96	128	141	107	33	6	-
Female	2	2	13	12	29	54	174	264	274	156	42	-
Black, both sexes	-	-	-	2	4	11	7	8	6	2	1	-
Male	-	-	-	1	1	6	4	4	1	-	-	-
Female	-	-	-	1	3	5	3	4	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	2	-	1	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	2	-	1	1	-	-
Michigan												
All races, both sexes	22	34	48	78	134	259	438	587	529	242	38	-
Male	17	24	24	40	62	130	195	214	148	55	5	-
Female	5	10	24	38	72	129	243	373	381	187	33	-
White, both sexes	15	20	39	65	112	227	402	558	500	227	33	-
Male	12	16	19	34	53	117	177	202	141	52	4	-
Female	3	4	20	31	59	110	225	356	359	175	29	-
Black, both sexes	7	13	9	12	19	30	34	27	27	15	4	-
Male	5	7	5	5	8	11	17	11	5	3	1	-
Female	2	6	4	7	11	19	17	16	22	12	3	-
American Indian 1/, both sexes	-	1	-	1	2	1	1	1	-	-	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	1	2	-	1	-
Male	-	-	-	-	1	1	1	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Minnesota												
All races, both sexes	6	3	11	12	38	99	178	272	277	134	41	-
Male	5	3	7	10	28	51	93	117	80	35	8	-
Female	1	-	4	2	10	48	85	155	197	99	33	-
White, both sexes	3	3	11	11	32	94	176	267	274	134	40	-
Male	3	3	7	9	25	47	92	114	79	35	8	-
Female	-	-	4	2	7	47	84	153	195	99	32	-
Black, both sexes	2	-	-	1	4	2	-	3	1	-	1	-
Male	2	-	-	1	2	2	-	1	-	-	-	-
Female	-	-	-	-	2	-	-	2	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	2	2	-	-	-
Male	-	-	-	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Mississippi												
All races, both sexes	19	45	33	74	77	165	219	269	239	113	30	-
Male	9	28	18	46	41	79	83	86	58	17	7	-
Female	10	17	15	28	36	86	136	183	181	96	23	-
White, both sexes	6	19	18	53	59	123	168	213	188	84	18	-
Male	3	13	10	34	34	63	68	66	48	13	4	-
Female	3	6	8	19	25	60	100	147	140	71	14	-
Black, both sexes	13	25	15	21	18	41	50	55	51	29	12	-
Male	6	14	8	12	7	15	14	20	10	4	3	-
Female	7	11	7	9	11	26	36	35	41	25	9	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Missouri												
All races, both sexes	14	19	26	50	91	146	264	328	302	130	41	-
Male	8	11	13	22	57	63	106	112	83	32	8	-
Female	6	8	13	28	34	83	158	216	219	98	33	-
White, both sexes	12	15	25	46	77	132	251	317	291	125	38	-
Male	7	9	12	21	45	60	104	107	79	30	8	-
Female	5	6	13	25	32	72	147	210	212	95	30	-
Black, both sexes	2	4	1	4	14	14	12	11	11	5	3	-
Male	1	2	1	1	12	3	2	5	4	2	-	-
Female	1	2	-	3	2	11	10	6	7	3	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	6	10	6	12	25	49	86	82	37	1	-
Male	-	3	9	5	4	13	23	40	27	9	-	-
Female	-	3	1	1	8	12	26	46	55	28	1	-
White, both sexes	-	6	9	6	10	24	48	86	82	37	1	-
Male	-	3	8	5	3	12	22	40	27	9	-	-
Female	-	3	1	1	7	12	26	46	55	28	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	1	1	-	-	-	-	-
Male	-	-	1	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Nebraska												
All races, both sexes	8	6	9	13	20	49	88	100	127	57	18	-
Male	5	4	4	6	7	22	39	28	37	12	2	-
Female	3	2	5	7	13	27	49	72	90	45	16	-
White, both sexes	7	5	8	11	19	46	86	98	125	57	16	-
Male	4	4	3	5	6	21	38	27	37	12	2	-
Female	3	1	5	6	13	25	48	71	88	45	14	-
Black, both sexes	1	1	1	1	-	2	2	2	2	-	2	-
Male	1	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	2	-	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	14	24	36	48	52	111	109	64	19	7	-
Male	1	7	10	27	28	29	49	46	20	5	2	-
Female	4	7	14	9	20	23	62	63	44	14	5	-
White, both sexes	5	14	19	30	45	44	106	100	64	18	7	-
Male	1	7	8	24	27	23	48	41	20	5	2	-
Female	4	7	11	6	18	21	58	59	44	13	5	-
Black, both sexes	-	-	4	3	3	4	1	5	-	1	-	-
Male	-	-	2	3	1	3	-	3	-	-	-	-
Female	-	-	2	-	2	1	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	4	2	3	-	-	-	-
Male	-	-	-	-	-	3	1	2	-	-	-	-
Female	-	-	-	2	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
New Hampshire												
All races, both sexes	2	-	3	3	10	24	50	62	66	26	7	-
Male	2	-	2	2	8	12	21	22	18	6	-	-
Female	-	-	1	1	2	12	29	40	48	20	7	-
White, both sexes	2	-	3	3	10	24	50	62	66	26	7	-
Male	2	-	2	2	8	12	21	22	18	6	-	-
Female	-	-	1	1	2	12	29	40	48	20	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	20	34	35	72	150	242	338	295	137	34	-
Male	5	10	22	17	39	73	110	134	89	27	4	-
Female	4	10	12	18	33	77	132	204	206	110	30	-
White, both sexes	5	13	23	25	58	134	220	318	283	128	33	-
Male	4	7	16	11	29	68	103	130	85	24	3	-
Female	1	6	7	14	29	66	117	188	198	104	30	-
Black, both sexes	4	7	11	9	12	15	22	17	8	9	1	-
Male	1	3	6	6	9	4	7	4	2	3	1	-
Female	3	4	5	3	3	11	15	13	6	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	-	2	4	-	-	-
Male	-	-	-	-	1	1	-	-	2	-	-	-
Female	-	-	-	1	1	-	-	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
New Mexico												
All races, both sexes	1	5	5	23	17	43	73	89	79	44	9	-
Male	-	4	4	12	8	23	29	28	25	10	2	-
Female	1	1	1	11	9	20	44	61	54	34	7	-
White, both sexes	1	5	5	23	16	41	72	87	76	44	8	-
Male	-	4	4	12	7	22	28	28	24	10	1	-
Female	1	1	1	11	9	19	44	59	52	34	7	-
Black, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	16	41	57	76	128	289	439	611	538	280	62	-
Male	12	24	30	43	60	157	171	250	162	68	9	-
Female	4	17	27	33	68	132	268	361	376	212	53	-
White, both sexes	10	31	45	60	108	261	405	577	515	271	57	-
Male	7	20	26	37	52	143	157	241	157	66	9	-
Female	3	11	19	23	56	118	248	336	358	205	48	-
Black, both sexes	5	8	12	16	17	24	29	31	19	7	4	-
Male	4	3	4	6	7	13	12	8	4	1	-	-
Female	1	5	8	10	10	11	17	23	15	6	4	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	2	4	4	3	4	2	1	-
Male	1	-	-	-	1	1	2	1	1	1	-	-
Female	-	1	-	-	1	3	2	2	3	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
North Carolina												
All races, both sexes	20	24	48	64	100	193	277	311	270	102	28	-
Male	11	12	26	34	47	87	130	90	79	22	2	-
Female	9	12	22	30	53	106	147	221	191	80	26	-
White, both sexes	8	14	27	48	81	164	231	272	244	86	25	-
Male	5	10	14	22	38	73	106	76	75	19	2	-
Female	3	4	13	26	43	91	125	196	169	67	23	-
Black, both sexes	11	10	21	15	19	28	45	38	24	16	3	-
Male	5	2	12	11	9	13	23	14	2	3	-	-
Female	6	8	9	4	10	15	22	24	22	13	3	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	2	-	-	-
Male	1	-	-	1	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	2	2	-	4	5	6	26	27	22	2	-
Male	-	1	-	-	2	3	3	12	4	1	-	-
Female	-	1	2	-	2	2	3	14	23	21	2	-
White, both sexes	-	1	2	-	2	3	4	25	27	22	2	-
Male	-	-	-	-	1	2	3	12	4	1	-	-
Female	-	1	2	-	1	1	1	13	23	21	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	2	2	2	1	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Ohio												
All races, both sexes	17	31	43	43	116	211	355	428	333	145	29	-
Male	7	19	25	21	56	104	144	170	102	28	4	-
Female	10	12	18	22	60	107	211	258	231	117	25	-
White, both sexes	8	25	35	35	105	189	331	407	321	134	23	-
Male	2	15	21	19	47	96	137	160	98	25	4	-
Female	6	10	14	16	58	93	194	247	223	109	19	-
Black, both sexes	9	6	8	8	11	21	24	21	11	10	6	-
Male	5	4	4	2	9	7	7	10	3	3	-	-
Female	4	2	4	6	2	14	17	11	8	7	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	21	31	51	65	100	147	230	256	220	84	23	-
Male	17	16	30	37	53	77	84	90	69	15	4	-
Female	4	15	21	28	47	70	146	166	151	69	19	-
White, both sexes	17	25	43	53	87	136	217	244	202	80	22	-
Male	14	13	26	29	48	73	79	85	66	14	4	-
Female	3	12	17	24	39	63	138	159	136	66	18	-
Black, both sexes	3	4	2	6	10	9	6	4	6	3	1	-
Male	3	1	2	3	3	2	2	1	-	1	-	-
Female	-	3	-	3	7	7	4	3	6	2	1	-
American Indian 1/, both sexes	1	2	5	6	3	2	7	8	10	1	-	-
Male	-	2	2	5	2	2	3	4	3	-	-	-
Female	1	-	3	1	1	-	4	4	7	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Oregon												
All races, both sexes	12	9	12	11	30	74	116	185	175	99	19	-
Male	4	6	10	7	19	37	50	93	68	24	6	-
Female	8	3	2	4	11	37	66	92	107	75	13	-
White, both sexes	11	9	11	10	30	72	112	178	172	99	18	-
Male	4	6	10	7	19	36	49	90	67	24	5	-
Female	7	3	1	3	11	36	63	88	105	75	13	-
Black, both sexes	-	-	-	-	-	-	3	3	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	1	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	3	1	-	1	-
Male	-	-	-	-	-	-	-	2	-	-	1	-
Female	-	-	-	1	-	1	1	1	1	-	-	-
Pennsylvania												
All races, both sexes	20	42	65	105	197	356	606	871	751	307	64	-
Male	13	23	40	59	102	160	245	312	215	52	10	-
Female	7	19	25	46	95	196	361	559	536	255	54	-
White, both sexes	15	32	54	91	179	322	573	837	722	297	60	-
Male	12	17	33	52	92	151	233	300	204	50	9	-
Female	3	15	21	39	87	171	340	537	518	247	51	-
Black, both sexes	5	10	11	13	18	31	33	32	28	10	4	-
Male	1	6	7	7	10	8	12	11	10	2	1	-
Female	4	4	4	6	8	23	21	21	18	8	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	3	-	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	1	-	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Rhode Island												
All races, both sexes	-	1	2	10	10	19	29	31	35	18	2	-
Male	-	1	2	6	6	8	13	12	7	7	-	-
Female	-	-	-	4	4	11	16	19	28	11	2	-
White, both sexes	-	1	2	9	10	18	28	29	35	18	2	-
Male	-	1	2	5	6	7	13	11	7	7	-	-
Female	-	-	-	4	4	11	15	18	28	11	2	-
Black, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	17	31	35	56	63	103	151	187	144	54	9	-
Male	11	18	22	33	29	54	66	70	47	13	1	-
Female	6	13	13	23	34	49	85	117	97	41	8	-
White, both sexes	8	23	20	36	54	82	117	156	112	45	6	-
Male	3	13	12	23	26	46	54	63	37	12	-	-
Female	5	10	8	13	28	36	63	93	75	33	6	-
Black, both sexes	9	8	15	19	9	21	32	30	32	9	3	-
Male	8	5	10	10	3	8	12	7	10	1	1	-
Female	1	3	5	9	6	13	20	23	22	8	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
South Dakota												
All races, both sexes	-	-	1	1	-	4	8	10	15	9	6	-
Male	-	-	-	1	-	3	5	2	6	1	1	-
Female	-	-	1	-	-	1	3	8	9	8	5	-
White, both sexes	-	-	1	1	-	4	7	10	15	9	6	-
Male	-	-	-	1	-	3	4	2	6	1	1	-
Female	-	-	1	-	-	1	3	8	9	8	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	15	13	30	32	50	82	137	121	121	56	8	-
Male	7	9	17	18	20	38	42	51	29	10	-	-
Female	8	4	13	14	30	44	95	70	92	46	8	-
White, both sexes	6	6	29	21	39	71	119	110	106	51	5	-
Male	2	5	17	13	17	33	38	46	24	10	-	-
Female	4	1	12	8	22	38	81	64	82	41	5	-
Black, both sexes	9	7	1	11	11	11	18	11	13	5	3	-
Male	5	4	-	5	3	5	4	5	5	-	-	-
Female	4	3	1	6	8	6	14	6	8	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Texas												
All races, both sexes	71	79	126	171	272	448	733	878	700	361	69	-
Male	44	45	71	100	146	208	316	338	196	78	12	-
Female	27	34	55	71	126	240	417	540	504	283	57	-
White, both sexes	52	59	101	148	236	399	665	811	638	331	58	-
Male	32	38	56	90	126	190	291	314	182	69	9	-
Female	20	21	45	58	110	209	374	497	456	262	49	-
Black, both sexes	18	19	24	22	35	46	64	62	58	29	11	-
Male	11	7	15	9	20	18	25	22	12	8	3	-
Female	7	12	9	13	15	28	39	40	46	21	8	-
American Indian 1/, both sexes	-	1	1	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	3	4	4	3	1	-	-
Male	1	-	-	-	-	-	-	1	1	1	-	-
Female	-	-	-	-	1	3	4	3	2	-	-	-
Utah												
All races, both sexes	3	6	5	15	33	47	120	166	126	48	5	1
Male	1	3	3	7	20	21	59	55	28	15	1	1
Female	2	3	2	8	13	26	61	111	98	33	4	-
White, both sexes	3	5	5	15	32	46	119	164	126	48	4	1
Male	1	2	3	7	19	20	58	55	28	15	1	1
Female	2	3	2	8	13	26	61	109	98	33	3	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	2	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Vermont												
All races, both sexes	-	-	1	1	1	7	9	18	13	14	2	-
Male	-	-	-	1	-	3	5	5	7	3	1	-
Female	-	-	1	-	1	4	4	13	6	11	1	-
White, both sexes	-	-	1	1	1	7	9	18	13	14	2	-
Male	-	-	-	1	-	3	5	5	7	3	1	-
Female	-	-	1	-	1	4	4	13	6	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	13	37	32	57	100	162	250	314	286	145	38	-
Male	10	27	17	33	54	80	114	104	82	23	3	-
Female	3	10	15	24	46	82	136	210	204	122	35	-
White, both sexes	5	18	22	32	75	139	210	279	255	133	31	-
Male	4	13	14	18	37	68	90	92	77	21	2	-
Female	1	5	8	14	38	71	120	187	178	112	29	-
Black, both sexes	8	18	9	24	24	20	38	33	30	10	6	-
Male	6	13	2	14	16	10	22	12	5	2	1	-
Female	2	5	7	10	8	10	16	21	25	8	5	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	3	2	1	1	1	1	-
Male	-	1	1	-	1	2	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Washington												
All races, both sexes	5	2	9	6	23	39	89	117	128	54	19	-
Male	2	-	4	5	11	15	38	56	49	19	4	-
Female	3	2	5	1	12	24	51	61	79	35	15	-
White, both sexes	3	1	9	5	21	38	85	114	126	54	19	-
Male	1	-	4	4	10	15	36	54	49	19	4	-
Female	2	1	5	1	11	23	49	60	77	35	15	-
Black, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	2	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	2	2	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	2	-	-	-
West Virginia												
All races, both sexes	3	5	18	13	36	68	95	114	104	41	12	-
Male	1	1	10	7	12	28	39	44	28	6	2	-
Female	2	4	8	6	24	40	56	70	76	35	10	-
White, both sexes	3	5	17	13	35	65	94	111	104	38	12	-
Male	1	1	9	7	12	27	39	43	28	4	2	-
Female	2	4	8	6	23	38	55	68	76	34	10	-
Black, both sexes	-	-	1	-	1	3	1	3	-	3	-	-
Male	-	-	1	-	-	1	-	1	-	2	-	-
Female	-	-	-	-	1	2	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Wisconsin												
All races, both sexes	11	16	28	35	56	143	256	358	358	180	47	-
Male	8	11	18	22	26	82	109	165	116	48	7	-
Female	3	5	10	13	30	61	147	193	242	132	40	-
White, both sexes	10	13	27	34	51	139	244	353	355	180	47	-
Male	7	10	17	22	25	79	105	163	115	48	7	-
Female	3	3	10	12	26	60	139	190	240	132	40	-
Black, both sexes	1	2	1	1	4	4	8	3	3	-	-	-
Male	1	-	1	-	1	3	3	-	1	-	-	-
Female	-	2	-	1	3	1	5	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	-	2	2	8	4	11	27	37	31	12	4	-
Male	-	1	1	4	2	6	17	18	11	2	2	-
Female	-	1	1	4	2	5	10	19	20	10	2	-
White, both sexes	-	1	2	8	3	11	27	37	31	12	4	-
Male	-	1	1	4	1	6	17	18	11	2	2	-
Female	-	-	1	4	2	5	10	19	20	10	2	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
United States												
All races, both sexes	4,414	14	4	2	5	6	7	6	15	23	53	87
Male	2,010	8	2	1	3	2	4	5	7	14	36	57
Female	2,404	6	2	1	2	4	3	1	8	9	17	30
White, both sexes	3,709	6	3	1	2	5	2	4	8	13	38	53
Male	1,653	2	1	-	2	2	1	3	4	6	25	37
Female	2,056	4	2	1	-	3	1	1	4	7	13	16
Black, both sexes	625	8	1	1	3	1	4	2	7	9	15	32
Male	325	6	1	1	1	-	3	2	3	7	11	19
Female	300	2	-	-	2	1	1	-	4	2	4	13
American Indian 1/, both sexes	20	-	-	-	-	-	1	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	60	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	237	1	-	-	-	1	-	-	2	3	9	18
Male	115	-	-	-	-	1	-	-	2	3	5	12
Female	122	1	-	-	-	-	-	-	-	-	4	6
White, both sexes	128	-	-	-	-	1	-	-	1	1	6	8
Male	54	-	-	-	-	1	-	-	1	1	4	6
Female	74	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	107	1	-	-	-	-	-	-	1	1	3	10
Male	60	-	-	-	-	-	-	-	1	1	1	6
Female	47	1	-	-	-	-	-	-	-	-	2	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	46	-	1	-	-	-	-	1	-	-	-	-
Male	22	-	-	-	-	-	-	1	-	-	-	-
Female	24	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	1	-	-	-	-	1	-	-	-	-
Male	20	-	-	-	-	-	-	1	-	-	-	-
Female	22	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Arkansas												
All races, both sexes	47	-	-	-	1	1	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	1	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	5	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	102	-	-	-	-	-	-	-	-	1	-	-
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	92	-	-	-	-	-	-	-	-	1	-	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Colorado												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	1	-
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	1	-
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	98	-	-	-	-	-	-	-	-	-	-	-
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	60	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	93	-	-	-	-	-	-	-	-	-	-	-
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Florida												
All races, both sexes	150	1	-	-	-	-	-	1	1	-	1	2
Male	66	1	-	-	-	-	-	1	-	-	-	1
Female	84	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	132	1	-	-	-	-	-	-	-	-	1	-
Male	55	1	-	-	-	-	-	-	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	17	-	-	-	-	-	-	1	1	-	-	2
Male	11	-	-	-	-	-	-	1	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	199	1	-	-	-	-	-	-	1	-	2	4
Male	87	1	-	-	-	-	-	-	-	-	2	2
Female	112	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	145	-	-	-	-	-	-	-	-	-	1	2
Male	63	-	-	-	-	-	-	-	-	-	1	1
Female	82	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	53	1	-	-	-	-	-	-	1	-	1	2
Male	24	1	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Hawaii												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Illinois												
All races, both sexes	196	3	-	1	-	-	-	1	-	2	-	4
Male	84	1	-	-	-	-	-	1	-	1	-	2
Female	112	2	-	1	-	-	-	-	-	1	-	2
White, both sexes	170	1	-	1	-	-	-	-	-	-	-	2
Male	73	-	-	-	-	-	-	-	-	-	-	2
Female	97	1	-	1	-	-	-	-	-	-	-	-
Black, both sexes	23	2	-	-	-	-	-	1	-	2	-	2
Male	9	1	-	-	-	-	-	1	-	1	-	-
Female	14	1	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	118	-	-	-	-	1	-	-	-	-	4	2
Male	64	-	-	-	-	1	-	-	-	-	3	2
Female	54	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	112	-	-	-	-	1	-	-	-	-	3	1
Male	60	-	-	-	-	1	-	-	-	-	2	1
Female	52	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Iowa												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Kentucky												
All races, both sexes	75	-	-	-	-	-	-	-	1	1	2	2
Male	31	-	-	-	-	-	-	-	1	1	2	2
Female	44	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	1	1	2	2
Male	31	-	-	-	-	-	-	-	1	1	2	2
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	58	-	1	-	-	-	-	-	-	1	1	2
Male	32	-	1	-	-	-	-	-	-	1	1	1
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	13	-	1	-	-	-	-	-	-	1	1	-
Male	8	-	1	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Maine												
All races, both sexes	24	-	-	-	-	1	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	1	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	24	-	-	-	-	-	-	-	1	1	-	2
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	1	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Massachusetts												
All races, both sexes	134	-	1	-	-	-	-	1	1	-	-	1
Male	54	-	1	-	-	-	-	1	1	-	-	1
Female	80	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	130	-	1	-	-	-	-	1	1	-	-	-
Male	53	-	1	-	-	-	-	1	1	-	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	153	-	-	-	-	-	1	-	-	1	1	1
Male	75	-	-	-	-	-	1	-	-	1	1	1
Female	78	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	130	-	-	-	-	-	-	-	-	1	-	-
Male	60	-	-	-	-	-	-	-	-	1	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	21	-	-	-	-	-	1	-	-	-	1	1
Male	13	-	-	-	-	-	1	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Minnesota												
All races, both sexes	100	-	-	-	-	-	-	-	-	-	1	1
Male	48	-	-	-	-	-	-	-	-	-	1	-
Female	52	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	96	-	-	-	-	-	-	-	-	-	1	1
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	50	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	126	-	-	-	-	-	1	-	-	-	3	1
Male	61	-	-	-	-	-	-	-	-	-	-	1
Female	65	-	-	-	-	-	1	-	-	-	3	-
White, both sexes	86	-	-	-	-	-	1	-	-	-	2	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	1	-	-	-	2	-
Black, both sexes	39	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Missouri												
All races, both sexes	74	-	-	-	-	-	-	-	-	1	1	3
Male	20	-	-	-	-	-	-	-	-	1	-	2
Female	54	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	66	-	-	-	-	-	-	-	-	1	1	3
Male	18	-	-	-	-	-	-	-	-	1	-	2
Female	48	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Nebraska												
All races, both sexes	76	-	-	-	-	-	-	-	-	2	-	4
Male	49	-	-	-	-	-	-	-	-	-	-	3
Female	27	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	76	-	-	-	-	-	-	-	-	2	-	4
Male	49	-	-	-	-	-	-	-	-	-	-	3
Female	27	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	34	1	-	-	-	-	-	-	1	-	-	1
Male	17	-	-	-	-	-	-	-	1	-	-	-
Female	17	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	30	1	-	-	-	-	-	-	1	-	-	-
Male	17	-	-	-	-	-	-	-	1	-	-	-
Female	13	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
New Hampshire												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	143	3	-	-	-	-	1	-	-	-	3	5
Male	64	2	-	-	-	-	-	-	-	-	3	3
Female	79	1	-	-	-	-	1	-	-	-	-	2
White, both sexes	127	1	-	-	-	-	-	-	-	-	3	4
Male	56	-	-	-	-	-	-	-	-	-	3	2
Female	71	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	16	2	-	-	-	-	1	-	-	-	-	1
Male	8	2	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
New Mexico												
All races, both sexes	24	-	-	-	-	-	-	-	1	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	1	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	229	-	-	-	-	-	-	1	1	1	2	3
Male	103	-	-	-	-	-	-	-	-	1	1	2
Female	126	-	-	-	-	-	-	1	1	-	1	1
White, both sexes	209	-	-	-	-	-	-	1	1	1	2	2
Male	95	-	-	-	-	-	-	-	-	1	1	2
Female	114	-	-	-	-	-	-	1	1	-	1	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
North Carolina												
All races, both sexes	147	2	1	-	-	-	-	-	1	-	5	3
Male	67	2	-	-	-	-	-	-	1	-	3	2
Female	80	-	1	-	-	-	-	-	-	-	2	1
White, both sexes	106	1	1	-	-	-	-	-	-	-	3	2
Male	50	1	-	-	-	-	-	-	-	-	2	2
Female	56	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	38	1	-	-	-	-	-	-	1	-	2	1
Male	15	1	-	-	-	-	-	-	1	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Ohio												
All races, both sexes	119	-	-	-	-	-	-	-	-	-	3	-
Male	43	-	-	-	-	-	-	-	-	-	3	-
Female	76	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	107	-	-	-	-	-	-	-	-	-	3	-
Male	41	-	-	-	-	-	-	-	-	-	3	-
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	82	-	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Oregon												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	270	-	-	1	-	2	1	-	-	2	1	3
Male	124	-	-	1	-	-	1	-	-	1	1	3
Female	146	-	-	-	-	2	-	-	-	1	-	-
White, both sexes	235	-	-	-	-	1	-	-	-	1	1	1
Male	102	-	-	-	-	-	-	-	-	-	1	1
Female	133	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	33	-	-	1	-	1	1	-	-	1	-	2
Male	20	-	-	1	-	-	1	-	-	1	-	2
Female	13	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Rhode Island												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	110	1	-	-	-	-	1	-	-	-	1	5
Male	61	-	-	-	-	-	1	-	-	-	1	5
Female	49	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	1	-	-	-	-	1	-	-	-	1	2
Male	39	-	-	-	-	-	1	-	-	-	1	2
Female	33	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	38	-	-	-	-	-	-	-	-	-	-	3
Male	22	-	-	-	-	-	-	-	-	-	-	3
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	56	-	-	-	1	-	-	-	-	-	1	2
Male	25	-	-	-	1	-	-	-	-	-	1	1
Female	31	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	51	-	-	-	-	-	-	-	-	-	1	2
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	1	-	-	-	-	-	-	-
Male	4	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Texas												
All races, both sexes	383	1	-	-	1	-	-	1	3	1	6	6
Male	185	1	-	-	1	-	-	1	-	1	3	3
Female	198	-	-	-	-	-	-	-	3	-	3	3
White, both sexes	326	-	-	-	1	-	-	1	2	1	4	5
Male	154	-	-	-	1	-	-	1	-	1	1	2
Female	172	-	-	-	-	-	-	-	2	-	3	3
Black, both sexes	51	1	-	-	-	-	-	-	1	-	2	1
Male	27	1	-	-	-	-	-	-	-	-	2	1
Female	24	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	38	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Vermont												
All races, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	264	-	-	-	1	-	1	-	1	3	3	4
Male	130	-	-	-	-	-	1	-	1	2	3	2
Female	134	-	-	-	1	-	-	-	-	1	-	2
White, both sexes	181	-	-	-	-	-	-	-	-	-	-	1
Male	84	-	-	-	-	-	-	-	-	-	-	1
Female	97	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	80	-	-	-	1	-	1	-	1	3	3	3
Male	45	-	-	-	-	-	1	-	1	2	3	1
Female	35	-	-	-	1	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Washington												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	122	-	-	-	-	-	-	-	-	2	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	120	-	-	-	-	-	-	-	-	2	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	81	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Wisconsin												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
United States												
All races, both sexes	123	173	187	241	346	522	741	804	652	303	100	-
Male	91	105	118	143	195	283	365	299	186	69	17	-
Female	32	68	69	98	151	239	376	505	466	234	83	-
White, both sexes	80	120	138	185	280	450	652	720	593	267	89	-
Male	59	79	87	114	153	241	322	272	170	57	16	-
Female	21	41	51	71	127	209	330	448	423	210	73	-
Black, both sexes	38	47	44	51	64	65	76	65	50	32	10	-
Male	29	22	29	26	41	38	36	24	15	10	1	-
Female	9	25	15	25	23	27	40	41	35	22	9	-
American Indian 1/, both sexes	1	1	2	-	1	3	4	3	2	-	-	-
Male	-	1	1	-	1	2	1	2	-	-	-	-
Female	1	-	1	-	-	1	3	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	4	5	3	5	1	4	9	16	7	4	1	-
Male	3	3	1	3	-	2	6	1	1	2	-	-
Female	1	2	2	2	1	2	3	15	6	2	1	-
Alabama												
All races, both sexes	18	18	18	19	27	19	30	23	21	8	2	-
Male	13	10	10	6	15	10	14	5	6	3	-	-
Female	5	8	8	13	12	9	16	18	15	5	2	-
White, both sexes	8	13	9	9	11	14	16	13	13	4	1	-
Male	4	7	3	3	4	6	8	2	3	1	-	-
Female	4	6	6	6	7	8	8	11	10	3	1	-
Black, both sexes	10	5	9	10	16	5	14	10	7	4	1	-
Male	9	3	7	3	11	4	6	3	3	2	-	-
Female	1	2	2	7	5	1	8	7	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Alaska												
All races, both sexes	-	-	-	-	-	1	1	1	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	3	2	2	3	10	6	6	9	2	-	-
Male	-	2	2	1	2	6	4	3	1	-	-	-
Female	1	1	-	1	1	4	2	3	8	2	-	-
White, both sexes	1	3	2	2	3	8	5	5	9	2	-	-
Male	-	2	2	1	2	4	4	3	1	-	-	-
Female	1	1	-	1	1	4	1	2	8	2	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Arkansas												
All races, both sexes	-	1	3	6	3	7	7	8	5	4	-	-
Male	-	-	3	5	1	3	3	3	1	1	-	-
Female	-	1	-	1	2	4	4	5	4	3	-	-
White, both sexes	-	1	2	4	3	7	7	8	4	4	-	-
Male	-	-	2	3	1	3	3	3	1	1	-	-
Female	-	1	-	1	2	4	4	5	3	3	-	-
Black, both sexes	-	-	1	2	-	-	-	-	1	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	5	7	2	13	13	16	22	12	8	2	-
Male	1	1	5	1	8	7	7	5	2	3	1	-
Female	-	4	2	1	5	6	9	17	10	5	1	-
White, both sexes	-	4	6	2	12	13	14	19	11	8	2	-
Male	-	1	4	1	7	7	7	4	2	3	1	-
Female	-	3	2	1	5	6	7	15	9	5	1	-
Black, both sexes	1	-	1	-	1	-	1	1	-	-	-	-
Male	1	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Colorado												
All races, both sexes	-	-	3	4	4	8	10	9	6	3	-	-
Male	-	-	3	1	3	5	7	5	1	1	-	-
Female	-	-	-	3	1	3	3	4	5	2	-	-
White, both sexes	-	-	3	4	4	8	10	9	6	3	-	-
Male	-	-	3	1	3	5	7	5	1	1	-	-
Female	-	-	-	3	1	3	3	4	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	2	2	5	2	10	25	19	18	7	6	-
Male	2	1	1	3	1	5	13	5	6	-	1	-
Female	-	1	1	2	1	5	12	14	12	7	5	-
White, both sexes	1	2	2	4	2	9	25	19	18	6	5	-
Male	1	1	1	3	1	4	13	5	6	-	-	-
Female	-	1	1	1	1	5	12	14	12	6	5	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Delaware												
All races, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	1	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Florida												
All races, both sexes	2	7	7	6	8	17	27	31	28	9	2	-
Male	1	3	5	3	6	10	13	11	8	3	-	-
Female	1	4	2	3	2	7	14	20	20	6	2	-
White, both sexes	1	6	3	5	7	16	24	31	26	9	2	-
Male	1	3	2	3	5	9	10	11	7	3	-	-
Female	-	3	1	2	2	7	14	20	19	6	2	-
Black, both sexes	1	1	4	1	1	1	3	-	1	-	-	-
Male	-	-	3	-	1	1	3	-	1	-	-	-
Female	1	1	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Georgia												
All races, both sexes	7	9	7	16	14	30	33	25	30	14	6	-
Male	4	5	5	8	8	11	17	6	13	4	1	-
Female	3	4	2	8	6	19	16	19	17	10	5	-
White, both sexes	3	5	4	7	9	22	30	18	27	12	5	-
Male	2	4	2	4	5	7	16	4	12	4	1	-
Female	1	1	2	3	4	15	14	14	15	8	4	-
Black, both sexes	4	4	3	9	5	8	3	6	3	2	1	-
Male	2	1	3	4	3	4	1	2	1	-	-	-
Female	2	3	-	5	2	4	2	4	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Hawaii												
All races, both sexes	-	-	2	3	-	1	4	7	1	5	1	-
Male	-	-	1	1	-	-	3	-	-	2	-	-
Female	-	-	1	2	-	1	1	7	1	3	1	-
White, both sexes	-	-	-	1	-	-	1	2	-	2	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	-	1	3	5	1	3	1	-
Male	-	-	1	-	-	-	2	-	-	2	-	-
Female	-	-	1	2	-	1	1	5	1	1	1	-
Idaho												
All races, both sexes	-	-	-	1	-	2	1	2	3	1	-	-
Male	-	-	-	1	-	-	1	1	1	1	-	-
Female	-	-	-	-	-	2	-	1	2	-	-	-
White, both sexes	-	-	-	1	-	2	1	2	3	1	-	-
Male	-	-	-	1	-	-	1	1	1	1	-	-
Female	-	-	-	-	-	2	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Illinois												
All races, both sexes	8	6	7	11	9	20	44	35	28	11	6	-
Male	7	4	3	3	6	9	21	14	7	4	1	-
Female	1	2	4	8	3	11	23	21	21	7	5	-
White, both sexes	8	2	6	10	8	16	41	33	27	9	6	-
Male	7	2	3	3	5	8	18	14	7	3	1	-
Female	1	-	3	7	3	8	23	19	20	6	5	-
Black, both sexes	-	4	1	1	1	4	1	2	-	2	-	-
Male	-	2	-	-	1	1	1	-	-	1	-	-
Female	-	2	1	1	-	3	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	1	2	5	12	16	11	24	18	10	10	2	-
Male	-	2	4	9	6	7	15	9	4	2	-	-
Female	1	-	1	3	10	4	9	9	6	8	2	-
White, both sexes	1	2	5	11	16	11	23	18	10	8	2	-
Male	-	2	4	8	6	7	15	9	4	1	-	-
Female	1	-	1	3	10	4	8	9	6	7	2	-
Black, both sexes	-	-	-	1	-	-	1	-	-	2	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Iowa												
All races, both sexes	-	-	1	-	2	2	7	7	4	6	5	-
Male	-	-	1	-	-	2	3	4	2	3	-	-
Female	-	-	-	-	2	-	4	3	2	3	5	-
White, both sexes	-	-	1	-	2	2	7	6	4	6	4	-
Male	-	-	1	-	-	2	3	4	2	3	-	-
Female	-	-	-	-	2	-	4	2	2	3	4	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	2	2	1	4	5	12	7	8	4	1	-
Male	2	2	2	1	2	2	7	6	3	-	-	-
Female	-	-	-	-	2	3	5	1	5	4	1	-
White, both sexes	1	2	2	1	3	5	11	5	8	3	1	-
Male	1	2	2	1	2	2	7	4	3	-	-	-
Female	-	-	-	-	1	3	4	1	5	3	1	-
Black, both sexes	1	-	-	-	1	-	-	2	-	1	-	-
Male	1	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Kentucky												
All races, both sexes	1	2	6	2	4	12	12	11	13	5	1	-
Male	1	-	2	1	1	6	6	3	4	1	-	-
Female	-	2	4	1	3	6	6	8	9	4	1	-
White, both sexes	1	1	6	2	4	12	12	10	13	5	1	-
Male	1	-	2	1	1	6	6	3	4	1	-	-
Female	-	1	4	1	3	6	6	7	9	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Louisiana												
All races, both sexes	1	5	4	3	4	6	6	9	10	4	1	-
Male	-	5	3	1	3	4	4	3	5	-	-	-
Female	1	-	1	2	1	2	2	6	5	4	1	-
White, both sexes	-	4	3	3	3	5	5	8	9	3	-	-
Male	-	4	2	1	3	3	3	2	5	-	-	-
Female	-	-	1	2	-	2	2	6	4	3	-	-
Black, both sexes	1	1	1	-	1	1	1	1	1	1	1	-
Male	-	1	1	-	-	1	1	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Maine												
All races, both sexes	1	2	3	2	1	1	6	2	3	2	-	-
Male	1	1	3	2	-	1	4	-	1	-	-	-
Female	-	1	-	-	1	-	2	2	2	2	-	-
White, both sexes	1	2	3	2	1	1	6	2	3	2	-	-
Male	1	1	3	2	-	1	4	-	1	-	-	-
Female	-	1	-	-	1	-	2	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	1	3	3	5	3	3	1	-	-
Male	-	-	-	1	3	1	2	2	-	-	-	-
Female	-	-	1	-	-	2	3	1	3	1	-	-
White, both sexes	-	-	1	1	3	3	5	3	3	1	-	-
Male	-	-	-	1	3	1	2	2	-	-	-	-
Female	-	-	1	-	-	2	3	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Massachusetts												
All races, both sexes	3	1	1	4	11	16	28	28	26	9	3	-
Male	2	1	1	3	4	5	13	13	5	2	1	-
Female	1	-	-	1	7	11	15	15	21	7	2	-
White, both sexes	2	1	1	4	11	16	28	28	24	9	3	-
Male	2	1	1	3	4	5	13	13	5	2	1	-
Female	-	-	-	1	7	11	15	15	19	7	2	-
Black, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	7	5	10	10	24	27	30	20	7	8	-
Male	1	4	4	5	7	16	12	16	3	2	1	-
Female	-	3	1	5	3	8	15	14	17	5	7	-
White, both sexes	-	5	4	9	9	19	24	27	18	6	8	-
Male	-	2	4	5	6	14	9	14	3	1	1	-
Female	-	3	-	4	3	5	15	13	15	5	7	-
Black, both sexes	1	2	1	1	1	4	2	3	2	1	-	-
Male	1	2	-	-	1	1	2	2	-	1	-	-
Female	-	-	1	1	-	3	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Minnesota												
All races, both sexes	1	1	3	-	7	10	16	25	24	9	2	-
Male	1	-	2	-	5	7	6	13	10	2	1	-
Female	-	1	1	-	2	3	10	12	14	7	1	-
White, both sexes	1	1	3	-	5	10	16	24	23	9	2	-
Male	1	-	2	-	3	7	6	13	10	2	1	-
Female	-	1	1	-	2	3	10	11	13	7	1	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Mississippi												
All races, both sexes	8	5	3	9	11	18	15	23	19	8	2	-
Male	8	3	3	9	6	9	6	6	8	2	-	-
Female	-	2	-	-	5	9	9	17	11	6	2	-
White, both sexes	4	4	1	4	8	11	14	19	14	3	-	-
Male	4	2	1	4	4	7	5	5	7	-	-	-
Female	-	2	-	-	4	4	9	14	7	3	-	-
Black, both sexes	4	1	2	5	3	7	1	3	5	5	2	-
Male	4	1	2	5	2	2	1	1	1	2	-	-
Female	-	-	-	-	1	5	-	2	4	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Missouri												
All races, both sexes	3	2	2	3	5	6	11	13	20	3	1	-
Male	1	2	-	2	1	2	3	3	3	-	-	-
Female	2	-	2	1	4	4	8	10	17	3	1	-
White, both sexes	2	1	2	3	4	4	9	12	20	3	1	-
Male	1	1	-	2	1	1	3	3	3	-	-	-
Female	1	-	2	1	3	3	6	9	17	3	1	-
Black, both sexes	1	1	-	-	1	2	2	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	2	-	-	1	2	5	6	2	-	-
Male	1	-	1	-	-	1	-	2	-	-	-	-
Female	-	-	1	-	-	-	2	3	6	2	-	-
White, both sexes	1	-	2	-	-	1	1	5	5	2	-	-
Male	1	-	1	-	-	1	-	2	-	-	-	-
Female	-	-	1	-	-	-	1	3	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Nebraska												
All races, both sexes	4	4	4	5	7	9	6	7	9	11	4	-
Male	3	2	3	4	5	7	5	5	4	6	2	-
Female	1	2	1	1	2	2	1	2	5	5	2	-
White, both sexes	4	4	4	5	7	9	6	7	9	11	4	-
Male	3	2	3	4	5	7	5	5	4	6	2	-
Female	1	2	1	1	2	2	1	2	5	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	2	-	3	2	5	6	9	3	1	-	-
Male	-	2	-	-	2	5	3	3	1	-	-	-
Female	-	-	-	3	-	-	3	6	2	1	-	-
White, both sexes	-	2	-	3	2	5	4	8	3	1	-	-
Male	-	2	-	-	2	5	3	3	1	-	-	-
Female	-	-	-	3	-	-	1	5	2	1	-	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
New Hampshire												
All races, both sexes	-	-	-	-	1	1	4	6	2	2	2	-
Male	-	-	-	-	-	-	3	3	1	-	-	-
Female	-	-	-	-	1	1	1	3	1	2	2	-
White, both sexes	-	-	-	-	1	1	4	6	2	2	2	-
Male	-	-	-	-	-	-	3	3	1	-	-	-
Female	-	-	-	-	1	1	1	3	1	2	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	1	4	11	23	24	24	25	14	2	-
Male	1	-	-	4	6	15	11	8	7	3	1	-
Female	-	2	1	-	5	8	13	16	18	11	1	-
White, both sexes	1	1	1	3	9	21	22	22	23	14	2	-
Male	1	-	-	3	5	13	11	7	7	3	1	-
Female	-	1	1	-	4	8	11	15	16	11	1	-
Black, both sexes	-	1	-	1	2	2	2	2	2	-	-	-
Male	-	-	-	1	1	2	-	1	-	-	-	-
Female	-	1	-	-	1	-	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
New Mexico												
All races, both sexes	-	-	2	-	4	3	4	4	5	1	-	-
Male	-	-	1	-	4	2	3	1	2	-	-	-
Female	-	-	1	-	-	1	1	3	3	1	-	-
White, both sexes	-	-	2	-	4	3	4	4	5	1	-	-
Male	-	-	1	-	4	2	3	1	2	-	-	-
Female	-	-	1	-	-	1	1	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	8	5	10	17	29	36	56	40	13	6	-
Male	-	4	5	6	10	17	21	19	13	2	2	-
Female	1	4	-	4	7	12	15	37	27	11	4	-
White, both sexes	1	6	5	5	17	27	34	53	37	12	5	-
Male	-	3	5	3	10	15	20	19	12	2	2	-
Female	1	3	-	2	7	12	14	34	25	10	3	-
Black, both sexes	-	2	-	3	-	2	2	3	2	1	1	-
Male	-	1	-	1	-	2	1	-	-	-	-	-
Female	-	1	-	2	-	-	1	3	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	-	-	1	-	-	-
Male	-	-	-	2	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
North Carolina												
All races, both sexes	6	8	8	11	15	15	21	22	16	11	2	-
Male	6	4	3	7	10	9	9	5	5	1	-	-
Female	-	4	5	4	5	6	12	17	11	10	2	-
White, both sexes	4	5	5	6	11	12	17	18	11	9	1	-
Male	4	3	2	4	8	8	7	5	3	1	-	-
Female	-	2	3	2	3	4	10	13	8	8	1	-
Black, both sexes	2	3	1	5	4	2	4	4	5	2	1	-
Male	2	1	-	3	2	-	2	-	2	-	-	-
Female	-	2	1	2	2	2	2	4	3	2	1	-
American Indian 1/, both sexes	-	-	2	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	2	-	4	2	3	-	-	-
Male	-	-	-	1	2	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	3	-	-	-
White, both sexes	-	-	-	1	2	-	4	2	3	-	-	-
Male	-	-	-	1	2	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Ohio												
All races, both sexes	3	3	3	11	13	13	19	27	14	7	3	-
Male	2	2	-	4	7	5	6	10	2	1	1	-
Female	1	1	3	7	6	8	13	17	12	6	2	-
White, both sexes	1	2	1	11	11	13	18	24	13	7	3	-
Male	1	2	-	4	7	5	6	9	2	1	1	-
Female	-	-	1	7	4	8	12	15	11	6	2	-
Black, both sexes	2	1	1	-	2	-	1	2	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	2	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	2	1	2	4	10	14	18	16	11	2	-
Male	-	2	1	2	1	7	6	7	4	1	-	-
Female	2	-	-	-	3	3	8	11	12	10	2	-
White, both sexes	1	2	1	2	2	8	12	16	14	10	2	-
Male	-	2	1	2	-	5	5	6	3	1	-	-
Female	1	-	-	-	2	3	7	10	11	9	2	-
Black, both sexes	-	-	-	-	2	2	1	1	1	1	-	-
Male	-	-	-	-	1	2	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Oregon												
All races, both sexes	2	-	1	3	5	1	5	9	8	1	1	-
Male	1	-	1	2	1	1	2	3	3	1	-	-
Female	1	-	-	1	4	-	3	6	5	-	1	-
White, both sexes	2	-	1	3	5	1	5	9	8	1	1	-
Male	1	-	1	2	1	1	2	3	3	1	-	-
Female	1	-	-	1	4	-	3	6	5	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	11	13	16	25	40	49	50	27	15	8	-
Male	2	7	9	11	12	25	18	21	8	4	-	-
Female	4	4	4	5	13	15	31	29	19	11	8	-
White, both sexes	5	9	10	12	22	35	44	47	24	15	8	-
Male	1	6	7	9	10	20	17	19	7	4	-	-
Female	4	3	3	3	12	15	27	28	17	11	8	-
Black, both sexes	-	2	3	4	3	5	4	3	3	-	-	-
Male	-	1	2	2	2	5	-	2	1	-	-	-
Female	-	1	1	2	1	-	4	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Rhode Island												
All races, both sexes	-	1	-	-	-	1	2	1	3	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	3	-	-	-
White, both sexes	-	1	-	-	-	1	2	1	3	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	7	9	10	8	8	9	21	15	12	3	-	-
Male	7	4	7	4	5	4	11	5	6	1	-	-
Female	-	5	3	4	3	5	10	10	6	2	-	-
White, both sexes	5	1	6	5	4	8	15	12	9	2	-	-
Male	5	1	4	3	2	4	7	5	3	1	-	-
Female	-	-	2	2	2	4	8	7	6	1	-	-
Black, both sexes	2	8	4	3	4	1	6	3	3	1	-	-
Male	2	3	3	1	3	-	4	-	3	-	-	-
Female	-	5	1	2	1	1	2	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
South Dakota												
All races, both sexes	-	-	-	-	1	1	1	1	2	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	2	-	-	-
White, both sexes	-	-	-	-	1	1	1	1	2	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	3	9	2	3	7	6	8	7	3	-	-
Male	3	2	5	1	2	2	5	2	-	-	-	-
Female	1	1	4	1	1	5	1	6	7	3	-	-
White, both sexes	3	2	8	2	3	6	6	8	7	3	-	-
Male	2	2	4	1	2	1	5	2	-	-	-	-
Female	1	-	4	1	1	5	1	6	7	3	-	-
Black, both sexes	1	1	1	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Texas												
All races, both sexes	10	18	20	20	27	53	47	78	53	30	8	-
Male	7	12	12	14	18	30	28	32	13	7	2	-
Female	3	6	8	6	9	23	19	46	40	23	6	-
White, both sexes	7	13	12	19	23	44	40	69	50	27	8	-
Male	5	9	8	13	15	26	24	27	13	6	2	-
Female	2	4	4	6	8	18	16	42	37	21	6	-
Black, both sexes	3	3	8	-	4	9	7	6	3	3	-	-
Male	2	1	4	-	3	4	4	4	-	1	-	-
Female	1	2	4	-	1	5	3	2	3	2	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	2	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Utah												
All races, both sexes	2	-	1	3	2	4	6	6	8	4	1	-
Male	1	-	1	1	1	1	2	2	1	-	-	-
Female	1	-	-	2	1	3	4	4	7	4	1	-
White, both sexes	1	-	1	3	2	3	6	6	8	4	1	-
Male	-	-	1	1	1	1	2	2	1	-	-	-
Female	1	-	-	2	1	2	4	4	7	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	10	15	11	13	25	23	51	45	40	15	3	-
Male	10	12	6	11	15	11	24	16	13	3	-	-
Female	-	3	5	2	10	12	27	29	27	12	3	-
White, both sexes	7	10	8	10	14	17	36	34	33	9	2	-
Male	7	8	5	8	8	7	17	11	11	1	-	-
Female	-	2	3	2	6	10	19	23	22	8	2	-
Black, both sexes	3	5	3	3	10	6	14	11	6	6	1	-
Male	3	4	1	3	7	4	6	5	2	2	-	-
Female	-	1	2	-	3	2	8	6	4	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Washington												
All races, both sexes	1	1	-	3	2	5	4	8	10	4	-	-
Male	1	1	-	3	1	4	-	4	4	2	-	-
Female	-	-	-	-	1	1	4	4	6	2	-	-
White, both sexes	1	1	-	3	2	3	4	7	10	4	-	-
Male	1	1	-	3	1	2	-	4	4	2	-	-
Female	-	-	-	-	1	1	4	3	6	2	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	1	2	1	3	9	13	28	31	15	12	4	-
Male	-	2	-	1	3	5	13	11	4	-	-	-
Female	1	-	1	2	6	8	15	20	11	12	4	-
White, both sexes	-	2	1	3	9	13	27	31	15	12	4	-
Male	-	2	-	1	3	5	13	11	4	-	-	-
Female	-	-	1	2	6	8	14	20	11	12	4	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Wisconsin												
All races, both sexes	1	3	1	-	1	2	4	8	4	2	-	-
Male	1	2	-	-	1	1	3	3	-	1	-	-
Female	-	1	1	-	-	1	1	5	4	1	-	-
White, both sexes	1	-	1	-	1	2	4	8	4	2	-	-
Male	1	-	-	-	1	1	3	3	-	1	-	-
Female	-	-	1	-	-	1	1	5	4	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	1	-	1	1	2	-
Male	-	-	-	-	-	1	-	-	1	-	1	-
Female	-	-	-	-	-	-	1	-	-	1	1	-
White, both sexes	-	-	-	-	-	1	1	-	1	1	2	-
Male	-	-	-	-	-	1	-	-	1	-	1	-
Female	-	-	-	-	-	-	1	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
United States												
All races, both sexes	678	28	21	11	20	33	36	48	38	41	51	70
Male	342	15	12	6	9	20	22	32	19	22	25	33
Female	336	13	9	5	11	13	14	16	19	19	26	37
White, both sexes	536	14	16	6	16	20	26	36	30	35	39	49
Male	275	8	10	2	9	14	18	25	15	17	20	22
Female	261	6	6	4	7	6	8	11	15	18	19	27
Black, both sexes	123	11	5	4	3	11	10	9	7	5	10	20
Male	56	6	2	3	-	5	4	5	3	4	3	10
Female	67	5	3	1	3	6	6	4	4	1	7	10
American Indian 1/, both sexes	4	1	-	-	1	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	1	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	2	-	1	-	1	-	3	1	1	2	-
Male	10	1	-	1	-	1	-	2	1	1	2	-
Female	5	1	-	-	-	-	-	1	-	-	-	-
Alabama												
All races, both sexes	13	-	-	1	1	1	-	3	1	-	2	2
Male	6	-	-	1	-	-	-	3	-	-	-	1
Female	7	-	-	-	1	1	-	-	1	-	2	1
White, both sexes	9	-	-	-	1	1	-	2	1	-	-	2
Male	4	-	-	-	-	-	-	2	-	-	-	1
Female	5	-	-	-	1	1	-	-	1	-	-	1
Black, both sexes	3	-	-	1	-	-	-	-	-	-	2	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	6	-	-	-	-	-	-	1	-	-	-	2
Male	3	-	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	6	-	-	-	-	-	-	1	-	-	-	2
Male	3	-	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Arkansas												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	40	2	1	1	2	2	1	1	3	6	1	5
Male	21	2	1	1	-	2	1	1	-	3	1	1
Female	19	-	-	-	2	-	-	-	3	3	-	4
White, both sexes	30	1	1	-	2	1	-	-	1	6	-	3
Male	14	1	1	-	-	1	-	-	-	3	-	-
Female	16	-	-	-	2	-	-	-	1	3	-	3
Black, both sexes	7	-	-	1	-	-	1	1	2	-	-	2
Male	4	-	-	1	-	-	1	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	1	-	-	-	-	1	-
Male	3	1	-	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Colorado												
All races, both sexes	13	1	-	2	-	-	-	2	-	-	1	2
Male	5	-	-	2	-	-	-	1	-	-	-	-
Female	8	1	-	-	-	-	-	1	-	-	1	2
White, both sexes	12	1	-	1	-	-	-	2	-	-	1	2
Male	4	-	-	1	-	-	-	1	-	-	-	-
Female	8	1	-	-	-	-	-	1	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	12	-	-	-	-	1	-	3	1	1	-	-
Male	8	-	-	-	-	-	-	2	1	1	-	-
Female	4	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	11	-	-	-	-	1	-	2	1	1	-	-
Male	7	-	-	-	-	-	-	1	1	1	-	-
Female	4	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Florida												
All races, both sexes	57	1	1	-	1	5	5	2	2	2	2	10
Male	32	-	1	-	1	4	4	1	2	2	1	6
Female	25	1	-	-	-	1	1	1	-	-	1	4
White, both sexes	39	-	1	-	1	3	3	-	1	1	1	5
Male	23	-	1	-	1	3	3	-	1	1	-	3
Female	16	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	15	1	-	-	-	2	2	1	1	1	-	4
Male	7	-	-	-	-	1	1	1	1	1	-	2
Female	8	1	-	-	-	1	1	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	29	-	-	1	-	1	1	1	1	-	3	2
Male	10	-	-	1	-	1	-	-	1	-	-	1
Female	19	-	-	-	-	-	1	1	-	-	3	1
White, both sexes	23	-	-	-	-	-	1	1	1	-	2	2
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	16	-	-	-	-	-	1	1	-	-	2	1
Black, both sexes	5	-	-	1	-	1	-	-	-	-	1	-
Male	3	-	-	1	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Illinois												
All races, both sexes	23	1	1	-	1	-	1	5	1	-	1	2
Male	10	1	1	-	-	-	-	3	-	-	-	1
Female	13	-	-	-	1	-	1	2	1	-	1	1
White, both sexes	19	1	-	-	1	-	-	4	1	-	-	2
Male	9	1	-	-	-	-	-	3	-	-	-	1
Female	10	-	-	-	1	-	-	1	1	-	-	1
Black, both sexes	4	-	1	-	-	-	1	1	-	-	1	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	23	1	1	-	-	2	2	2	2	-	2	3
Male	10	1	-	-	-	-	2	-	-	-	1	2
Female	13	-	1	-	-	2	-	2	2	-	1	1
White, both sexes	17	1	1	-	-	-	2	1	2	-	2	1
Male	8	1	-	-	-	-	2	-	-	-	1	1
Female	9	-	1	-	-	-	-	1	2	-	1	-
Black, both sexes	6	-	-	-	-	2	-	1	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	2	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Iowa												
All races, both sexes	6	1	-	-	-	1	-	-	-	2	-	-
Male	3	1	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	4	-	-	-	-	1	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	2	1	-	-	-	-	-	-	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	1	-	1	-	-	1	-	-	-	1	-
Male	5	-	-	-	-	-	1	-	-	-	1	-
Female	4	1	-	1	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	1	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Kentucky												
All races, both sexes	7	-	1	-	-	-	1	-	-	1	-	-
Male	4	-	1	-	-	-	1	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	1	-	-	-	1	-	-	1	-	-
Male	4	-	1	-	-	-	1	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	12	2	1	1	-	-	1	-	1	1	-	3
Male	7	1	1	-	-	-	1	-	-	1	-	1
Female	5	1	-	1	-	-	-	-	1	-	-	2
White, both sexes	8	-	1	-	-	-	1	-	1	1	-	2
Male	5	-	1	-	-	-	1	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	4	2	-	1	-	-	-	-	-	-	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	1
Female	2	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Maine												
All races, both sexes	9	-	1	-	-	-	1	2	1	1	3	-
Male	6	-	1	-	-	-	-	2	1	-	2	-
Female	3	-	-	-	-	-	1	-	-	1	1	-
White, both sexes	9	-	1	-	-	-	1	2	1	1	3	-
Male	6	-	1	-	-	-	-	2	1	-	2	-
Female	3	-	-	-	-	-	1	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	18	-	1	-	-	1	1	3	2	1	2	1
Male	9	-	1	-	-	1	1	2	2	-	-	-
Female	9	-	-	-	-	-	-	1	-	1	2	1
White, both sexes	10	-	1	-	-	1	1	1	1	1	-	1
Male	6	-	1	-	-	1	1	1	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	8	-	-	-	-	-	-	2	1	-	2	-
Male	3	-	-	-	-	-	-	1	1	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Massachusetts												
All races, both sexes	14	-	-	-	-	1	-	1	1	1	3	-
Male	7	-	-	-	-	1	-	-	-	1	2	-
Female	7	-	-	-	-	-	-	1	1	-	1	-
White, both sexes	14	-	-	-	-	1	-	1	1	1	3	-
Male	7	-	-	-	-	1	-	-	-	1	2	-
Female	7	-	-	-	-	-	-	1	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	37	-	1	-	1	2	-	1	2	2	2	6
Male	18	-	-	-	1	2	-	1	1	1	1	2
Female	19	-	1	-	-	-	-	-	1	1	1	4
White, both sexes	33	-	1	-	1	1	-	1	2	2	2	4
Male	17	-	-	-	1	1	-	1	1	1	1	2
Female	16	-	1	-	-	-	-	-	1	1	1	2
Black, both sexes	4	-	-	-	-	1	-	-	-	-	-	2
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Minnesota												
All races, both sexes	20	2	-	2	-	2	4	1	1	2	-	2
Male	8	-	-	-	-	1	2	-	-	1	-	2
Female	12	2	-	2	-	1	2	1	1	1	-	-
White, both sexes	15	1	-	2	-	-	3	1	1	1	-	2
Male	5	-	-	-	-	-	1	-	-	-	-	2
Female	10	1	-	2	-	-	2	1	1	1	-	-
Black, both sexes	3	-	-	-	-	2	1	-	-	-	-	-
Male	2	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Missouri												
All races, both sexes	9	-	-	-	1	1	2	1	-	1	-	-
Male	4	-	-	-	1	-	2	1	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	6	-	-	-	1	1	2	-	-	-	-	-
Male	3	-	-	-	1	-	2	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Nebraska												
All races, both sexes	11	-	-	-	1	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	1	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	19	-	1	-	-	-	2	1	-	-	5	2
Male	12	-	-	-	-	-	2	1	-	-	3	1
Female	7	-	1	-	-	-	-	-	-	-	2	1
White, both sexes	14	-	1	-	-	-	2	1	-	-	3	-
Male	9	-	-	-	-	-	2	1	-	-	2	-
Female	5	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	2
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Mexico												
All races, both sexes	4	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	34	1	1	1	1	2	2	4	2	3	2	2
Male	19	1	1	1	1	-	1	3	2	2	-	1
Female	15	-	-	-	-	2	1	1	-	1	2	1
White, both sexes	29	1	1	1	1	2	-	4	1	3	2	2
Male	16	1	1	1	1	-	-	3	1	2	-	1
Female	13	-	-	-	-	2	-	1	-	1	2	1
Black, both sexes	4	-	-	-	-	-	2	-	1	-	-	-
Male	3	-	-	-	-	-	1	-	1	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
North Carolina												
All races, both sexes	19	3	2	-	1	-	2	-	3	-	-	1
Male	5	1	-	-	1	-	-	-	1	-	-	-
Female	14	2	2	-	-	-	2	-	2	-	-	1
White, both sexes	13	1	-	-	1	-	1	-	3	-	-	1
Male	5	1	-	-	1	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	1	-	2	-	-	1
Black, both sexes	5	1	2	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	2	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Ohio												
All races, both sexes	39	2	-	-	1	2	3	1	2	2	2	7
Male	22	1	-	-	1	1	2	1	1	2	-	4
Female	17	1	-	-	-	1	1	-	1	-	2	3
White, both sexes	32	1	-	-	1	1	3	1	2	1	2	6
Male	19	-	-	-	1	1	2	1	1	1	-	4
Female	13	1	-	-	-	-	1	-	1	-	2	2
Black, both sexes	7	1	-	-	-	1	-	-	-	1	-	1
Male	3	1	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	9	-	1	-	2	1	-	-	-	-	2	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	7	-	1	-	2	-	-	-	-	-	2	-
White, both sexes	9	-	1	-	2	1	-	-	-	-	2	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	7	-	1	-	2	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Oregon												
All races, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	25	-	-	-	-	-	2	3	4	2	4	3
Male	12	-	-	-	-	-	1	1	2	-	2	1
Female	13	-	-	-	-	-	1	2	2	2	2	2
White, both sexes	22	-	-	-	-	-	2	3	3	2	4	2
Male	10	-	-	-	-	-	1	1	2	-	2	-
Female	12	-	-	-	-	-	1	2	1	2	2	2
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Rhode Island												
All races, both sexes	5	-	-	-	1	1	-	1	-	-	1	-
Male	3	-	-	-	-	1	-	1	-	-	1	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	1	-	1	-	-	1	-
Male	3	-	-	-	-	1	-	1	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	11	1	-	-	-	-	-	2	-	-	2	1
Male	6	1	-	-	-	-	-	2	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	2	-	-	2	-
Male	5	-	-	-	-	-	-	2	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	-	1
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
South Dakota												
All races, both sexes	2	-	-	-	1	-	-	1	-	-	-	-
Male	2	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	1	-	-	1	-	-	-	-
Male	2	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	13	1	-	-	-	-	1	1	1	1	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	8	1	-	-	-	-	1	-	1	1	-	1
White, both sexes	10	-	-	-	-	-	1	1	-	1	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	3	1	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Texas												
All races, both sexes	57	6	4	1	3	3	2	3	1	3	4	7
Male	30	4	2	-	1	2	-	2	1	2	4	4
Female	27	2	2	1	2	1	2	1	-	1	-	3
White, both sexes	42	4	3	1	2	2	-	2	1	2	4	5
Male	22	2	2	-	1	2	-	1	1	1	4	2
Female	20	2	1	1	1	-	-	1	-	1	-	3
Black, both sexes	14	2	1	-	-	1	2	1	-	1	-	2
Male	8	2	-	-	-	-	-	1	-	1	-	2
Female	6	-	1	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	14	-	1	-	1	1	1	-	1	4	1	3
Male	9	-	-	-	1	1	1	-	1	1	-	3
Female	5	-	1	-	-	-	-	-	-	3	1	-
White, both sexes	14	-	1	-	1	1	1	-	1	4	1	3
Male	9	-	-	-	1	1	1	-	1	1	-	3
Female	5	-	1	-	-	-	-	-	-	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	15	-	2	-	-	2	-	-	1	1	2	1
Male	15	-	2	-	-	2	-	-	1	1	2	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	1	-	-	1	-	-	-	1	1	-
Male	9	-	1	-	-	1	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	1	-	-	1	-	-	-	-	1	1
Male	5	-	1	-	-	1	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Washington												
All races, both sexes	9	-	-	-	-	1	-	-	1	1	-	-
Male	3	-	-	-	-	-	-	-	1	1	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	1	1	-	-
Male	3	-	-	-	-	-	-	-	1	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	5	1	-	-	-	-	-	-	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	5	1	-	-	-	-	-	-	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Wisconsin												
All races, both sexes	9	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
United States												
All races, both sexes	58	36	28	18	20	20	22	35	23	16	5	-
Male	29	16	15	7	13	11	8	12	9	5	2	-
Female	29	20	13	11	7	9	14	23	14	11	3	-
White, both sexes	50	31	25	13	19	18	21	32	19	16	5	-
Male	26	14	13	6	12	10	7	12	8	5	2	-
Female	24	17	12	7	7	8	14	20	11	11	3	-
Black, both sexes	7	5	2	4	1	1	1	3	4	-	-	-
Male	3	2	1	1	1	1	1	-	1	-	-	-
Female	4	3	1	3	-	-	-	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	3	1	-	2	-	1	5	-	-	-	-
Male	1	2	1	-	2	-	1	1	-	-	-	-
Female	2	1	-	-	-	-	-	4	-	-	-	-
White, both sexes	3	3	1	-	2	-	1	5	-	-	-	-
Male	1	2	1	-	2	-	1	1	-	-	-	-
Female	2	1	-	-	-	-	-	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Colorado												
All races, both sexes	2	-	2	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	2	-	2	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	-	-	1	1	1	-	-	1	-	-
Male	-	1	-	-	-	1	1	-	-	1	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	1	1	1	-	-	1	-	-
Male	-	1	-	-	-	1	1	-	-	1	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Delaware												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Florida												
All races, both sexes	3	2	2	1	3	1	3	6	2	2	1	-
Male	1	1	1	-	2	1	-	3	-	1	-	-
Female	2	1	1	1	1	-	3	3	2	1	1	-
White, both sexes	2	2	2	1	3	1	3	6	-	2	1	-
Male	1	1	1	-	2	1	-	3	-	1	-	-
Female	1	1	1	1	1	-	3	3	-	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4	1	1	2	-	1	2	2	4	2	-	-
Male	2	-	-	-	-	1	1	-	1	1	-	-
Female	2	1	1	2	-	-	1	2	3	1	-	-
White, both sexes	3	1	1	1	-	1	1	2	4	2	-	-
Male	2	-	-	-	-	1	-	-	1	1	-	-
Female	1	1	1	1	-	-	1	2	3	1	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Illinois												
All races, both sexes	2	-	1	-	-	-	2	3	2	-	-	-
Male	1	-	1	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	1	2	2	-	-	-
White, both sexes	2	-	1	-	-	-	2	3	2	-	-	-
Male	1	-	1	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	1	1	2	-	-	-	-	1	1	-	-
Male	1	-	-	1	-	-	-	-	1	1	-	-
Female	1	1	1	1	-	-	-	-	-	-	-	-
White, both sexes	1	1	1	2	-	-	-	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	1	1	-	-
Female	1	1	1	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Iowa												
All races, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	2	-	-	-	-	-	-	1	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	2	-	-	-	-	-	-	1	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Kentucky												
All races, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
White, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	2	1	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Massachusetts												
All races, both sexes	2	3	-	-	-	1	-	1	-	-	-	-
Male	-	3	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	2	3	-	-	-	1	-	1	-	-	-	-
Male	-	3	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7	2	3	-	2	1	-	3	-	2	-	-
Male	3	1	2	-	-	1	-	1	-	1	-	-
Female	4	1	1	-	2	-	-	2	-	1	-	-
White, both sexes	6	2	3	-	2	1	-	3	-	2	-	-
Male	3	1	2	-	-	1	-	1	-	1	-	-
Female	3	1	1	-	2	-	-	2	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Minnesota												
All races, both sexes	-	-	1	-	-	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Missouri												
All races, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Nebraska												
All races, both sexes	2	1	1	-	1	-	1	-	1	1	-	-
Male	2	-	1	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	1	-	1	-	-	1	-	-
White, both sexes	2	1	1	-	1	-	1	-	1	1	-	-
Male	2	-	1	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	1	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	1	1	2	1	1	1	-	-	-	-	-
Male	-	1	1	2	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	1	1	1	1	1	1	1	-	-	-	-	-
Male	-	1	1	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Mexico												
All races, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	2	2	2	-	3	1	-	-	1	-	-
Male	2	1	-	1	-	2	-	-	-	-	-	-
Female	-	1	2	1	-	1	1	-	-	1	-	-
White, both sexes	2	1	2	2	-	2	1	-	-	1	-	-
Male	2	-	-	1	-	2	-	-	-	-	-	-
Female	-	1	2	1	-	-	1	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
North Carolina												
All races, both sexes	1	3	-	-	-	-	-	-	1	2	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	1	2	-	-	-	-	-	-	-	2	-	-
White, both sexes	1	2	-	-	-	-	-	-	1	2	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	1	1	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Ohio												
All races, both sexes	2	3	-	1	3	1	3	2	1	-	1	-
Male	-	1	-	-	3	1	2	1	-	-	1	-
Female	2	2	-	1	-	-	1	1	1	-	-	-
White, both sexes	2	2	-	-	2	1	3	2	1	-	1	-
Male	-	1	-	-	2	1	2	1	-	-	1	-
Female	2	1	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Oregon												
All races, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	1	3	1	-	-	-	-	1	-	-	-
Male	1	-	2	1	-	-	-	-	1	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	2	1	-	-	-	-	1	-	-	-
Male	1	-	1	1	-	-	-	-	1	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Rhode Island												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	2	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	1	-	-	1	-	-	1	1	-	1	-
Male	2	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	1	-
White, both sexes	2	-	-	-	1	-	-	1	1	-	1	-
Male	2	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Texas												
All races, both sexes	6	2	2	2	1	1	-	4	2	-	-	-
Male	4	-	-	-	1	-	-	2	1	-	-	-
Female	2	2	2	2	-	1	-	2	1	-	-	-
White, both sexes	5	2	2	1	1	1	-	3	1	-	-	-
Male	3	-	-	-	1	-	-	2	-	-	-	-
Female	2	2	2	1	-	1	-	1	1	-	-	-
Black, both sexes	1	-	-	1	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	1	-	-	1	1	-	-	-	-	1	-
Male	2	1	-	-	1	1	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	1	-	-	-	-	-	1	-
Male	2	1	-	-	1	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Washington												
All races, both sexes	2	1	-	-	-	-	1	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	1	1	-	1	-	-
White, both sexes	1	1	-	-	-	-	1	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Wisconsin												
All races, both sexes	1	1	1	1	1	-	-	1	-	1	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	1	-	1	-	-
White, both sexes	1	1	1	1	1	-	-	1	-	1	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
United States												
All races, both sexes	10,986	22	15	7	17	31	62	92	132	202	332	438
Male	5,578	9	9	5	9	22	53	67	111	140	214	306
Female	5,408	13	6	2	8	9	9	25	21	62	118	132
White, both sexes	9,320	15	12	6	13	17	40	57	94	134	258	333
Male	4,677	7	8	5	8	12	38	44	83	92	173	241
Female	4,643	8	4	1	5	5	2	13	11	42	85	92
Black, both sexes	1,469	5	3	1	4	14	19	33	34	61	66	90
Male	800	2	1	-	1	10	13	22	26	44	36	56
Female	669	3	2	1	3	4	6	11	8	17	30	34
American Indian 1/, both sexes	48	1	-	-	-	-	1	1	2	4	2	1
Male	28	-	-	-	-	-	-	1	1	2	1	1
Female	20	1	-	-	-	-	1	-	1	2	1	-
Asian or Pacific Islander 2/, both sexes	149	1	-	-	-	-	2	1	2	3	6	14
Male	73	-	-	-	-	-	2	-	1	2	4	8
Female	76	1	-	-	-	-	-	1	1	1	2	6
Alabama												
All races, both sexes	397	-	-	-	1	1	-	6	2	4	7	21
Male	214	-	-	-	1	-	-	5	1	3	5	15
Female	183	-	-	-	-	1	-	1	1	1	2	6
White, both sexes	319	-	-	-	-	1	-	2	1	1	6	16
Male	164	-	-	-	-	-	-	2	1	1	4	12
Female	155	-	-	-	-	1	-	-	-	-	2	4
Black, both sexes	77	-	-	-	1	-	-	4	1	3	1	5
Male	49	-	-	-	1	-	-	3	-	2	1	3
Female	28	-	-	-	-	-	-	1	1	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Alaska												
All races, both sexes	29	-	-	-	-	-	-	-	-	1	-	-
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	249	-	1	1	2	1	-	2	5	8	8	12
Male	146	-	1	-	1	-	-	1	5	7	5	7
Female	103	-	-	1	1	1	-	1	-	1	3	5
White, both sexes	229	-	1	1	2	1	-	1	3	7	7	11
Male	130	-	1	-	1	-	-	1	3	6	4	6
Female	99	-	-	1	1	1	-	-	-	1	3	5
Black, both sexes	12	-	-	-	-	-	-	1	1	1	-	1
Male	9	-	-	-	-	-	-	-	1	1	-	1
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Arkansas												
All races, both sexes	217	-	3	-	1	-	-	1	-	1	3	4
Male	92	-	2	-	-	-	-	-	-	-	1	4
Female	125	-	1	-	1	-	-	1	-	1	2	-
White, both sexes	189	-	2	-	-	-	-	1	-	-	2	4
Male	81	-	2	-	-	-	-	-	-	-	1	4
Female	108	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	28	-	1	-	1	-	-	-	-	1	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	1	-	1	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,047	4	-	-	1	2	5	8	22	26	42	60
Male	503	2	-	-	-	2	3	5	19	21	31	44
Female	544	2	-	-	1	-	2	3	3	5	11	16
White, both sexes	907	3	-	-	1	2	3	7	18	17	38	48
Male	440	2	-	-	-	2	2	5	17	13	30	39
Female	467	1	-	-	1	-	1	2	1	4	8	9
Black, both sexes	85	-	-	-	-	-	2	1	3	5	3	7
Male	42	-	-	-	-	-	1	-	2	5	1	2
Female	43	-	-	-	-	-	1	1	1	-	2	5
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	2	-	-
Male	4	-	-	-	-	-	-	-	-	2	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	1	-	-	-	-	-	-	1	2	1	5
Male	17	-	-	-	-	-	-	-	-	1	-	3
Female	32	1	-	-	-	-	-	-	1	1	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Colorado												
All races, both sexes	78	-	1	-	-	1	1	-	-	-	4	4
Male	34	-	1	-	-	1	-	-	-	-	3	2
Female	44	-	-	-	-	-	1	-	-	-	1	2
White, both sexes	71	-	1	-	-	1	-	-	-	-	4	4
Male	32	-	1	-	-	1	-	-	-	-	3	2
Female	39	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	80	-	1	-	-	-	1	1	1	1	-	3
Male	39	-	-	-	-	-	1	1	1	1	-	2
Female	41	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	78	-	1	-	-	-	1	1	1	1	-	3
Male	37	-	-	-	-	-	1	1	1	1	-	2
Female	41	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Delaware												
All races, both sexes	33	-	-	-	-	-	-	-	2	2	2	-
Male	18	-	-	-	-	-	-	-	2	2	2	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	2	1	1	-
Male	13	-	-	-	-	-	-	-	2	1	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Florida												
All races, both sexes	839	2	1	-	4	-	3	7	7	11	8	11
Male	417	1	1	-	1	-	1	5	7	8	5	7
Female	422	1	-	-	3	-	2	2	-	3	3	4
White, both sexes	751	2	1	-	3	-	-	3	5	6	7	9
Male	369	1	1	-	1	-	-	2	5	4	5	5
Female	382	1	-	-	2	-	-	1	-	2	2	4
Black, both sexes	85	-	-	-	1	-	3	4	2	5	1	2
Male	47	-	-	-	-	-	1	3	2	4	-	2
Female	38	-	-	-	1	-	2	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	794	-	-	2	2	5	3	5	9	9	23	43
Male	455	-	-	2	1	3	3	4	8	8	13	33
Female	339	-	-	-	1	2	-	1	1	1	10	10
White, both sexes	578	-	-	2	2	2	1	2	6	5	17	29
Male	333	-	-	2	1	1	1	2	5	4	8	23
Female	245	-	-	-	1	1	-	-	1	1	9	6
Black, both sexes	209	-	-	-	-	3	1	3	3	4	6	14
Male	118	-	-	-	-	2	1	2	3	4	5	10
Female	91	-	-	-	-	1	-	1	-	-	1	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Hawaii												
All races, both sexes	66	-	-	-	-	-	-	-	1	1	4	11
Male	42	-	-	-	-	-	-	-	1	1	3	6
Female	24	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	27	-	-	-	-	-	-	-	-	1	1	2
Male	17	-	-	-	-	-	-	-	-	1	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	1	-	3	9
Male	25	-	-	-	-	-	-	-	1	-	2	5
Female	14	-	-	-	-	-	-	-	-	-	1	4
Idaho												
All races, both sexes	28	-	-	-	-	-	1	2	-	1	2	3
Male	17	-	-	-	-	-	1	2	-	1	-	3
Female	11	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	28	-	-	-	-	-	1	2	-	1	2	3
Male	17	-	-	-	-	-	1	2	-	1	-	3
Female	11	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Illinois												
All races, both sexes	458	1	1	-	-	1	2	3	6	3	11	17
Male	211	-	-	-	-	1	1	2	5	2	8	12
Female	247	1	1	-	-	-	1	1	1	1	3	5
White, both sexes	389	1	1	-	-	1	2	2	3	3	9	16
Male	175	-	-	-	-	1	1	2	2	2	6	11
Female	214	1	1	-	-	-	1	-	1	1	3	5
Black, both sexes	68	-	-	-	-	-	-	1	3	-	2	1
Male	35	-	-	-	-	-	-	-	3	-	2	1
Female	33	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	170	1	1	-	-	-	2	1	4	6	9	7
Male	93	-	1	-	-	-	2	1	4	3	7	5
Female	77	1	-	-	-	-	-	-	-	3	2	2
White, both sexes	155	1	1	-	-	-	2	1	4	3	8	7
Male	84	-	1	-	-	-	2	1	4	1	6	5
Female	71	1	-	-	-	-	-	-	-	2	2	2
Black, both sexes	14	-	-	-	-	-	-	-	-	3	1	-
Male	9	-	-	-	-	-	-	-	-	2	1	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Iowa												
All races, both sexes	68	-	-	-	-	-	-	-	-	1	1	-
Male	33	-	-	-	-	-	-	-	-	1	-	-
Female	35	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	65	-	-	-	-	-	-	-	-	1	1	-
Male	32	-	-	-	-	-	-	-	-	1	-	-
Female	33	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	82	-	-	-	-	-	1	2	-	3	3	7
Male	42	-	-	-	-	-	1	-	-	-	3	4
Female	40	-	-	-	-	-	-	2	-	3	-	3
White, both sexes	78	-	-	-	-	-	1	1	-	3	3	6
Male	40	-	-	-	-	-	1	-	-	-	3	4
Female	38	-	-	-	-	-	-	1	-	3	-	2
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Kentucky												
All races, both sexes	252	-	-	1	-	1	1	3	1	4	7	12
Male	140	-	-	1	-	-	1	3	1	2	4	8
Female	112	-	-	-	-	1	-	-	-	2	3	4
White, both sexes	234	-	-	1	-	-	1	3	1	4	6	10
Male	132	-	-	1	-	-	1	3	1	2	3	7
Female	102	-	-	-	-	-	-	-	-	2	3	3
Black, both sexes	14	-	-	-	-	1	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	307	3	-	-	-	-	1	-	3	5	8	8
Male	145	-	-	-	-	-	1	-	3	4	7	6
Female	162	3	-	-	-	-	-	-	-	1	1	2
White, both sexes	232	2	-	-	-	-	1	-	2	2	4	3
Male	106	-	-	-	-	-	1	-	2	2	3	3
Female	126	2	-	-	-	-	-	-	-	-	1	-
Black, both sexes	74	1	-	-	-	-	-	-	1	3	3	5
Male	38	-	-	-	-	-	-	-	1	2	3	3
Female	36	1	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Maine												
All races, both sexes	40	-	-	-	-	-	1	-	-	-	2	1
Male	24	-	-	-	-	-	1	-	-	-	2	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	1	-	-	-	2	1
Male	23	-	-	-	-	-	1	-	-	-	2	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	109	-	-	-	2	2	-	2	2	7	17	8
Male	67	-	-	-	1	2	-	2	2	4	12	5
Female	42	-	-	-	1	-	-	-	-	3	5	3
White, both sexes	79	-	-	-	1	-	-	2	1	4	13	5
Male	48	-	-	-	1	-	-	2	1	2	11	3
Female	31	-	-	-	-	-	-	-	-	2	2	2
Black, both sexes	29	-	-	-	1	2	-	-	1	3	4	3
Male	19	-	-	-	-	2	-	-	1	2	1	2
Female	10	-	-	-	1	-	-	-	-	1	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Massachusetts												
All races, both sexes	175	-	-	-	1	-	2	-	2	3	4	5
Male	82	-	-	-	1	-	2	-	2	2	3	3
Female	93	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	168	-	-	-	1	-	2	-	2	3	4	5
Male	79	-	-	-	1	-	2	-	2	2	3	3
Female	89	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	270	1	1	-	-	1	-	9	1	5	7	6
Male	142	1	-	-	-	-	-	6	1	3	7	4
Female	128	-	1	-	-	1	-	3	-	2	-	2
White, both sexes	232	1	-	-	-	1	-	5	-	4	4	5
Male	122	1	-	-	-	-	-	4	-	3	4	3
Female	110	-	-	-	-	1	-	1	-	1	-	2
Black, both sexes	35	-	1	-	-	-	-	4	1	1	3	-
Male	18	-	-	-	-	-	-	2	1	-	3	-
Female	17	-	1	-	-	-	-	2	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Minnesota												
All races, both sexes	124	-	1	-	-	-	1	-	2	5	4	1
Male	65	-	1	-	-	-	1	-	1	3	3	1
Female	59	-	-	-	-	-	-	-	1	2	1	-
White, both sexes	112	-	1	-	-	-	1	-	1	2	3	1
Male	59	-	1	-	-	-	1	-	1	1	2	1
Female	53	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	3	1	-
Male	5	-	-	-	-	-	-	-	-	2	1	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	294	-	-	1	-	-	2	-	1	3	2	10
Male	137	-	-	-	-	-	1	-	1	2	1	7
Female	157	-	-	1	-	-	1	-	-	1	1	3
White, both sexes	197	-	-	-	-	-	1	-	1	-	1	6
Male	97	-	-	-	-	-	1	-	1	-	1	5
Female	100	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	97	-	-	1	-	-	1	-	-	3	1	4
Male	40	-	-	-	-	-	-	-	-	2	-	2
Female	57	-	-	1	-	-	1	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Missouri												
All races, both sexes	219	1	-	-	-	1	3	-	4	5	10	3
Male	107	-	-	-	-	1	3	-	4	4	5	3
Female	112	1	-	-	-	-	-	-	-	1	5	-
White, both sexes	192	1	-	-	-	1	3	-	3	2	9	3
Male	91	-	-	-	-	1	3	-	3	1	5	3
Female	101	1	-	-	-	-	-	-	-	1	4	-
Black, both sexes	27	-	-	-	-	-	-	-	1	3	1	-
Male	16	-	-	-	-	-	-	-	1	3	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	36	-	-	-	-	-	-	-	-	1	1	1
Male	24	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	33	-	-	-	-	-	-	-	-	1	1	1
Male	22	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Nebraska												
All races, both sexes	102	-	-	-	-	-	-	1	1	3	4	3
Male	53	-	-	-	-	-	-	1	1	3	2	2
Female	49	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	96	-	-	-	-	-	-	1	1	2	4	2
Male	49	-	-	-	-	-	-	1	1	2	2	1
Female	47	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	107	-	-	-	-	1	-	-	1	-	2	2
Male	45	-	-	-	-	1	-	-	1	-	1	1
Female	62	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	95	-	-	-	-	1	-	-	1	-	2	2
Male	42	-	-	-	-	1	-	-	1	-	1	1
Female	53	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Hampshire												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	1	4
Male	38	-	-	-	-	-	-	-	-	-	1	4
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	1	4
Male	38	-	-	-	-	-	-	-	-	-	1	4
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	239	1	-	-	2	-	2	1	-	6	5	8
Male	111	-	-	-	2	-	2	1	-	4	4	5
Female	128	1	-	-	-	-	-	-	-	2	1	3
White, both sexes	207	-	-	-	2	-	-	1	-	4	3	5
Male	92	-	-	-	2	-	-	1	-	3	3	4
Female	115	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	29	1	-	-	-	-	2	-	-	2	2	3
Male	18	-	-	-	-	-	2	-	-	1	1	1
Female	11	1	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Mexico												
All races, both sexes	54	-	-	-	-	-	-	1	-	-	-	2
Male	28	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	1	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	470	-	1	1	-	1	3	6	6	14	25	27
Male	252	-	-	1	-	1	3	4	5	11	15	18
Female	218	-	1	-	-	-	-	2	1	3	10	9
White, both sexes	401	-	1	1	-	-	2	4	4	9	18	18
Male	211	-	-	1	-	-	2	2	3	7	11	13
Female	190	-	1	-	-	-	-	2	1	2	7	5
Black, both sexes	64	-	-	-	-	1	1	2	2	5	7	9
Male	37	-	-	-	-	1	1	2	2	4	4	5
Female	27	-	-	-	-	-	-	-	-	1	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
North Carolina												
All races, both sexes	305	2	-	-	-	1	1	2	1	4	7	11
Male	139	1	-	-	-	-	1	2	1	2	3	8
Female	166	1	-	-	-	1	-	-	-	2	4	3
White, both sexes	224	1	-	-	-	-	-	1	1	4	4	5
Male	98	-	-	-	-	-	-	1	1	2	2	4
Female	126	1	-	-	-	-	-	-	-	2	2	1
Black, both sexes	77	1	-	-	-	1	1	1	-	-	3	6
Male	38	1	-	-	-	-	1	1	-	-	1	4
Female	39	-	-	-	-	1	-	-	-	-	2	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Ohio												
All races, both sexes	291	-	-	-	-	1	2	1	7	4	7	10
Male	142	-	-	-	-	1	2	1	5	4	3	6
Female	149	-	-	-	-	-	-	-	2	-	4	4
White, both sexes	256	-	-	-	-	-	2	-	3	3	6	10
Male	125	-	-	-	-	-	2	-	2	3	3	6
Female	131	-	-	-	-	-	-	-	1	-	3	4
Black, both sexes	35	-	-	-	-	1	-	1	4	1	1	-
Male	17	-	-	-	-	1	-	1	3	1	-	-
Female	18	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	122	-	1	-	-	1	-	3	-	4	7	3
Male	48	-	-	-	-	-	-	2	-	1	3	3
Female	74	-	1	-	-	1	-	1	-	3	4	-
White, both sexes	111	-	1	-	-	-	-	-	-	3	7	2
Male	44	-	-	-	-	-	-	-	-	1	3	2
Female	67	-	1	-	-	-	-	-	-	2	4	-
Black, both sexes	6	-	-	-	-	1	-	2	-	-	-	1
Male	3	-	-	-	-	-	-	2	-	-	-	1
Female	3	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Oregon												
All races, both sexes	85	1	-	1	-	-	-	2	2	2	2	1
Male	46	-	-	1	-	-	-	2	1	1	2	1
Female	39	1	-	-	-	-	-	-	1	1	-	-
White, both sexes	80	-	-	1	-	-	-	2	1	2	2	1
Male	42	-	-	1	-	-	-	2	-	1	2	1
Female	38	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	450	1	-	-	-	-	2	2	4	3	11	14
Male	221	1	-	-	-	-	2	1	4	1	8	9
Female	229	-	-	-	-	-	-	1	-	2	3	5
White, both sexes	412	1	-	-	-	-	1	2	3	3	8	14
Male	196	1	-	-	-	-	1	1	3	1	6	9
Female	216	-	-	-	-	-	-	1	-	2	2	5
Black, both sexes	38	-	-	-	-	-	1	-	1	-	3	-
Male	25	-	-	-	-	-	1	-	1	-	2	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Rhode Island												
All races, both sexes	17	-	-	-	-	-	-	-	2	2	-	1
Male	12	-	-	-	-	-	-	-	2	-	-	1
Female	5	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	15	-	-	-	-	-	-	-	2	2	-	-
Male	10	-	-	-	-	-	-	-	2	-	-	-
Female	5	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	324	1	-	-	-	2	5	9	7	10	23	28
Male	205	-	-	-	-	2	5	8	4	7	16	19
Female	119	1	-	-	-	-	-	1	3	3	7	9
White, both sexes	203	-	-	-	-	-	3	4	2	5	16	14
Male	119	-	-	-	-	-	3	3	1	4	13	10
Female	84	-	-	-	-	-	-	1	1	1	3	4
Black, both sexes	120	1	-	-	-	2	2	5	5	5	7	14
Male	85	-	-	-	-	2	2	5	3	3	3	9
Female	35	1	-	-	-	-	-	-	2	2	4	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
South Dakota												
All races, both sexes	36	-	-	-	-	-	1	-	-	-	4	3
Male	20	-	-	-	-	-	-	-	-	-	4	3
Female	16	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	4	3
Male	19	-	-	-	-	-	-	-	-	-	4	3
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	205	1	-	-	-	1	2	3	6	8	8	10
Male	104	1	-	-	-	1	2	3	5	5	2	6
Female	101	-	-	-	-	-	-	-	1	3	6	4
White, both sexes	172	-	-	-	-	1	1	3	5	6	7	8
Male	90	-	-	-	-	1	1	3	4	5	2	5
Female	82	-	-	-	-	-	-	-	1	1	5	3
Black, both sexes	32	1	-	-	-	-	1	-	1	2	1	2
Male	14	1	-	-	-	-	1	-	1	-	-	1
Female	18	-	-	-	-	-	-	-	-	2	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Texas												
All races, both sexes	875	2	2	-	1	4	9	6	20	17	19	29
Male	405	2	2	-	1	3	8	3	14	11	8	17
Female	470	-	-	-	-	1	1	3	6	6	11	12
White, both sexes	770	2	1	-	1	3	8	3	17	12	12	26
Male	355	2	1	-	1	2	8	2	14	8	6	14
Female	415	-	-	-	-	1	-	1	3	4	6	12
Black, both sexes	100	-	1	-	-	1	1	3	3	5	7	3
Male	48	-	1	-	-	1	-	1	-	3	2	3
Female	52	-	-	-	-	-	1	2	3	2	5	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	159	-	-	-	-	-	1	-	-	1	2	3
Male	91	-	-	-	-	-	1	-	-	1	2	2
Female	68	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	149	-	-	-	-	-	-	-	-	1	1	3
Male	83	-	-	-	-	-	-	-	-	1	1	2
Female	66	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Vermont												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	220	-	-	-	-	3	3	-	-	2	6	10
Male	118	-	-	-	-	3	3	-	-	2	3	7
Female	102	-	-	-	-	-	-	-	-	-	3	3
White, both sexes	171	-	-	-	-	2	1	-	-	1	3	5
Male	93	-	-	-	-	2	1	-	-	1	1	4
Female	78	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	48	-	-	-	-	1	2	-	-	1	3	5
Male	25	-	-	-	-	1	2	-	-	1	2	3
Female	23	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Washington												
All races, both sexes	113	-	-	-	-	-	-	1	-	3	-	2
Male	49	-	-	-	-	-	-	1	-	1	-	2
Female	64	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	107	-	-	-	-	-	-	-	-	3	-	2
Male	46	-	-	-	-	-	-	-	-	1	-	2
Female	61	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Wisconsin												
All races, both sexes	152	-	-	-	-	-	1	2	-	2	6	4
Male	84	-	-	-	-	-	1	1	-	1	5	3
Female	68	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	148	-	-	-	-	-	1	2	-	1	6	4
Male	84	-	-	-	-	-	1	1	-	1	5	3
Female	64	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	41	-	-	-	-	-	-	-	-	1	2	3
Male	21	-	-	-	-	-	-	-	-	1	1	2
Female	20	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	40	-	-	-	-	-	-	-	-	1	1	3
Male	21	-	-	-	-	-	-	-	-	1	1	2
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
United States												
All races, both sexes	476	573	610	632	820	1,120	1,632	1,765	1,307	581	120	-
Male	335	381	419	415	488	601	752	697	387	134	24	-
Female	141	192	191	217	332	519	880	1,068	920	447	96	-
White, both sexes	352	440	485	521	682	967	1,446	1,606	1,213	530	99	-
Male	251	297	334	338	410	518	676	639	363	121	19	-
Female	101	143	151	183	272	449	770	967	850	409	80	-
Black, both sexes	114	122	109	93	118	139	173	125	84	42	20	-
Male	76	76	73	68	69	77	72	42	22	10	4	-
Female	38	46	36	25	49	62	101	83	62	32	16	-
American Indian 1/, both sexes	5	1	8	5	3	6	2	3	1	2	-	-
Male	4	1	6	4	1	3	1	1	1	-	-	-
Female	1	-	2	1	2	3	1	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes	5	10	8	13	17	8	11	31	9	7	1	-
Male	4	7	6	5	8	3	3	15	1	3	1	-
Female	1	3	2	8	9	5	8	16	8	4	-	-
Alabama												
All races, both sexes	18	40	31	36	40	49	50	54	23	11	3	-
Male	12	25	23	24	23	22	22	23	7	2	1	-
Female	6	15	8	12	17	27	28	31	16	9	2	-
White, both sexes	13	36	26	31	34	39	45	42	16	9	1	-
Male	8	23	18	20	17	16	19	17	4	1	1	-
Female	5	13	8	11	17	23	26	25	12	8	-	-
Black, both sexes	4	4	5	5	6	10	5	12	7	2	2	-
Male	3	2	5	4	6	6	3	6	3	1	-	-
Female	1	2	-	1	-	4	2	6	4	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Alaska												
All races, both sexes	4	-	6	3	5	1	3	4	1	1	-	-
Male	4	-	6	1	2	1	-	3	1	-	-	-
Female	-	-	-	2	3	-	3	1	-	1	-	-
White, both sexes	4	-	4	2	3	-	2	2	1	1	-	-
Male	4	-	4	1	2	-	-	2	1	-	-	-
Female	-	-	-	1	1	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	1	-	1	2	-	-	-	-
Male	-	-	2	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	10	17	30	24	25	30	26	26	10	2	-
Male	8	5	12	20	18	16	16	13	9	1	1	-
Female	1	5	5	10	6	9	14	13	17	9	1	-
White, both sexes	9	9	15	27	23	22	29	25	25	10	1	-
Male	8	4	10	18	17	14	15	12	9	1	-	-
Female	1	5	5	9	6	8	14	13	16	9	1	-
Black, both sexes	-	1	-	3	-	2	1	-	1	-	-	-
Male	-	1	-	2	-	2	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Arkansas												
All races, both sexes	1	11	14	7	19	22	36	43	28	17	6	-
Male	1	7	8	6	13	8	15	17	6	2	2	-
Female	-	4	6	1	6	14	21	26	22	15	4	-
White, both sexes	1	8	10	5	17	21	33	39	25	15	6	-
Male	1	4	6	4	11	8	14	16	6	2	2	-
Female	-	4	4	1	6	13	19	23	19	13	4	-
Black, both sexes	-	3	4	2	2	1	3	4	3	2	-	-
Male	-	3	2	2	2	-	1	1	-	-	-	-
Female	-	-	2	-	-	1	2	3	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	54	44	28	40	48	77	143	189	165	78	11	-
Male	40	25	23	22	27	39	65	73	45	16	1	-
Female	14	19	5	18	21	38	78	116	120	62	10	-
White, both sexes	45	37	22	32	35	61	131	170	157	71	9	-
Male	35	23	18	18	22	30	62	63	44	14	1	-
Female	10	14	4	14	13	31	69	107	113	57	8	-
Black, both sexes	8	4	4	3	9	11	6	10	4	3	2	-
Male	4	1	4	2	5	6	2	5	1	1	-	-
Female	4	3	-	1	4	5	4	5	3	2	2	-
American Indian 1/, both sexes	-	-	-	1	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	3	2	4	4	3	6	9	4	3	-	-
Male	1	1	1	2	-	1	1	5	-	1	-	-
Female	-	2	1	2	4	2	5	4	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Colorado												
All races, both sexes	4	3	3	5	5	10	12	13	9	3	-	-
Male	2	3	2	4	1	5	7	2	1	-	-	-
Female	2	-	1	1	4	5	5	11	8	3	-	-
White, both sexes	4	3	1	4	3	9	12	13	9	3	-	-
Male	2	3	-	4	1	5	7	2	1	-	-	-
Female	2	-	1	-	2	4	5	11	8	3	-	-
Black, both sexes	-	-	1	-	2	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	5	3	4	4	8	13	22	6	7	-	-
Male	-	3	2	3	1	4	3	14	1	2	-	-
Female	-	2	1	1	3	4	10	8	5	5	-	-
White, both sexes	-	5	3	4	4	8	12	21	6	7	-	-
Male	-	3	2	3	1	4	2	13	1	2	-	-
Female	-	2	1	1	3	4	10	8	5	5	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Delaware												
All races, both sexes	3	-	5	-	2	3	6	1	5	2	-	-
Male	2	-	5	-	1	-	2	-	2	-	-	-
Female	1	-	-	-	1	3	4	1	3	2	-	-
White, both sexes	2	-	4	-	1	3	4	1	5	2	-	-
Male	1	-	4	-	1	-	1	-	2	-	-	-
Female	1	-	-	-	-	3	3	1	3	2	-	-
Black, both sexes	1	-	1	-	1	-	2	-	-	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	1	1	1	3	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	1	-	2	1	-	-
White, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Florida												
All races, both sexes	16	18	20	34	43	91	159	187	144	63	10	-
Male	10	13	15	23	29	49	87	83	50	19	3	-
Female	6	5	5	11	14	42	72	104	94	44	7	-
White, both sexes	11	14	17	31	36	79	143	172	139	63	10	-
Male	7	11	13	20	25	45	78	77	47	19	3	-
Female	4	3	4	11	11	34	65	95	92	44	7	-
Black, both sexes	5	4	3	3	7	12	16	12	5	-	-	-
Male	3	2	2	3	4	4	9	5	3	-	-	-
Female	2	2	1	-	3	8	7	7	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Georgia												
All races, both sexes	51	81	82	62	76	65	99	93	58	19	7	-
Male	31	58	54	40	48	46	52	28	15	6	2	-
Female	20	23	28	22	28	19	47	65	43	13	5	-
White, both sexes	29	48	56	51	58	49	79	73	49	14	6	-
Male	17	40	40	31	39	33	43	24	13	4	2	-
Female	12	8	16	20	19	16	36	49	36	10	4	-
Black, both sexes	21	32	25	11	16	16	20	19	9	5	1	-
Male	13	17	14	9	8	13	9	4	2	2	-	-
Female	8	15	11	2	8	3	11	15	7	3	1	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Hawaii												
All races, both sexes	8	9	1	5	4	3	4	9	3	3	-	-
Male	6	7	1	4	2	-	3	5	1	2	-	-
Female	2	2	-	1	2	3	1	4	2	1	-	-
White, both sexes	5	6	-	2	2	3	2	2	1	-	-	-
Male	4	5	-	2	1	-	1	1	-	-	-	-
Female	1	1	-	-	1	3	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	1	3	2	-	2	7	2	3	-	-
Male	2	2	1	2	1	-	2	4	1	2	-	-
Female	1	1	-	1	1	-	-	3	1	1	-	-
Idaho												
All races, both sexes	3	-	4	1	1	1	4	4	-	1	-	-
Male	2	-	3	1	1	1	-	2	-	-	-	-
Female	1	-	1	-	-	-	4	2	-	1	-	-
White, both sexes	3	-	4	1	1	1	4	4	-	1	-	-
Male	2	-	3	1	1	1	-	2	-	-	-	-
Female	1	-	1	-	-	-	4	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Illinois												
All races, both sexes	21	22	20	20	32	50	73	77	63	31	4	-
Male	15	18	14	12	15	24	34	22	17	9	-	-
Female	6	4	6	8	17	26	39	55	46	22	4	-
White, both sexes	17	16	16	16	25	42	58	72	58	28	3	-
Male	11	12	11	8	12	20	28	22	17	9	-	-
Female	6	4	5	8	13	22	30	50	41	19	3	-
Black, both sexes	4	6	4	4	6	8	15	5	5	3	1	-
Male	4	6	3	4	2	4	6	-	-	-	-	-
Female	-	-	1	-	4	4	9	5	5	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	8	7	7	8	12	17	29	25	17	6	3	-
Male	5	7	7	6	9	8	13	9	3	2	1	-
Female	3	-	-	2	3	9	16	16	14	4	2	-
White, both sexes	7	5	6	8	11	13	28	25	16	6	3	-
Male	5	5	6	6	8	6	13	9	3	2	1	-
Female	2	-	-	2	3	7	15	16	13	4	2	-
Black, both sexes	1	2	1	-	1	3	1	-	1	-	-	-
Male	-	2	1	-	1	2	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Iowa												
All races, both sexes	7	4	2	4	3	4	9	13	12	5	3	-
Male	3	2	2	3	1	1	6	7	5	2	-	-
Female	4	2	-	1	2	3	3	6	7	3	3	-
White, both sexes	5	4	2	4	3	4	8	13	12	5	3	-
Male	2	2	2	3	1	1	6	7	5	2	-	-
Female	3	2	-	1	2	3	2	6	7	3	3	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	12	7	3	5	5	3	10	9	6	1	-
Male	2	11	6	2	2	2	2	3	4	-	-	-
Female	3	1	1	1	3	3	1	7	5	6	1	-
White, both sexes	5	12	6	2	5	5	3	10	9	6	1	-
Male	2	11	5	1	2	2	2	3	4	-	-	-
Female	3	1	1	1	3	3	1	7	5	6	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Kentucky												
All races, both sexes	19	16	23	16	20	26	32	35	21	12	2	-
Male	17	13	19	11	9	12	12	16	8	2	1	-
Female	2	3	4	5	11	14	20	19	13	10	1	-
White, both sexes	17	14	22	16	18	23	30	34	21	11	2	-
Male	16	12	18	11	9	12	10	15	8	2	1	-
Female	1	2	4	5	9	11	20	19	13	9	1	-
Black, both sexes	1	2	-	-	1	3	2	1	-	1	-	-
Male	-	1	-	-	-	-	2	1	-	-	-	-
Female	1	1	-	-	1	3	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	12	13	17	22	21	37	53	50	36	17	1	-
Male	8	7	11	13	11	18	20	19	13	3	1	-
Female	4	6	6	9	10	19	33	31	23	14	-	-
White, both sexes	6	9	10	15	16	28	46	43	31	14	-	-
Male	4	3	6	10	9	14	17	17	13	2	-	-
Female	2	6	4	5	7	14	29	26	18	12	-	-
Black, both sexes	6	4	7	7	5	9	7	7	5	3	1	-
Male	4	4	5	3	2	4	3	2	-	1	1	-
Female	2	-	2	4	3	5	4	5	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Maine												
All races, both sexes	2	4	8	2	4	2	3	4	4	2	1	-
Male	2	3	4	1	2	2	1	4	1	-	-	-
Female	-	1	4	1	2	-	2	-	3	2	1	-
White, both sexes	1	4	8	2	4	2	3	4	4	2	1	-
Male	1	3	4	1	2	2	1	4	1	-	-	-
Female	-	1	4	1	2	-	2	-	3	2	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	5	3	5	3	13	11	13	10	3	1	-
Male	2	3	2	2	3	10	6	5	4	2	-	-
Female	-	2	1	3	-	3	5	8	6	1	1	-
White, both sexes	1	5	1	4	3	9	8	9	9	3	1	-
Male	1	3	-	2	3	6	5	2	4	2	-	-
Female	-	2	1	2	-	3	3	7	5	1	1	-
Black, both sexes	1	-	2	1	-	4	3	3	1	-	-	-
Male	1	-	2	-	-	4	1	3	-	-	-	-
Female	-	-	-	1	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Massachusetts												
All races, both sexes	2	7	2	9	17	19	35	38	18	9	2	-
Male	1	5	2	5	7	10	16	18	3	2	-	-
Female	1	2	-	4	10	9	19	20	15	7	2	-
White, both sexes	2	6	2	9	16	18	33	36	18	9	2	-
Male	1	4	2	5	7	9	16	17	3	2	-	-
Female	1	2	-	4	9	9	17	19	15	7	2	-
Black, both sexes	-	-	-	-	-	1	2	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Michigan												
All races, both sexes	6	14	13	15	23	30	43	47	31	15	2	-
Male	5	10	11	9	13	17	23	20	9	3	-	-
Female	1	4	2	6	10	13	20	27	22	12	2	-
White, both sexes	3	9	9	11	20	29	39	45	31	15	1	-
Male	3	7	8	7	12	17	21	20	9	3	-	-
Female	-	2	1	4	8	12	18	25	22	12	1	-
Black, both sexes	3	5	4	2	3	1	4	2	-	-	1	-
Male	2	3	3	1	1	-	2	-	-	-	-	-
Female	1	2	1	1	2	1	2	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Minnesota												
All races, both sexes	5	12	5	4	7	14	19	24	13	4	3	-
Male	4	8	2	3	3	7	11	11	3	1	2	-
Female	1	4	3	1	4	7	8	13	10	3	1	-
White, both sexes	4	11	4	4	6	14	17	23	13	4	3	-
Male	3	8	1	3	3	7	11	10	3	1	2	-
Female	1	3	3	1	3	7	6	13	10	3	1	-
Black, both sexes	1	1	1	-	1	-	1	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	16	11	24	19	28	31	46	49	32	12	7	-
Male	12	5	19	12	17	13	15	19	8	2	3	-
Female	4	6	5	7	11	18	31	30	24	10	4	-
White, both sexes	7	4	18	10	21	19	34	40	26	8	1	-
Male	5	2	15	6	13	9	14	17	6	2	-	-
Female	2	2	3	4	8	10	20	23	20	6	1	-
Black, both sexes	9	7	6	9	7	12	12	9	6	4	6	-
Male	7	3	4	6	4	4	1	2	2	-	3	-
Female	2	4	2	3	3	8	11	7	4	4	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Missouri												
All races, both sexes	10	13	8	18	12	17	36	32	27	14	5	-
Male	8	8	5	13	8	7	15	8	12	3	-	-
Female	2	5	3	5	4	10	21	24	15	11	5	-
White, both sexes	5	11	8	15	12	15	31	30	25	14	4	-
Male	4	6	5	11	8	6	14	8	10	3	-	-
Female	1	5	3	4	4	9	17	22	15	11	4	-
Black, both sexes	5	2	-	3	-	2	5	2	2	-	1	-
Male	4	2	-	2	-	1	1	-	2	-	-	-
Female	1	-	-	1	-	1	4	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	3	3	2	1	8	8	4	1	1	-
Male	2	-	1	3	2	1	5	4	3	1	-	-
Female	-	-	2	-	-	-	3	4	1	-	1	-
White, both sexes	1	-	2	2	2	1	8	8	4	1	1	-
Male	1	-	1	2	2	1	5	4	3	1	-	-
Female	-	-	1	-	-	-	3	4	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Nebraska												
All races, both sexes	2	5	4	3	8	15	14	21	7	9	2	-
Male	2	4	2	2	5	10	8	7	1	3	-	-
Female	-	1	2	1	3	5	6	14	6	6	2	-
White, both sexes	2	4	4	2	8	15	13	20	7	9	2	-
Male	2	3	2	1	5	10	8	7	1	3	-	-
Female	-	1	2	1	3	5	5	13	6	6	2	-
Black, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	2	4	7	10	7	11	17	19	18	5	1	-
Male	-	2	5	7	4	8	7	3	5	-	-	-
Female	2	2	2	3	3	3	10	16	13	5	1	-
White, both sexes	2	3	5	10	7	9	13	18	16	5	1	-
Male	-	2	4	7	4	7	7	3	4	-	-	-
Female	2	1	1	3	3	2	6	15	12	5	1	-
Black, both sexes	-	1	1	-	-	2	3	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Hampshire												
All races, both sexes	2	3	5	2	7	5	7	12	14	3	-	-
Male	2	2	2	2	5	4	5	5	6	-	-	-
Female	-	1	3	-	2	1	2	7	8	3	-	-
White, both sexes	2	3	5	2	7	5	7	12	14	3	-	-
Male	2	2	2	2	5	4	5	5	6	-	-	-
Female	-	1	3	-	2	1	2	7	8	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	12	7	10	11	33	40	48	33	15	1	-
Male	2	8	6	5	8	18	14	19	6	7	-	-
Female	2	4	1	5	3	15	26	29	27	8	1	-
White, both sexes	2	10	5	7	10	31	37	45	30	14	1	-
Male	2	6	4	3	7	16	12	18	5	6	-	-
Female	-	4	1	4	3	15	25	27	25	8	1	-
Black, both sexes	2	2	2	2	1	2	3	2	2	1	-	-
Male	-	2	2	1	1	2	2	1	1	1	-	-
Female	2	-	-	1	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Mexico												
All races, both sexes	1	1	5	2	1	7	9	11	7	6	1	-
Male	-	1	3	1	-	5	7	5	2	2	-	-
Female	1	-	2	1	1	2	2	6	5	4	1	-
White, both sexes	1	1	5	2	1	7	8	10	7	6	1	-
Male	-	1	3	1	-	5	6	5	2	2	-	-
Female	1	-	2	1	1	2	2	5	5	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	20	19	23	22	36	47	77	69	44	26	3	-
Male	16	12	10	17	21	27	38	29	15	9	-	-
Female	4	7	13	5	15	20	39	40	29	17	3	-
White, both sexes	16	13	21	19	30	42	70	65	41	24	3	-
Male	13	10	9	15	16	23	36	26	15	9	-	-
Female	3	3	12	4	14	19	34	39	26	15	3	-
Black, both sexes	4	6	1	2	5	4	7	3	3	2	-	-
Male	3	2	-	2	4	3	2	2	-	-	-	-
Female	1	4	1	-	1	1	5	1	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	1	-	-	-	-
Male	-	-	1	-	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
North Carolina												
All races, both sexes	18	13	22	19	30	35	48	37	44	9	1	-
Male	15	8	12	15	14	18	14	12	10	3	-	-
Female	3	5	10	4	16	17	34	25	34	6	1	-
White, both sexes	10	9	15	14	23	31	35	29	35	7	-	-
Male	8	5	9	11	8	17	10	10	9	1	-	-
Female	2	4	6	3	15	14	25	19	26	6	-	-
Black, both sexes	7	4	6	5	6	3	13	8	9	2	1	-
Male	6	3	2	4	5	1	4	2	1	2	-	-
Female	1	1	4	1	1	2	9	6	8	-	1	-
American Indian 1/, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	2	1	1	-	2	1	3	-	-	-
Male	-	-	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	2	-	3	-	-	-
White, both sexes	-	-	2	1	1	-	2	1	3	-	-	-
Male	-	-	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	2	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Ohio												
All races, both sexes	16	19	18	22	32	30	38	40	30	12	2	-
Male	8	10	10	16	19	15	11	18	10	3	-	-
Female	8	9	8	6	13	15	27	22	20	9	2	-
White, both sexes	13	18	16	19	24	28	35	38	28	12	1	-
Male	7	10	9	14	13	15	11	18	9	3	-	-
Female	6	8	7	5	11	13	24	20	19	9	1	-
Black, both sexes	3	1	2	3	8	2	3	2	2	-	1	-
Male	1	-	1	2	6	-	-	-	1	-	-	-
Female	2	1	1	1	2	2	3	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	10	3	5	15	16	18	25	7	3	-
Male	-	-	6	1	3	7	10	6	5	1	-	-
Female	1	-	4	2	2	8	6	12	20	6	3	-
White, both sexes	-	-	10	3	4	13	16	18	24	7	3	-
Male	-	-	6	1	3	6	10	6	5	1	-	-
Female	-	-	4	2	1	7	6	12	19	6	3	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Oregon												
All races, both sexes	4	6	2	6	3	10	13	6	19	3	2	-
Male	3	4	2	3	3	6	7	2	8	-	-	-
Female	1	2	-	3	-	4	6	4	11	3	2	-
White, both sexes	4	5	1	5	3	10	13	6	19	3	2	-
Male	3	3	1	2	3	6	7	2	8	-	-	-
Female	1	2	-	3	-	4	6	4	11	3	2	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	19	18	33	25	34	53	63	81	58	23	6	-
Male	13	11	26	17	23	27	28	27	18	4	1	-
Female	6	7	7	8	11	26	35	54	40	19	5	-
White, both sexes	14	16	24	23	32	47	58	80	57	23	6	-
Male	11	11	19	15	21	22	25	27	18	4	1	-
Female	3	5	5	8	11	25	33	53	39	19	5	-
Black, both sexes	5	2	9	2	2	6	5	1	1	-	-	-
Male	2	-	7	2	2	5	3	-	-	-	-	-
Female	3	2	2	-	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Rhode Island												
All races, both sexes	-	-	-	-	3	3	-	2	3	1	-	-
Male	-	-	-	-	3	3	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
White, both sexes	-	-	-	-	2	3	-	2	3	1	-	-
Male	-	-	-	-	2	3	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	20	30	31	29	23	30	32	23	12	7	2	-
Male	16	24	22	21	13	20	17	7	2	2	-	-
Female	4	6	9	8	10	10	15	16	10	5	2	-
White, both sexes	11	19	21	15	15	22	25	16	9	5	1	-
Male	9	14	14	10	7	12	11	6	1	1	-	-
Female	2	5	7	5	8	10	14	10	8	4	1	-
Black, both sexes	9	10	10	14	8	8	7	7	3	2	1	-
Male	7	9	8	11	6	8	6	1	1	1	-	-
Female	2	1	2	3	2	-	1	6	2	1	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
South Dakota												
All races, both sexes	1	-	1	3	2	4	2	5	8	-	2	-
Male	1	-	1	2	1	1	1	3	2	-	1	-
Female	-	-	-	1	1	3	1	2	6	-	1	-
White, both sexes	1	-	1	2	2	4	2	5	8	-	2	-
Male	1	-	1	1	1	1	1	3	2	-	1	-
Female	-	-	-	1	1	3	1	2	6	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10	9	12	10	22	22	35	20	20	5	1	-
Male	8	3	6	6	17	10	16	6	6	1	-	-
Female	2	6	6	4	5	12	19	14	14	4	1	-
White, both sexes	9	6	11	6	19	20	28	19	18	4	1	-
Male	7	2	6	4	14	10	14	5	6	1	-	-
Female	2	4	5	2	5	10	14	14	12	3	1	-
Black, both sexes	1	3	1	4	3	2	7	1	1	1	-	-
Male	1	1	-	2	3	-	2	1	-	-	-	-
Female	-	2	1	2	-	2	5	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Texas												
All races, both sexes	34	39	36	48	61	87	125	151	109	65	11	-
Male	24	23	22	28	35	48	52	61	34	6	3	-
Female	10	16	14	20	26	39	73	90	75	59	8	-
White, both sexes	28	28	31	43	54	79	112	142	102	56	10	-
Male	19	16	19	24	32	45	45	57	31	6	3	-
Female	9	12	12	19	22	34	67	85	71	50	7	-
Black, both sexes	6	10	4	4	7	8	12	9	7	8	1	-
Male	5	6	2	4	3	3	7	4	3	-	-	-
Female	1	4	2	-	4	5	5	5	4	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Utah												
All races, both sexes	5	7	3	12	17	14	38	35	13	8	-	-
Male	3	4	3	9	9	9	25	17	3	3	-	-
Female	2	3	-	3	8	5	13	18	10	5	-	-
White, both sexes	5	7	3	12	14	14	37	31	13	8	-	-
Male	3	4	3	9	6	9	24	15	3	3	-	-
Female	2	3	-	3	8	5	13	16	10	5	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	4	-	-	-	-
Male	-	-	-	-	2	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Vermont												
All races, both sexes	1	-	2	1	1	1	3	2	3	1	1	-
Male	-	-	2	1	1	-	3	2	1	1	-	-
Female	1	-	-	-	-	1	-	-	2	-	1	-
White, both sexes	1	-	2	1	1	1	3	2	3	1	1	-
Male	-	-	2	1	1	-	3	2	1	1	-	-
Female	1	-	-	-	-	1	-	-	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	9	9	11	14	24	28	35	26	29	8	3	-
Male	6	7	5	9	15	18	17	10	9	3	1	-
Female	3	2	6	5	9	10	18	16	20	5	2	-
White, both sexes	5	4	7	13	17	22	33	24	26	6	2	-
Male	4	2	4	9	14	16	15	10	7	2	1	-
Female	1	2	3	4	3	6	18	14	19	4	1	-
Black, both sexes	4	5	4	1	6	6	2	2	3	2	1	-
Male	2	5	1	-	1	2	2	-	2	1	-	-
Female	2	-	3	1	5	4	-	2	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Washington												
All races, both sexes	5	8	4	3	6	11	24	27	9	8	2	-
Male	4	5	2	1	6	5	7	10	4	1	-	-
Female	1	3	2	2	-	6	17	17	5	7	2	-
White, both sexes	5	8	4	3	5	9	24	26	9	7	2	-
Male	4	5	2	1	5	4	7	10	4	1	-	-
Female	1	3	2	2	-	5	17	16	5	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	1	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
West Virginia												
All races, both sexes	1	1	5	3	9	11	11	5	8	2	-	-
Male	-	1	2	3	3	4	3	-	1	1	-	-
Female	1	-	3	-	6	7	8	5	7	1	-	-
White, both sexes	1	1	5	3	8	11	11	5	7	2	-	-
Male	-	1	2	3	2	4	3	-	1	1	-	-
Female	1	-	3	-	6	7	8	5	6	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Wisconsin												
All races, both sexes	11	2	8	6	11	22	20	32	20	4	1	-
Male	8	1	6	4	9	13	10	16	5	1	-	-
Female	3	1	2	2	2	9	10	16	15	3	1	-
White, both sexes	11	2	8	5	10	22	19	32	20	4	1	-
Male	8	1	6	4	9	13	10	16	5	1	-	-
Female	3	1	2	1	1	9	9	16	15	3	1	-
Black, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	7	3	3	6	4	3	3	3	1	-	-
Male	-	5	3	1	3	2	1	1	1	-	-	-
Female	2	2	-	2	3	2	2	2	2	1	-	-
White, both sexes	2	7	3	3	6	4	3	3	3	1	-	-
Male	-	5	3	1	3	2	1	1	1	-	-	-
Female	2	2	-	2	3	2	2	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
United States												
All races, both sexes	137,119	142	54	45	50	73	137	183	344	717	1,504	2,557
Male	54,524	83	26	26	30	36	79	106	184	394	769	1,353
Female	82,595	59	28	19	20	37	58	77	160	323	735	1,204
White, both sexes	115,864	89	39	30	37	59	100	124	221	436	953	1,670
Male	45,198	48	16	19	27	31	60	72	123	239	510	900
Female	70,666	41	23	11	10	28	40	52	98	197	443	770
Black, both sexes	17,045	49	12	12	9	13	31	51	100	233	464	766
Male	7,424	33	8	5	2	5	15	28	47	129	213	380
Female	9,621	16	4	7	7	8	16	23	53	104	251	386
American Indian 1/, both sexes	548	2	2	2	-	1	2	-	5	5	26	22
Male	231	1	2	1	-	-	1	-	3	2	19	14
Female	317	1	-	1	-	1	1	-	2	3	7	8
Asian or Pacific Islander 2/, both sexes	3,662	2	1	1	4	-	4	8	18	43	61	99
Male	1,671	1	-	1	1	-	3	6	11	24	27	59
Female	1,991	1	1	-	3	-	1	2	7	19	34	40
Alabama												
All races, both sexes	2,740	3	-	1	1	-	-	4	10	16	43	60
Male	1,114	2	-	1	1	-	-	2	5	9	19	31
Female	1,626	1	-	-	-	-	-	2	5	7	24	29
White, both sexes	1,949	1	-	-	1	-	-	2	5	6	19	31
Male	752	1	-	-	1	-	-	-	3	4	8	14
Female	1,197	-	-	-	-	-	-	2	2	2	11	17
Black, both sexes	779	1	-	1	-	-	-	2	5	10	24	29
Male	357	1	-	1	-	-	-	2	2	5	11	17
Female	422	-	-	-	-	-	-	-	3	5	13	12
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Alaska												
All races, both sexes	177	-	-	-	-	1	1	-	-	1	4	3
Male	80	-	-	-	-	-	1	-	-	-	4	1
Female	97	-	-	-	-	1	-	-	-	1	-	2
White, both sexes	124	-	-	-	-	1	-	-	-	-	2	2
Male	57	-	-	-	-	-	-	-	-	-	2	-
Female	67	-	-	-	-	1	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	35	-	-	-	-	-	1	-	-	-	1	1
Male	16	-	-	-	-	-	1	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,226	2	1	2	-	1	5	7	7	21	29	35
Male	904	1	1	-	-	1	2	1	4	16	18	18
Female	1,322	1	-	2	-	-	3	6	3	5	11	17
White, both sexes	2,095	2	-	1	-	1	5	5	5	17	25	30
Male	838	1	-	-	-	1	2	1	3	13	16	16
Female	1,257	1	-	1	-	-	3	4	2	4	9	14
Black, both sexes	72	-	-	-	-	-	-	2	1	2	-	2
Male	40	-	-	-	-	-	-	-	-	1	-	1
Female	32	-	-	-	-	-	-	2	1	1	-	1
American Indian 1/, both sexes	44	-	1	1	-	-	-	-	1	1	3	3
Male	23	-	1	-	-	-	-	-	1	1	2	1
Female	21	-	-	1	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Arkansas												
All races, both sexes	1,884	3	-	-	-	1	2	1	12	15	14	31
Male	724	-	-	-	-	-	2	1	6	9	9	18
Female	1,160	3	-	-	-	1	-	-	6	6	5	13
White, both sexes	1,550	1	-	-	-	-	1	1	9	6	6	14
Male	588	-	-	-	-	-	1	1	5	3	4	9
Female	962	1	-	-	-	-	-	-	4	3	2	5
Black, both sexes	316	2	-	-	-	-	1	-	2	9	8	16
Male	131	-	-	-	-	-	1	-	-	6	5	9
Female	185	2	-	-	-	-	-	-	2	3	3	7
American Indian 1/, both sexes	12	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	1	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	15,039	16	2	6	8	9	17	23	34	88	191	303
Male	6,101	10	2	4	4	5	11	16	20	57	101	166
Female	8,938	6	-	2	4	4	6	7	14	31	90	137
White, both sexes	12,194	11	1	5	4	9	14	18	23	61	141	222
Male	4,846	6	1	4	3	5	9	13	14	40	75	119
Female	7,348	5	-	1	1	4	5	5	9	21	66	103
Black, both sexes	1,129	5	1	1	2	-	2	2	6	11	27	42
Male	474	4	1	-	-	-	1	1	3	6	14	20
Female	655	1	-	1	2	-	1	1	3	5	13	22
American Indian 1/, both sexes	52	-	-	-	-	-	-	-	-	-	2	2
Male	25	-	-	-	-	-	-	-	-	-	2	2
Female	27	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1,664	-	-	-	2	-	1	3	5	16	21	37
Male	756	-	-	-	1	-	1	2	3	11	10	25
Female	908	-	-	-	1	-	-	1	2	5	11	12

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Colorado												
All races, both sexes	1,532	2	1	1	1	-	3	2	3	5	22	24
Male	559	-	1	1	1	-	1	1	1	2	9	12
Female	973	2	-	-	-	-	2	1	2	3	13	12
White, both sexes	1,445	2	1	1	1	-	3	1	3	3	19	21
Male	514	-	1	1	1	-	1	-	1	-	9	10
Female	931	2	-	-	-	-	2	1	2	3	10	11
Black, both sexes	57	-	-	-	-	-	-	1	-	2	2	2
Male	32	-	-	-	-	-	-	1	-	2	-	2
Female	25	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	1
Connecticut												
All races, both sexes	1,547	-	-	1	-	-	1	2	3	7	11	19
Male	586	-	-	1	-	-	1	-	3	5	7	13
Female	961	-	-	-	-	-	-	2	-	2	4	6
White, both sexes	1,430	-	-	1	-	-	1	2	2	7	7	11
Male	527	-	-	1	-	-	1	-	2	5	4	7
Female	903	-	-	-	-	-	-	2	-	2	3	4
Black, both sexes	100	-	-	-	-	-	-	-	1	-	4	8
Male	51	-	-	-	-	-	-	-	1	-	3	6
Female	49	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Delaware												
All races, both sexes	384	-	1	-	-	-	-	1	3	2	3	5
Male	137	-	1	-	-	-	-	-	1	2	1	2
Female	247	-	-	-	-	-	-	1	2	-	2	3
White, both sexes	322	-	-	-	-	-	-	1	1	1	3	3
Male	108	-	-	-	-	-	-	-	-	1	1	1
Female	214	-	-	-	-	-	-	1	1	-	2	2
Black, both sexes	62	-	1	-	-	-	-	-	2	1	-	2
Male	29	-	1	-	-	-	-	-	1	1	-	1
Female	33	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	221	-	1	-	1	1	-	-	1	2	4	5
Male	95	-	1	-	1	1	-	-	1	-	2	3
Female	126	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	57	-	-	-	-	-	-	-	-	1	1	-
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	34	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	163	-	1	-	1	1	-	-	1	1	3	5
Male	71	-	1	-	1	1	-	-	1	-	1	3
Female	92	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Florida												
All races, both sexes	8,925	10	2	4	3	4	8	13	25	38	75	159
Male	3,772	5	1	2	2	4	2	9	16	14	30	83
Female	5,153	5	1	2	1	-	6	4	9	24	45	76
White, both sexes	7,571	5	2	3	3	2	4	7	20	27	45	97
Male	3,210	3	1	2	2	2	2	4	12	10	19	55
Female	4,361	2	1	1	1	-	2	3	8	17	26	42
Black, both sexes	1,259	5	-	1	-	2	4	5	4	10	27	58
Male	516	2	-	-	-	2	-	4	3	4	9	24
Female	743	3	-	1	-	-	4	1	1	6	18	34
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	83	-	-	-	-	-	-	1	1	1	3	3
Male	40	-	-	-	-	-	-	1	1	-	2	3
Female	43	-	-	-	-	-	-	-	-	1	1	-
Georgia												
All races, both sexes	3,889	6	2	-	2	2	4	10	14	39	81	112
Male	1,601	3	1	-	2	2	2	5	6	18	36	52
Female	2,288	3	1	-	-	-	2	5	8	21	45	60
White, both sexes	2,677	2	2	-	1	2	3	6	3	16	38	57
Male	1,059	1	1	-	1	2	2	4	1	6	23	31
Female	1,618	1	1	-	-	-	1	2	2	10	15	26
Black, both sexes	1,156	4	-	-	1	-	1	4	11	22	40	54
Male	520	2	-	-	1	-	-	1	5	12	12	21
Female	636	2	-	-	-	-	1	3	6	10	28	33
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	-	-	-	1	2	1
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	1	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Hawaii												
All races, both sexes	665	-	-	-	-	-	1	-	3	4	8	15
Male	307	-	-	-	-	-	-	-	-	1	2	8
Female	358	-	-	-	-	-	1	-	3	3	6	7
White, both sexes	144	-	-	-	-	-	-	-	-	1	3	1
Male	62	-	-	-	-	-	-	-	-	-	1	1
Female	82	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	518	-	-	-	-	-	1	-	3	3	5	14
Male	244	-	-	-	-	-	-	-	-	1	1	7
Female	274	-	-	-	-	-	1	-	3	2	4	7
Idaho												
All races, both sexes	725	-	-	-	-	-	2	-	1	3	5	6
Male	307	-	-	-	-	-	2	-	1	1	3	2
Female	418	-	-	-	-	-	-	-	-	2	2	4
White, both sexes	711	-	-	-	-	-	2	-	1	3	5	6
Male	300	-	-	-	-	-	2	-	1	1	3	2
Female	411	-	-	-	-	-	-	-	-	2	2	4
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Illinois												
All races, both sexes	5,989	7	4	1	5	1	4	6	8	18	57	109
Male	2,329	6	1	-	4	1	-	3	5	13	27	64
Female	3,660	1	3	1	1	-	4	3	3	5	30	45
White, both sexes	5,047	4	2	1	4	1	3	2	5	7	38	67
Male	1,920	4	1	-	4	1	-	1	3	5	20	43
Female	3,127	-	1	1	-	-	3	1	2	2	18	24
Black, both sexes	850	3	2	-	1	-	1	4	3	11	17	40
Male	371	2	-	-	-	-	-	2	2	8	7	20
Female	479	1	2	-	1	-	1	2	1	3	10	20
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	91	-	-	-	-	-	-	-	-	-	1	2
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	3,238	3	-	1	1	-	4	8	13	14	38	51
Male	1,269	2	-	1	-	-	3	5	6	8	16	31
Female	1,969	1	-	-	1	-	1	3	7	6	22	20
White, both sexes	2,975	2	-	1	1	-	2	8	10	8	32	41
Male	1,145	1	-	1	-	-	2	5	5	4	14	23
Female	1,830	1	-	-	1	-	-	3	5	4	18	18
Black, both sexes	255	1	-	-	-	-	2	-	3	5	6	10
Male	120	1	-	-	-	-	1	-	1	3	2	8
Female	135	-	-	-	-	-	1	-	2	2	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Iowa												
All races, both sexes	1,718	1	-	-	-	-	-	2	2	3	6	17
Male	630	-	-	-	-	-	-	2	1	-	4	6
Female	1,088	1	-	-	-	-	-	-	1	3	2	11
White, both sexes	1,692	1	-	-	-	-	-	2	2	2	6	14
Male	619	-	-	-	-	-	-	2	1	-	4	5
Female	1,073	1	-	-	-	-	-	-	1	2	2	9
Black, both sexes	21	-	-	-	-	-	-	-	-	1	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,489	1	1	-	1	-	2	1	3	8	15	24
Male	572	1	-	-	-	-	-	-	1	6	10	15
Female	917	-	1	-	1	-	2	1	2	2	5	9
White, both sexes	1,394	-	1	-	-	-	2	-	3	6	13	16
Male	529	-	-	-	-	-	-	-	1	4	8	11
Female	865	-	1	-	-	-	2	-	2	2	5	5
Black, both sexes	76	1	-	-	1	-	-	-	-	2	2	5
Male	34	1	-	-	-	-	-	-	-	2	2	3
Female	42	-	-	-	1	-	-	-	-	-	-	2
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Kentucky												
All races, both sexes	2,197	-	2	1	1	1	1	4	8	12	21	47
Male	885	-	1	-	-	-	1	3	5	6	14	26
Female	1,312	-	1	1	1	1	-	1	3	6	7	21
White, both sexes	2,042	-	2	1	1	1	1	2	8	8	18	38
Male	817	-	1	-	-	-	1	1	5	5	11	23
Female	1,225	-	1	1	1	1	-	1	3	3	7	15
Black, both sexes	147	-	-	-	-	-	-	2	-	3	3	9
Male	66	-	-	-	-	-	-	2	-	1	3	3
Female	81	-	-	-	-	-	-	-	-	2	-	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
Louisiana												
All races, both sexes	2,195	1	2	1	1	3	2	4	10	15	29	61
Male	876	1	1	-	-	1	-	2	4	7	12	31
Female	1,319	-	1	1	1	2	2	2	6	8	17	30
White, both sexes	1,462	-	1	-	-	3	1	2	6	4	7	25
Male	555	-	1	-	-	1	-	1	3	2	4	14
Female	907	-	-	-	-	2	1	1	3	2	3	11
Black, both sexes	719	1	1	1	1	-	1	2	4	11	21	35
Male	313	1	-	-	-	-	-	1	1	5	8	16
Female	406	-	1	1	1	-	1	1	3	6	13	19
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Maine												
All races, both sexes	670	-	1	1	-	-	2	2	2	2	5	7
Male	252	-	1	1	-	-	1	-	-	2	3	5
Female	418	-	-	-	-	-	1	2	2	-	2	2
White, both sexes	669	-	1	1	-	-	2	2	2	2	5	7
Male	252	-	1	1	-	-	1	-	-	2	3	5
Female	417	-	-	-	-	-	1	2	2	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,365	5	2	1	2	1	3	3	4	25	23	53
Male	950	2	-	-	1	1	3	2	2	8	8	31
Female	1,415	3	2	1	1	-	-	1	2	17	15	22
White, both sexes	1,710	4	1	-	1	-	1	1	3	10	11	22
Male	651	2	-	-	1	-	1	1	2	3	4	15
Female	1,059	2	1	-	-	-	-	-	1	7	7	7
Black, both sexes	597	1	1	1	1	1	2	2	1	15	12	31
Male	273	-	-	-	-	1	2	1	-	5	4	16
Female	324	1	1	1	1	-	-	1	1	10	8	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	56	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Massachusetts												
All races, both sexes	2,880	-	-	1	-	1	3	-	5	8	19	30
Male	1,060	-	-	1	-	-	2	-	3	4	8	18
Female	1,820	-	-	-	-	1	1	-	2	4	11	12
White, both sexes	2,690	-	-	1	-	1	2	-	3	3	15	22
Male	984	-	-	1	-	-	1	-	2	1	7	14
Female	1,706	-	-	-	-	1	1	-	1	2	8	8
Black, both sexes	130	-	-	-	-	-	1	-	2	5	4	7
Male	53	-	-	-	-	-	1	-	1	3	1	3
Female	77	-	-	-	-	-	-	-	1	2	3	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4,752	2	-	1	-	5	6	5	14	19	40	66
Male	1,878	1	-	-	-	4	4	1	7	8	17	26
Female	2,874	1	-	1	-	1	2	4	7	11	23	40
White, both sexes	4,066	2	-	1	-	4	2	3	7	11	25	45
Male	1,588	1	-	-	-	4	2	1	5	5	10	19
Female	2,478	1	-	1	-	-	-	2	2	6	15	26
Black, both sexes	641	-	-	-	-	1	3	2	6	7	14	20
Male	274	-	-	-	-	-	1	-	2	3	7	6
Female	367	-	-	-	-	1	2	2	4	4	7	14
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	1	-	1	1	1	1
Male	9	-	-	-	-	-	1	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Minnesota												
All races, both sexes	2,219	3	1	-	-	2	5	3	2	10	17	30
Male	861	2	-	-	-	1	3	1	-	3	11	16
Female	1,358	1	1	-	-	1	2	2	2	7	6	14
White, both sexes	2,133	2	-	-	-	2	3	2	2	8	15	23
Male	825	1	-	-	-	1	3	1	-	2	9	14
Female	1,308	1	-	-	-	1	-	1	2	6	6	9
Black, both sexes	32	1	-	-	-	-	1	1	-	-	2	2
Male	11	1	-	-	-	-	-	-	-	-	2	1
Female	21	-	-	-	-	-	1	1	-	-	-	1
American Indian 1/, both sexes	23	-	-	-	-	-	1	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	31	-	1	-	-	-	-	-	-	1	-	5
Male	17	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	1	-	-	-	-	-	-	-	-	4
Mississippi												
All races, both sexes	1,585	2	-	1	-	3	1	5	3	8	29	60
Male	650	2	-	-	-	2	1	3	2	5	14	33
Female	935	-	-	1	-	1	-	2	1	3	15	27
White, both sexes	1,018	1	-	-	-	2	-	2	3	3	6	21
Male	410	1	-	-	-	2	-	2	2	2	4	13
Female	608	-	-	-	-	-	-	-	1	1	2	8
Black, both sexes	564	1	-	1	-	1	1	3	-	5	23	38
Male	239	1	-	-	-	-	1	1	-	3	10	19
Female	325	-	-	1	-	1	-	2	-	2	13	19
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Missouri												
All races, both sexes	3,247	4	1	2	1	-	2	2	6	17	34	58
Male	1,225	3	-	1	-	-	2	1	4	9	16	26
Female	2,022	1	1	1	1	-	-	1	2	8	18	32
White, both sexes	2,892	2	1	2	1	-	2	2	4	9	26	42
Male	1,075	2	-	1	-	-	2	1	2	4	13	20
Female	1,817	-	1	1	1	-	-	1	2	5	13	22
Black, both sexes	334	2	-	-	-	-	-	-	2	7	7	14
Male	142	1	-	-	-	-	-	-	2	5	3	4
Female	192	1	-	-	-	-	-	-	-	2	4	10
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	1	1	-
Montana												
All races, both sexes	461	-	-	-	-	-	-	-	-	-	4	5
Male	182	-	-	-	-	-	-	-	-	-	3	2
Female	279	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	440	-	-	-	-	-	-	-	-	-	2	4
Male	173	-	-	-	-	-	-	-	-	-	1	1
Female	267	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	2	1
Male	8	-	-	-	-	-	-	-	-	-	2	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Nebraska												
All races, both sexes	922	-	2	-	-	1	-	-	-	5	15	15
Male	359	-	2	-	-	1	-	-	-	4	8	8
Female	563	-	-	-	-	-	-	-	-	1	7	7
White, both sexes	883	-	1	-	-	1	-	-	-	4	13	8
Male	340	-	1	-	-	1	-	-	-	3	6	5
Female	543	-	-	-	-	-	-	-	-	1	7	3
Black, both sexes	30	-	1	-	-	-	-	-	-	1	-	5
Male	15	-	1	-	-	-	-	-	-	1	-	2
Female	15	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	2	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Nevada												
All races, both sexes	847	2	-	-	1	1	1	1	5	10	16	27
Male	374	-	-	-	-	1	1	1	3	5	8	15
Female	473	2	-	-	1	-	-	-	2	5	8	12
White, both sexes	734	2	-	-	-	1	1	1	2	8	14	19
Male	322	-	-	-	-	1	1	1	1	5	8	11
Female	412	2	-	-	-	-	-	-	1	3	6	8
Black, both sexes	63	-	-	-	-	-	-	-	1	-	1	3
Male	31	-	-	-	-	-	-	-	-	-	-	2
Female	32	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	1	-	-	-	1	2	1	5
Male	17	-	-	-	-	-	-	-	1	-	-	2
Female	26	-	-	-	1	-	-	-	-	2	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
New Hampshire												
All races, both sexes	494	3	-	-	-	1	1	-	2	1	3	7
Male	193	2	-	-	-	-	-	-	-	1	3	5
Female	301	1	-	-	-	1	1	-	2	-	-	2
White, both sexes	489	3	-	-	-	1	1	-	2	1	3	7
Male	189	2	-	-	-	-	-	-	-	1	3	5
Female	300	1	-	-	-	1	1	-	2	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3,468	3	-	3	2	2	2	9	12	23	39	76
Male	1,426	1	-	1	1	2	-	6	7	15	20	49
Female	2,042	2	-	2	1	-	2	3	5	8	19	27
White, both sexes	2,882	2	-	2	2	1	2	2	9	14	18	53
Male	1,160	-	-	1	1	1	-	2	6	9	11	39
Female	1,722	2	-	1	1	-	2	-	3	5	7	14
Black, both sexes	476	1	-	1	-	1	-	6	3	8	16	20
Male	218	1	-	-	-	1	-	3	1	5	5	7
Female	258	-	-	1	-	-	-	3	2	3	11	13
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	106	-	-	-	-	-	-	1	-	1	5	2
Male	47	-	-	-	-	-	-	1	-	1	4	2
Female	59	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
New Mexico												
All races, both sexes	739	1	1	1	-	1	1	-	3	2	5	19
Male	281	1	-	1	-	-	1	-	2	1	3	5
Female	458	-	1	-	-	1	-	-	1	1	2	14
White, both sexes	687	1	1	-	-	1	1	-	2	2	3	18
Male	258	1	-	-	-	-	1	-	1	1	1	5
Female	429	-	1	-	-	1	-	-	1	1	2	13
Black, both sexes	18	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	-	1	-	-	-	-	-	-	2	1
Male	14	-	-	1	-	-	-	-	-	-	2	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6,398	11	2	2	4	2	5	5	22	28	70	129
Male	2,552	7	1	2	2	2	4	4	11	17	38	64
Female	3,846	4	1	-	2	-	1	1	11	11	32	65
White, both sexes	5,414	5	2	1	4	2	4	3	12	12	34	93
Male	2,116	3	1	1	2	2	4	2	5	9	19	45
Female	3,298	2	1	-	2	-	-	1	7	3	15	48
Black, both sexes	799	5	-	1	-	-	1	1	8	13	35	29
Male	352	4	-	1	-	-	-	1	4	6	18	13
Female	447	1	-	-	-	-	1	-	4	7	17	16
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	182	1	-	-	-	-	-	1	2	3	1	7
Male	84	-	-	-	-	-	-	1	2	2	1	6
Female	98	1	-	-	-	-	-	-	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
North Carolina												
All races, both sexes	4,572	4	2	-	1	5	6	5	8	36	68	84
Male	1,834	4	-	-	1	1	4	4	4	22	37	48
Female	2,738	-	2	-	-	4	2	1	4	14	31	36
White, both sexes	3,438	2	2	-	1	3	5	3	4	13	32	39
Male	1,331	2	-	-	1	1	3	3	2	7	20	18
Female	2,107	-	2	-	-	2	2	-	2	6	12	21
Black, both sexes	1,068	2	-	-	-	2	1	2	4	22	30	44
Male	467	2	-	-	-	-	1	1	2	15	13	29
Female	601	-	-	-	-	2	-	1	2	7	17	15
American Indian 1/, both sexes	42	-	-	-	-	-	-	-	-	1	5	1
Male	22	-	-	-	-	-	-	-	-	-	3	1
Female	20	-	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	428	-	1	1	-	-	-	-	2	1	1	6
Male	177	-	1	-	-	-	-	-	-	-	-	4
Female	251	-	-	1	-	-	-	-	2	1	1	2
White, both sexes	421	-	-	1	-	-	-	-	1	1	1	6
Male	173	-	-	-	-	-	-	-	-	-	-	4
Female	248	-	-	1	-	-	-	-	1	1	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	1	-	-	-	-	-	1	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Ohio												
All races, both sexes	5,828	4	5	3	3	4	3	9	10	18	47	100
Male	2,235	2	2	3	3	1	2	2	5	12	21	49
Female	3,593	2	3	-	-	3	1	7	5	6	26	51
White, both sexes	5,143	3	4	2	3	4	1	8	9	11	35	80
Male	1,931	2	1	2	3	1	-	2	5	8	16	42
Female	3,212	1	3	-	-	3	1	6	4	3	19	38
Black, both sexes	660	1	1	1	-	-	2	1	1	7	12	19
Male	294	-	1	1	-	-	2	-	-	4	5	7
Female	366	1	-	-	-	-	-	1	1	3	7	12
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Oklahoma												
All races, both sexes	2,085	4	2	-	-	1	2	1	3	6	17	34
Male	819	2	1	-	-	-	1	-	2	5	9	19
Female	1,266	2	1	-	-	1	1	1	1	1	8	15
White, both sexes	1,835	3	1	-	-	1	2	-	1	6	12	29
Male	715	2	-	-	-	-	1	-	1	5	7	16
Female	1,120	1	1	-	-	1	1	-	-	1	5	13
Black, both sexes	153	1	1	-	-	-	-	1	2	-	3	4
Male	67	-	1	-	-	-	-	-	1	-	-	3
Female	86	1	-	-	-	-	-	1	1	-	3	1
American Indian 1/, both sexes	81	-	-	-	-	-	-	-	-	-	2	1
Male	30	-	-	-	-	-	-	-	-	-	2	-
Female	51	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Oregon												
All races, both sexes	1,978	-	-	-	1	-	3	2	2	6	16	22
Male	830	-	-	-	1	-	1	1	1	4	9	11
Female	1,148	-	-	-	-	-	2	1	1	2	7	11
White, both sexes	1,897	-	-	-	1	-	2	2	2	5	13	18
Male	793	-	-	-	1	-	-	1	1	3	7	8
Female	1,104	-	-	-	-	-	2	1	1	2	6	10
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	-	1	-	-	-	2	2
Male	13	-	-	-	-	-	1	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	1	1
Pennsylvania												
All races, both sexes	7,151	5	4	4	1	3	6	8	12	27	68	104
Male	2,722	1	1	2	-	-	3	5	6	18	37	65
Female	4,429	4	3	2	1	3	3	3	6	9	31	39
White, both sexes	6,421	4	4	3	-	2	4	7	11	19	48	64
Male	2,419	-	1	1	-	-	2	4	5	13	28	41
Female	4,002	4	3	2	-	2	2	3	6	6	20	23
Black, both sexes	677	1	-	1	1	1	2	1	1	7	17	38
Male	281	1	-	1	-	-	1	1	1	4	8	23
Female	396	-	-	-	1	1	1	-	-	3	9	15
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	52	-	-	-	-	-	-	-	-	1	3	2
Male	21	-	-	-	-	-	-	-	-	1	1	1
Female	31	-	-	-	-	-	-	-	-	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Rhode Island												
All races, both sexes	421	-	1	-	-	-	-	-	1	1	2	6
Male	163	-	1	-	-	-	-	-	-	1	1	4
Female	258	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	409	-	1	-	-	-	-	-	1	-	2	6
Male	157	-	1	-	-	-	-	-	-	-	1	4
Female	252	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,291	3	-	-	-	-	1	-	7	25	36	70
Male	918	2	-	-	-	-	-	-	3	10	25	34
Female	1,373	1	-	-	-	-	1	-	4	15	11	36
White, both sexes	1,552	-	-	-	-	-	1	-	3	13	14	34
Male	611	-	-	-	-	-	-	-	1	7	8	16
Female	941	-	-	-	-	-	1	-	2	6	6	18
Black, both sexes	717	3	-	-	-	-	-	-	4	11	22	34
Male	298	2	-	-	-	-	-	-	2	3	17	18
Female	419	1	-	-	-	-	-	-	2	8	5	16
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
South Dakota												
All races, both sexes	442	1	-	-	-	-	-	-	2	-	2	7
Male	185	1	-	-	-	-	-	-	2	-	-	3
Female	257	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	422	-	-	-	-	-	-	-	1	-	1	6
Male	174	-	-	-	-	-	-	-	1	-	-	2
Female	248	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	1	-	-	-	-	-	-	1	-	1	1
Male	11	1	-	-	-	-	-	-	1	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3,407	4	1	1	1	-	6	4	11	20	36	78
Male	1,355	2	-	1	1	-	6	3	10	9	16	42
Female	2,052	2	1	-	-	-	-	1	1	11	20	36
White, both sexes	2,829	2	1	1	1	-	4	2	7	17	20	45
Male	1,094	-	-	1	1	-	4	1	7	6	8	24
Female	1,735	2	1	-	-	-	-	1	-	11	12	21
Black, both sexes	567	2	-	-	-	-	2	2	4	3	16	33
Male	254	2	-	-	-	-	2	2	3	3	8	18
Female	313	-	-	-	-	-	-	-	1	-	8	15
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Texas												
All races, both sexes	9,366	16	6	4	3	6	15	20	25	61	137	227
Male	3,782	8	3	3	2	2	8	14	11	32	73	111
Female	5,584	8	3	1	1	4	7	6	14	29	64	116
White, both sexes	7,928	13	5	1	3	4	12	16	14	47	91	157
Male	3,165	6	2	1	2	2	7	11	7	23	53	78
Female	4,763	7	3	-	1	2	5	5	7	24	38	79
Black, both sexes	1,279	2	1	2	-	2	3	4	10	12	39	65
Male	544	1	1	1	-	-	1	3	4	8	18	30
Female	735	1	-	1	-	2	2	1	6	4	21	35
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	1	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	148	1	-	1	-	-	-	-	-	2	7	3
Male	70	1	-	1	-	-	-	-	-	1	2	2
Female	78	-	-	-	-	-	-	-	-	1	5	1
Utah												
All races, both sexes	674	1	-	-	-	1	2	-	2	2	4	6
Male	270	-	-	-	-	1	1	-	2	2	3	4
Female	404	1	-	-	-	-	1	-	-	-	1	2
White, both sexes	650	1	-	-	-	1	2	-	1	2	4	5
Male	268	-	-	-	-	1	1	-	1	2	3	4
Female	382	1	-	-	-	-	1	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Vermont												
All races, both sexes	264	1	-	-	-	-	-	-	1	1	1	2
Male	107	1	-	-	-	-	-	-	1	1	1	1
Female	157	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	264	1	-	-	-	-	-	-	1	1	1	2
Male	107	1	-	-	-	-	-	-	1	1	1	1
Female	157	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,523	4	2	-	2	5	3	8	8	10	40	55
Male	1,362	4	1	-	-	2	3	6	4	5	22	25
Female	2,161	-	1	-	2	3	-	2	4	5	18	30
White, both sexes	2,650	2	1	-	1	4	3	6	2	5	17	26
Male	989	2	-	-	-	2	3	4	1	1	7	12
Female	1,661	-	1	-	1	2	-	2	1	4	10	14
Black, both sexes	796	2	1	-	-	1	-	1	5	4	22	25
Male	333	2	1	-	-	-	-	1	2	3	15	11
Female	463	-	-	-	-	1	-	-	3	1	7	14
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	73	-	-	-	1	-	-	1	1	1	1	4
Male	38	-	-	-	-	-	-	1	1	1	-	2
Female	35	-	-	-	1	-	-	-	-	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Washington												
All races, both sexes	2,725	2	-	-	1	2	1	-	7	13	19	33
Male	1,059	2	-	-	1	-	-	-	5	7	13	20
Female	1,666	-	-	-	-	2	1	-	2	6	6	13
White, both sexes	2,464	2	-	-	1	2	1	-	4	9	11	25
Male	935	2	-	-	1	-	-	-	3	5	7	14
Female	1,529	-	-	-	-	2	1	-	1	4	4	11
Black, both sexes	82	-	-	-	-	-	-	-	1	-	2	5
Male	37	-	-	-	-	-	-	-	1	-	1	4
Female	45	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	2	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	150	-	-	-	-	-	-	-	2	4	4	2
Male	76	-	-	-	-	-	-	-	1	2	3	1
Female	74	-	-	-	-	-	-	-	1	2	1	1
West Virginia												
All races, both sexes	1,072	-	-	-	-	-	-	1	-	4	8	16
Male	413	-	-	-	-	-	-	1	-	2	5	7
Female	659	-	-	-	-	-	-	-	-	2	3	9
White, both sexes	1,040	-	-	-	-	-	-	1	-	4	8	15
Male	398	-	-	-	-	-	-	1	-	2	5	6
Female	642	-	-	-	-	-	-	-	-	2	3	9
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Wisconsin												
All races, both sexes	2,829	2	1	-	2	2	-	2	3	15	24	36
Male	1,108	2	1	-	2	-	-	1	1	7	12	19
Female	1,721	-	-	-	-	2	-	1	2	8	12	17
White, both sexes	2,680	1	1	-	2	2	-	2	2	10	22	30
Male	1,033	1	1	-	2	-	-	1	1	5	12	13
Female	1,647	-	-	-	-	2	-	1	1	5	10	17
Black, both sexes	113	1	-	-	-	-	-	-	1	3	2	6
Male	56	1	-	-	-	-	-	-	-	1	-	6
Female	57	-	-	-	-	-	-	-	1	2	2	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	2	-	-
Male	15	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
Wyoming												
All races, both sexes	236	-	-	-	-	-	-	-	-	2	3	3
Male	94	-	-	-	-	-	-	-	-	1	1	2
Female	142	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	233	-	-	-	-	-	-	-	-	2	3	3
Male	93	-	-	-	-	-	-	-	-	1	1	2
Female	140	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
United States												
All races, both sexes	3,784	4,811	5,707	7,334	10,889	17,739	25,980	27,647	18,996	7,135	1,290	1
Male	2,129	2,757	3,140	3,963	5,400	8,097	10,210	9,252	5,003	1,327	160	-
Female	1,655	2,054	2,567	3,371	5,489	9,642	15,770	18,395	13,993	5,808	1,130	1
White, both sexes	2,465	3,271	4,146	5,546	8,684	15,070	22,914	25,037	17,341	6,504	1,127	1
Male	1,379	1,848	2,262	3,002	4,310	6,946	9,095	8,411	4,552	1,206	142	-
Female	1,086	1,423	1,884	2,544	4,374	8,124	13,819	16,626	12,789	5,298	985	1
Black, both sexes	1,153	1,303	1,310	1,439	1,770	2,085	2,357	1,962	1,269	508	149	-
Male	664	775	748	765	879	894	847	578	306	87	16	-
Female	489	528	562	674	891	1,191	1,510	1,384	963	421	133	-
American Indian 1/, both sexes	30	32	34	49	56	65	82	67	52	11	3	-
Male	19	21	23	24	26	22	26	15	10	2	-	-
Female	11	11	11	25	30	43	56	52	42	9	3	-
Asian or Pacific Islander 2/, both sexes	136	205	217	300	379	519	627	581	334	112	11	-
Male	67	113	107	172	185	235	242	248	135	32	2	-
Female	69	92	110	128	194	284	385	333	199	80	9	-
Alabama												
All races, both sexes	108	129	123	173	268	388	491	514	294	93	21	-
Male	52	82	73	97	139	177	186	156	65	14	3	-
Female	56	47	50	76	129	211	305	358	229	79	18	-
White, both sexes	61	60	80	122	177	279	374	416	227	75	13	-
Male	31	35	43	68	90	131	136	129	46	9	3	-
Female	30	25	37	54	87	148	238	287	181	66	10	-
Black, both sexes	47	67	43	50	88	107	117	97	65	18	8	-
Male	21	45	30	29	47	46	50	27	18	5	-	-
Female	26	22	13	21	41	61	67	70	47	13	8	-
American Indian 1/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	3	-	-	1	2	-	-	-
Male	-	2	-	-	2	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Alaska												
All races, both sexes	10	21	9	13	17	22	34	25	12	3	1	-
Male	4	11	8	8	10	11	9	8	5	-	-	-
Female	6	10	1	5	7	11	25	17	7	3	1	-
White, both sexes	6	16	6	8	11	16	24	18	11	2	1	-
Male	2	9	5	5	9	7	6	7	5	-	-	-
Female	4	7	1	3	2	9	18	11	6	2	1	-
Black, both sexes	1	1	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	3	3	1	3	5	5	6	5	1	-	-	-
Male	2	1	1	1	1	4	2	1	-	-	-	-
Female	1	2	-	2	4	1	4	4	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	2	-	1	4	2	-	-	-	-
Male	-	1	2	2	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	3	2	-	-	-	-
Arizona												
All races, both sexes	55	77	97	111	186	282	441	460	288	102	17	-
Male	34	39	50	62	99	137	169	154	76	21	1	-
Female	21	38	47	49	87	145	272	306	212	81	16	-
White, both sexes	48	70	86	104	175	261	422	445	279	97	17	-
Male	30	35	43	59	92	127	161	147	72	18	1	-
Female	18	35	43	45	83	134	261	298	207	79	16	-
Black, both sexes	6	6	5	4	8	10	10	9	4	3	-	-
Male	3	4	3	2	6	7	4	5	2	2	-	-
Female	3	2	2	2	2	3	6	4	2	1	-	-
American Indian 1/, both sexes	1	1	3	3	2	5	7	6	4	2	-	-
Male	1	-	3	1	1	2	4	2	2	1	-	-
Female	-	1	-	2	1	3	3	4	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	-	1	6	2	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	1	5	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Arkansas												
All races, both sexes	52	65	83	117	164	247	349	383	245	84	16	-
Male	22	36	42	55	70	105	141	126	68	13	1	-
Female	30	29	41	62	94	142	208	257	177	71	15	-
White, both sexes	38	42	64	92	127	207	299	340	217	76	10	-
Male	16	21	33	47	54	93	119	109	61	12	-	-
Female	22	21	31	45	73	114	180	231	156	64	10	-
Black, both sexes	13	22	18	25	36	38	46	42	24	8	6	-
Male	6	14	8	8	16	12	22	16	6	1	1	-
Female	7	8	10	17	20	26	24	26	18	7	5	-
American Indian 1/, both sexes	1	-	1	-	-	1	3	1	3	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	3	-	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
California												
All races, both sexes	436	536	604	794	1,126	1,882	2,819	3,085	2,060	836	164	-
Male	232	304	312	436	551	836	1,120	1,089	627	182	16	-
Female	204	232	292	358	575	1,046	1,699	1,996	1,433	654	148	-
White, both sexes	293	356	418	573	847	1,490	2,383	2,649	1,794	735	147	-
Male	160	199	223	309	420	665	955	929	525	159	13	-
Female	133	157	195	264	427	825	1,428	1,720	1,269	576	134	-
Black, both sexes	81	85	89	97	115	123	144	141	99	44	12	-
Male	43	49	43	52	53	57	54	38	27	7	1	-
Female	38	36	46	45	62	66	90	103	72	37	11	-
American Indian 1/, both sexes	3	6	5	4	5	5	6	11	2	1	-	-
Male	2	4	-	3	2	2	2	5	1	-	-	-
Female	1	2	5	1	3	3	4	6	1	1	-	-
Asian or Pacific Islander 2/, both sexes	59	89	92	120	159	264	286	284	165	56	5	-
Male	27	52	46	72	76	112	109	117	74	16	2	-
Female	32	37	46	48	83	152	177	167	91	40	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Colorado												
All races, both sexes	45	43	55	93	125	182	292	312	223	83	15	-
Male	29	22	21	46	55	72	111	99	64	9	2	-
Female	16	21	34	47	70	110	181	213	159	74	13	-
White, both sexes	40	38	44	84	113	173	282	303	218	80	15	-
Male	24	19	15	40	50	67	106	95	63	9	2	-
Female	16	19	29	44	63	106	176	208	155	71	13	-
Black, both sexes	4	5	8	6	6	5	4	6	4	2	-	-
Male	4	3	5	4	2	3	3	2	1	-	-	-
Female	-	2	3	2	4	2	1	4	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	1	-	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	3	4	4	5	3	-	1	-	-
Male	1	-	1	2	1	2	2	2	-	-	-	-
Female	-	-	2	1	3	2	3	1	-	1	-	-
Connecticut												
All races, both sexes	31	48	42	64	97	182	275	393	227	113	31	-
Male	20	26	27	39	55	80	110	134	50	15	-	-
Female	11	22	15	25	42	102	165	259	177	98	31	-
White, both sexes	23	38	33	51	84	172	258	386	214	111	29	-
Male	16	19	22	31	44	76	104	132	48	15	-	-
Female	7	19	11	20	40	96	154	254	166	96	29	-
Black, both sexes	6	6	8	10	11	10	14	6	12	2	2	-
Male	3	4	5	7	9	4	5	2	2	-	-	-
Female	3	2	3	3	2	6	9	4	10	2	2	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	-	3	1	-	2	1	1	-	-	-
Male	1	3	-	1	1	-	-	-	-	-	-	-
Female	1	1	-	2	-	-	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Delaware												
All races, both sexes	8	10	13	30	38	39	77	87	46	19	2	-
Male	5	5	6	19	16	22	22	22	12	1	-	-
Female	3	5	7	11	22	17	55	65	34	18	2	-
White, both sexes	4	7	10	24	26	35	69	78	41	17	2	-
Male	3	4	3	15	11	21	19	19	10	-	-	-
Female	1	3	7	9	15	14	50	59	31	17	2	-
Black, both sexes	4	3	3	6	12	4	8	9	5	2	-	-
Male	2	1	3	4	5	1	3	3	2	1	-	-
Female	2	2	-	2	7	3	5	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	14	9	13	21	23	17	40	29	26	12	2	-
Male	7	6	4	10	11	10	16	12	8	2	-	-
Female	7	3	9	11	12	7	24	17	18	10	2	-
White, both sexes	-	-	4	4	5	5	9	10	11	6	1	-
Male	-	-	1	1	3	4	2	7	4	-	-	-
Female	-	-	3	3	2	1	7	3	7	6	1	-
Black, both sexes	14	9	9	17	18	12	31	19	15	5	1	-
Male	7	6	3	9	8	6	14	5	4	1	-	-
Female	7	3	6	8	10	6	17	14	11	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Florida												
All races, both sexes	264	300	425	489	744	1,189	1,714	1,755	1,185	439	80	-
Male	157	166	234	268	382	549	717	660	361	92	18	-
Female	107	134	191	221	362	640	997	1,095	824	347	62	-
White, both sexes	170	208	308	366	588	1,028	1,530	1,607	1,084	397	70	-
Male	101	112	176	208	311	487	645	616	341	84	17	-
Female	69	96	132	158	277	541	885	991	743	313	53	-
Black, both sexes	92	89	105	117	139	150	168	137	94	42	10	-
Male	54	52	53	58	65	54	65	39	19	8	1	-
Female	38	37	52	59	74	96	103	98	75	34	9	-
American Indian 1/, both sexes	1	1	1	-	2	2	1	1	2	-	-	-
Male	1	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	11	6	15	9	15	10	5	-	-	-
Male	1	1	4	2	5	7	7	5	1	-	-	-
Female	-	1	7	4	10	2	8	5	4	-	-	-
Georgia												
All races, both sexes	156	190	255	288	342	517	643	654	401	150	21	-
Male	94	109	144	149	175	237	248	202	87	27	2	-
Female	62	81	111	139	167	280	395	452	314	123	19	-
White, both sexes	64	81	144	184	221	348	510	540	324	117	14	-
Male	34	47	72	97	107	157	209	165	76	22	1	-
Female	30	34	72	87	114	191	301	375	248	95	13	-
Black, both sexes	86	105	110	96	115	155	127	111	75	32	7	-
Male	55	60	72	50	66	76	37	34	10	5	1	-
Female	31	45	38	46	49	79	90	77	65	27	6	-
American Indian 1/, both sexes	-	1	-	-	-	1	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	3	1	8	6	13	5	3	1	1	-	-
Male	5	1	-	2	2	4	2	3	1	-	-	-
Female	1	2	1	6	4	9	3	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Hawaii												
All races, both sexes	16	33	30	39	56	74	118	129	105	30	4	-
Male	7	19	20	21	33	38	49	54	45	10	-	-
Female	9	14	10	18	23	36	69	75	60	20	4	-
White, both sexes	-	5	8	7	13	21	28	23	24	9	1	-
Male	-	3	5	4	11	12	7	6	10	2	-	-
Female	-	2	3	3	2	9	21	17	14	7	1	-
Black, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	27	22	31	43	53	90	105	81	21	3	-
Male	7	16	15	16	22	26	42	48	35	8	-	-
Female	9	11	7	15	21	27	48	57	46	13	3	-
Idaho												
All races, both sexes	14	18	16	44	53	102	148	155	114	40	4	-
Male	5	10	11	23	33	48	62	55	39	11	1	-
Female	9	8	5	21	20	54	86	100	75	29	3	-
White, both sexes	14	16	16	42	53	99	144	153	113	40	4	-
Male	5	9	11	22	33	47	59	54	39	11	1	-
Female	9	7	5	20	20	52	85	99	74	29	3	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	2	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	3	1	-	-	-	-
Male	-	-	-	-	-	1	2	1	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Illinois												
All races, both sexes	144	174	229	308	456	756	1,123	1,257	918	336	68	-
Male	71	91	126	173	226	346	445	425	240	50	12	-
Female	73	83	103	135	230	410	678	832	678	286	56	-
White, both sexes	78	121	155	211	356	647	977	1,159	845	307	57	-
Male	36	58	83	120	181	300	391	389	224	45	11	-
Female	42	63	72	91	175	347	586	770	621	262	46	-
Black, both sexes	61	49	69	89	93	100	123	82	66	25	11	-
Male	32	30	42	48	41	44	45	27	16	4	1	-
Female	29	19	27	41	52	56	78	55	50	21	10	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	4	5	8	7	9	23	16	7	4	-	-
Male	3	3	1	5	4	2	9	9	-	1	-	-
Female	2	1	4	3	3	7	14	7	7	3	-	-
Indiana												
All races, both sexes	88	106	138	189	257	437	618	649	429	153	41	-
Male	45	72	77	102	142	186	232	213	100	23	5	-
Female	43	34	61	87	115	251	386	436	329	130	36	-
White, both sexes	69	80	112	158	237	405	592	620	411	149	37	-
Male	33	51	64	87	129	174	222	205	97	23	5	-
Female	36	29	48	71	108	231	370	415	314	126	32	-
Black, both sexes	19	25	26	30	19	31	25	27	18	4	4	-
Male	12	20	13	14	12	12	10	8	3	-	-	-
Female	7	5	13	16	7	19	15	19	15	4	4	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	1	2	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Iowa												
All races, both sexes	29	45	51	58	121	174	315	420	324	123	27	-
Male	16	27	23	28	59	76	145	130	87	21	5	-
Female	13	18	28	30	62	98	170	290	237	102	22	-
White, both sexes	28	42	49	56	115	173	312	417	324	122	27	-
Male	16	27	21	27	56	75	144	128	87	21	5	-
Female	12	15	28	29	59	98	168	289	237	101	22	-
Black, both sexes	1	1	2	2	4	1	3	2	-	1	-	-
Male	-	-	2	1	2	1	1	1	-	-	-	-
Female	1	1	-	1	2	-	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	2	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	30	41	35	65	110	176	286	339	235	99	17	-
Male	17	23	14	32	58	92	105	110	63	22	3	-
Female	13	18	21	33	52	84	181	229	172	77	14	-
White, both sexes	26	33	29	60	100	161	270	329	232	96	17	-
Male	15	19	10	29	53	85	100	106	63	22	3	-
Female	11	14	19	31	47	76	170	223	169	74	14	-
Black, both sexes	3	5	4	4	9	12	14	9	3	2	-	-
Male	1	2	2	2	5	6	4	4	-	-	-	-
Female	2	3	2	2	4	6	10	5	3	2	-	-
American Indian 1/, both sexes	1	1	1	-	-	1	1	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	2	1	1	-	1	-	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	-	1	-	-	1	1	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Kentucky												
All races, both sexes	61	77	105	165	179	281	414	397	313	93	14	-
Male	41	40	57	79	97	136	162	116	81	18	2	-
Female	20	37	48	86	82	145	252	281	232	75	12	-
White, both sexes	50	65	99	150	165	264	388	378	300	90	13	-
Male	34	35	53	68	91	128	155	112	76	17	1	-
Female	16	30	46	82	74	136	233	266	224	73	12	-
Black, both sexes	11	12	5	14	13	16	24	18	13	3	1	-
Male	7	5	4	11	5	8	6	4	5	1	1	-
Female	4	7	1	3	8	8	18	14	8	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	2	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	1	1	-	-	-	-
Louisiana												
All races, both sexes	87	94	97	152	210	327	394	385	236	77	7	-
Male	59	45	56	74	113	133	147	120	54	14	2	-
Female	28	49	41	78	97	194	247	265	182	63	5	-
White, both sexes	34	45	54	85	122	240	284	295	193	58	3	-
Male	21	23	27	42	62	101	107	94	42	9	1	-
Female	13	22	27	43	60	139	177	201	151	49	2	-
Black, both sexes	53	47	42	66	86	86	110	86	42	19	4	-
Male	38	21	28	31	50	32	40	24	11	5	1	-
Female	15	26	14	35	36	54	70	62	31	14	3	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	2	-	-	4	1	-	-	-
Male	-	-	1	1	1	-	-	2	1	-	-	-
Female	-	1	-	-	1	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Maine												
All races, both sexes	14	20	24	25	61	94	96	153	112	46	3	-
Male	8	14	14	10	27	45	40	46	22	13	-	-
Female	6	6	10	15	34	49	56	107	90	33	3	-
White, both sexes	14	20	24	25	61	93	96	153	112	46	3	-
Male	8	14	14	10	27	45	40	46	22	13	-	-
Female	6	6	10	15	34	48	56	107	90	33	3	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	67	102	108	123	176	306	432	499	294	115	21	-
Male	41	61	58	65	90	149	158	163	78	25	4	-
Female	26	41	50	58	86	157	274	336	216	90	17	-
White, both sexes	26	44	55	73	111	219	344	420	249	97	18	-
Male	14	24	31	34	56	105	122	141	68	23	4	-
Female	12	20	24	39	55	114	222	279	181	74	14	-
Black, both sexes	40	56	51	44	60	77	77	70	35	16	3	-
Male	27	35	26	27	30	38	31	20	9	1	-	-
Female	13	21	25	17	30	39	46	50	26	15	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	6	5	10	11	8	10	2	-	-
Male	-	2	-	4	4	6	5	2	1	1	-	-
Female	1	-	1	2	1	4	6	6	9	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Massachusetts												
All races, both sexes	35	67	77	117	178	372	601	641	520	160	45	-
Male	20	36	42	64	95	182	229	199	120	32	5	-
Female	15	31	35	53	83	190	372	442	400	128	40	-
White, both sexes	33	63	68	101	163	340	566	618	501	148	42	-
Male	19	33	38	57	86	164	219	189	118	30	5	-
Female	14	30	30	44	77	176	347	429	383	118	37	-
Black, both sexes	-	2	6	10	10	22	21	15	16	7	2	-
Male	-	2	3	4	6	12	7	7	2	1	-	-
Female	-	-	3	6	4	10	14	8	14	6	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	3	6	4	9	13	8	3	5	1	-
Male	1	1	1	3	3	6	3	3	-	1	-	-
Female	1	1	2	3	1	3	10	5	3	4	1	-
Michigan												
All races, both sexes	121	147	183	237	354	648	936	1,001	672	248	47	-
Male	64	92	97	133	181	291	380	350	169	45	8	-
Female	57	55	86	104	173	357	556	651	503	203	39	-
White, both sexes	72	95	130	192	293	552	826	906	629	230	41	-
Male	38	58	67	106	152	253	342	315	160	42	8	-
Female	34	37	63	86	141	299	484	591	469	188	33	-
Black, both sexes	47	51	51	43	56	88	104	87	38	18	5	-
Male	25	34	29	26	27	33	36	33	9	3	-	-
Female	22	17	22	17	29	55	68	54	29	15	5	-
American Indian 1/, both sexes	1	-	1	-	1	3	4	2	4	-	1	-
Male	1	-	1	-	-	3	2	-	-	-	-	-
Female	-	-	-	-	1	-	2	2	4	-	1	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	4	5	2	6	1	-	-	-
Male	-	-	-	1	2	2	-	2	-	-	-	-
Female	1	1	1	1	2	3	2	4	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Minnesota												
All races, both sexes	41	60	72	89	143	254	398	482	403	178	26	-
Male	28	41	39	50	70	119	163	174	104	34	2	-
Female	13	19	33	39	73	135	235	308	299	144	24	-
White, both sexes	35	55	63	76	136	247	392	472	401	173	26	-
Male	26	38	36	42	65	117	160	171	104	33	2	-
Female	9	17	27	34	71	130	232	301	297	140	24	-
Black, both sexes	3	2	5	4	1	-	2	2	2	4	-	-
Male	1	1	2	-	1	-	1	-	-	1	-	-
Female	2	1	3	4	-	-	1	2	2	3	-	-
American Indian 1/, both sexes	2	3	2	1	3	5	-	4	-	1	-	-
Male	1	2	1	1	2	1	-	-	-	-	-	-
Female	1	1	1	-	1	4	-	4	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	8	3	2	4	4	-	-	-	-
Male	-	-	-	7	2	1	2	3	-	-	-	-
Female	1	-	2	1	1	1	2	1	-	-	-	-
Mississippi												
All races, both sexes	57	84	94	113	155	243	260	231	167	62	7	-
Male	28	53	55	62	71	109	89	71	39	10	1	-
Female	29	31	39	51	84	134	171	160	128	52	6	-
White, both sexes	26	33	46	64	95	180	192	176	121	41	6	-
Male	10	16	26	41	47	87	72	51	27	6	1	-
Female	16	17	20	23	48	93	120	125	94	35	5	-
Black, both sexes	31	51	48	49	59	63	68	54	46	21	1	-
Male	18	37	29	21	24	22	17	20	12	4	-	-
Female	13	14	19	28	35	41	51	34	34	17	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Missouri												
All races, both sexes	89	101	121	179	267	441	611	629	474	171	37	-
Male	52	64	61	107	125	194	218	182	120	33	7	-
Female	37	37	60	72	142	247	393	447	354	138	30	-
White, both sexes	57	63	90	151	239	403	559	592	451	160	36	-
Male	31	41	49	90	112	181	201	175	112	31	7	-
Female	26	22	41	61	127	222	358	417	339	129	29	-
Black, both sexes	31	35	30	26	28	36	50	34	20	11	1	-
Male	21	22	12	16	13	12	17	7	5	2	-	-
Female	10	13	18	10	15	24	33	27	15	9	1	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	2	-	1	2	1	2	-	-	-
Male	-	1	-	1	-	-	-	-	2	-	-	-
Female	1	1	1	1	-	1	2	1	-	-	-	-
Montana												
All races, both sexes	13	11	11	25	32	48	90	113	75	29	5	-
Male	7	4	5	12	16	23	42	38	23	7	-	-
Female	6	7	6	13	16	25	48	75	52	22	5	-
White, both sexes	10	11	10	23	27	45	88	112	74	29	5	-
Male	5	4	4	12	15	22	41	38	23	7	-	-
Female	5	7	6	11	12	23	47	74	51	22	5	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	1	2	4	3	2	1	1	-	-	-
Male	1	-	1	-	1	1	1	-	-	-	-	-
Female	1	-	-	2	3	2	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Nebraska												
All races, both sexes	25	31	22	27	56	109	175	207	141	86	5	-
Male	14	24	14	9	30	58	73	59	40	15	-	-
Female	11	7	8	18	26	51	102	148	101	71	5	-
White, both sexes	20	30	18	24	53	107	172	203	138	86	5	-
Male	13	23	11	7	29	57	71	59	39	15	-	-
Female	7	7	7	17	24	50	101	144	99	71	5	-
Black, both sexes	5	1	4	1	3	1	2	3	3	-	-	-
Male	1	1	3	1	1	1	2	-	1	-	-	-
Female	4	-	1	-	2	-	-	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Nevada												
All races, both sexes	32	45	48	69	83	102	163	130	85	25	1	-
Male	19	26	24	39	42	52	63	48	20	7	-	-
Female	13	19	24	30	41	50	100	82	65	18	1	-
White, both sexes	24	37	36	52	68	91	151	120	81	25	1	-
Male	14	23	18	30	31	46	59	46	20	7	-	-
Female	10	14	18	22	37	45	92	74	61	18	1	-
Black, both sexes	5	4	7	9	10	7	7	8	1	-	-	-
Male	2	1	5	7	8	3	1	2	-	-	-	-
Female	3	3	2	2	2	4	6	6	1	-	-	-
American Indian 1/, both sexes	-	2	-	2	-	-	1	-	1	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	5	6	5	4	4	2	2	-	-	-
Male	3	1	1	1	3	3	2	-	-	-	-	-
Female	-	1	4	5	2	1	2	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
New Hampshire												
All races, both sexes	10	10	16	19	33	61	89	118	76	40	4	-
Male	5	4	12	15	17	33	39	34	16	5	2	-
Female	5	6	4	4	16	28	50	84	60	35	2	-
White, both sexes	7	10	16	18	33	61	89	117	76	40	4	-
Male	3	4	12	14	17	33	39	33	16	5	2	-
Female	4	6	4	4	16	28	50	84	60	35	2	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	88	129	143	154	256	441	698	693	497	169	29	-
Male	51	71	89	89	136	195	278	256	139	18	2	-
Female	37	58	54	65	120	246	420	437	358	151	27	-
White, both sexes	58	86	92	113	191	370	604	636	452	149	26	-
Male	36	47	54	69	103	162	243	234	125	15	2	-
Female	22	39	38	44	88	208	361	402	327	134	24	-
Black, both sexes	25	34	44	33	52	59	74	43	37	16	3	-
Male	14	19	30	16	26	29	29	16	13	3	-	-
Female	11	15	14	17	26	30	45	27	24	13	3	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	9	7	8	12	12	20	14	7	4	-	-
Male	1	5	5	4	7	4	6	6	1	-	-	-
Female	3	4	2	4	5	8	14	8	6	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
New Mexico												
All races, both sexes	18	20	32	40	66	97	144	136	110	34	8	-
Male	9	7	18	21	35	49	59	38	23	6	2	-
Female	9	13	14	19	31	48	85	98	87	28	6	-
White, both sexes	18	18	27	35	62	92	137	126	102	34	7	-
Male	9	6	16	18	33	45	56	38	19	6	2	-
Female	9	12	11	17	29	47	81	88	83	28	5	-
Black, both sexes	-	1	2	1	2	3	3	3	2	-	-	-
Male	-	1	-	1	1	2	1	-	1	-	-	-
Female	-	-	2	-	1	1	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	3	3	1	2	3	7	6	-	1	-
Male	-	-	2	2	1	2	1	-	3	-	-	-
Female	-	-	1	1	-	-	2	7	3	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	1	-	-	-	-	-	-	-
New York												
All races, both sexes	162	256	274	336	542	806	1,231	1,296	822	328	65	-
Male	95	153	148	184	264	357	498	432	210	56	3	-
Female	67	103	126	152	278	449	733	864	612	272	62	-
White, both sexes	106	186	189	242	423	695	1,090	1,171	769	314	57	-
Male	57	109	98	136	210	314	451	393	200	53	2	-
Female	49	77	91	106	213	381	639	778	569	261	55	-
Black, both sexes	51	60	71	72	96	85	109	97	45	12	8	-
Male	35	39	46	37	43	35	35	24	8	2	1	-
Female	16	21	25	35	53	50	74	73	37	10	7	-
American Indian 1/, both sexes	-	-	-	2	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	10	14	20	23	26	31	28	8	2	-	-
Male	3	5	4	11	11	8	12	15	2	1	-	-
Female	2	5	10	9	12	18	19	13	6	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
North Carolina												
All races, both sexes	155	173	219	279	373	640	849	869	564	194	38	-
Male	78	110	133	160	190	296	303	278	122	33	6	-
Female	77	63	86	119	183	344	546	591	442	161	32	-
White, both sexes	81	96	127	173	262	486	689	750	481	162	27	-
Male	40	61	76	100	129	228	251	252	104	29	4	-
Female	41	35	51	73	133	258	438	498	377	133	23	-
Black, both sexes	68	74	85	100	103	148	150	113	80	30	10	-
Male	33	46	52	55	57	66	47	25	17	4	2	-
Female	35	28	33	45	46	82	103	88	63	26	8	-
American Indian 1/, both sexes	4	3	4	4	4	6	6	2	-	2	-	-
Male	3	3	3	3	2	2	2	-	-	-	-	-
Female	1	-	1	1	2	4	4	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes	2	-	3	2	4	-	4	4	3	-	1	-
Male	2	-	2	2	2	-	3	1	1	-	-	-
Female	-	-	1	-	2	-	1	3	2	-	1	-
North Dakota												
All races, both sexes	5	5	8	16	22	48	89	93	87	35	8	-
Male	2	5	6	9	11	24	47	40	21	7	-	-
Female	3	-	2	7	11	24	42	53	66	28	8	-
White, both sexes	5	5	8	14	21	48	87	93	87	35	8	-
Male	2	5	6	8	10	24	46	40	21	7	-	-
Female	3	-	2	6	11	24	41	53	66	28	8	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	1	-	1	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Ohio												
All races, both sexes	158	198	219	305	459	761	1,180	1,225	798	275	44	-
Male	86	108	115	162	214	355	456	389	198	47	3	-
Female	72	90	104	143	245	406	724	836	600	228	41	-
White, both sexes	109	141	178	249	383	661	1,081	1,145	740	258	38	-
Male	55	75	95	122	175	311	419	367	185	43	2	-
Female	54	66	83	127	208	350	662	778	555	215	36	-
Black, both sexes	47	55	37	53	71	98	96	78	57	17	6	-
Male	31	31	19	38	37	44	35	21	13	4	1	-
Female	16	24	18	15	34	54	61	57	44	13	5	-
American Indian 1/, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	4	2	5	1	3	2	1	-	-	-
Male	-	1	1	2	2	-	2	1	-	-	-	-
Female	2	-	3	-	3	1	1	1	1	-	-	-
Oklahoma												
All races, both sexes	62	65	98	112	187	282	389	394	296	105	25	-
Male	32	43	57	64	88	125	153	127	68	18	5	-
Female	30	22	41	48	99	157	236	267	228	87	20	-
White, both sexes	43	42	80	88	169	253	357	357	273	98	20	-
Male	21	26	49	49	79	116	143	116	62	18	4	-
Female	22	16	31	39	90	137	214	241	211	80	16	-
Black, both sexes	12	19	14	15	11	17	15	19	12	4	3	-
Male	5	14	5	10	5	8	5	5	4	-	1	-
Female	7	5	9	5	6	9	10	14	8	4	2	-
American Indian 1/, both sexes	5	2	3	6	6	9	16	17	11	2	1	-
Male	5	1	3	3	4	-	5	5	2	-	-	-
Female	-	1	-	3	2	9	11	12	9	2	1	-
Asian or Pacific Islander 2/, both sexes	2	2	1	3	1	3	1	1	-	1	1	-
Male	1	2	-	2	-	1	-	1	-	-	-	-
Female	1	-	1	1	1	2	1	-	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Oregon												
All races, both sexes	44	62	65	82	133	242	411	442	292	127	26	-
Male	26	34	42	49	72	114	180	150	90	41	4	-
Female	18	28	23	33	61	128	231	292	202	86	22	-
White, both sexes	40	53	59	76	125	233	399	431	287	125	26	-
Male	24	31	40	46	67	109	177	144	89	41	4	-
Female	16	22	19	30	58	124	222	287	198	84	22	-
Black, both sexes	3	5	1	2	3	7	4	2	1	1	-	-
Male	2	3	-	1	3	5	-	2	1	-	-	-
Female	1	2	1	1	-	2	4	-	-	1	-	-
American Indian 1/, both sexes	1	-	3	2	1	-	1	1	2	-	-	-
Male	-	-	2	-	-	-	-	1	-	-	-	-
Female	1	-	1	2	1	-	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	2	2	4	2	7	8	2	1	-	-
Male	-	-	-	2	2	-	3	3	-	-	-	-
Female	-	4	2	-	2	2	4	5	2	1	-	-
Pennsylvania												
All races, both sexes	155	210	253	310	511	905	1,404	1,631	1,049	418	63	-
Male	90	122	157	177	248	394	548	515	261	67	5	-
Female	65	88	96	133	263	511	856	1,116	788	351	58	-
White, both sexes	121	156	208	255	442	808	1,300	1,528	990	390	57	-
Male	74	85	130	151	213	358	518	481	247	62	5	-
Female	47	71	78	104	229	450	782	1,047	743	328	52	-
Black, both sexes	33	51	43	47	64	93	95	94	55	26	6	-
Male	16	36	25	21	30	34	27	33	14	5	-	-
Female	17	15	18	26	34	59	68	61	41	21	6	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	8	5	4	9	9	4	2	-	-
Male	-	-	2	5	5	2	3	1	-	-	-	-
Female	1	2	-	3	-	2	6	8	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Rhode Island												
All races, both sexes	12	9	9	16	24	52	97	100	65	21	5	-
Male	9	7	6	10	13	24	35	33	14	4	1	-
Female	3	2	3	6	11	28	62	67	51	17	4	-
White, both sexes	12	9	8	14	23	52	94	100	62	20	5	-
Male	9	7	5	9	13	24	33	33	13	4	1	-
Female	3	2	3	5	10	28	61	67	49	16	4	-
Black, both sexes	-	-	-	-	1	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
South Carolina												
All races, both sexes	74	113	130	160	176	304	446	371	266	85	24	-
Male	44	55	72	95	77	146	161	116	64	13	1	-
Female	30	58	58	65	99	158	285	255	202	72	23	-
White, both sexes	32	55	81	92	111	222	332	277	204	67	14	-
Male	21	27	45	57	52	114	116	85	53	9	-	-
Female	11	28	36	35	59	108	216	192	151	58	14	-
Black, both sexes	42	56	49	66	65	78	109	91	60	17	10	-
Male	23	26	27	36	25	31	45	29	10	3	1	-
Female	19	30	22	30	40	47	64	62	50	14	9	-
American Indian 1/, both sexes	-	-	-	-	-	-	3	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	2	-	4	2	2	-	-	-	-
Male	-	2	-	2	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	3	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
South Dakota												
All races, both sexes	7	7	11	8	24	51	84	97	86	47	8	-
Male	5	2	6	5	11	25	41	49	23	11	1	-
Female	2	5	5	3	13	26	43	48	63	36	7	-
White, both sexes	6	5	11	7	18	49	82	97	84	47	8	-
Male	4	-	6	4	9	23	41	49	23	11	1	-
Female	2	5	5	3	9	26	41	48	61	36	7	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	2	-	1	6	2	2	-	1	-	-	-
Male	1	2	-	1	2	2	-	-	-	-	-	-
Female	-	-	-	-	4	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	103	135	174	208	312	455	614	608	432	181	23	-
Male	57	79	102	109	133	204	223	220	106	27	5	-
Female	46	56	72	99	179	251	391	388	326	154	18	-
White, both sexes	57	88	121	156	260	391	548	547	388	157	16	-
Male	32	54	66	85	109	174	198	202	94	24	4	-
Female	25	34	55	71	151	217	350	345	294	133	12	-
Black, both sexes	46	45	52	52	50	62	65	59	43	24	7	-
Male	25	24	36	24	22	29	24	17	11	3	1	-
Female	21	21	16	28	28	33	41	42	32	21	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	2	2	1	2	1	-	-	-
Male	-	1	-	-	2	1	1	1	1	-	-	-
Female	-	1	1	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Texas												
All races, both sexes	335	393	448	530	796	1,274	1,730	1,662	1,191	418	68	1
Male	188	223	244	259	397	578	687	530	319	78	12	-
Female	147	170	204	271	399	696	1,043	1,132	872	340	56	1
White, both sexes	232	293	347	412	647	1,080	1,533	1,504	1,077	378	61	1
Male	132	172	194	203	316	491	612	485	286	71	11	-
Female	100	121	153	209	331	589	921	1,019	791	307	50	1
Black, both sexes	95	86	91	105	131	162	172	147	106	37	7	-
Male	52	47	44	49	73	65	69	41	30	6	1	-
Female	43	39	47	56	58	97	103	106	76	31	6	-
American Indian 1/, both sexes	-	1	1	2	-	1	1	-	1	1	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	8	13	9	11	18	31	24	11	7	2	-	-
Male	4	4	5	6	8	22	6	4	3	1	-	-
Female	4	9	4	5	10	9	18	7	4	1	-	-
Utah												
All races, both sexes	10	16	29	30	56	99	132	157	94	33	-	-
Male	7	9	13	13	25	45	49	60	27	9	-	-
Female	3	7	16	17	31	54	83	97	67	24	-	-
White, both sexes	10	13	28	28	52	94	129	154	93	33	-	-
Male	7	9	13	13	25	44	49	60	27	9	-	-
Female	3	4	15	15	27	50	80	94	66	24	-	-
Black, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	3	2	2	3	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	2	1	-	3	1	2	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Vermont												
All races, both sexes	3	5	6	7	19	41	53	56	50	13	5	-
Male	-	3	3	4	9	19	23	20	17	4	-	-
Female	3	2	3	3	10	22	30	36	33	9	5	-
White, both sexes	3	5	6	7	19	41	53	56	50	13	5	-
Male	-	3	3	4	9	19	23	20	17	4	-	-
Female	3	2	3	3	10	22	30	36	33	9	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	93	124	140	176	321	470	653	704	497	180	28	-
Male	53	71	75	94	153	215	246	224	128	29	2	-
Female	40	53	65	82	168	255	407	480	369	151	26	-
White, both sexes	46	64	84	118	217	369	511	573	432	151	18	-
Male	22	38	45	57	107	168	199	186	109	24	2	-
Female	24	26	39	61	110	201	312	387	323	127	16	-
Black, both sexes	44	56	53	53	93	86	129	123	61	27	10	-
Male	29	30	27	34	41	40	40	35	18	4	-	-
Female	15	26	26	19	52	46	89	88	43	23	10	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	2	5	10	15	12	7	4	2	-	-
Male	2	3	2	3	5	7	6	3	1	1	-	-
Female	1	1	-	2	5	8	6	4	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Washington												
All races, both sexes	58	79	102	129	198	317	528	593	449	171	23	-
Male	36	38	54	65	80	150	210	214	121	41	2	-
Female	22	41	48	64	118	167	318	379	328	130	21	-
White, both sexes	43	67	82	108	164	285	477	569	428	163	23	-
Male	27	32	40	56	65	131	194	205	112	39	2	-
Female	16	35	42	52	99	154	283	364	316	124	21	-
Black, both sexes	7	7	6	7	11	10	15	3	5	3	-	-
Male	7	2	4	2	4	6	3	-	2	1	-	-
Female	-	5	2	5	7	4	12	3	3	2	-	-
American Indian 1/, both sexes	2	-	-	2	5	3	7	2	5	-	-	-
Male	-	-	-	1	3	1	3	-	-	-	-	-
Female	2	-	-	1	2	2	4	2	5	-	-	-
Asian or Pacific Islander 2/, both sexes	6	5	14	12	18	19	29	19	11	5	-	-
Male	2	4	10	6	8	12	10	9	7	1	-	-
Female	4	1	4	6	10	7	19	10	4	4	-	-
West Virginia												
All races, both sexes	34	40	45	67	92	162	202	210	123	57	11	-
Male	23	24	21	36	50	80	66	60	29	7	2	-
Female	11	16	24	31	42	82	136	150	94	50	9	-
White, both sexes	33	39	43	64	89	155	195	205	122	57	10	-
Male	22	23	21	33	49	75	64	60	29	7	1	-
Female	11	16	22	31	40	80	131	145	93	50	9	-
Black, both sexes	1	1	2	2	3	7	7	5	1	-	1	-
Male	1	1	-	2	1	5	2	-	-	-	1	-
Female	-	-	2	-	2	2	5	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Wisconsin												
All races, both sexes	52	71	91	118	196	311	562	631	490	186	34	-
Male	32	36	45	66	99	141	240	237	126	39	2	-
Female	20	35	46	52	97	170	322	394	364	147	32	-
White, both sexes	36	59	77	101	177	295	545	616	487	181	34	-
Male	24	30	36	56	86	134	232	233	125	39	2	-
Female	12	29	41	45	91	161	313	383	362	142	32	-
Black, both sexes	14	12	12	9	14	14	9	10	2	4	-	-
Male	7	6	8	5	9	5	6	2	-	-	-	-
Female	7	6	4	4	5	9	3	8	2	4	-	-
American Indian 1/, both sexes	1	-	1	1	2	-	5	-	-	-	-	-
Male	-	-	-	1	2	-	1	-	-	-	-	-
Female	1	-	1	-	-	-	4	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	7	3	2	3	5	1	1	-	-
Male	1	-	1	4	2	2	1	2	1	-	-	-
Female	-	-	-	3	1	-	2	3	-	1	-	-
Wyoming												
All races, both sexes	4	7	15	15	16	29	42	47	38	13	2	-
Male	3	5	8	5	7	15	17	15	14	1	-	-
Female	1	2	7	10	9	14	25	32	24	12	2	-
White, both sexes	4	7	15	14	15	29	41	47	38	13	2	-
Male	3	5	8	4	7	15	17	15	14	1	-	-
Female	1	2	7	10	8	14	24	32	24	12	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
United States												
All races, both sexes	5,846	2	-	9	13	26	49	61	112	211	421	589
Male	2,089	1	-	4	6	11	26	35	60	94	164	246
Female	3,757	1	-	5	7	15	23	26	52	117	257	343
White, both sexes	4,689	1	-	6	8	22	41	50	83	143	288	446
Male	1,710	-	-	4	6	10	22	30	48	59	121	199
Female	2,979	1	-	2	2	12	19	20	35	84	167	247
Black, both sexes	850	1	-	2	4	4	7	11	23	53	102	113
Male	283	1	-	-	-	1	4	5	9	26	32	35
Female	567	-	-	2	4	3	3	6	14	27	70	78
American Indian 1/, both sexes	30	-	-	1	-	-	1	-	1	2	7	5
Male	9	-	-	-	-	-	-	-	1	-	4	1
Female	21	-	-	1	-	-	1	-	-	2	3	4
Asian or Pacific Islander 2/, both sexes	277	-	-	-	1	-	-	-	5	13	24	25
Male	87	-	-	-	-	-	-	-	2	9	7	11
Female	190	-	-	-	1	-	-	-	3	4	17	14
Alabama												
All races, both sexes	80	-	-	-	-	-	-	-	4	-	6	5
Male	23	-	-	-	-	-	-	-	2	-	4	3
Female	57	-	-	-	-	-	-	-	2	-	2	2
White, both sexes	58	-	-	-	-	-	-	-	2	-	4	4
Male	16	-	-	-	-	-	-	-	1	-	3	2
Female	42	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	22	-	-	-	-	-	-	-	2	-	2	1
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	15	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Alaska												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	110	-	-	1	-	1	3	2	3	3	3	10
Male	35	-	-	-	-	1	1	-	2	2	1	4
Female	75	-	-	1	-	-	2	2	1	1	2	6
White, both sexes	100	-	-	-	-	1	3	1	1	3	3	8
Male	32	-	-	-	-	1	1	-	1	2	1	4
Female	68	-	-	-	-	-	2	1	-	1	2	4
Black, both sexes	6	-	-	-	-	-	-	1	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	4	-	-	1	-	-	-	-	1	-	-	2
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	1	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Arkansas												
All races, both sexes	40	-	-	-	-	-	-	1	2	2	1	5
Male	16	-	-	-	-	-	-	1	1	1	-	2
Female	24	-	-	-	-	-	-	-	1	1	1	3
White, both sexes	32	-	-	-	-	-	-	1	2	1	-	3
Male	15	-	-	-	-	-	-	1	1	-	-	2
Female	17	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	658	-	-	3	2	5	4	8	9	17	59	66
Male	230	-	-	2	1	3	2	5	7	12	21	36
Female	428	-	-	1	1	2	2	3	2	5	38	30
White, both sexes	490	-	-	3	1	5	4	7	7	10	43	56
Male	177	-	-	2	1	3	2	5	5	7	16	30
Female	313	-	-	1	-	2	2	2	2	3	27	26
Black, both sexes	51	-	-	-	1	-	-	1	1	2	6	2
Male	14	-	-	-	-	-	-	-	1	1	1	-
Female	37	-	-	-	1	-	-	1	-	1	5	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	114	-	-	-	-	-	-	-	1	5	9	7
Male	37	-	-	-	-	-	-	-	1	4	3	5
Female	77	-	-	-	-	-	-	-	-	1	6	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Colorado												
All races, both sexes	83	-	-	-	-	-	2	1	2	1	9	4
Male	27	-	-	-	-	-	1	-	1	-	1	2
Female	56	-	-	-	-	-	1	1	1	1	8	2
White, both sexes	77	-	-	-	-	-	2	1	2	1	8	4
Male	25	-	-	-	-	-	1	-	1	-	1	2
Female	52	-	-	-	-	-	1	1	1	1	7	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	69	-	-	-	-	-	-	-	2	2	3	5
Male	28	-	-	-	-	-	-	-	2	1	2	4
Female	41	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	59	-	-	-	-	-	-	-	2	2	1	4
Male	23	-	-	-	-	-	-	-	2	1	-	3
Female	36	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	2	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Delaware												
All races, both sexes	17	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	16	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	22	-	-	-	-	-	-	-	1	2	3	2
Male	8	-	-	-	-	-	-	-	1	-	1	1
Female	14	-	-	-	-	-	-	-	-	2	2	1
White, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	19	-	-	-	-	-	-	-	1	1	3	2
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	12	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Florida												
All races, both sexes	404	-	-	3	-	2	3	-	8	14	27	34
Male	147	-	-	1	-	2	1	-	3	2	8	15
Female	257	-	-	2	-	-	2	-	5	12	19	19
White, both sexes	341	-	-	2	-	1	1	-	8	11	18	23
Male	132	-	-	1	-	1	1	-	3	2	7	14
Female	209	-	-	1	-	-	-	-	5	9	11	9
Black, both sexes	58	-	-	1	-	1	2	-	-	3	8	11
Male	13	-	-	-	-	1	-	-	-	-	-	1
Female	45	-	-	1	-	-	2	-	-	3	8	10
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	173	-	-	-	-	-	1	5	4	9	17	19
Male	49	-	-	-	-	-	1	1	2	4	4	7
Female	124	-	-	-	-	-	-	4	2	5	13	12
White, both sexes	124	-	-	-	-	-	1	3	3	4	8	12
Male	39	-	-	-	-	-	1	1	1	1	4	4
Female	85	-	-	-	-	-	-	2	2	3	4	8
Black, both sexes	45	-	-	-	-	-	-	2	1	5	7	7
Male	10	-	-	-	-	-	-	-	1	3	-	3
Female	35	-	-	-	-	-	-	2	-	2	7	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Hawaii												
All races, both sexes	47	-	-	-	-	-	-	-	1	2	3	5
Male	15	-	-	-	-	-	-	-	-	1	-	3
Female	32	-	-	-	-	-	-	-	1	1	3	2
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	-	-	-	1	1	3	4
Male	13	-	-	-	-	-	-	-	-	1	-	2
Female	24	-	-	-	-	-	-	-	1	-	3	2
Idaho												
All races, both sexes	25	-	-	-	-	-	-	-	1	-	1	2
Male	14	-	-	-	-	-	-	-	1	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	1	-	1	2
Male	13	-	-	-	-	-	-	-	1	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Illinois												
All races, both sexes	224	-	-	-	2	-	1	1	3	6	13	32
Male	80	-	-	-	1	-	-	1	3	3	5	15
Female	144	-	-	-	1	-	1	-	-	3	8	17
White, both sexes	175	-	-	-	1	-	1	1	2	3	9	23
Male	60	-	-	-	1	-	-	1	2	1	4	12
Female	115	-	-	-	-	-	1	-	-	2	5	11
Black, both sexes	45	-	-	-	1	-	-	-	1	3	3	9
Male	19	-	-	-	-	-	-	-	1	2	1	3
Female	26	-	-	-	1	-	-	-	-	1	2	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	142	-	-	-	-	-	-	6	6	5	9	14
Male	67	-	-	-	-	-	-	5	3	2	2	10
Female	75	-	-	-	-	-	-	1	3	3	7	4
White, both sexes	126	-	-	-	-	-	-	6	5	3	8	11
Male	57	-	-	-	-	-	-	5	3	1	2	8
Female	69	-	-	-	-	-	-	1	2	2	6	3
Black, both sexes	14	-	-	-	-	-	-	-	1	2	1	3
Male	9	-	-	-	-	-	-	-	-	1	-	2
Female	5	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Iowa												
All races, both sexes	54	-	-	-	-	-	-	2	2	3	2	5
Male	22	-	-	-	-	-	-	2	1	-	1	2
Female	32	-	-	-	-	-	-	-	1	3	1	3
White, both sexes	51	-	-	-	-	-	-	2	2	2	2	5
Male	21	-	-	-	-	-	-	2	1	-	1	2
Female	30	-	-	-	-	-	-	-	1	2	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	59	-	-	-	1	-	1	-	1	4	6	8
Male	26	-	-	-	-	-	-	-	-	2	3	6
Female	33	-	-	-	1	-	1	-	1	2	3	2
White, both sexes	51	-	-	-	-	-	1	-	1	4	4	7
Male	21	-	-	-	-	-	-	-	-	2	1	5
Female	30	-	-	-	-	-	1	-	1	2	3	2
Black, both sexes	8	-	-	-	1	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	2	1
Female	3	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Kentucky												
All races, both sexes	87	-	-	-	-	1	-	1	2	2	7	9
Male	40	-	-	-	-	-	-	1	2	2	4	5
Female	47	-	-	-	-	1	-	-	-	-	3	4
White, both sexes	81	-	-	-	-	1	-	1	2	2	7	9
Male	39	-	-	-	-	-	-	1	2	2	4	5
Female	42	-	-	-	-	1	-	-	-	-	3	4
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	81	-	-	1	1	1	-	1	-	4	4	13
Male	29	-	-	-	-	-	-	1	-	1	2	4
Female	52	-	-	1	1	1	-	-	-	3	2	9
White, both sexes	46	-	-	-	-	1	-	1	-	3	3	8
Male	15	-	-	-	-	-	-	1	-	1	1	3
Female	31	-	-	-	-	1	-	-	-	2	2	5
Black, both sexes	33	-	-	1	1	-	-	-	-	1	1	5
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	20	-	-	1	1	-	-	-	-	1	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Maine												
All races, both sexes	31	-	-	-	-	-	1	2	2	1	1	2
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	20	-	-	-	-	-	1	2	2	-	1	1
White, both sexes	31	-	-	-	-	-	1	2	2	1	1	2
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	20	-	-	-	-	-	1	2	2	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	116	-	-	-	-	-	1	-	3	9	6	15
Male	51	-	-	-	-	-	1	-	2	5	1	8
Female	65	-	-	-	-	-	-	-	1	4	5	7
White, both sexes	80	-	-	-	-	-	-	-	2	6	2	7
Male	37	-	-	-	-	-	-	-	2	3	-	5
Female	43	-	-	-	-	-	-	-	-	3	2	2
Black, both sexes	33	-	-	-	-	-	1	-	1	3	4	8
Male	14	-	-	-	-	-	1	-	-	2	1	3
Female	19	-	-	-	-	-	-	-	1	1	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Massachusetts												
All races, both sexes	112	-	-	-	-	-	-	-	1	2	10	6
Male	37	-	-	-	-	-	-	-	1	1	4	3
Female	75	-	-	-	-	-	-	-	-	1	6	3
White, both sexes	101	-	-	-	-	-	-	-	-	-	8	6
Male	33	-	-	-	-	-	-	-	-	-	3	3
Female	68	-	-	-	-	-	-	-	-	-	5	3
Black, both sexes	9	-	-	-	-	-	-	-	1	2	2	-
Male	4	-	-	-	-	-	-	-	1	1	1	-
Female	5	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	194	-	-	-	-	2	-	1	5	7	10	20
Male	63	-	-	-	-	2	-	-	1	1	4	3
Female	131	-	-	-	-	-	-	1	4	6	6	17
White, both sexes	147	-	-	-	-	2	-	1	2	5	5	15
Male	50	-	-	-	-	2	-	-	1	1	1	3
Female	97	-	-	-	-	-	-	1	1	4	4	12
Black, both sexes	41	-	-	-	-	-	-	-	2	2	4	5
Male	12	-	-	-	-	-	-	-	-	-	3	-
Female	29	-	-	-	-	-	-	-	2	2	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Minnesota												
All races, both sexes	117	-	-	-	-	1	3	3	1	6	7	13
Male	45	-	-	-	-	-	2	1	-	1	5	4
Female	72	-	-	-	-	1	1	2	1	5	2	9
White, both sexes	107	-	-	-	-	1	2	2	1	4	6	10
Male	42	-	-	-	-	-	2	1	-	-	4	4
Female	65	-	-	-	-	1	-	1	1	4	2	6
Black, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	3
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	3
Mississippi												
All races, both sexes	62	1	-	-	-	-	-	1	1	5	3	7
Male	19	1	-	-	-	-	-	1	-	3	1	1
Female	43	-	-	-	-	-	-	-	1	2	2	6
White, both sexes	38	-	-	-	-	-	-	1	1	3	-	3
Male	14	-	-	-	-	-	-	1	-	2	-	1
Female	24	-	-	-	-	-	-	-	1	1	-	2
Black, both sexes	24	1	-	-	-	-	-	-	-	2	3	4
Male	5	1	-	-	-	-	-	-	-	1	1	-
Female	19	-	-	-	-	-	-	-	-	1	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Missouri												
All races, both sexes	139	-	-	-	-	-	2	1	1	7	12	22
Male	40	-	-	-	-	-	2	1	1	4	3	7
Female	99	-	-	-	-	-	-	-	-	3	9	15
White, both sexes	118	-	-	-	-	-	2	1	-	5	11	16
Male	35	-	-	-	-	-	2	1	-	3	3	6
Female	83	-	-	-	-	-	-	-	-	2	8	10
Black, both sexes	19	-	-	-	-	-	-	-	1	2	1	5
Male	4	-	-	-	-	-	-	-	1	1	-	-
Female	15	-	-	-	-	-	-	-	-	1	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Nebraska												
All races, both sexes	37	-	-	-	-	-	-	-	-	2	5	4
Male	18	-	-	-	-	-	-	-	-	1	4	1
Female	19	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	35	-	-	-	-	-	-	-	-	2	5	3
Male	17	-	-	-	-	-	-	-	-	1	4	1
Female	18	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Nevada												
All races, both sexes	54	-	-	-	1	1	-	-	2	2	7	7
Male	19	-	-	-	-	1	-	-	1	1	4	1
Female	35	-	-	-	1	-	-	-	1	1	3	6
White, both sexes	40	-	-	-	-	1	-	-	1	1	7	5
Male	17	-	-	-	-	1	-	-	1	1	4	1
Female	23	-	-	-	-	-	-	-	-	-	3	4
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	1	-	-	-	-	1	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	1	-	-	-	-	1	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
New Hampshire												
All races, both sexes	18	-	-	-	-	1	1	-	1	1	2	2
Male	11	-	-	-	-	-	-	-	-	1	2	2
Female	7	-	-	-	-	1	1	-	1	-	-	-
White, both sexes	18	-	-	-	-	1	1	-	1	1	2	2
Male	11	-	-	-	-	-	-	-	-	1	2	2
Female	7	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	160	-	-	-	2	-	1	3	6	3	13	25
Male	69	-	-	-	1	-	-	2	5	1	4	13
Female	91	-	-	-	1	-	1	1	1	2	9	12
White, both sexes	122	-	-	-	2	-	1	1	6	3	6	18
Male	60	-	-	-	1	-	-	1	5	1	2	11
Female	62	-	-	-	1	-	1	-	1	2	4	7
Black, both sexes	31	-	-	-	-	-	-	2	-	-	6	6
Male	7	-	-	-	-	-	-	1	-	-	1	1
Female	24	-	-	-	-	-	-	1	-	-	5	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
New Mexico												
All races, both sexes	31	-	-	-	-	-	-	-	1	1	2	6
Male	11	-	-	-	-	-	-	-	-	-	2	1
Female	20	-	-	-	-	-	-	-	1	1	-	5
White, both sexes	28	-	-	-	-	-	-	-	1	1	1	6
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	1	1	-	5
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	353	-	-	-	-	2	3	2	12	8	21	34
Male	132	-	-	-	-	2	3	1	4	7	7	13
Female	221	-	-	-	-	-	-	1	8	1	14	21
White, both sexes	265	-	-	-	-	2	3	2	7	4	11	29
Male	95	-	-	-	-	2	3	1	2	3	4	11
Female	170	-	-	-	-	-	-	1	5	1	7	18
Black, both sexes	63	-	-	-	-	-	-	-	4	3	9	3
Male	25	-	-	-	-	-	-	-	1	3	2	1
Female	38	-	-	-	-	-	-	-	3	-	7	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	1	1	1	2
Male	12	-	-	-	-	-	-	-	1	1	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
North Carolina												
All races, both sexes	191	-	-	-	1	1	2	3	2	11	14	19
Male	61	-	-	-	1	-	2	3	1	4	4	9
Female	130	-	-	-	-	1	-	-	1	7	10	10
White, both sexes	135	-	-	-	1	-	2	2	2	5	7	9
Male	42	-	-	-	1	-	2	2	1	2	2	4
Female	93	-	-	-	-	-	-	-	1	3	5	5
Black, both sexes	48	-	-	-	-	1	-	1	-	5	6	10
Male	16	-	-	-	-	-	-	1	-	2	2	5
Female	32	-	-	-	-	1	-	-	-	3	4	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Ohio												
All races, both sexes	240	-	-	-	1	1	3	4	2	4	14	26
Male	75	-	-	-	1	-	2	1	1	2	3	9
Female	165	-	-	-	-	1	1	3	1	2	11	17
White, both sexes	209	-	-	-	1	1	1	4	2	3	9	23
Male	64	-	-	-	1	-	-	1	1	2	2	8
Female	145	-	-	-	-	1	1	3	1	1	7	15
Black, both sexes	30	-	-	-	-	-	2	-	-	1	5	3
Male	11	-	-	-	-	-	2	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	1	4	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	76	1	-	-	-	-	2	-	-	1	8	5
Male	29	-	-	-	-	-	1	-	-	1	4	2
Female	47	1	-	-	-	-	1	-	-	-	4	3
White, both sexes	65	1	-	-	-	-	2	-	-	1	5	5
Male	26	-	-	-	-	-	1	-	-	1	3	2
Female	39	1	-	-	-	-	1	-	-	-	2	3
Black, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Oregon												
All races, both sexes	76	-	-	-	-	-	2	-	1	1	7	5
Male	23	-	-	-	-	-	-	-	1	-	5	2
Female	53	-	-	-	-	-	2	-	-	1	2	3
White, both sexes	73	-	-	-	-	-	2	-	1	1	6	5
Male	22	-	-	-	-	-	-	-	1	-	4	2
Female	51	-	-	-	-	-	2	-	-	1	2	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	273	-	-	1	-	1	3	1	3	12	17	20
Male	94	-	-	1	-	-	1	1	1	7	8	7
Female	179	-	-	-	-	1	2	-	2	5	9	13
White, both sexes	231	-	-	1	-	1	2	1	3	7	14	14
Male	84	-	-	1	-	-	1	1	1	3	7	5
Female	147	-	-	-	-	1	1	-	2	4	7	9
Black, both sexes	37	-	-	-	-	-	1	-	-	4	2	6
Male	8	-	-	-	-	-	-	-	-	3	1	2
Female	29	-	-	-	-	-	1	-	-	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	2	2
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	14	-	-	-	-	-	-	-	-	-	2	2
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	95	-	-	-	-	-	1	-	2	6	6	11
Male	22	-	-	-	-	-	-	-	1	4	3	-
Female	73	-	-	-	-	-	1	-	1	2	3	11
White, both sexes	60	-	-	-	-	-	1	-	1	2	2	7
Male	13	-	-	-	-	-	-	-	1	2	-	-
Female	47	-	-	-	-	-	1	-	-	-	2	7
Black, both sexes	33	-	-	-	-	-	-	-	1	4	4	2
Male	9	-	-	-	-	-	-	-	-	2	3	-
Female	24	-	-	-	-	-	-	-	1	2	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
South Dakota												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	153	-	-	-	-	-	2	2	4	7	8	20
Male	60	-	-	-	-	-	2	1	4	2	2	7
Female	93	-	-	-	-	-	-	1	-	5	6	13
White, both sexes	120	-	-	-	-	-	2	1	4	6	5	12
Male	49	-	-	-	-	-	2	-	4	1	2	3
Female	71	-	-	-	-	-	-	1	-	5	3	9
Black, both sexes	32	-	-	-	-	-	-	1	-	1	3	8
Male	11	-	-	-	-	-	-	1	-	1	-	4
Female	21	-	-	-	-	-	-	-	-	-	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Texas												
All races, both sexes	425	-	-	-	2	2	5	6	5	24	40	37
Male	161	-	-	-	1	-	3	4	2	7	19	17
Female	264	-	-	-	1	2	2	2	3	17	21	20
White, both sexes	353	-	-	-	2	1	4	4	2	20	28	32
Male	133	-	-	-	1	-	2	2	1	6	15	14
Female	220	-	-	-	1	1	2	2	1	14	13	18
Black, both sexes	56	-	-	-	-	1	1	2	3	3	8	4
Male	22	-	-	-	-	-	1	2	1	1	3	2
Female	34	-	-	-	-	1	-	-	2	2	5	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	1	4	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	1	3	-
Utah												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	1	3
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	132	-	-	-	-	2	1	3	2	2	11	12
Male	46	-	-	-	-	-	1	1	1	1	5	4
Female	86	-	-	-	-	2	-	2	1	1	6	8
White, both sexes	96	-	-	-	-	1	1	3	1	1	7	7
Male	29	-	-	-	-	-	1	1	1	-	2	3
Female	67	-	-	-	-	1	-	2	-	1	5	4
Black, both sexes	31	-	-	-	-	1	-	-	1	1	3	4
Male	15	-	-	-	-	-	-	-	-	1	3	1
Female	16	-	-	-	-	1	-	-	1	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Washington												
All races, both sexes	115	-	-	-	-	2	1	-	3	3	6	9
Male	35	-	-	-	-	-	-	-	1	2	2	5
Female	80	-	-	-	-	2	1	-	2	1	4	4
White, both sexes	100	-	-	-	-	2	1	-	2	2	4	7
Male	30	-	-	-	-	-	-	-	1	1	2	4
Female	70	-	-	-	-	2	1	-	1	1	2	3
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	1	1	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	1	-	1	1
West Virginia												
All races, both sexes	57	-	-	-	-	-	-	-	-	2	4	2
Male	21	-	-	-	-	-	-	-	-	2	2	-
Female	36	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	55	-	-	-	-	-	-	-	-	2	4	2
Male	20	-	-	-	-	-	-	-	-	2	2	-
Female	35	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Wisconsin												
All races, both sexes	92	-	-	-	-	-	-	1	-	5	9	10
Male	24	-	-	-	-	-	-	1	-	-	3	3
Female	68	-	-	-	-	-	-	-	-	5	6	7
White, both sexes	84	-	-	-	-	-	-	1	-	3	8	10
Male	23	-	-	-	-	-	-	1	-	-	3	3
Female	61	-	-	-	-	-	-	-	-	3	5	7
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Wyoming												
All races, both sexes	12	-	-	-	-	-	-	-	-	2	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	2	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
United States												
All races, both sexes	648	620	517	418	445	547	525	418	167	42	6	-
Male	257	240	177	140	142	184	152	108	36	6	-	-
Female	391	380	340	278	303	363	373	310	131	36	6	-
White, both sexes	484	488	421	346	367	474	458	366	154	38	5	-
Male	201	197	144	111	120	166	135	99	34	4	-	-
Female	283	291	277	235	247	308	323	267	120	34	5	-
Black, both sexes	132	95	65	53	53	43	45	31	10	3	-	-
Male	47	32	23	24	13	12	9	7	2	1	-	-
Female	85	63	42	29	40	31	36	24	8	2	-	-
American Indian 1/, both sexes	2	4	1	1	-	2	3	-	-	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	2	2	-	1	-	2	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	33	30	18	25	28	19	21	3	1	1	-
Male	9	9	9	5	9	6	8	2	-	1	-	-
Female	21	24	21	13	16	22	11	19	3	-	1	-
Alabama												
All races, both sexes	6	13	5	5	7	12	3	11	2	1	-	-
Male	2	4	1	1	1	1	1	3	-	-	-	-
Female	4	9	4	4	6	11	2	8	2	1	-	-
White, both sexes	6	7	2	5	5	9	3	8	2	1	-	-
Male	2	2	-	1	1	1	1	2	-	-	-	-
Female	4	5	2	4	4	8	2	6	2	1	-	-
Black, both sexes	-	6	3	-	2	3	-	3	-	-	-	-
Male	-	2	1	-	-	-	-	1	-	-	-	-
Female	-	4	2	-	2	3	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Alaska												
All races, both sexes	3	3	-	1	1	2	3	1	-	-	-	-
Male	-	3	-	-	1	1	1	-	-	-	-	-
Female	3	-	-	1	-	1	2	1	-	-	-	-
White, both sexes	2	3	-	1	1	2	1	-	-	-	-	-
Male	-	3	-	-	1	1	1	-	-	-	-	-
Female	2	-	-	1	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	12	21	7	5	5	12	9	11	2	-	-	-
Male	4	6	1	1	1	5	1	4	1	-	-	-
Female	8	15	6	4	4	7	8	7	1	-	-	-
White, both sexes	10	19	7	5	5	12	9	11	2	-	-	-
Male	3	5	1	1	1	5	1	4	1	-	-	-
Female	7	14	6	4	4	7	8	7	1	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Arkansas												
All races, both sexes	5	5	3	2	5	4	2	2	1	-	-	-
Male	3	4	2	-	1	1	-	-	-	-	-	-
Female	2	1	1	2	4	3	2	2	1	-	-	-
White, both sexes	5	5	2	1	4	4	1	2	1	-	-	-
Male	3	4	2	-	1	1	-	-	-	-	-	-
Female	2	1	-	1	3	3	1	2	1	-	-	-
Black, both sexes	-	-	1	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	86	72	57	40	40	73	59	43	11	3	1	-
Male	31	25	15	13	14	17	19	6	-	1	-	-
Female	55	47	42	27	26	56	40	37	11	2	1	-
White, both sexes	60	53	39	35	28	56	44	28	9	2	-	-
Male	22	20	9	11	11	15	13	5	-	-	-	-
Female	38	33	30	24	17	41	31	23	9	2	-	-
Black, both sexes	10	7	6	3	5	2	3	2	-	-	-	-
Male	5	1	2	1	1	-	1	-	-	-	-	-
Female	5	6	4	2	4	2	2	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	11	12	2	7	15	12	13	2	1	1	-
Male	4	4	4	1	2	2	5	1	-	1	-	-
Female	12	7	8	1	5	13	7	12	2	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Colorado												
All races, both sexes	15	10	4	5	6	6	8	5	3	2	-	-
Male	9	4	1	1	2	2	1	1	-	1	-	-
Female	6	6	3	4	4	4	7	4	3	1	-	-
White, both sexes	14	8	2	5	6	6	8	5	3	2	-	-
Male	8	3	1	1	2	2	1	1	-	1	-	-
Female	6	5	1	4	4	4	7	4	3	1	-	-
Black, both sexes	1	2	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	7	7	6	5	5	10	9	6	1	1	-	-
Male	4	4	-	1	2	5	2	1	-	-	-	-
Female	3	3	6	4	3	5	7	5	1	1	-	-
White, both sexes	7	6	4	4	5	10	7	5	1	1	-	-
Male	4	3	-	1	2	5	2	-	-	-	-	-
Female	3	3	4	3	3	5	5	5	1	1	-	-
Black, both sexes	-	1	2	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	2	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Delaware												
All races, both sexes	2	-	1	4	1	4	2	-	-	-	-	-
Male	1	-	1	2	1	3	-	-	-	-	-	-
Female	1	-	-	2	-	1	2	-	-	-	-	-
White, both sexes	1	-	1	4	1	4	2	-	-	-	-	-
Male	1	-	1	2	1	3	-	-	-	-	-	-
Female	-	-	-	2	-	1	2	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	1	1	3	2	-	1	1	-	-	-	-
Male	2	1	1	-	-	-	-	1	-	-	-	-
Female	3	-	-	3	2	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	5	1	1	2	2	-	1	-	-	-	-	-
Male	2	1	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	2	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Florida												
All races, both sexes	44	39	50	29	30	38	37	30	9	7	-	-
Male	17	14	13	14	14	16	13	12	2	-	-	-
Female	27	25	37	15	16	22	24	18	7	7	-	-
White, both sexes	30	37	42	23	30	37	36	27	8	7	-	-
Male	12	14	12	9	14	15	13	12	2	-	-	-
Female	18	23	30	14	16	22	23	15	6	7	-	-
Black, both sexes	14	1	5	6	-	1	1	3	1	-	-	-
Male	5	-	-	5	-	1	-	-	-	-	-	-
Female	9	1	5	1	-	-	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	2	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	22	16	19	19	9	9	14	6	4	-	-	-
Male	7	6	3	4	2	4	3	1	-	-	-	-
Female	15	10	16	15	7	5	11	5	4	-	-	-
White, both sexes	16	11	17	17	7	5	11	6	3	-	-	-
Male	6	5	3	4	2	3	3	1	-	-	-	-
Female	10	6	14	13	5	2	8	5	3	-	-	-
Black, both sexes	6	4	2	2	1	4	3	-	1	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	5	3	2	2	1	3	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Hawaii												
All races, both sexes	5	7	5	6	5	2	1	4	1	-	-	-
Male	1	3	3	1	2	-	-	1	-	-	-	-
Female	4	4	2	5	3	2	1	3	1	-	-	-
White, both sexes	-	1	2	1	1	1	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	1	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	6	3	5	4	1	1	3	-	-	-	-
Male	1	3	2	1	2	-	-	1	-	-	-	-
Female	4	3	1	4	2	1	1	2	-	-	-	-
Idaho												
All races, both sexes	5	2	1	2	3	2	4	2	-	-	-	-
Male	2	1	1	1	2	-	2	2	-	-	-	-
Female	3	1	-	1	1	2	2	-	-	-	-	-
White, both sexes	5	2	1	2	3	2	3	2	-	-	-	-
Male	2	1	1	1	2	-	1	2	-	-	-	-
Female	3	1	-	1	1	2	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Illinois												
All races, both sexes	23	23	23	15	13	18	23	18	8	2	-	-
Male	8	12	7	5	2	4	9	4	1	-	-	-
Female	15	11	16	10	11	14	14	14	7	2	-	-
White, both sexes	16	17	19	13	11	14	19	17	7	2	-	-
Male	3	9	6	4	2	3	8	3	1	-	-	-
Female	13	8	13	9	9	11	11	14	6	2	-	-
Black, both sexes	7	5	3	2	2	4	3	1	1	-	-	-
Male	5	3	1	1	-	1	-	1	-	-	-	-
Female	2	2	2	1	2	3	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	13	8	12	9	19	13	16	9	2	1	-	-
Male	4	5	5	5	10	3	6	4	2	1	-	-
Female	9	3	7	4	9	10	10	5	-	-	-	-
White, both sexes	13	7	12	5	17	11	16	9	2	1	-	-
Male	4	4	5	2	8	2	6	4	2	1	-	-
Female	9	3	7	3	9	9	10	5	-	-	-	-
Black, both sexes	-	1	-	4	1	1	-	-	-	-	-	-
Male	-	1	-	3	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Iowa												
All races, both sexes	5	7	3	5	5	2	7	5	-	1	-	-
Male	1	2	2	3	1	1	4	2	-	-	-	-
Female	4	5	1	2	4	1	3	3	-	1	-	-
White, both sexes	5	6	3	4	5	2	7	5	-	1	-	-
Male	1	2	2	2	1	1	4	2	-	-	-	-
Female	4	4	1	2	4	1	3	3	-	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	7	7	4	5	1	4	5	3	2	-	-	-
Male	3	1	2	2	-	4	2	1	-	-	-	-
Female	4	6	2	3	1	-	3	2	2	-	-	-
White, both sexes	6	7	3	5	1	3	4	3	2	-	-	-
Male	3	1	2	2	-	3	1	1	-	-	-	-
Female	3	6	1	3	1	-	3	2	2	-	-	-
Black, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Kentucky												
All races, both sexes	16	7	9	9	2	7	8	3	4	-	-	-
Male	7	3	5	2	1	4	2	-	2	-	-	-
Female	9	4	4	7	1	3	6	3	2	-	-	-
White, both sexes	14	5	9	8	1	7	8	3	4	-	-	-
Male	6	3	5	2	1	4	2	-	2	-	-	-
Female	8	2	4	6	-	3	6	3	2	-	-	-
Black, both sexes	2	2	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	2	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	9	10	7	3	2	12	6	6	1	-	-	-
Male	3	2	4	1	1	6	2	2	-	-	-	-
Female	6	8	3	2	1	6	4	4	1	-	-	-
White, both sexes	5	4	4	1	1	8	3	3	1	-	-	-
Male	1	1	1	-	-	4	1	1	-	-	-	-
Female	4	3	3	1	1	4	2	2	1	-	-	-
Black, both sexes	4	6	2	2	1	4	3	2	-	-	-	-
Male	2	1	2	1	1	2	1	1	-	-	-	-
Female	2	5	-	1	-	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Maine												
All races, both sexes	4	4	2	1	3	3	3	1	1	-	-	-
Male	2	2	2	-	-	1	1	1	-	-	-	-
Female	2	2	-	1	3	2	2	-	1	-	-	-
White, both sexes	4	4	2	1	3	3	3	1	1	-	-	-
Male	2	2	2	-	-	1	1	1	-	-	-	-
Female	2	2	-	1	3	2	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	11	13	9	12	12	8	9	5	3	-	-	-
Male	3	7	6	4	5	4	1	2	2	-	-	-
Female	8	6	3	8	7	4	8	3	1	-	-	-
White, both sexes	7	10	9	8	10	6	8	4	1	-	-	-
Male	1	6	6	2	4	4	1	2	1	-	-	-
Female	6	4	3	6	6	2	7	2	-	-	-	-
Black, both sexes	4	3	-	3	2	1	1	1	1	-	-	-
Male	2	1	-	2	1	-	-	-	1	-	-	-
Female	2	2	-	1	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Massachusetts												
All races, both sexes	9	9	7	6	10	18	12	12	6	4	-	-
Male	5	5	2	3	4	3	1	3	1	1	-	-
Female	4	4	5	3	6	15	11	9	5	3	-	-
White, both sexes	9	7	7	6	10	16	11	11	6	4	-	-
Male	5	4	2	3	4	3	1	3	1	1	-	-
Female	4	3	5	3	6	13	10	8	5	3	-	-
Black, both sexes	-	1	-	-	-	2	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	18	17	20	20	12	20	17	18	6	-	1	-
Male	7	8	7	4	3	8	5	7	3	-	-	-
Female	11	9	13	16	9	12	12	11	3	-	1	-
White, both sexes	10	10	17	17	9	18	13	16	6	-	1	-
Male	5	4	7	2	3	7	4	7	3	-	-	-
Female	5	6	10	15	6	11	9	9	3	-	1	-
Black, both sexes	8	7	3	2	2	1	4	1	-	-	-	-
Male	2	4	-	1	-	1	1	-	-	-	-	-
Female	6	3	3	1	2	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Minnesota												
All races, both sexes	9	12	13	7	11	10	10	9	1	-	1	-
Male	6	5	5	2	5	6	3	-	-	-	-	-
Female	3	7	8	5	6	4	7	9	1	-	1	-
White, both sexes	9	12	13	6	11	9	10	9	1	-	1	-
Male	6	5	5	2	5	5	3	-	-	-	-	-
Female	3	7	8	4	6	4	7	9	1	-	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	10	2	6	7	2	8	3	4	2	-	-	-
Male	-	1	-	3	1	2	2	2	1	-	-	-
Female	10	1	6	4	1	6	1	2	1	-	-	-
White, both sexes	4	1	5	3	1	8	3	4	1	-	-	-
Male	-	1	-	1	1	2	2	2	1	-	-	-
Female	4	-	5	2	-	6	1	2	-	-	-	-
Black, both sexes	6	1	1	4	1	-	-	-	1	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	6	1	1	2	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Missouri												
All races, both sexes	11	10	8	7	7	16	11	14	5	4	1	-
Male	2	4	3	3	1	6	-	3	-	-	-	-
Female	9	6	5	4	6	10	11	11	5	4	1	-
White, both sexes	8	6	8	7	7	15	10	13	4	4	1	-
Male	2	2	3	3	1	6	-	3	-	-	-	-
Female	6	4	5	4	6	9	10	10	4	4	1	-
Black, both sexes	3	4	-	-	-	1	-	1	1	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	3	2	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	1	4	1	4	-	4	2	-	-	-	-	-
Male	-	1	1	1	-	2	1	-	-	-	-	-
Female	1	3	-	3	-	2	1	-	-	-	-	-
White, both sexes	1	4	1	4	-	4	2	-	-	-	-	-
Male	-	1	1	1	-	2	1	-	-	-	-	-
Female	1	3	-	3	-	2	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Nebraska												
All races, both sexes	2	5	3	2	7	1	5	1	-	-	-	-
Male	1	3	3	-	3	-	2	-	-	-	-	-
Female	1	2	-	2	4	1	3	1	-	-	-	-
White, both sexes	2	5	2	2	7	1	5	1	-	-	-	-
Male	1	3	2	-	3	-	2	-	-	-	-	-
Female	1	2	-	2	4	1	3	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	7	6	6	2	4	5	1	-	-	-	-
Male	2	2	1	3	1	1	1	-	-	-	-	-
Female	1	5	5	3	1	3	4	1	-	-	-	-
White, both sexes	2	6	4	3	1	4	5	-	-	-	-	-
Male	2	2	1	2	-	1	1	-	-	-	-	-
Female	-	4	3	1	1	3	4	-	-	-	-	-
Black, both sexes	1	1	1	1	1	-	-	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	1	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
New Hampshire												
All races, both sexes	1	3	2	-	3	-	-	1	-	-	-	-
Male	1	1	2	-	2	-	-	-	-	-	-	-
Female	-	2	-	-	1	-	-	1	-	-	-	-
White, both sexes	1	3	2	-	3	-	-	1	-	-	-	-
Male	1	1	2	-	2	-	-	-	-	-	-	-
Female	-	2	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	19	15	5	12	15	15	9	5	1	-	-
Male	2	6	6	4	6	7	7	4	1	-	-	-
Female	9	13	9	1	6	8	8	5	4	1	-	-
White, both sexes	7	13	10	5	10	11	15	9	5	-	-	-
Male	2	6	5	4	4	6	7	4	1	-	-	-
Female	5	7	5	1	6	5	8	5	4	-	-	-
Black, both sexes	2	4	5	-	2	3	-	-	-	1	-	-
Male	-	-	1	-	2	1	-	-	-	-	-	-
Female	2	4	4	-	-	2	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
New Mexico												
All races, both sexes	2	3	4	2	1	2	4	2	1	-	-	-
Male	1	-	2	1	1	2	1	-	-	-	-	-
Female	1	3	2	1	-	-	3	2	1	-	-	-
White, both sexes	2	2	3	2	1	2	4	2	1	-	-	-
Male	1	-	1	1	1	2	1	-	-	-	-	-
Female	1	2	2	1	-	-	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	36	52	33	29	23	24	37	28	9	-	-	-
Male	18	21	11	14	3	8	12	6	2	-	-	-
Female	18	31	22	15	20	16	25	22	7	-	-	-
White, both sexes	27	38	23	23	15	21	29	23	8	-	-	-
Male	12	16	5	11	3	7	9	5	1	-	-	-
Female	15	22	18	12	12	14	20	18	7	-	-	-
Black, both sexes	7	10	6	3	6	1	6	4	1	-	-	-
Male	4	4	5	1	-	-	2	1	1	-	-	-
Female	3	6	1	2	6	1	4	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	4	3	2	2	2	1	-	-	-	-
Male	2	1	1	2	-	1	1	-	-	-	-	-
Female	-	3	3	1	2	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
North Carolina												
All races, both sexes	27	16	16	16	20	14	11	16	2	-	-	-
Male	6	8	5	3	3	4	3	4	1	-	-	-
Female	21	8	11	13	17	10	8	12	1	-	-	-
White, both sexes	18	10	9	13	18	13	9	15	2	-	-	-
Male	4	5	2	3	3	4	2	4	1	-	-	-
Female	14	5	7	10	15	9	7	11	1	-	-	-
Black, both sexes	7	4	6	3	1	1	2	1	-	-	-	-
Male	1	1	3	-	-	-	1	-	-	-	-	-
Female	6	3	3	3	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	1	1	-	2	4	-	1	1	-	-
Male	1	-	1	-	-	1	3	-	-	-	-	-
Female	2	-	-	1	-	1	1	-	1	1	-	-
White, both sexes	3	-	1	1	-	2	4	-	1	1	-	-
Male	1	-	1	-	-	1	3	-	-	-	-	-
Female	2	-	-	1	-	1	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Ohio												
All races, both sexes	27	32	15	15	18	23	22	23	10	-	-	-
Male	13	11	2	4	6	5	5	7	3	-	-	-
Female	14	21	13	11	12	18	17	16	7	-	-	-
White, both sexes	22	27	14	15	15	20	21	22	9	-	-	-
Male	10	9	2	4	5	4	5	7	3	-	-	-
Female	12	18	12	11	10	16	16	15	6	-	-	-
Black, both sexes	5	5	1	-	2	3	1	1	1	-	-	-
Male	3	2	-	-	1	1	-	-	-	-	-	-
Female	2	3	1	-	1	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	12	10	8	3	4	7	7	5	3	-	-	-
Male	5	6	3	1	-	3	1	1	1	-	-	-
Female	7	4	5	2	4	4	6	4	2	-	-	-
White, both sexes	9	8	8	3	4	5	6	5	3	-	-	-
Male	5	4	3	1	-	3	1	1	1	-	-	-
Female	4	4	5	2	4	2	5	4	2	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Oregon												
All races, both sexes	9	12	6	4	10	5	4	5	5	-	-	-
Male	3	2	2	1	2	1	1	2	1	-	-	-
Female	6	10	4	3	8	4	3	3	4	-	-	-
White, both sexes	9	12	5	4	9	5	4	5	5	-	-	-
Male	3	2	2	1	2	1	1	2	1	-	-	-
Female	6	10	3	3	7	4	3	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	21	24	23	18	25	30	26	27	17	4	-	-
Male	11	9	7	8	9	10	5	5	4	-	-	-
Female	10	15	16	10	16	20	21	22	13	4	-	-
White, both sexes	18	22	19	16	20	24	25	25	16	3	-	-
Male	11	8	7	8	8	9	5	5	4	-	-	-
Female	7	14	12	8	12	15	20	20	12	3	-	-
Black, both sexes	3	2	4	1	4	5	1	2	1	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	3	1	4	1	4	4	1	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Rhode Island												
All races, both sexes	2	-	1	3	-	1	2	1	-	-	1	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	2	-	-	2	-	1	1	1	-	-	1	-
White, both sexes	2	-	1	2	-	1	2	1	-	-	1	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	2	-	-	1	-	1	1	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	8	5	8	8	8	8	12	8	2	2	-	-
Male	2	-	-	3	1	3	2	1	1	1	-	-
Female	6	5	8	5	7	5	10	7	1	1	-	-
White, both sexes	4	5	7	5	5	7	6	5	2	1	-	-
Male	2	-	-	2	1	3	1	-	1	-	-	-
Female	2	5	7	3	4	4	5	5	1	1	-	-
Black, both sexes	4	-	1	3	3	1	6	3	-	1	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	4	-	1	2	3	1	5	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
South Dakota												
All races, both sexes	3	1	5	1	-	2	3	1	2	1	-	-
Male	2	-	2	1	-	1	1	-	1	1	-	-
Female	1	1	3	-	-	1	2	1	1	-	-	-
White, both sexes	3	1	5	1	-	2	3	1	2	1	-	-
Male	2	-	2	1	-	1	1	-	1	1	-	-
Female	1	1	3	-	-	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	20	17	19	5	16	15	6	10	1	1	-	-
Male	9	8	8	2	4	6	2	3	-	-	-	-
Female	11	9	11	3	12	9	4	7	1	1	-	-
White, both sexes	15	11	15	5	14	14	6	8	1	1	-	-
Male	8	8	6	2	3	6	2	2	-	-	-	-
Female	7	3	9	3	11	8	4	6	1	1	-	-
Black, both sexes	5	5	4	-	2	1	-	2	-	-	-	-
Male	1	-	2	-	1	-	-	1	-	-	-	-
Female	4	5	2	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Texas												
All races, both sexes	55	39	34	30	39	35	31	25	15	1	-	-
Male	25	15	17	7	14	9	11	7	3	-	-	-
Female	30	24	17	23	25	26	20	18	12	1	-	-
White, both sexes	42	34	31	25	28	31	30	24	14	1	-	-
Male	19	14	15	6	9	8	11	7	3	-	-	-
Female	23	20	16	19	19	23	19	17	11	1	-	-
Black, both sexes	10	4	2	5	8	2	1	1	1	-	-	-
Male	5	1	2	1	3	-	-	-	-	-	-	-
Female	5	3	-	4	5	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	-	3	2	-	-	-	-	-	-
Male	1	-	-	-	2	1	-	-	-	-	-	-
Female	2	1	1	-	1	1	-	-	-	-	-	-
Utah												
All races, both sexes	1	4	2	2	3	1	2	2	-	1	-	-
Male	1	2	1	2	-	-	-	1	-	-	-	-
Female	-	2	1	-	3	1	2	1	-	1	-	-
White, both sexes	1	2	2	2	3	1	2	2	-	1	-	-
Male	1	2	1	2	-	-	-	1	-	-	-	-
Female	-	-	1	-	3	1	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Vermont												
All races, both sexes	2	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	10	12	11	12	13	16	5	9	-	-	-
Male	5	5	6	3	5	4	4	-	1	-	-	-
Female	6	5	6	8	7	9	12	5	8	-	-	-
White, both sexes	4	7	9	8	10	11	13	4	9	-	-	-
Male	1	4	4	1	4	3	3	-	1	-	-	-
Female	3	3	5	7	6	8	10	4	8	-	-	-
Black, both sexes	7	2	3	3	1	1	3	1	-	-	-	-
Male	4	-	2	2	-	1	1	-	-	-	-	-
Female	3	2	1	1	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Washington												
All races, both sexes	14	14	9	7	9	10	10	8	7	2	1	-
Male	8	2	2	2	2	4	2	2	1	-	-	-
Female	6	12	7	5	7	6	8	6	6	2	1	-
White, both sexes	12	13	8	5	8	9	9	8	7	2	1	-
Male	7	2	2	1	2	3	2	2	1	-	-	-
Female	5	11	6	4	6	6	7	6	6	2	1	-
Black, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	1	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	7	5	4	5	6	10	8	4	-	-	-	-
Male	4	2	1	2	-	4	3	1	-	-	-	-
Female	3	3	3	3	6	6	5	3	-	-	-	-
White, both sexes	6	5	4	5	6	10	7	4	-	-	-	-
Male	3	2	1	2	-	4	3	1	-	-	-	-
Female	3	3	3	3	6	6	4	3	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Wisconsin												
All races, both sexes	7	12	7	8	9	7	8	5	2	2	-	-
Male	3	3	1	1	3	2	2	1	1	-	-	-
Female	4	9	6	7	6	5	6	4	1	2	-	-
White, both sexes	5	11	7	8	7	7	8	5	2	2	-	-
Male	3	3	1	1	2	2	2	1	1	-	-	-
Female	2	8	6	7	5	5	6	4	1	2	-	-
Black, both sexes	2	1	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	2	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	1	1	-	1	2	2	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	-	1	1	2	-	-	-	-
White, both sexes	1	1	1	1	-	1	2	2	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
United States												
All races, both sexes	27,216	65	12	11	15	28	52	69	129	284	604	1,059
Male	12,763	37	5	5	11	14	30	34	69	179	336	606
Female	14,453	28	7	6	4	14	22	35	60	105	268	453
White, both sexes	22,053	44	8	7	12	22	33	44	78	157	356	628
Male	10,155	23	2	3	9	11	19	23	39	102	205	370
Female	11,898	21	6	4	3	11	14	21	39	55	151	258
Black, both sexes	3,924	21	2	3	2	6	15	20	39	103	203	362
Male	1,986	14	2	1	2	3	8	7	22	64	105	188
Female	1,938	7	-	2	-	3	7	13	17	39	98	174
American Indian 1/, both sexes	144	-	1	1	-	-	1	-	2	2	14	9
Male	73	-	1	1	-	-	1	-	1	2	11	7
Female	71	-	-	-	-	-	-	-	1	-	3	2
Asian or Pacific Islander 2/, both sexes	1,095	-	1	-	1	-	3	5	10	22	31	60
Male	549	-	-	-	-	-	2	4	7	11	15	41
Female	546	-	1	-	1	-	1	1	3	11	16	19
Alabama												
All races, both sexes	563	1	-	-	-	-	-	2	3	7	19	22
Male	254	1	-	-	-	-	-	-	1	4	8	17
Female	309	-	-	-	-	-	-	2	2	3	11	5
White, both sexes	399	1	-	-	-	-	-	2	2	1	6	7
Male	171	1	-	-	-	-	-	-	1	1	1	6
Female	228	-	-	-	-	-	-	2	1	-	5	1
Black, both sexes	161	-	-	-	-	-	-	-	1	6	13	15
Male	82	-	-	-	-	-	-	-	-	3	7	11
Female	79	-	-	-	-	-	-	-	1	3	6	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Alaska												
All races, both sexes	46	-	-	-	-	1	1	-	-	-	2	-
Male	20	-	-	-	-	-	1	-	-	-	2	-
Female	26	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	1	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	1	-	-	-	1	-
Male	5	-	-	-	-	-	1	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	394	1	1	1	-	-	1	3	2	8	12	17
Male	197	-	1	-	-	-	1	-	1	7	6	8
Female	197	1	-	1	-	-	-	3	1	1	6	9
White, both sexes	357	1	-	1	-	-	1	2	2	5	8	15
Male	179	-	-	-	-	-	1	-	1	4	4	7
Female	178	1	-	1	-	-	-	2	1	1	4	8
Black, both sexes	15	-	-	-	-	-	-	1	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	17	-	1	-	-	-	-	-	-	1	3	1
Male	11	-	1	-	-	-	-	-	-	1	2	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Arkansas												
All races, both sexes	303	1	-	-	-	-	1	-	5	6	3	13
Male	128	-	-	-	-	-	1	-	3	4	1	8
Female	175	1	-	-	-	-	-	-	2	2	2	5
White, both sexes	247	1	-	-	-	-	1	-	5	1	1	6
Male	103	-	-	-	-	-	1	-	3	1	1	3
Female	144	1	-	-	-	-	-	-	2	-	-	3
Black, both sexes	52	-	-	-	-	-	-	-	-	5	2	7
Male	23	-	-	-	-	-	-	-	-	3	-	5
Female	29	-	-	-	-	-	-	-	-	2	2	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,826	5	-	1	3	4	7	6	11	38	88	126
Male	1,394	4	-	1	2	2	5	4	7	30	48	72
Female	1,432	1	-	-	1	2	2	2	4	8	40	54
White, both sexes	2,126	3	-	1	2	4	4	6	7	29	69	85
Male	1,036	2	-	1	2	2	3	4	4	23	39	48
Female	1,090	1	-	-	-	2	1	2	3	6	30	37
Black, both sexes	241	2	-	-	-	-	2	-	1	2	9	19
Male	120	2	-	-	-	-	1	-	1	1	4	8
Female	121	-	-	-	-	-	1	-	-	1	5	11
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	449	-	-	-	1	-	1	-	3	7	10	21
Male	232	-	-	-	-	-	1	-	2	6	5	15
Female	217	-	-	-	1	-	-	-	1	1	5	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Colorado												
All races, both sexes	312	1	-	-	-	-	1	1	1	2	8	13
Male	139	-	-	-	-	-	-	1	-	-	4	6
Female	173	1	-	-	-	-	1	-	1	2	4	7
White, both sexes	292	1	-	-	-	-	1	-	1	2	7	12
Male	127	-	-	-	-	-	-	-	-	-	4	6
Female	165	1	-	-	-	-	1	-	1	2	3	6
Black, both sexes	13	-	-	-	-	-	-	1	-	-	1	-
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Connecticut												
All races, both sexes	321	-	-	-	-	-	1	2	1	2	3	6
Male	137	-	-	-	-	-	1	-	1	2	1	4
Female	184	-	-	-	-	-	-	2	-	-	2	2
White, both sexes	292	-	-	-	-	-	1	2	-	2	1	2
Male	121	-	-	-	-	-	1	-	-	2	-	1
Female	171	-	-	-	-	-	-	2	-	-	1	1
Black, both sexes	21	-	-	-	-	-	-	-	1	-	2	4
Male	12	-	-	-	-	-	-	-	1	-	1	3
Female	9	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Delaware												
All races, both sexes	80	-	-	-	-	-	-	-	2	1	-	1
Male	27	-	-	-	-	-	-	-	-	1	-	-
Female	53	-	-	-	-	-	-	-	2	-	-	1
White, both sexes	61	-	-	-	-	-	-	-	1	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	55	-	1	-	1	1	-	-	-	-	-	1
Male	29	-	1	-	1	1	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	45	-	1	-	1	1	-	-	-	-	-	1
Male	25	-	1	-	1	1	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Florida												
All races, both sexes	1,772	6	-	-	1	2	4	6	10	9	24	69
Male	853	3	-	-	-	2	1	3	7	4	13	41
Female	919	3	-	-	1	-	3	3	3	5	11	28
White, both sexes	1,492	4	-	-	1	1	3	4	9	7	14	41
Male	706	2	-	-	-	1	1	2	6	4	7	24
Female	786	2	-	-	1	-	2	2	3	3	7	17
Black, both sexes	247	2	-	-	-	1	1	1	1	1	8	25
Male	128	1	-	-	-	1	-	-	1	-	5	14
Female	119	1	-	-	-	-	1	1	-	1	3	11
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	1	-	1	2	3
Male	18	-	-	-	-	-	-	1	-	-	1	3
Female	14	-	-	-	-	-	-	-	-	1	1	-
Georgia												
All races, both sexes	839	2	1	-	2	1	2	4	5	18	28	55
Male	401	1	-	-	2	1	-	3	2	8	15	27
Female	438	1	1	-	-	-	2	1	3	10	13	28
White, both sexes	493	1	1	-	1	1	1	3	-	8	10	23
Male	227	1	-	-	1	1	-	3	-	4	7	15
Female	266	-	1	-	-	-	1	-	-	4	3	8
Black, both sexes	323	1	-	-	1	-	1	1	5	9	18	31
Male	166	-	-	-	1	-	-	-	2	4	8	12
Female	157	1	-	-	-	-	1	1	3	5	10	19
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Hawaii												
All races, both sexes	160	-	-	-	-	-	1	-	2	2	2	6
Male	81	-	-	-	-	-	-	-	-	-	1	3
Female	79	-	-	-	-	-	1	-	2	2	1	3
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	132	-	-	-	-	-	1	-	2	2	2	6
Male	69	-	-	-	-	-	-	-	-	-	1	3
Female	63	-	-	-	-	-	1	-	2	2	1	3
Idaho												
All races, both sexes	91	-	-	-	-	-	-	-	-	2	3	2
Male	41	-	-	-	-	-	-	-	-	1	1	1
Female	50	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	90	-	-	-	-	-	-	-	-	2	3	2
Male	40	-	-	-	-	-	-	-	-	1	1	1
Female	50	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Illinois												
All races, both sexes	1,156	5	-	-	1	-	3	5	1	8	19	48
Male	519	4	-	-	1	-	-	2	-	6	9	30
Female	637	1	-	-	-	-	3	3	1	2	10	18
White, both sexes	938	3	-	-	1	-	2	1	1	1	12	23
Male	423	3	-	-	1	-	-	-	-	1	7	17
Female	515	-	-	-	-	-	2	1	1	-	5	6
Black, both sexes	194	2	-	-	-	-	1	4	-	7	6	23
Male	87	1	-	-	-	-	-	2	-	5	2	12
Female	107	1	-	-	-	-	1	2	-	2	4	11
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	593	3	-	1	-	-	1	2	6	6	14	19
Male	281	2	-	1	-	-	1	-	3	5	7	13
Female	312	1	-	-	-	-	-	2	3	1	7	6
White, both sexes	526	2	-	1	-	-	-	2	4	3	12	14
Male	242	1	-	1	-	-	-	-	2	2	6	8
Female	284	1	-	-	-	-	-	2	2	1	6	6
Black, both sexes	64	1	-	-	-	-	1	-	2	2	2	5
Male	38	1	-	-	-	-	1	-	1	2	1	5
Female	26	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Iowa												
All races, both sexes	273	1	-	-	-	-	-	-	-	-	1	6
Male	119	-	-	-	-	-	-	-	-	-	1	1
Female	154	1	-	-	-	-	-	-	-	-	-	5
White, both sexes	265	1	-	-	-	-	-	-	-	-	1	5
Male	115	-	-	-	-	-	-	-	-	-	1	-
Female	150	1	-	-	-	-	-	-	-	-	-	5
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	257	-	-	-	-	-	1	1	-	3	6	10
Male	128	-	-	-	-	-	-	-	-	3	6	6
Female	129	-	-	-	-	-	1	1	-	-	-	4
White, both sexes	231	-	-	-	-	-	1	-	-	1	6	7
Male	116	-	-	-	-	-	-	-	-	1	6	6
Female	115	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	2	-	1
Male	7	-	-	-	-	-	-	-	-	2	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Kentucky												
All races, both sexes	494	-	1	1	-	-	-	1	4	5	9	19
Male	214	-	-	-	-	-	-	-	2	1	5	11
Female	280	-	1	1	-	-	-	1	2	4	4	8
White, both sexes	453	-	1	1	-	-	-	1	4	3	7	13
Male	195	-	-	-	-	-	-	-	2	1	3	8
Female	258	-	1	1	-	-	-	1	2	2	4	5
Black, both sexes	39	-	-	-	-	-	-	-	-	1	2	6
Male	18	-	-	-	-	-	-	-	-	-	2	3
Female	21	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Louisiana												
All races, both sexes	422	-	-	-	-	2	1	2	7	5	13	23
Male	192	-	-	-	-	1	-	1	2	3	7	12
Female	230	-	-	-	-	1	1	1	5	2	6	11
White, both sexes	255	-	-	-	-	2	1	-	4	-	3	7
Male	101	-	-	-	-	1	-	-	1	-	2	4
Female	154	-	-	-	-	1	1	-	3	-	1	3
Black, both sexes	163	-	-	-	-	-	-	2	3	5	10	15
Male	87	-	-	-	-	-	-	1	1	3	5	7
Female	76	-	-	-	-	-	-	1	2	2	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Maine												
All races, both sexes	114	-	-	-	-	-	1	-	-	1	4	2
Male	55	-	-	-	-	-	1	-	-	1	3	2
Female	59	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	113	-	-	-	-	-	1	-	-	1	4	2
Male	55	-	-	-	-	-	1	-	-	1	3	2
Female	58	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	503	3	-	1	1	1	2	2	-	9	13	23
Male	246	1	-	-	1	1	2	1	-	2	6	13
Female	257	2	-	1	-	-	-	1	-	7	7	10
White, both sexes	334	2	-	-	1	-	1	-	-	3	7	8
Male	153	1	-	-	1	-	1	-	-	-	3	4
Female	181	1	-	-	-	-	-	-	-	3	4	4
Black, both sexes	153	1	-	1	-	1	1	2	-	6	6	15
Male	84	-	-	-	-	1	1	1	-	2	3	9
Female	69	1	-	1	-	-	-	1	-	4	3	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Massachusetts												
All races, both sexes	582	-	-	-	-	-	2	-	1	5	2	17
Male	249	-	-	-	-	-	2	-	-	3	1	9
Female	333	-	-	-	-	-	-	-	1	2	1	8
White, both sexes	535	-	-	-	-	-	1	-	1	2	2	11
Male	233	-	-	-	-	-	1	-	-	1	1	7
Female	302	-	-	-	-	-	-	-	1	1	1	4
Black, both sexes	25	-	-	-	-	-	1	-	-	3	-	5
Male	6	-	-	-	-	-	1	-	-	2	-	1
Female	19	-	-	-	-	-	-	-	-	1	-	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	956	1	-	-	-	1	3	3	4	6	14	19
Male	456	1	-	-	-	1	1	-	2	3	7	10
Female	500	-	-	-	-	-	2	3	2	3	7	9
White, both sexes	788	1	-	-	-	1	-	1	2	2	8	10
Male	370	1	-	-	-	1	-	-	1	1	5	5
Female	418	-	-	-	-	-	-	1	1	1	3	5
Black, both sexes	150	-	-	-	-	-	3	2	2	3	6	8
Male	78	-	-	-	-	-	1	-	1	2	2	4
Female	72	-	-	-	-	-	2	2	1	1	4	4
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Minnesota												
All races, both sexes	424	2	1	-	-	-	1	-	-	4	7	10
Male	198	1	-	-	-	-	-	-	-	2	5	8
Female	226	1	1	-	-	-	1	-	-	2	2	2
White, both sexes	404	2	-	-	-	-	-	-	-	4	6	9
Male	190	1	-	-	-	-	-	-	-	2	4	7
Female	214	1	-	-	-	-	-	-	-	2	2	2
Black, both sexes	9	-	-	-	-	-	1	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	1	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	1	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	275	-	-	1	-	2	1	-	-	-	13	21
Male	127	-	-	-	-	1	1	-	-	-	9	10
Female	148	-	-	1	-	1	-	-	-	-	4	11
White, both sexes	167	-	-	-	-	1	-	-	-	-	4	6
Male	76	-	-	-	-	1	-	-	-	-	4	3
Female	91	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	106	-	-	1	-	1	1	-	-	-	9	14
Male	50	-	-	-	-	-	1	-	-	-	5	6
Female	56	-	-	1	-	1	-	-	-	-	4	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Missouri												
All races, both sexes	624	2	-	-	1	-	-	1	3	9	7	17
Male	283	2	-	-	-	-	-	-	1	4	4	10
Female	341	-	-	-	1	-	-	1	2	5	3	7
White, both sexes	534	2	-	-	1	-	-	1	3	3	4	11
Male	236	2	-	-	-	-	-	-	1	-	3	7
Female	298	-	-	-	1	-	-	1	2	3	1	4
Black, both sexes	82	-	-	-	-	-	-	-	-	5	2	5
Male	44	-	-	-	-	-	-	-	-	4	1	2
Female	38	-	-	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	1	-
Montana												
All races, both sexes	91	-	-	-	-	-	-	-	-	-	1	2
Male	35	-	-	-	-	-	-	-	-	-	1	2
Female	56	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Nebraska												
All races, both sexes	157	-	2	-	-	1	-	-	-	2	7	5
Male	79	-	2	-	-	1	-	-	-	2	2	2
Female	78	-	-	-	-	-	-	-	-	-	5	3
White, both sexes	144	-	1	-	-	1	-	-	-	1	5	1
Male	71	-	1	-	-	1	-	-	-	1	-	1
Female	73	-	-	-	-	-	-	-	-	-	5	-
Black, both sexes	10	-	1	-	-	-	-	-	-	1	-	4
Male	5	-	1	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	2	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	187	2	-	-	-	-	1	-	2	5	8	11
Male	100	-	-	-	-	-	1	-	2	3	3	8
Female	87	2	-	-	-	-	-	-	-	2	5	3
White, both sexes	158	2	-	-	-	-	1	-	-	5	6	6
Male	84	-	-	-	-	-	1	-	-	3	3	5
Female	74	2	-	-	-	-	-	-	-	2	3	1
Black, both sexes	13	-	-	-	-	-	-	-	-	-	1	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	1	-	1	3
Male	7	-	-	-	-	-	-	-	1	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Hampshire												
All races, both sexes	112	1	-	-	-	-	-	-	-	-	1	3
Male	58	1	-	-	-	-	-	-	-	-	1	2
Female	54	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	108	1	-	-	-	-	-	-	-	-	1	3
Male	55	1	-	-	-	-	-	-	-	-	1	2
Female	53	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	885	3	-	-	-	1	1	4	4	14	15	32
Male	435	1	-	-	-	1	-	3	2	10	11	23
Female	450	2	-	-	-	-	1	1	2	4	4	9
White, both sexes	718	2	-	-	-	1	1	1	2	8	10	23
Male	342	-	-	-	-	1	-	1	1	6	8	18
Female	376	2	-	-	-	-	1	-	1	2	2	5
Black, both sexes	126	1	-	-	-	-	-	2	2	5	1	8
Male	72	1	-	-	-	-	-	1	1	3	-	4
Female	54	-	-	-	-	-	-	1	1	2	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	41	-	-	-	-	-	-	1	-	1	4	1
Male	21	-	-	-	-	-	-	1	-	1	3	1
Female	20	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Mexico												
All races, both sexes	125	1	-	1	-	-	-	-	-	1	-	5
Male	57	1	-	1	-	-	-	-	-	1	-	1
Female	68	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	115	1	-	-	-	-	-	-	-	1	-	4
Male	53	1	-	-	-	-	-	-	-	1	-	1
Female	62	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	1	-	-	-	-	-	-	-	1
Male	3	-	-	1	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,554	4	-	-	-	-	1	1	5	12	32	51
Male	731	4	-	-	-	-	1	1	4	9	20	28
Female	823	-	-	-	-	-	-	-	1	3	12	23
White, both sexes	1,260	1	-	-	-	-	1	-	1	6	15	33
Male	574	1	-	-	-	-	1	-	-	5	11	18
Female	686	-	-	-	-	-	-	-	1	1	4	15
Black, both sexes	237	3	-	-	-	-	-	-	3	4	17	13
Male	131	3	-	-	-	-	-	-	3	3	9	5
Female	106	-	-	-	-	-	-	-	-	1	8	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	56	-	-	-	-	-	-	1	1	2	-	5
Male	26	-	-	-	-	-	-	1	1	1	-	5
Female	30	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
North Carolina												
All races, both sexes	910	1	1	-	-	3	2	2	3	18	29	41
Male	441	1	-	-	-	-	1	1	1	13	19	25
Female	469	-	1	-	-	3	1	1	2	5	10	16
White, both sexes	645	-	1	-	-	2	2	1	1	6	8	18
Male	302	-	-	-	-	-	1	1	1	4	6	9
Female	343	-	1	-	-	2	1	-	-	2	2	9
Black, both sexes	244	1	-	-	-	1	-	1	2	12	16	23
Male	123	1	-	-	-	-	-	-	-	9	9	16
Female	121	-	-	-	-	1	-	1	2	3	7	7
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	4	-
Male	10	-	-	-	-	-	-	-	-	-	3	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	61	-	-	1	-	-	-	-	1	1	1	2
Male	31	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	1	-	-	-	-	1	1	1	1
White, both sexes	61	-	-	1	-	-	-	-	1	1	1	2
Male	31	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	1	-	-	-	-	1	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Ohio												
All races, both sexes	1,170	-	1	2	1	2	-	1	5	6	17	40
Male	523	-	1	2	1	-	-	-	3	4	10	20
Female	647	-	-	-	-	2	-	1	2	2	7	20
White, both sexes	1,008	-	1	1	1	2	-	-	5	3	14	29
Male	448	-	1	1	1	-	-	-	3	3	8	18
Female	560	-	-	-	-	2	-	-	2	-	6	11
Black, both sexes	153	-	-	1	-	-	-	1	-	3	3	10
Male	73	-	-	1	-	-	-	-	-	1	2	2
Female	80	-	-	-	-	-	-	1	-	2	1	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Oklahoma												
All races, both sexes	282	2	-	-	-	-	-	-	1	2	4	10
Male	127	1	-	-	-	-	-	-	-	2	1	7
Female	155	1	-	-	-	-	-	-	1	-	3	3
White, both sexes	244	1	-	-	-	-	-	-	-	2	3	8
Male	113	1	-	-	-	-	-	-	-	2	1	6
Female	131	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	22	1	-	-	-	-	-	-	1	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	13	1	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Oregon												
All races, both sexes	331	-	-	-	-	-	1	1	-	4	4	10
Male	150	-	-	-	-	-	1	1	-	4	1	4
Female	181	-	-	-	-	-	-	-	-	-	3	6
White, both sexes	313	-	-	-	-	-	-	1	-	3	3	7
Male	141	-	-	-	-	-	-	1	-	3	1	2
Female	172	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	1	-	-	-	1	2
Male	4	-	-	-	-	-	1	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	1
Pennsylvania												
All races, both sexes	1,556	2	1	1	-	1	2	5	7	5	31	47
Male	731	1	-	-	-	-	1	3	5	2	16	33
Female	825	1	1	1	-	1	1	2	2	3	15	14
White, both sexes	1,353	1	1	1	-	-	1	5	6	4	21	26
Male	624	-	-	-	-	-	-	3	4	2	12	21
Female	729	1	1	1	-	-	1	2	2	2	9	5
Black, both sexes	183	1	-	-	-	1	1	-	1	1	9	19
Male	97	1	-	-	-	-	1	-	1	-	4	11
Female	86	-	-	-	-	1	-	-	-	1	5	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Rhode Island												
All races, both sexes	106	-	-	-	-	-	-	-	1	1	-	2
Male	52	-	-	-	-	-	-	-	-	1	-	2
Female	54	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	104	-	-	-	-	-	-	-	1	-	-	2
Male	50	-	-	-	-	-	-	-	-	-	-	2
Female	54	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	524	2	-	-	-	-	-	-	4	10	19	33
Male	246	1	-	-	-	-	-	-	2	4	13	20
Female	278	1	-	-	-	-	-	-	2	6	6	13
White, both sexes	354	-	-	-	-	-	-	-	1	6	7	14
Male	165	-	-	-	-	-	-	-	-	3	4	9
Female	189	-	-	-	-	-	-	-	1	3	3	5
Black, both sexes	162	2	-	-	-	-	-	-	3	4	12	19
Male	75	1	-	-	-	-	-	-	2	1	9	11
Female	87	1	-	-	-	-	-	-	1	3	3	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
South Dakota												
All races, both sexes	70	-	-	-	-	-	-	-	-	-	2	3
Male	28	-	-	-	-	-	-	-	-	-	-	3
Female	42	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	63	-	-	-	-	-	-	-	-	-	1	2
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	39	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	663	3	1	-	1	-	2	1	3	9	17	32
Male	304	1	-	-	1	-	2	1	2	6	9	17
Female	359	2	1	-	-	-	-	-	1	3	8	15
White, both sexes	530	2	1	-	1	-	1	1	1	7	8	19
Male	237	-	-	-	1	-	1	1	1	4	4	9
Female	293	2	1	-	-	-	-	-	-	3	4	10
Black, both sexes	130	1	-	-	-	-	1	-	2	2	9	13
Male	66	1	-	-	-	-	1	-	1	2	5	8
Female	64	-	-	-	-	-	-	-	1	-	4	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Texas												
All races, both sexes	1,858	7	-	-	-	2	7	10	15	23	62	103
Male	890	2	-	-	-	1	5	7	8	16	34	53
Female	968	5	-	-	-	1	2	3	7	7	28	50
White, both sexes	1,549	6	-	-	-	2	7	8	9	15	38	69
Male	742	2	-	-	-	1	5	6	5	10	23	38
Female	807	4	-	-	-	1	2	2	4	5	15	31
Black, both sexes	247	1	-	-	-	-	-	2	5	8	21	31
Male	118	-	-	-	-	-	-	1	3	6	10	13
Female	129	1	-	-	-	-	-	1	2	2	11	18
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	-	3	2
Male	27	-	-	-	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	2	1
Utah												
All races, both sexes	99	-	-	-	-	-	-	-	2	-	1	1
Male	44	-	-	-	-	-	-	-	2	-	1	-
Female	55	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	92	-	-	-	-	-	-	-	1	-	1	1
Male	42	-	-	-	-	-	-	-	1	-	1	-
Female	50	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Vermont												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	716	2	1	-	1	2	-	1	4	6	18	21
Male	327	2	-	-	-	1	-	1	3	4	11	10
Female	389	-	1	-	1	1	-	-	1	2	7	11
White, both sexes	506	1	1	-	1	2	-	-	1	3	4	9
Male	225	1	-	-	-	1	-	-	-	1	2	3
Female	281	-	1	-	1	1	-	-	1	2	2	6
Black, both sexes	182	1	-	-	-	-	-	-	2	2	14	9
Male	91	1	-	-	-	-	-	-	2	2	9	5
Female	91	-	-	-	-	-	-	-	-	-	5	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	1	1	1	-	3
Male	11	-	-	-	-	-	-	1	1	1	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Washington												
All races, both sexes	531	1	-	-	-	-	-	-	2	3	7	15
Male	246	1	-	-	-	-	-	-	2	1	6	8
Female	285	-	-	-	-	-	-	-	-	2	1	7
White, both sexes	444	1	-	-	-	-	-	-	-	1	2	11
Male	205	1	-	-	-	-	-	-	-	1	1	5
Female	239	-	-	-	-	-	-	-	-	-	1	6
Black, both sexes	20	-	-	-	-	-	-	-	1	-	-	2
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	2	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	56	-	-	-	-	-	-	-	1	2	3	1
Male	25	-	-	-	-	-	-	-	1	-	3	1
Female	31	-	-	-	-	-	-	-	-	2	-	-
West Virginia												
All races, both sexes	206	-	-	-	-	-	-	1	-	1	1	6
Male	97	-	-	-	-	-	-	1	-	-	-	3
Female	109	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	200	-	-	-	-	-	-	1	-	1	1	5
Male	92	-	-	-	-	-	-	1	-	-	-	2
Female	108	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Wisconsin												
All races, both sexes	501	-	-	-	2	1	-	1	2	5	12	17
Male	225	-	-	-	2	-	-	-	1	3	7	10
Female	276	-	-	-	-	1	-	1	1	2	5	7
White, both sexes	461	-	-	-	2	1	-	1	2	4	11	14
Male	202	-	-	-	2	-	-	-	1	3	7	7
Female	259	-	-	-	-	1	-	1	1	1	4	7
Black, both sexes	31	-	-	-	-	-	-	-	-	1	1	3
Male	17	-	-	-	-	-	-	-	-	-	-	3
Female	14	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
United States												
All races, both sexes	1,609	1,806	1,747	1,936	2,670	3,902	4,818	4,023	1,880	442	55	-
Male	975	1,050	987	1,050	1,361	1,807	2,004	1,486	606	102	9	-
Female	634	756	760	886	1,309	2,095	2,814	2,537	1,274	340	46	-
White, both sexes	973	1,143	1,276	1,480	2,204	3,432	4,305	3,701	1,717	385	48	-
Male	594	642	707	820	1,128	1,604	1,834	1,372	553	88	7	-
Female	379	501	569	660	1,076	1,828	2,471	2,329	1,164	297	41	-
Black, both sexes	552	555	370	316	328	318	344	212	107	41	5	-
Male	330	341	225	155	169	131	114	71	26	7	1	-
Female	222	214	145	161	159	187	230	141	81	34	4	-
American Indian 1/, both sexes	18	14	8	16	12	13	18	6	6	3	-	-
Male	14	8	5	8	5	3	4	-	1	1	-	-
Female	4	6	3	8	7	10	14	6	5	2	-	-
Asian or Pacific Islander 2/, both sexes	66	94	93	124	126	139	151	104	50	13	2	-
Male	37	59	50	67	59	69	52	43	26	6	1	-
Female	29	35	43	57	67	70	99	61	24	7	1	-
Alabama												
All races, both sexes	46	46	37	39	64	77	84	76	32	7	1	-
Male	21	27	22	24	28	34	32	26	7	1	1	-
Female	25	19	15	15	36	43	52	50	25	6	-	-
White, both sexes	25	21	26	30	48	61	68	66	28	6	1	-
Male	10	11	15	19	21	29	26	22	6	1	1	-
Female	15	10	11	11	27	32	42	44	22	5	-	-
Black, both sexes	21	24	11	8	16	15	16	10	4	1	-	-
Male	11	15	7	5	7	5	6	4	1	-	-	-
Female	10	9	4	3	9	10	10	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Alaska												
All races, both sexes	5	8	5	3	5	5	7	4	-	-	-	-
Male	3	3	4	2	3	1	1	-	-	-	-	-
Female	2	5	1	1	2	4	6	4	-	-	-	-
White, both sexes	3	5	3	2	4	3	2	3	-	-	-	-
Male	1	2	2	2	3	1	-	-	-	-	-	-
Female	2	3	1	-	1	2	2	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	-	1	1	1	2	1	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	1	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	1	3	-	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Arizona												
All races, both sexes	22	20	33	24	47	59	61	62	17	2	1	-
Male	15	12	17	18	29	32	23	21	5	1	-	-
Female	7	8	16	6	18	27	38	41	12	1	1	-
White, both sexes	18	17	29	23	45	56	57	61	15	-	1	-
Male	13	11	15	17	28	31	22	21	4	-	-	-
Female	5	6	14	6	17	25	35	40	11	-	1	-
Black, both sexes	3	2	1	1	1	2	2	-	-	-	-	-
Male	1	1	-	1	1	1	-	-	-	-	-	-
Female	2	1	1	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	1	1	2	-	-	1	1	1	2	2	-	-
Male	1	-	2	-	-	-	1	-	1	1	-	-
Female	-	1	-	-	-	1	-	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Arkansas												
All races, both sexes	16	10	18	22	31	49	60	46	16	5	1	-
Male	5	7	7	5	14	21	27	17	6	2	-	-
Female	11	3	11	17	17	28	33	29	10	3	1	-
White, both sexes	10	8	11	21	30	43	50	40	15	3	1	-
Male	3	5	4	5	13	20	23	14	6	1	-	-
Female	7	3	7	16	17	23	27	26	9	2	1	-
Black, both sexes	6	1	6	1	1	5	9	6	1	2	-	-
Male	2	1	2	-	1	1	4	3	-	1	-	-
Female	4	-	4	1	-	4	5	3	1	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
California												
All races, both sexes	195	198	198	232	258	339	462	435	177	35	8	-
Male	122	121	110	132	131	161	190	173	70	7	2	-
Female	73	77	88	100	127	178	272	262	107	28	6	-
White, both sexes	120	130	135	158	187	252	374	374	151	29	6	-
Male	77	77	78	90	95	121	156	153	56	4	1	-
Female	43	53	57	68	92	131	218	221	95	25	5	-
Black, both sexes	44	31	30	22	19	19	21	15	4	1	-	-
Male	27	17	15	12	10	8	8	6	-	-	-	-
Female	17	14	15	10	9	11	13	9	4	1	-	-
American Indian 1/, both sexes	3	3	-	1	-	1	1	-	-	-	-	-
Male	2	2	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	34	33	51	52	67	66	46	22	5	2	-
Male	16	25	17	29	26	32	26	14	14	3	1	-
Female	12	9	16	22	26	35	40	32	8	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Colorado												
All races, both sexes	16	16	23	27	33	47	58	39	18	8	-	-
Male	13	8	10	15	17	21	19	15	9	1	-	-
Female	3	8	13	12	16	26	39	24	9	7	-	-
White, both sexes	13	15	20	25	29	47	56	38	18	7	-	-
Male	10	7	8	14	15	21	18	14	9	1	-	-
Female	3	8	12	11	14	26	38	24	9	6	-	-
Black, both sexes	2	1	2	1	2	-	2	-	-	1	-	-
Male	2	1	2	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	12	20	17	19	22	45	59	67	37	6	2	-
Male	9	9	14	9	13	24	20	20	9	1	-	-
Female	3	11	3	10	9	21	39	47	28	5	2	-
White, both sexes	9	16	14	13	20	44	58	67	35	6	2	-
Male	8	6	12	6	12	23	20	20	9	1	-	-
Female	1	10	2	7	8	21	38	47	26	5	2	-
Black, both sexes	1	1	2	4	2	1	1	-	2	-	-	-
Male	-	1	2	2	1	1	-	-	-	-	-	-
Female	1	-	-	2	1	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	-	2	-	-	-	-	-	-	-	-
Male	1	2	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Delaware												
All races, both sexes	5	5	2	6	12	9	15	16	3	3	-	-
Male	3	2	1	3	5	5	3	4	-	-	-	-
Female	2	3	1	3	7	4	12	12	3	3	-	-
White, both sexes	2	3	2	5	8	7	12	15	3	3	-	-
Male	1	1	1	3	4	4	3	4	-	-	-	-
Female	1	2	1	2	4	3	9	11	3	3	-	-
Black, both sexes	3	2	-	1	4	2	3	1	-	-	-	-
Male	2	1	-	-	1	1	-	-	-	-	-	-
Female	1	1	-	1	3	1	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	3	6	11	7	2	6	6	3	1	-	-
Male	3	3	1	4	6	1	4	1	2	-	-	-
Female	3	-	5	7	1	1	2	5	1	1	-	-
White, both sexes	-	-	2	3	1	-	1	1	1	1	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	-	2	2	-	-	-	1	-	1	-	-
Black, both sexes	6	3	4	8	6	2	5	5	2	-	-	-
Male	3	3	1	3	5	1	3	1	1	-	-	-
Female	3	-	3	5	1	1	2	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Florida												
All races, both sexes	109	111	131	116	188	263	343	237	115	25	3	-
Male	74	61	83	68	107	121	141	85	34	5	-	-
Female	35	50	48	48	81	142	202	152	81	20	3	-
White, both sexes	68	74	100	97	158	236	313	227	109	23	3	-
Male	44	39	66	60	91	113	127	82	33	4	-	-
Female	24	35	34	37	67	123	186	145	76	19	3	-
Black, both sexes	39	36	25	18	21	25	28	8	5	2	-	-
Male	28	21	14	7	13	7	12	2	1	1	-	-
Female	11	15	11	11	8	18	16	6	4	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	6	1	9	2	2	2	1	-	-	-
Male	1	1	3	1	3	1	2	1	-	-	-	-
Female	-	-	3	-	6	1	-	1	1	-	-	-
Georgia												
All races, both sexes	70	81	71	70	81	115	101	92	35	5	-	-
Male	47	42	38	37	42	50	44	36	6	-	-	-
Female	23	39	33	33	39	65	57	56	29	5	-	-
White, both sexes	17	27	36	44	47	82	84	77	27	3	-	-
Male	11	13	16	22	23	34	40	31	5	-	-	-
Female	6	14	20	22	24	48	44	46	22	3	-	-
Black, both sexes	49	54	35	21	30	28	15	15	7	2	-	-
Male	33	29	22	13	18	15	4	5	-	-	-	-
Female	16	25	13	8	12	13	11	10	7	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	5	4	4	1	-	1	-	-	-
Male	3	-	-	2	1	1	-	-	1	-	-	-
Female	1	-	-	3	3	3	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Hawaii												
All races, both sexes	5	16	8	14	12	22	29	25	13	3	-	-
Male	3	9	5	6	6	16	13	10	7	2	-	-
Female	2	7	3	8	6	6	16	15	6	1	-	-
White, both sexes	-	2	1	2	2	6	7	3	4	-	-	-
Male	-	1	-	2	1	5	2	-	1	-	-	-
Female	-	1	1	-	1	1	5	3	3	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	13	7	12	10	16	22	22	9	3	-	-
Male	3	8	5	4	5	11	11	10	6	2	-	-
Female	2	5	2	8	5	5	11	12	3	1	-	-
Idaho												
All races, both sexes	2	3	1	15	8	11	16	17	11	-	-	-
Male	-	2	-	7	6	6	9	5	3	-	-	-
Female	2	1	1	8	2	5	7	12	8	-	-	-
White, both sexes	2	3	1	15	8	11	15	17	11	-	-	-
Male	-	2	-	7	6	6	8	5	3	-	-	-
Female	2	1	1	8	2	5	7	12	8	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Illinois												
All races, both sexes	66	73	60	74	117	162	208	193	90	18	5	-
Male	32	37	31	40	50	72	93	74	30	6	2	-
Female	34	36	29	34	67	90	115	119	60	12	3	-
White, both sexes	30	49	39	56	99	147	183	187	82	17	5	-
Male	17	22	19	35	44	65	84	72	28	6	2	-
Female	13	27	20	21	55	82	99	115	54	11	3	-
Black, both sexes	32	22	19	16	16	15	17	6	7	1	-	-
Male	13	13	12	3	6	7	7	2	2	-	-	-
Female	19	9	7	13	10	8	10	4	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	2	2	2	-	8	-	1	-	-	-
Male	2	2	-	2	-	-	2	-	-	-	-	-
Female	2	-	2	-	2	-	6	-	1	-	-	-
Indiana												
All races, both sexes	33	40	37	44	66	89	93	107	27	5	-	-
Male	19	26	24	24	34	39	42	35	5	1	-	-
Female	14	14	13	20	32	50	51	72	22	4	-	-
White, both sexes	22	30	29	37	63	84	91	101	26	5	-	-
Male	13	16	20	21	33	38	42	34	4	1	-	-
Female	9	14	9	16	30	46	49	67	22	4	-	-
Black, both sexes	11	10	8	7	3	5	1	5	1	-	-	-
Male	6	10	4	3	1	1	-	1	1	-	-	-
Female	5	-	4	4	2	4	1	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Iowa												
All races, both sexes	10	16	15	17	31	38	55	52	22	9	-	-
Male	5	12	6	7	15	18	31	16	7	-	-	-
Female	5	4	9	10	16	20	24	36	15	9	-	-
White, both sexes	9	16	15	16	30	38	54	50	22	8	-	-
Male	5	12	6	7	14	18	30	15	7	-	-	-
Female	4	4	9	9	16	20	24	35	15	8	-	-
Black, both sexes	1	-	-	1	-	-	1	2	-	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	15	13	8	18	35	26	44	46	24	7	-	-
Male	9	9	3	8	17	16	16	18	14	3	-	-
Female	6	4	5	10	18	10	28	28	10	4	-	-
White, both sexes	13	10	7	16	33	22	39	45	24	7	-	-
Male	7	7	2	6	17	15	15	17	14	3	-	-
Female	6	3	5	10	16	7	24	28	10	4	-	-
Black, both sexes	1	-	-	1	1	3	5	1	-	-	-	-
Male	1	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	2	4	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	1	-	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Kentucky												
All races, both sexes	19	32	30	54	50	65	85	72	34	13	-	-
Male	15	17	13	25	31	24	31	24	10	5	-	-
Female	4	15	17	29	19	41	54	48	24	8	-	-
White, both sexes	13	27	29	49	47	63	82	69	32	12	-	-
Male	11	15	12	22	29	24	31	24	9	4	-	-
Female	2	12	17	27	18	39	51	45	23	8	-	-
Black, both sexes	6	5	1	5	2	2	3	3	2	1	-	-
Male	4	2	1	3	1	-	-	-	1	1	-	-
Female	2	3	-	2	1	2	3	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	29	29	27	31	43	68	60	47	29	5	1	-
Male	19	16	15	13	25	25	28	14	10	1	-	-
Female	10	13	12	18	18	43	32	33	19	4	1	-
White, both sexes	10	14	13	16	26	53	46	32	23	4	1	-
Male	5	7	6	5	14	19	20	9	8	-	-	-
Female	5	7	7	11	12	34	26	23	15	4	1	-
Black, both sexes	19	15	14	14	17	15	14	14	5	1	-	-
Male	14	9	9	7	11	6	8	4	1	1	-	-
Female	5	6	5	7	6	9	6	10	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Maine												
All races, both sexes	7	5	6	6	13	21	13	24	8	3	-	-
Male	5	3	4	2	6	10	8	8	1	1	-	-
Female	2	2	2	4	7	11	5	16	7	2	-	-
White, both sexes	7	5	6	6	13	20	13	24	8	3	-	-
Male	5	3	4	2	6	10	8	8	1	1	-	-
Female	2	2	2	4	7	10	5	16	7	2	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	37	37	34	29	43	65	94	76	27	5	1	-
Male	24	23	21	12	26	38	38	31	6	-	-	-
Female	13	14	13	17	17	27	56	45	21	5	1	-
White, both sexes	13	11	22	17	28	48	78	69	23	3	-	-
Male	8	6	13	5	16	27	33	29	6	-	-	-
Female	5	5	9	12	12	21	45	40	17	3	-	-
Black, both sexes	24	25	11	10	13	13	13	6	2	2	1	-
Male	16	16	8	5	8	8	4	2	-	-	-	-
Female	8	9	3	5	5	5	9	4	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	2	4	3	1	2	-	-	-
Male	-	1	-	2	2	3	1	-	-	-	-	-
Female	-	-	1	-	-	1	2	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Massachusetts												
All races, both sexes	14	25	24	32	51	110	125	109	52	13	-	-
Male	10	16	10	14	24	48	52	36	19	5	-	-
Female	4	9	14	18	27	62	73	73	33	8	-	-
White, both sexes	12	25	21	29	46	100	118	104	52	11	-	-
Male	9	16	9	12	23	45	51	35	19	4	-	-
Female	3	9	12	17	23	55	67	69	33	7	-	-
Black, both sexes	-	-	-	1	2	5	4	3	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	2	4	3	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	3	2	2	4	3	2	-	1	-	-
Male	1	-	1	2	1	2	-	1	-	1	-	-
Female	1	-	2	-	1	2	3	1	-	-	-	-
Michigan												
All races, both sexes	51	61	51	67	96	149	200	128	79	22	1	-
Male	26	42	28	40	55	63	91	56	25	4	1	-
Female	25	19	23	27	41	86	109	72	54	18	-	-
White, both sexes	29	36	38	57	81	132	181	115	73	20	1	-
Male	12	22	19	34	46	58	87	49	24	4	1	-
Female	17	14	19	23	35	74	94	66	49	16	-	-
Black, both sexes	20	24	12	9	13	14	16	11	5	2	-	-
Male	13	20	9	6	7	3	3	6	1	-	-	-
Female	7	4	3	3	6	11	13	5	4	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	2	1	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	2	2	1	1	1	-	-	-
Male	-	-	-	-	2	1	-	1	-	-	-	-
Female	1	1	1	1	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Minnesota												
All races, both sexes	16	20	25	29	31	65	81	74	51	6	1	-
Male	13	12	14	14	11	29	33	30	26	-	-	-
Female	3	8	11	15	20	36	48	44	25	6	1	-
White, both sexes	14	18	23	24	30	65	79	72	51	6	1	-
Male	11	12	14	11	11	29	33	29	26	-	-	-
Female	3	6	9	13	19	36	46	43	25	6	1	-
Black, both sexes	1	1	2	2	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	2	2	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	-	-	1	2	-	-	-	-
Male	-	-	-	3	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Mississippi												
All races, both sexes	21	29	11	15	32	42	44	22	16	4	1	-
Male	12	14	9	7	13	20	16	10	3	2	-	-
Female	9	15	2	8	19	22	28	12	13	2	1	-
White, both sexes	10	9	7	7	26	35	35	15	8	3	1	-
Male	5	2	5	4	12	16	15	5	2	2	-	-
Female	5	7	2	3	14	19	20	10	6	1	1	-
Black, both sexes	11	20	4	8	5	7	9	7	8	1	-	-
Male	7	12	4	3	1	4	1	5	1	-	-	-
Female	4	8	-	5	4	3	8	2	7	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Missouri												
All races, both sexes	34	35	40	51	60	105	101	84	58	15	1	-
Male	23	21	24	34	25	38	43	28	23	3	-	-
Female	11	14	16	17	35	67	58	56	35	12	1	-
White, both sexes	18	21	28	43	54	99	92	82	57	14	1	-
Male	12	13	17	30	23	37	39	27	22	3	-	-
Female	6	8	11	13	31	62	53	55	35	11	1	-
Black, both sexes	15	12	12	7	6	5	9	2	1	1	-	-
Male	11	7	7	4	2	-	4	1	1	-	-	-
Female	4	5	5	3	4	5	5	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	7	3	4	3	7	14	20	17	13	-	-	-
Male	5	-	1	1	2	7	9	4	3	-	-	-
Female	2	3	3	2	5	7	11	13	10	-	-	-
White, both sexes	5	3	4	2	4	14	19	17	13	-	-	-
Male	4	-	1	1	2	7	9	4	3	-	-	-
Female	1	3	3	1	2	7	10	13	10	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	3	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	3	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Nebraska												
All races, both sexes	10	11	8	4	12	29	30	23	8	4	1	-
Male	6	9	4	1	8	15	13	11	2	1	-	-
Female	4	2	4	3	4	14	17	12	6	3	1	-
White, both sexes	7	11	7	3	12	29	30	23	8	4	1	-
Male	5	9	3	-	8	15	13	11	2	1	-	-
Female	2	2	4	3	4	14	17	12	6	3	1	-
Black, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	21	13	14	18	20	28	13	12	2	1	-
Male	10	13	9	10	7	10	16	3	4	1	-	-
Female	6	8	4	4	11	10	12	10	8	1	1	-
White, both sexes	12	17	9	12	17	17	26	13	12	2	1	-
Male	8	11	6	9	7	8	15	3	4	1	-	-
Female	4	6	3	3	10	9	11	10	8	1	1	-
Black, both sexes	3	1	2	-	1	2	1	-	-	-	-	-
Male	1	-	2	-	-	1	-	-	-	-	-	-
Female	2	1	-	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	1	-	1	-	-	-	-	-	-
Male	1	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Hampshire												
All races, both sexes	5	3	3	11	10	16	19	23	12	3	2	-
Male	3	-	3	9	5	9	10	10	4	-	1	-
Female	2	3	-	2	5	7	9	13	8	3	1	-
White, both sexes	2	3	3	10	10	16	19	23	12	3	2	-
Male	1	-	3	8	5	9	10	10	4	-	1	-
Female	1	3	-	2	5	7	9	13	8	3	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	48	54	48	57	82	114	184	131	80	13	-	-
Male	31	29	36	30	37	58	78	55	25	5	-	-
Female	17	25	12	27	45	56	106	76	55	8	-	-
White, both sexes	31	31	31	38	63	101	163	125	76	11	-	-
Male	21	14	21	22	29	52	71	51	22	4	-	-
Female	10	17	10	16	34	49	92	74	54	7	-	-
Black, both sexes	15	16	12	16	15	10	14	4	3	2	-	-
Male	9	10	11	7	6	5	7	3	3	1	-	-
Female	6	6	1	9	9	5	7	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	7	5	3	4	3	7	2	1	-	-	-
Male	1	5	4	1	2	1	-	1	-	-	-	-
Female	1	2	1	2	2	2	7	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Mexico												
All races, both sexes	7	8	8	10	22	20	21	12	8	1	-	-
Male	3	4	6	6	13	9	9	2	1	-	-	-
Female	4	4	2	4	9	11	12	10	7	1	-	-
White, both sexes	7	8	6	7	21	19	21	12	7	1	-	-
Male	3	4	5	5	13	8	9	2	1	-	-	-
Female	4	4	1	2	8	11	12	10	6	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	2	-	-	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
New York												
All races, both sexes	70	95	103	107	171	241	276	246	104	32	3	-
Male	41	60	54	61	87	113	122	90	30	6	-	-
Female	29	35	49	46	84	128	154	156	74	26	3	-
White, both sexes	45	63	69	76	136	202	252	234	94	29	3	-
Male	24	36	34	45	70	97	114	85	28	5	-	-
Female	21	27	35	31	66	105	138	149	66	24	3	-
Black, both sexes	24	28	29	23	26	32	19	7	6	3	-	-
Male	17	22	18	13	13	14	7	2	1	1	-	-
Female	7	6	11	10	13	18	12	5	5	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	5	7	9	7	5	5	4	-	-	-
Male	-	2	2	3	4	2	1	3	1	-	-	-
Female	1	2	3	4	5	5	4	2	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
North Carolina												
All races, both sexes	70	71	60	62	100	144	132	103	55	12	1	-
Male	35	47	39	36	55	68	49	37	12	2	-	-
Female	35	24	21	26	45	76	83	66	43	10	1	-
White, both sexes	29	34	38	40	82	123	113	90	46	11	-	-
Male	17	22	24	22	48	56	43	36	10	2	-	-
Female	12	12	14	18	34	67	70	54	36	9	-	-
Black, both sexes	38	37	21	19	17	20	16	10	8	1	1	-
Male	15	25	14	11	6	11	4	1	1	-	-	-
Female	23	12	7	8	11	9	12	9	7	1	1	-
American Indian 1/, both sexes	3	-	-	1	1	1	2	1	-	-	-	-
Male	3	-	-	1	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	-	1	2	1	-	-	-
Male	-	-	1	2	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
North Dakota												
All races, both sexes	1	2	1	4	5	9	14	8	10	-	1	-
Male	-	2	1	3	2	5	8	5	4	-	-	-
Female	1	-	-	1	3	4	6	3	6	-	1	-
White, both sexes	1	2	1	4	5	9	14	8	10	-	1	-
Male	-	2	1	3	2	5	8	5	4	-	-	-
Female	1	-	-	1	3	4	6	3	6	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Ohio												
All races, both sexes	64	72	79	93	124	157	219	186	78	21	2	-
Male	32	42	43	50	57	69	86	71	26	6	-	-
Female	32	30	36	43	67	88	133	115	52	15	2	-
White, both sexes	39	45	63	77	110	145	201	178	72	20	2	-
Male	17	26	36	39	52	64	81	69	23	6	-	-
Female	22	19	27	38	58	81	120	109	49	14	2	-
Black, both sexes	25	27	13	15	12	11	17	8	6	1	-	-
Male	15	16	6	11	4	5	5	2	3	-	-	-
Female	10	11	7	4	8	6	12	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	-	2	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	2	-	1	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	13	18	18	10	30	35	66	44	26	3	-	-
Male	9	15	10	5	14	13	32	13	5	-	-	-
Female	4	3	8	5	16	22	34	31	21	3	-	-
White, both sexes	9	11	16	8	27	31	61	41	24	2	-	-
Male	6	9	10	4	12	13	32	12	5	-	-	-
Female	3	2	6	4	15	18	29	29	19	2	-	-
Black, both sexes	2	6	1	-	3	1	1	1	2	-	-	-
Male	1	5	-	-	2	-	-	-	-	-	-	-
Female	1	1	1	-	1	1	1	1	2	-	-	-
American Indian 1/, both sexes	2	1	-	1	-	3	3	1	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	3	3	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Oregon												
All races, both sexes	18	20	14	24	24	45	78	60	17	8	3	-
Male	12	7	10	13	16	23	32	17	5	4	-	-
Female	6	13	4	11	8	22	46	43	12	4	3	-
White, both sexes	15	16	14	24	23	44	77	58	17	8	3	-
Male	10	7	10	13	15	23	31	16	5	4	-	-
Female	5	9	4	11	8	21	46	42	12	4	3	-
Black, both sexes	3	1	-	-	1	1	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	3	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	79	85	88	94	145	242	303	279	103	28	8	-
Male	46	48	56	62	80	117	128	98	28	7	-	-
Female	33	37	32	32	65	125	175	181	75	21	8	-
White, both sexes	59	55	69	75	130	224	282	262	98	26	7	-
Male	35	28	42	50	68	110	122	93	27	7	-	-
Female	24	27	27	25	62	114	160	169	71	19	7	-
Black, both sexes	19	27	17	15	13	17	19	16	4	2	1	-
Male	11	19	12	9	10	6	6	5	1	-	-	-
Female	8	8	5	6	3	11	13	11	3	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	4	2	1	2	1	1	-	-	-
Male	-	-	2	3	2	1	-	-	-	-	-	-
Female	1	2	-	1	-	-	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Rhode Island												
All races, both sexes	6	5	1	3	10	17	23	22	10	5	-	-
Male	6	4	-	2	5	8	8	11	3	2	-	-
Female	-	1	1	1	5	9	15	11	7	3	-	-
White, both sexes	6	5	1	3	10	17	23	22	9	5	-	-
Male	6	4	-	2	5	8	8	11	2	2	-	-
Female	-	1	1	1	5	9	15	11	7	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	33	54	33	46	34	64	96	58	31	6	1	-
Male	21	26	19	25	14	28	38	22	13	-	-	-
Female	12	28	14	21	20	36	58	36	18	6	1	-
White, both sexes	18	23	22	30	27	47	83	45	27	4	-	-
Male	12	10	13	18	11	22	33	17	13	-	-	-
Female	6	13	9	12	16	25	50	28	14	4	-	-
Black, both sexes	15	29	11	15	7	14	13	11	4	2	1	-
Male	9	14	6	6	3	5	5	3	-	-	-	-
Female	6	15	5	9	4	9	8	8	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	-	3	-	2	-	-	-	-
Male	-	2	-	1	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
South Dakota												
All races, both sexes	2	3	4	1	8	11	13	14	6	3	-	-
Male	1	1	2	1	2	4	6	6	2	-	-	-
Female	1	2	2	-	6	7	7	8	4	3	-	-
White, both sexes	2	2	4	-	6	11	12	14	6	3	-	-
Male	1	-	2	-	1	4	6	6	2	-	-	-
Female	1	2	2	-	5	7	6	8	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	2	-	1	-	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	46	54	58	50	72	91	102	62	44	13	2	-
Male	26	32	29	24	33	41	34	28	16	-	2	-
Female	20	22	29	26	39	50	68	34	28	13	-	-
White, both sexes	26	34	37	42	63	89	92	56	40	9	1	-
Male	17	21	15	20	27	40	32	28	15	-	1	-
Female	9	13	22	22	36	49	60	28	25	9	-	-
Black, both sexes	20	20	20	8	8	2	10	5	4	4	1	-
Male	9	11	14	4	5	1	2	-	1	-	1	-
Female	11	9	6	4	3	1	8	5	3	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Texas												
All races, both sexes	150	143	147	113	154	270	296	223	106	25	2	-
Male	94	82	78	51	80	139	117	79	37	7	-	-
Female	56	61	69	62	74	131	179	144	69	18	2	-
White, both sexes	111	103	118	91	130	247	267	206	97	23	2	-
Male	73	63	65	45	64	124	107	70	34	7	-	-
Female	38	40	53	46	66	123	160	136	63	16	2	-
Black, both sexes	36	32	23	15	18	10	20	15	9	1	-	-
Male	20	17	9	3	13	6	6	8	3	-	-	-
Female	16	15	14	12	5	4	14	7	6	1	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	8	5	6	6	13	8	2	-	1	-	-
Male	1	2	3	2	3	9	4	1	-	-	-	-
Female	2	6	2	4	3	4	4	1	-	1	-	-
Utah												
All races, both sexes	4	3	8	10	7	20	16	18	9	-	-	-
Male	3	3	3	4	5	8	6	7	2	-	-	-
Female	1	-	5	6	2	12	10	11	7	-	-	-
White, both sexes	4	3	7	8	6	19	15	18	9	-	-	-
Male	3	3	3	4	5	7	6	7	2	-	-	-
Female	1	-	4	4	1	12	9	11	7	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Vermont												
All races, both sexes	1	4	1	-	4	12	12	5	6	-	-	-
Male	-	2	-	-	1	6	6	3	2	-	-	-
Female	1	2	1	-	3	6	6	2	4	-	-	-
White, both sexes	1	4	1	-	4	12	12	5	6	-	-	-
Male	-	2	-	-	1	6	6	3	2	-	-	-
Female	1	2	1	-	3	6	6	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	42	55	42	45	76	101	120	118	43	18	-	-
Male	27	27	26	24	37	48	46	41	13	6	-	-
Female	15	28	16	21	39	53	74	77	30	12	-	-
White, both sexes	17	22	29	26	53	88	100	99	37	13	-	-
Male	9	9	19	14	29	44	41	36	11	5	-	-
Female	8	13	10	12	24	44	59	63	26	8	-	-
Black, both sexes	24	31	13	17	21	9	16	14	5	4	-	-
Male	17	17	7	10	8	3	4	3	2	1	-	-
Female	7	14	6	7	13	6	12	11	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	2	2	4	4	4	1	1	-	-
Male	1	1	-	-	-	1	1	2	-	-	-	-
Female	-	1	-	2	2	3	3	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Washington												
All races, both sexes	20	29	39	51	56	81	97	79	44	7	-	-
Male	13	19	21	29	25	32	44	31	13	1	-	-
Female	7	10	18	22	31	49	53	48	31	6	-	-
White, both sexes	12	22	27	41	44	77	86	77	38	5	-	-
Male	8	14	13	25	19	31	44	30	12	1	-	-
Female	4	8	14	16	25	46	42	47	26	4	-	-
Black, both sexes	3	3	3	2	2	1	3	-	-	-	-	-
Male	3	2	1	-	1	1	-	-	-	-	-	-
Female	-	1	2	2	1	-	3	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	-	1	-	3	-	-	-
Male	-	-	-	1	2	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	3	-	-	-
Asian or Pacific Islander 2/, both sexes	4	4	9	7	8	3	7	2	3	2	-	-
Male	2	3	7	3	3	-	-	1	1	-	-	-
Female	2	1	2	4	5	3	7	1	2	2	-	-
West Virginia												
All races, both sexes	12	8	12	22	15	38	39	31	17	3	-	-
Male	8	7	4	14	9	22	12	12	5	-	-	-
Female	4	1	8	8	6	16	27	19	12	3	-	-
White, both sexes	12	7	12	20	15	37	39	30	17	3	-	-
Male	8	6	4	12	9	21	12	12	5	-	-	-
Female	4	1	8	8	6	16	27	18	12	3	-	-
Black, both sexes	-	1	-	1	-	1	-	1	-	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Wisconsin												
All races, both sexes	24	32	31	32	47	60	103	81	40	10	1	-
Male	13	16	16	18	28	20	44	32	12	3	-	-
Female	11	16	15	14	19	40	59	49	28	7	1	-
White, both sexes	16	26	25	27	42	57	103	80	40	9	1	-
Male	8	13	11	16	25	19	44	31	12	3	-	-
Female	8	13	14	11	17	38	59	49	28	6	1	-
Black, both sexes	6	6	5	2	4	2	-	-	-	1	-	-
Male	4	3	5	-	2	-	-	-	-	-	-	-
Female	2	3	-	2	2	2	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	3	1	1	-	1	-	-	-	-
Male	1	-	-	2	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	6	5	1	3	7	11	4	-	-	-
Male	-	1	3	1	-	2	3	5	2	-	-	-
Female	-	-	3	4	1	1	4	6	2	-	-	-
White, both sexes	-	1	6	5	1	3	7	11	4	-	-	-
Male	-	1	3	1	-	2	3	5	2	-	-	-
Female	-	-	3	4	1	1	4	6	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
United States												
All races, both sexes	7,037	11	6	1	2	1	6	13	27	43	93	137
Male	2,981	7	5	-	2	-	4	10	17	24	53	80
Female	4,056	4	1	1	-	1	2	3	10	19	40	57
White, both sexes	6,055	10	5	1	2	-	6	11	17	27	69	91
Male	2,505	6	4	-	2	-	4	8	11	14	40	48
Female	3,550	4	1	1	-	-	2	3	6	13	29	43
Black, both sexes	815	1	1	-	-	1	-	2	7	14	24	42
Male	383	1	1	-	-	-	-	2	4	9	13	28
Female	432	-	-	-	-	1	-	-	3	5	11	14
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	-	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	3
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	145	-	-	-	-	-	-	-	3	2	-	1
Male	81	-	-	-	-	-	-	-	2	1	-	1
Female	64	-	-	-	-	-	-	-	1	1	-	-
Alabama												
All races, both sexes	131	-	-	-	1	-	-	-	1	-	3	5
Male	54	-	-	-	1	-	-	-	1	-	1	1
Female	77	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	101	-	-	-	1	-	-	-	-	-	2	3
Male	37	-	-	-	1	-	-	-	-	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	30	-	-	-	-	-	-	-	1	-	1	2
Male	17	-	-	-	-	-	-	-	1	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Alaska												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	116	1	-	-	-	-	-	1	1	2	2	1
Male	55	1	-	-	-	-	-	-	-	1	2	1
Female	61	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	111	1	-	-	-	-	-	1	1	2	2	1
Male	53	1	-	-	-	-	-	-	-	1	2	1
Female	58	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Arkansas												
All races, both sexes	87	-	-	-	-	-	-	-	2	1	2	1
Male	39	-	-	-	-	-	-	-	2	1	2	-
Female	48	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	73	-	-	-	-	-	-	-	1	-	1	1
Male	30	-	-	-	-	-	-	-	1	-	1	-
Female	43	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	637	2	-	-	-	-	2	1	2	6	9	12
Male	279	2	-	-	-	-	1	1	-	3	6	4
Female	358	-	-	-	-	-	1	-	2	3	3	8
White, both sexes	536	2	-	-	-	-	2	1	1	4	8	9
Male	229	2	-	-	-	-	1	1	-	2	5	3
Female	307	-	-	-	-	-	1	-	1	2	3	6
Black, both sexes	43	-	-	-	-	-	-	-	-	2	1	3
Male	17	-	-	-	-	-	-	-	-	1	1	1
Female	26	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	56	-	-	-	-	-	-	-	1	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Colorado												
All races, both sexes	53	-	1	-	-	-	-	-	-	-	1	1
Male	25	-	1	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	1	-	-	-	-	-	-	-	1	1
Male	24	-	1	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	107	-	-	-	-	-	-	-	-	-	2	2
Male	47	-	-	-	-	-	-	-	-	-	1	1
Female	60	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	98	-	-	-	-	-	-	-	-	-	2	1
Male	40	-	-	-	-	-	-	-	-	-	1	-
Female	58	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Delaware												
All races, both sexes	22	-	-	-	-	-	-	1	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	1	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Florida												
All races, both sexes	522	-	-	-	-	-	-	2	4	3	7	5
Male	219	-	-	-	-	-	-	2	4	2	3	3
Female	303	-	-	-	-	-	-	-	-	1	4	2
White, both sexes	447	-	-	-	-	-	-	2	2	2	5	5
Male	185	-	-	-	-	-	-	2	2	1	2	3
Female	262	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	69	-	-	-	-	-	-	-	1	1	2	-
Male	32	-	-	-	-	-	-	-	1	1	1	-
Female	37	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	150	1	-	-	-	-	-	-	1	5	6	5
Male	71	1	-	-	-	-	-	-	-	3	3	4
Female	79	-	-	-	-	-	-	-	1	2	3	1
White, both sexes	112	-	-	-	-	-	-	-	-	3	5	4
Male	52	-	-	-	-	-	-	-	-	1	3	4
Female	60	-	-	-	-	-	-	-	-	2	2	-
Black, both sexes	37	1	-	-	-	-	-	-	1	2	1	1
Male	18	1	-	-	-	-	-	-	-	2	-	-
Female	19	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Hawaii												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	29	-	-	-	-	-	1	-	-	-	-	1
Male	12	-	-	-	-	-	1	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	-	-	-	-	-	1	-	-	-	-	1
Male	11	-	-	-	-	-	1	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Illinois												
All races, both sexes	316	-	-	1	-	-	-	-	-	1	9	2
Male	138	-	-	-	-	-	-	-	-	1	5	2
Female	178	-	-	1	-	-	-	-	-	-	4	-
White, both sexes	256	-	-	1	-	-	-	-	-	-	6	2
Male	105	-	-	-	-	-	-	-	-	-	3	2
Female	151	-	-	1	-	-	-	-	-	-	3	-
Black, both sexes	55	-	-	-	-	-	-	-	-	1	3	-
Male	29	-	-	-	-	-	-	-	-	1	2	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	156	-	-	-	-	-	1	-	-	-	1	3
Male	70	-	-	-	-	-	1	-	-	-	1	2
Female	86	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	146	-	-	-	-	-	1	-	-	-	1	2
Male	65	-	-	-	-	-	1	-	-	-	1	1
Female	81	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Iowa												
All races, both sexes	104	-	-	-	-	-	-	-	-	-	-	1
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	104	-	-	-	-	-	-	-	-	-	-	1
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	69	-	-	-	-	-	-	-	1	-	-	3
Male	31	-	-	-	-	-	-	-	1	-	-	3
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	1	-	-	-
Male	27	-	-	-	-	-	-	-	1	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Kentucky												
All races, both sexes	101	-	-	-	-	-	-	1	-	-	-	3
Male	41	-	-	-	-	-	-	1	-	-	-	2
Female	60	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	96	-	-	-	-	-	-	-	-	-	-	3
Male	37	-	-	-	-	-	-	-	-	-	-	2
Female	59	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	79	-	1	-	-	-	-	-	-	-	1	2
Male	30	-	1	-	-	-	-	-	-	-	-	1
Female	49	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	57	-	1	-	-	-	-	-	-	-	-	-
Male	21	-	1	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Maine												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	183	-	-	-	-	-	-	-	1	-	-	3
Male	68	-	-	-	-	-	-	-	-	-	-	2
Female	115	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	143	-	-	-	-	-	-	-	1	-	-	2
Male	54	-	-	-	-	-	-	-	-	-	-	2
Female	89	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Massachusetts												
All races, both sexes	154	-	-	-	-	-	-	-	1	1	4	2
Male	61	-	-	-	-	-	-	-	1	-	3	2
Female	93	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	144	-	-	-	-	-	-	-	1	1	3	1
Male	56	-	-	-	-	-	-	-	1	-	3	1
Female	88	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	318	1	-	-	-	-	-	-	-	5	4	4
Male	126	-	-	-	-	-	-	-	-	4	2	1
Female	192	1	-	-	-	-	-	-	-	1	2	3
White, both sexes	281	1	-	-	-	-	-	-	-	3	3	2
Male	108	-	-	-	-	-	-	-	-	3	1	1
Female	173	1	-	-	-	-	-	-	-	-	2	1
Black, both sexes	37	-	-	-	-	-	-	-	-	2	1	2
Male	18	-	-	-	-	-	-	-	-	1	1	-
Female	19	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Minnesota												
All races, both sexes	130	-	-	-	-	-	-	-	-	-	1	2
Male	52	-	-	-	-	-	-	-	-	-	-	2
Female	78	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	123	-	-	-	-	-	-	-	-	-	1	1
Male	49	-	-	-	-	-	-	-	-	-	-	1
Female	74	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	84	-	-	-	-	-	-	-	1	-	1	3
Male	36	-	-	-	-	-	-	-	1	-	-	2
Female	48	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	62	-	-	-	-	-	-	-	1	-	1	2
Male	29	-	-	-	-	-	-	-	1	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Missouri												
All races, both sexes	185	-	-	-	-	-	-	-	1	1	2	4
Male	83	-	-	-	-	-	-	-	1	1	1	2
Female	102	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	164	-	-	-	-	-	-	-	1	1	1	3
Male	72	-	-	-	-	-	-	-	1	1	1	1
Female	92	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Nebraska												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	1	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	25	-	-	-	-	-	-	1	-	1	-	-
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	22	-	-	-	-	-	-	1	-	-	-	-
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
New Hampshire												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	144	-	-	-	-	-	-	-	2	-	3	5
Male	57	-	-	-	-	-	-	-	-	-	-	5
Female	87	-	-	-	-	-	-	-	2	-	3	-
White, both sexes	114	-	-	-	-	-	-	-	1	-	1	5
Male	46	-	-	-	-	-	-	-	-	-	-	5
Female	68	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	23	-	-	-	-	-	-	-	1	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
New Mexico												
All races, both sexes	27	-	1	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	26	-	1	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	384	-	-	-	1	-	-	1	1	2	3	7
Male	164	-	-	-	1	-	-	1	1	-	2	2
Female	220	-	-	-	-	-	-	-	-	2	1	5
White, both sexes	338	-	-	-	1	-	-	1	1	1	2	4
Male	139	-	-	-	1	-	-	1	1	-	1	-
Female	199	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	42	-	-	-	-	-	-	-	-	1	1	3
Male	22	-	-	-	-	-	-	-	-	-	1	2
Female	20	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
North Carolina												
All races, both sexes	267	-	-	-	-	-	-	-	2	1	6	5
Male	109	-	-	-	-	-	-	-	1	-	5	4
Female	158	-	-	-	-	-	-	-	1	1	1	1
White, both sexes	211	-	-	-	-	-	-	-	1	-	5	2
Male	85	-	-	-	-	-	-	-	-	-	5	1
Female	126	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	53	-	-	-	-	-	-	-	1	1	1	3
Male	24	-	-	-	-	-	-	-	1	-	-	3
Female	29	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Ohio												
All races, both sexes	410	-	-	-	-	-	-	2	-	2	4	6
Male	162	-	-	-	-	-	-	1	-	2	2	4
Female	248	-	-	-	-	-	-	1	-	-	2	2
White, both sexes	361	-	-	-	-	-	-	2	-	-	3	4
Male	135	-	-	-	-	-	-	1	-	-	2	2
Female	226	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	46	-	-	-	-	-	-	-	-	2	1	2
Male	26	-	-	-	-	-	-	-	-	2	-	2
Female	20	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	91	-	1	-	-	-	-	-	-	-	1	2
Male	31	-	1	-	-	-	-	-	-	-	-	2
Female	60	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	1	2
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	55	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Oregon												
All races, both sexes	84	-	-	-	-	-	-	-	-	1	2	1
Male	37	-	-	-	-	-	-	-	-	-	1	1
Female	47	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	84	-	-	-	-	-	-	-	-	1	2	1
Male	37	-	-	-	-	-	-	-	-	-	1	1
Female	47	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	409	2	-	-	-	-	-	-	1	2	1	2
Male	161	-	-	-	-	-	-	-	-	2	-	1
Female	248	2	-	-	-	-	-	-	1	-	1	1
White, both sexes	375	2	-	-	-	-	-	-	1	2	-	1
Male	152	-	-	-	-	-	-	-	-	2	-	1
Female	223	2	-	-	-	-	-	-	1	-	-	-
Black, both sexes	33	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	120	-	-	-	-	-	-	-	-	3	3	2
Male	47	-	-	-	-	-	-	-	-	-	3	1
Female	73	-	-	-	-	-	-	-	-	3	-	1
White, both sexes	80	-	-	-	-	-	-	-	-	2	1	1
Male	30	-	-	-	-	-	-	-	-	-	1	-
Female	50	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	40	-	-	-	-	-	-	-	-	1	2	1
Male	17	-	-	-	-	-	-	-	-	-	2	1
Female	23	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
South Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	191	-	-	-	-	-	1	1	1	-	1	4
Male	80	-	-	-	-	-	1	1	1	-	1	3
Female	111	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	161	-	-	-	-	-	1	-	-	-	-	1
Male	63	-	-	-	-	-	1	-	-	-	-	1
Female	98	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	30	-	-	-	-	-	-	1	1	-	1	3
Male	17	-	-	-	-	-	-	1	1	-	1	2
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Texas												
All races, both sexes	395	3	1	-	-	1	1	1	1	-	5	17
Male	177	2	1	-	-	-	-	1	1	-	3	10
Female	218	1	-	-	-	1	1	-	-	-	2	7
White, both sexes	334	3	1	-	-	-	1	1	1	-	4	12
Male	147	2	1	-	-	-	-	1	1	-	2	6
Female	187	1	-	-	-	-	1	-	-	-	2	6
Black, both sexes	55	-	-	-	-	1	-	-	-	-	1	5
Male	29	-	-	-	-	-	-	-	-	-	1	4
Female	26	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Vermont												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	161	-	-	-	-	-	-	1	1	-	1	2
Male	65	-	-	-	-	-	-	1	-	-	1	-
Female	96	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	127	-	-	-	-	-	-	1	-	-	-	-
Male	51	-	-	-	-	-	-	1	-	-	-	-
Female	76	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	30	-	-	-	-	-	-	-	1	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Washington												
All races, both sexes	159	1	-	-	-	-	-	-	1	3	3	2
Male	75	1	-	-	-	-	-	-	1	2	2	-
Female	84	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	147	1	-	-	-	-	-	-	1	2	3	2
Male	67	1	-	-	-	-	-	-	1	1	2	-
Female	80	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	-	4
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	48	-	-	-	-	-	-	-	-	1	-	4
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	24	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Wisconsin												
All races, both sexes	132	-	1	-	-	-	-	-	-	2	-	2
Male	62	-	1	-	-	-	-	-	-	2	-	2
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	122	-	1	-	-	-	-	-	-	2	-	-
Male	57	-	1	-	-	-	-	-	-	2	-	-
Female	65	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
United States												
All races, both sexes	198	262	341	412	566	945	1,305	1,354	927	324	63	-
Male	123	168	201	233	308	449	523	474	236	60	4	-
Female	75	94	140	179	258	496	782	880	691	264	59	-
White, both sexes	132	201	259	329	473	820	1,152	1,239	856	303	52	-
Male	80	127	154	188	264	385	462	432	214	59	3	-
Female	52	74	105	141	209	435	690	807	642	244	49	-
Black, both sexes	59	55	71	66	76	96	126	87	57	19	11	-
Male	40	36	40	34	34	47	47	31	15	-	1	-
Female	19	19	31	32	42	49	79	56	42	19	10	-
American Indian 1/, both sexes	2	1	1	3	3	4	1	3	1	-	-	-
Male	1	-	1	1	2	2	1	1	-	-	-	-
Female	1	1	-	2	1	2	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	5	10	14	14	25	26	25	13	2	-	-
Male	2	5	6	10	8	15	13	10	7	1	-	-
Female	3	-	4	4	6	10	13	15	6	1	-	-
Alabama												
All races, both sexes	7	6	8	6	15	10	25	27	10	5	2	-
Male	4	5	7	4	6	2	11	8	2	1	-	-
Female	3	1	1	2	9	8	14	19	8	4	2	-
White, both sexes	3	3	4	6	13	7	21	24	8	5	1	-
Male	2	2	3	4	5	2	9	7	1	1	-	-
Female	1	1	1	2	8	5	12	17	7	4	1	-
Black, both sexes	4	3	4	-	2	3	4	3	2	-	1	-
Male	2	3	4	-	1	-	2	1	1	-	-	-
Female	2	-	-	-	1	3	2	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Alaska												
All races, both sexes	-	-	-	2	-	2	1	-	2	-	-	-
Male	-	-	-	2	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	4	5	11	8	14	23	21	12	4	1	-
Male	5	3	4	6	4	6	8	11	3	-	-	-
Female	-	1	1	5	4	8	15	10	9	4	1	-
White, both sexes	5	4	4	10	8	13	22	20	12	4	1	-
Male	5	3	3	6	4	6	8	10	3	-	-	-
Female	-	1	1	4	4	7	14	10	9	4	1	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Arkansas												
All races, both sexes	4	3	3	6	9	18	13	15	8	2	-	-
Male	1	-	-	5	7	8	3	8	2	-	-	-
Female	3	3	3	1	2	10	10	7	6	2	-	-
White, both sexes	3	1	3	6	7	17	11	14	6	2	-	-
Male	-	-	-	5	6	7	2	7	1	-	-	-
Female	3	1	3	1	1	10	9	7	5	2	-	-
Black, both sexes	1	2	-	-	2	1	2	1	1	-	-	-
Male	1	-	-	-	1	1	1	1	-	-	-	-
Female	-	2	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12	18	30	44	51	96	111	124	88	26	3	-
Male	8	15	16	24	30	40	44	53	24	7	1	-
Female	4	3	14	20	21	56	67	71	64	19	2	-
White, both sexes	9	16	19	32	44	78	102	106	77	24	2	-
Male	6	13	11	16	26	32	41	44	19	6	1	-
Female	3	3	8	16	18	46	61	62	58	18	1	-
Black, both sexes	2	1	6	6	2	4	5	4	5	1	1	-
Male	1	1	2	4	1	1	2	1	1	-	-	-
Female	1	-	4	2	1	3	3	3	4	1	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	5	5	5	14	4	13	6	1	-	-
Male	1	1	3	3	3	7	1	7	4	1	-	-
Female	-	-	2	2	2	7	3	6	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Colorado												
All races, both sexes	1	6	3	3	4	3	10	11	6	2	1	-
Male	1	3	1	1	2	3	3	5	2	1	-	-
Female	-	3	2	2	2	-	7	6	4	1	1	-
White, both sexes	1	6	2	3	4	3	10	11	6	2	1	-
Male	1	3	-	1	2	3	3	5	2	1	-	-
Female	-	3	2	2	2	-	7	6	4	1	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	3	2	5	11	11	15	29	16	8	1	-
Male	1	3	1	4	8	6	5	11	4	2	-	-
Female	1	-	1	1	3	5	10	18	12	6	1	-
White, both sexes	-	3	-	5	8	10	15	29	16	8	1	-
Male	-	3	-	4	5	5	5	11	4	2	-	-
Female	-	-	-	1	3	5	10	18	12	6	1	-
Black, both sexes	2	-	2	-	3	1	-	-	-	-	-	-
Male	1	-	1	-	3	1	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Delaware												
All races, both sexes	-	3	2	3	1	3	3	2	2	-	1	-
Male	-	3	1	1	-	1	1	1	1	-	-	-
Female	-	-	1	2	1	2	2	1	1	-	1	-
White, both sexes	-	3	1	2	1	2	3	2	2	-	1	-
Male	-	3	-	-	-	1	1	1	1	-	-	-
Female	-	-	1	2	1	1	2	1	1	-	1	-
Black, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	2	1	1	2	1	1	1	-	-
Male	-	-	1	2	-	1	2	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
White, both sexes	-	-	1	-	1	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
Black, both sexes	-	-	-	2	-	1	2	1	1	-	-	-
Male	-	-	-	2	-	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Florida												
All races, both sexes	12	19	34	19	52	76	106	91	65	23	4	-
Male	8	12	17	10	24	33	43	34	18	6	-	-
Female	4	7	17	9	28	43	63	57	47	17	4	-
White, both sexes	11	13	28	15	36	66	94	81	61	22	4	-
Male	7	9	14	7	19	28	36	31	18	6	-	-
Female	4	4	14	8	17	38	58	50	43	16	4	-
Black, both sexes	1	6	5	4	14	10	12	8	4	1	-	-
Male	1	3	2	3	5	5	7	3	-	-	-	-
Female	-	3	3	1	9	5	5	5	4	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	2	-	-	-	-
Georgia												
All races, both sexes	5	12	11	12	7	16	21	27	13	7	1	-
Male	4	6	6	4	6	12	8	10	3	1	-	-
Female	1	6	5	8	1	4	13	17	10	6	1	-
White, both sexes	2	8	8	10	5	12	15	24	11	4	1	-
Male	1	3	4	4	5	8	6	9	3	1	-	-
Female	1	5	4	6	-	4	9	15	8	3	1	-
Black, both sexes	3	4	3	2	2	3	6	3	2	3	-	-
Male	3	3	2	-	1	3	2	1	-	-	-	-
Female	-	1	1	2	1	-	4	2	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Hawaii												
All races, both sexes	1	2	2	3	5	2	10	5	5	-	-	-
Male	1	2	-	3	3	1	4	2	3	-	-	-
Female	-	-	2	-	2	1	6	3	2	-	-	-
White, both sexes	-	-	-	-	2	1	2	1	1	-	-	-
Male	-	-	-	-	2	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	3	3	1	8	4	4	-	-	-
Male	1	2	-	3	1	1	4	1	2	-	-	-
Female	-	-	2	-	2	-	4	3	2	-	-	-
Idaho												
All races, both sexes	-	1	-	2	1	2	4	10	6	1	-	-
Male	-	-	-	1	1	2	-	5	1	1	-	-
Female	-	1	-	1	-	-	4	5	5	-	-	-
White, both sexes	-	1	-	2	1	2	4	9	6	1	-	-
Male	-	-	-	1	1	2	-	4	1	1	-	-
Female	-	1	-	1	-	-	4	5	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Illinois												
All races, both sexes	8	12	14	17	23	41	59	69	44	12	4	-
Male	5	6	12	12	11	20	28	21	12	2	1	-
Female	3	6	2	5	12	21	31	48	32	10	3	-
White, both sexes	4	8	9	13	17	32	48	64	39	11	2	-
Male	2	2	7	10	9	17	22	19	10	2	-	-
Female	2	6	2	3	8	15	26	45	29	9	2	-
Black, both sexes	4	4	4	4	6	9	7	5	5	1	2	-
Male	3	4	4	2	2	3	3	2	2	-	1	-
Female	1	-	-	2	4	6	4	3	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	4	-	-	-	-	-
Male	-	-	1	-	-	-	3	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	4	8	14	11	8	20	35	23	23	5	-	-
Male	2	7	9	4	6	5	14	9	9	1	-	-
Female	2	1	5	7	2	15	21	14	14	4	-	-
White, both sexes	3	7	11	9	8	18	35	23	23	5	-	-
Male	1	6	8	3	6	5	14	9	9	1	-	-
Female	2	1	3	6	2	13	21	14	14	4	-	-
Black, both sexes	1	1	3	2	-	2	-	-	-	-	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Iowa												
All races, both sexes	2	2	4	1	9	9	18	31	25	2	-	-
Male	2	1	3	-	4	3	6	13	5	-	-	-
Female	-	1	1	1	5	6	12	18	20	2	-	-
White, both sexes	2	2	4	1	9	9	18	31	25	2	-	-
Male	2	1	3	-	4	3	6	13	5	-	-	-
Female	-	1	1	1	5	6	12	18	20	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	3	3	4	4	9	11	13	13	2	-	-
Male	2	2	1	3	2	3	3	4	6	1	-	-
Female	1	1	2	1	2	6	8	9	7	1	-	-
White, both sexes	3	2	2	3	3	9	10	13	13	2	-	-
Male	2	2	1	3	1	3	3	4	6	1	-	-
Female	1	-	1	-	2	6	7	9	7	1	-	-
Black, both sexes	-	1	1	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Kentucky												
All races, both sexes	-	-	5	9	13	9	18	24	16	3	-	-
Male	-	-	3	4	8	3	7	10	3	-	-	-
Female	-	-	2	5	5	6	11	14	13	3	-	-
White, both sexes	-	-	5	9	12	8	17	23	16	3	-	-
Male	-	-	3	4	7	3	6	9	3	-	-	-
Female	-	-	2	5	5	5	11	14	13	3	-	-
Black, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	2	4	11	5	8	20	11	10	1	-	-
Male	2	-	3	4	1	5	8	4	-	1	-	-
Female	1	2	1	7	4	3	12	7	10	-	-	-
White, both sexes	2	1	3	6	3	8	12	10	10	1	-	-
Male	1	-	2	3	1	5	4	3	-	1	-	-
Female	1	1	1	3	2	3	8	7	10	-	-	-
Black, both sexes	1	1	1	5	2	-	8	1	-	-	-	-
Male	1	-	1	1	-	-	4	1	-	-	-	-
Female	-	1	-	4	2	-	4	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Maine												
All races, both sexes	1	-	3	1	2	5	7	3	1	2	-	-
Male	-	-	1	-	2	3	4	1	-	1	-	-
Female	1	-	2	1	-	2	3	2	1	1	-	-
White, both sexes	1	-	3	1	2	5	7	3	1	2	-	-
Male	-	-	1	-	2	3	4	1	-	1	-	-
Female	1	-	2	1	-	2	3	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	5	5	14	4	21	33	40	33	15	5	-
Male	4	2	2	9	2	7	16	12	9	3	-	-
Female	-	3	3	5	2	14	17	28	24	12	5	-
White, both sexes	1	2	2	7	4	19	27	34	27	13	4	-
Male	1	2	2	5	2	7	12	10	8	3	-	-
Female	-	-	-	2	2	12	15	24	19	10	4	-
Black, both sexes	3	3	3	5	-	2	5	5	6	2	1	-
Male	3	-	-	2	-	-	3	1	1	-	-	-
Female	-	3	3	3	-	2	2	4	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	1	1	-	-	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Massachusetts												
All races, both sexes	3	4	2	5	6	19	38	33	25	8	3	-
Male	-	2	2	3	4	9	15	11	5	4	-	-
Female	3	2	-	2	2	10	23	22	20	4	3	-
White, both sexes	3	3	2	3	5	19	34	33	25	8	3	-
Male	-	1	2	2	3	9	14	11	5	4	-	-
Female	3	2	-	1	2	10	20	22	20	4	3	-
Black, both sexes	-	-	-	1	1	-	4	-	-	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	6	9	11	22	30	44	53	67	46	11	5	-
Male	4	2	6	17	19	19	16	26	8	2	-	-
Female	2	7	5	5	11	25	37	41	38	9	5	-
White, both sexes	4	7	9	18	27	38	47	64	43	10	5	-
Male	2	1	5	13	17	16	15	25	7	2	-	-
Female	2	6	4	5	10	22	32	39	36	8	5	-
Black, both sexes	2	2	2	4	3	6	6	3	3	1	-	-
Male	2	1	1	4	2	3	1	1	1	-	-	-
Female	-	1	1	-	1	3	5	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Minnesota												
All races, both sexes	2	4	5	2	13	13	26	31	16	12	3	-
Male	2	4	4	1	10	5	8	10	4	2	-	-
Female	-	-	1	1	3	8	18	21	12	10	3	-
White, both sexes	2	4	4	1	12	13	26	29	16	11	3	-
Male	2	4	3	1	9	5	8	10	4	2	-	-
Female	-	-	1	-	3	8	18	19	12	9	3	-
Black, both sexes	-	-	1	1	-	-	-	1	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	7	6	8	8	12	9	17	6	2	1	-
Male	1	4	3	5	3	8	1	8	-	-	-	-
Female	2	3	3	3	5	4	8	9	6	2	1	-
White, both sexes	3	5	4	4	5	11	5	14	5	1	1	-
Male	1	2	3	3	2	8	1	7	-	-	-	-
Female	2	3	1	1	3	3	4	7	5	1	1	-
Black, both sexes	-	2	2	4	3	1	4	3	1	1	-	-
Male	-	2	-	2	1	-	-	1	-	-	-	-
Female	-	-	2	2	2	1	4	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Missouri												
All races, both sexes	7	7	12	15	18	32	42	21	14	7	2	-
Male	4	3	7	7	11	15	23	4	4	-	-	-
Female	3	4	5	8	7	17	19	17	10	7	2	-
White, both sexes	3	5	9	14	17	30	40	18	13	7	2	-
Male	1	3	5	6	10	14	22	3	4	-	-	-
Female	2	2	4	8	7	16	18	15	9	7	2	-
Black, both sexes	4	1	3	-	1	1	2	3	1	-	-	-
Male	3	-	2	-	1	1	1	1	-	-	-	-
Female	1	1	1	-	-	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	2	3	1	2	3	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	1	-	-	-	1	3	1	1	3	-	-	-
White, both sexes	1	-	-	-	1	2	1	2	3	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	2	1	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Nebraska												
All races, both sexes	2	3	-	2	6	4	7	2	5	4	-	-
Male	2	3	-	-	5	2	3	1	3	-	-	-
Female	-	-	-	2	1	2	4	1	2	4	-	-
White, both sexes	2	3	-	2	5	4	7	2	5	4	-	-
Male	2	3	-	-	5	2	3	1	3	-	-	-
Female	-	-	-	2	-	2	4	1	2	4	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	2	5	1	7	2	3	2	-	-	-
Male	-	-	1	1	1	2	1	1	1	-	-	-
Female	-	1	1	4	-	5	1	2	1	-	-	-
White, both sexes	-	-	2	4	1	7	2	3	2	-	-	-
Male	-	-	1	1	1	2	1	1	1	-	-	-
Female	-	-	1	3	-	5	1	2	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
New Hampshire												
All races, both sexes	1	-	1	1	3	2	6	8	6	-	-	-
Male	1	-	-	-	1	1	4	4	1	-	-	-
Female	-	-	1	1	2	1	2	4	5	-	-	-
White, both sexes	1	-	1	1	3	2	6	8	6	-	-	-
Male	1	-	-	-	1	1	4	4	1	-	-	-
Female	-	-	1	1	2	1	2	4	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	6	7	6	13	25	30	21	15	8	-	-
Male	2	3	5	4	9	11	11	5	1	1	-	-
Female	1	3	2	2	4	14	19	16	14	7	-	-
White, both sexes	2	5	4	3	9	20	24	20	13	7	-	-
Male	2	2	4	2	7	9	8	5	1	1	-	-
Female	-	3	-	1	2	11	16	15	12	6	-	-
Black, both sexes	-	1	3	3	3	3	5	1	1	-	-	-
Male	-	1	1	2	1	1	2	-	-	-	-	-
Female	-	-	2	1	2	2	3	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	-	1	1	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
New Mexico												
All races, both sexes	1	-	2	1	3	5	3	4	4	-	-	-
Male	1	-	1	-	3	3	1	2	-	-	-	-
Female	-	-	1	1	-	2	2	2	4	-	-	-
White, both sexes	1	-	2	1	3	5	3	3	4	-	-	-
Male	1	-	1	-	3	3	1	2	-	-	-	-
Female	-	-	1	1	-	2	2	1	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	18	14	24	34	47	70	89	42	21	4	-
Male	3	14	10	18	16	26	26	28	10	6	-	-
Female	3	4	4	6	18	21	44	61	32	15	4	-
White, both sexes	3	15	9	18	29	43	65	81	40	21	4	-
Male	1	11	7	15	14	23	24	24	10	6	-	-
Female	2	4	2	3	15	20	41	57	30	15	4	-
Black, both sexes	3	3	5	6	4	3	3	8	2	-	-	-
Male	2	3	3	3	1	2	1	4	-	-	-	-
Female	1	-	2	3	3	1	2	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
North Carolina												
All races, both sexes	8	8	9	19	16	42	46	56	36	11	2	-
Male	6	5	7	10	7	18	18	20	7	-	1	-
Female	2	3	2	9	9	24	28	36	29	11	1	-
White, both sexes	5	5	5	13	13	31	41	49	30	10	1	-
Male	4	3	4	8	6	13	18	17	5	-	1	-
Female	1	2	1	5	7	18	23	32	25	10	-	-
Black, both sexes	3	3	4	5	2	11	4	7	6	1	1	-
Male	2	2	3	2	1	5	-	3	2	-	-	-
Female	1	1	1	3	1	6	4	4	4	1	1	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	2	3	1	1	1	4	2	1	-	-
Male	-	1	1	1	1	1	-	2	-	-	-	-
Female	-	-	1	2	-	-	1	2	2	1	-	-
White, both sexes	-	1	2	3	1	1	1	4	2	1	-	-
Male	-	1	1	1	1	1	-	2	-	-	-	-
Female	-	-	1	2	-	-	1	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Ohio												
All races, both sexes	14	10	15	17	33	52	91	86	54	20	4	-
Male	7	8	9	6	17	36	32	22	14	2	-	-
Female	7	2	6	11	16	16	59	64	40	18	4	-
White, both sexes	8	8	13	15	29	42	81	82	52	19	3	-
Male	4	6	8	6	15	27	28	21	13	2	-	-
Female	4	2	5	9	14	15	53	61	39	17	3	-
Black, both sexes	5	2	2	2	3	10	10	3	2	1	1	-
Male	3	2	1	-	1	9	4	1	1	-	-	-
Female	2	-	1	2	2	1	6	2	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	5	1	6	5	4	11	10	16	23	3	3	-
Male	3	-	6	3	1	3	4	4	4	-	-	-
Female	2	1	-	2	3	8	6	12	19	3	3	-
White, both sexes	4	1	6	5	4	11	7	15	21	3	2	-
Male	2	-	6	3	1	3	2	4	4	-	-	-
Female	2	1	-	2	3	8	5	11	17	3	2	-
Black, both sexes	-	-	-	-	-	-	2	1	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Oregon												
All races, both sexes	-	2	5	2	6	16	20	17	8	4	-	-
Male	-	1	3	2	3	8	10	5	2	1	-	-
Female	-	1	2	-	3	8	10	12	6	3	-	-
White, both sexes	-	2	5	2	6	16	20	17	8	4	-	-
Male	-	1	3	2	3	8	10	5	2	1	-	-
Female	-	1	2	-	3	8	10	12	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	12	14	15	17	32	47	81	90	60	28	5	-
Male	7	9	10	11	20	21	34	29	14	3	-	-
Female	5	5	5	6	12	26	47	61	46	25	5	-
White, both sexes	11	13	14	16	29	44	71	84	56	27	4	-
Male	7	8	9	11	18	20	32	28	13	3	-	-
Female	4	5	5	5	11	24	39	56	43	24	4	-
Black, both sexes	1	1	1	1	3	3	9	6	4	1	1	-
Male	-	1	1	-	2	1	1	1	1	-	-	-
Female	1	-	-	1	1	2	8	5	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Rhode Island												
All races, both sexes	-	-	3	-	1	2	3	1	4	1	-	-
Male	-	-	2	-	-	2	2	-	-	1	-	-
Female	-	-	1	-	1	-	1	1	4	-	-	-
White, both sexes	-	-	2	-	1	2	3	1	4	1	-	-
Male	-	-	1	-	-	2	2	-	-	1	-	-
Female	-	-	1	-	1	-	1	1	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	8	6	7	3	17	23	21	19	3	2	-
Male	2	5	4	4	1	6	7	8	6	-	-	-
Female	1	3	2	3	2	11	16	13	13	3	2	-
White, both sexes	-	4	5	4	1	13	15	17	15	1	1	-
Male	-	2	3	2	1	5	4	7	5	-	-	-
Female	-	2	2	2	-	8	11	10	10	1	1	-
Black, both sexes	3	4	1	3	2	4	8	4	4	2	1	-
Male	2	3	1	2	-	1	3	1	1	-	-	-
Female	1	1	-	1	2	3	5	3	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
South Dakota												
All races, both sexes	-	-	-	1	1	3	1	3	3	2	-	-
Male	-	-	-	1	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	3	3	2	-	-
White, both sexes	-	-	-	1	1	3	1	3	3	2	-	-
Male	-	-	-	1	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	3	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	10	9	17	21	20	26	33	24	16	1	-
Male	3	6	7	7	8	9	13	12	6	2	-	-
Female	3	4	2	10	13	11	13	21	18	14	1	-
White, both sexes	3	7	7	14	19	17	22	31	23	16	-	-
Male	1	4	5	6	8	7	11	12	5	2	-	-
Female	2	3	2	8	11	10	11	19	18	14	-	-
Black, both sexes	3	3	2	3	2	3	4	2	1	-	1	-
Male	2	2	2	1	-	2	2	-	1	-	-	-
Female	1	1	-	2	2	1	2	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Texas												
All races, both sexes	17	16	23	19	33	68	63	61	43	19	3	-
Male	11	10	11	12	19	30	30	14	18	3	1	-
Female	6	6	12	7	14	38	33	47	25	16	2	-
White, both sexes	8	11	15	18	27	59	58	54	39	19	3	-
Male	6	6	8	11	15	26	30	12	16	3	1	-
Female	2	5	7	7	12	33	28	42	23	16	2	-
Black, both sexes	9	5	8	1	6	8	3	5	3	-	-	-
Male	5	4	3	1	4	3	-	2	2	-	-	-
Female	4	1	5	-	2	5	3	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	1	-	-	-
Utah												
All races, both sexes	2	1	1	-	2	5	4	2	2	2	-	-
Male	1	1	-	-	-	-	1	1	-	2	-	-
Female	1	-	1	-	2	5	3	1	2	-	-	-
White, both sexes	2	1	1	-	2	5	4	1	2	2	-	-
Male	1	1	-	-	-	-	1	1	-	2	-	-
Female	1	-	1	-	2	5	3	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Vermont												
All races, both sexes	-	1	1	1	3	5	2	1	5	-	-	-
Male	-	1	-	1	2	4	1	1	2	-	-	-
Female	-	-	1	-	1	1	1	-	3	-	-	-
White, both sexes	-	1	1	1	3	5	2	1	5	-	-	-
Male	-	1	-	1	2	4	1	1	2	-	-	-
Female	-	-	1	-	1	1	1	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	4	10	9	16	23	36	28	15	7	1	-
Male	3	3	2	7	9	13	11	8	7	-	-	-
Female	4	1	8	2	7	10	25	20	8	7	1	-
White, both sexes	4	4	9	9	12	17	27	24	14	5	1	-
Male	2	3	2	7	7	8	8	7	6	-	-	-
Female	2	1	7	2	5	9	19	17	8	5	1	-
Black, both sexes	2	-	1	-	3	4	9	4	1	2	-	-
Male	1	-	-	-	1	3	3	1	1	-	-	-
Female	1	-	1	-	2	1	6	3	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Washington												
All races, both sexes	7	7	10	5	14	21	34	23	20	8	-	-
Male	3	5	6	1	6	13	21	10	1	3	-	-
Female	4	2	4	4	8	8	13	13	19	5	-	-
White, both sexes	5	5	9	5	12	18	33	23	20	8	-	-
Male	2	4	5	1	5	11	20	10	1	3	-	-
Female	3	1	4	4	7	7	13	13	19	5	-	-
Black, both sexes	1	1	-	-	2	1	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	-	2	1	-	-	-	-	-
Male	-	1	1	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	4	3	5	5	7	7	8	2	1	-	-
Male	2	3	2	3	2	6	1	2	1	-	-	-
Female	-	1	1	2	3	1	6	6	1	1	-	-
White, both sexes	2	4	3	5	4	7	7	8	2	1	-	-
Male	2	3	2	3	2	6	1	2	1	-	-	-
Female	-	1	1	2	2	1	6	6	1	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Wisconsin												
All races, both sexes	5	6	8	8	5	15	24	28	23	4	1	-
Male	4	4	4	5	1	10	10	11	8	-	-	-
Female	1	2	4	3	4	5	14	17	15	4	1	-
White, both sexes	4	6	7	8	4	15	24	25	22	3	1	-
Male	4	4	3	5	-	10	10	10	8	-	-	-
Female	-	2	4	3	4	5	14	15	14	3	1	-
Black, both sexes	1	-	1	-	1	-	-	3	1	1	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	-	-	-	1	1	1	1	-	-	-
Male	1	1	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	1	1	1	1	-	-	-
Male	1	1	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
United States												
All races, both sexes	72,476	9	2	1	4	7	12	27	60	130	291	586
Male	27,241	6	1	1	1	4	9	19	30	79	166	330
Female	45,235	3	1	-	3	3	3	8	30	51	125	256
White, both sexes	62,065	6	2	1	1	7	8	10	33	81	179	376
Male	22,844	3	1	1	1	4	7	6	20	53	110	219
Female	39,221	3	1	-	-	3	1	4	13	28	69	157
Black, both sexes	8,582	3	-	-	2	-	4	15	26	45	105	196
Male	3,589	3	-	-	-	-	2	11	10	25	50	105
Female	4,993	-	-	-	2	-	2	4	16	20	55	91
American Indian 1/, both sexes	253	-	-	-	-	-	-	-	1	1	3	4
Male	102	-	-	-	-	-	-	-	-	-	3	2
Female	151	-	-	-	-	-	-	-	1	1	-	2
Asian or Pacific Islander 2/, both sexes	1,576	-	-	-	1	-	-	2	-	3	4	10
Male	706	-	-	-	-	-	-	2	-	1	3	4
Female	870	-	-	-	1	-	-	-	-	2	1	6
Alabama												
All races, both sexes	1,636	1	-	-	-	-	-	2	2	7	15	21
Male	659	1	-	-	-	-	-	2	1	5	6	8
Female	977	-	-	-	-	-	-	-	1	2	9	13
White, both sexes	1,161	-	-	-	-	-	-	-	1	4	7	12
Male	440	-	-	-	-	-	-	-	1	3	4	5
Female	721	-	-	-	-	-	-	-	-	1	3	7
Black, both sexes	468	1	-	-	-	-	-	2	1	3	8	9
Male	215	1	-	-	-	-	-	2	-	2	2	3
Female	253	-	-	-	-	-	-	-	1	1	6	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Alaska												
All races, both sexes	83	-	-	-	-	-	-	-	-	1	2	1
Male	41	-	-	-	-	-	-	-	-	-	2	-
Female	42	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	62	-	-	-	-	-	-	-	-	-	2	1
Male	32	-	-	-	-	-	-	-	-	-	2	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,258	-	-	-	-	-	-	-	1	7	7	7
Male	473	-	-	-	-	-	-	-	1	5	7	5
Female	785	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	1,198	-	-	-	-	-	-	-	1	6	7	6
Male	440	-	-	-	-	-	-	-	1	5	7	4
Female	758	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	40	-	-	-	-	-	-	-	-	1	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Arkansas												
All races, both sexes	1,135	-	-	-	-	-	1	-	2	3	8	11
Male	422	-	-	-	-	-	1	-	-	3	6	7
Female	713	-	-	-	-	-	-	-	2	-	2	4
White, both sexes	940	-	-	-	-	-	-	-	-	2	4	3
Male	338	-	-	-	-	-	-	-	-	2	2	3
Female	602	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	186	-	-	-	-	-	1	-	2	1	4	8
Male	83	-	-	-	-	-	1	-	-	1	4	4
Female	103	-	-	-	-	-	-	-	2	-	-	4
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,155	-	-	-	1	-	1	5	7	17	22	64
Male	2,738	-	-	-	-	-	1	5	4	9	16	34
Female	4,417	-	-	-	1	-	-	-	3	8	6	30
White, both sexes	5,905	-	-	-	-	-	1	2	5	12	12	48
Male	2,205	-	-	-	-	-	1	2	3	6	8	23
Female	3,700	-	-	-	-	-	-	-	2	6	4	25
Black, both sexes	524	-	-	-	1	-	-	1	2	2	8	10
Male	211	-	-	-	-	-	-	1	1	2	6	8
Female	313	-	-	-	1	-	-	-	1	-	2	2
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	702	-	-	-	-	-	-	2	-	3	1	6
Male	310	-	-	-	-	-	-	2	-	1	1	3
Female	392	-	-	-	-	-	-	-	-	2	-	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Colorado												
All races, both sexes	790	1	-	-	-	-	-	-	-	2	3	5
Male	257	-	-	-	-	-	-	-	-	2	2	3
Female	533	1	-	-	-	-	-	-	-	-	1	2
White, both sexes	742	1	-	-	-	-	-	-	-	-	2	3
Male	234	-	-	-	-	-	-	-	-	-	2	1
Female	508	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	32	-	-	-	-	-	-	-	-	2	-	2
Male	16	-	-	-	-	-	-	-	-	2	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	786	-	-	-	-	-	-	-	-	2	3	5
Male	287	-	-	-	-	-	-	-	-	2	3	4
Female	499	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	731	-	-	-	-	-	-	-	-	2	3	3
Male	259	-	-	-	-	-	-	-	-	2	3	3
Female	472	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	50	-	-	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Delaware												
All races, both sexes	212	-	-	-	-	-	-	-	-	-	1	1
Male	72	-	-	-	-	-	-	-	-	-	-	-
Female	140	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	182	-	-	-	-	-	-	-	-	-	1	1
Male	55	-	-	-	-	-	-	-	-	-	-	-
Female	127	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	107	-	-	-	-	-	-	-	-	-	1	1
Male	41	-	-	-	-	-	-	-	-	-	1	-
Female	66	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	78	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Florida												
All races, both sexes	4,484	-	-	1	-	-	-	4	3	8	14	35
Male	1,858	-	-	1	-	-	-	3	2	5	5	17
Female	2,626	-	-	-	-	-	-	1	1	3	9	18
White, both sexes	3,835	-	-	1	-	-	-	1	1	4	7	17
Male	1,594	-	-	1	-	-	-	-	1	2	3	9
Female	2,241	-	-	-	-	-	-	1	-	2	4	8
Black, both sexes	609	-	-	-	-	-	-	3	2	4	7	17
Male	245	-	-	-	-	-	-	3	1	3	2	7
Female	364	-	-	-	-	-	-	-	1	1	5	10
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,201	-	-	-	-	-	-	-	3	5	27	28
Male	868	-	-	-	-	-	-	-	1	2	13	12
Female	1,333	-	-	-	-	-	-	-	2	3	14	16
White, both sexes	1,569	-	-	-	-	-	-	-	-	1	14	16
Male	602	-	-	-	-	-	-	-	-	-	8	7
Female	967	-	-	-	-	-	-	-	-	1	6	9
Black, both sexes	605	-	-	-	-	-	-	-	3	4	12	12
Male	253	-	-	-	-	-	-	-	1	2	4	5
Female	352	-	-	-	-	-	-	-	2	2	8	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Hawaii												
All races, both sexes	352	-	-	-	-	-	-	-	-	-	2	3
Male	157	-	-	-	-	-	-	-	-	-	-	1
Female	195	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	87	-	-	-	-	-	-	-	-	-	2	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	263	-	-	-	-	-	-	-	-	-	-	3
Male	120	-	-	-	-	-	-	-	-	-	-	1
Female	143	-	-	-	-	-	-	-	-	-	-	2
Idaho												
All races, both sexes	407	-	-	-	-	-	-	-	-	-	-	1
Male	175	-	-	-	-	-	-	-	-	-	-	-
Female	232	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	399	-	-	-	-	-	-	-	-	-	-	1
Male	173	-	-	-	-	-	-	-	-	-	-	-
Female	226	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Illinois												
All races, both sexes	3,246	-	-	-	1	-	-	-	4	3	12	21
Male	1,165	-	-	-	1	-	-	-	2	3	8	13
Female	2,081	-	-	-	-	-	-	-	2	-	4	8
White, both sexes	2,814	-	-	-	1	-	-	-	2	3	9	14
Male	994	-	-	-	1	-	-	-	1	3	6	9
Female	1,820	-	-	-	-	-	-	-	1	-	3	5
Black, both sexes	387	-	-	-	-	-	-	-	2	-	3	7
Male	153	-	-	-	-	-	-	-	1	-	2	4
Female	234	-	-	-	-	-	-	-	1	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,838	-	-	-	-	-	2	-	1	2	7	11
Male	654	-	-	-	-	-	1	-	-	-	2	5
Female	1,184	-	-	-	-	-	1	-	1	2	5	6
White, both sexes	1,730	-	-	-	-	-	1	-	1	1	6	11
Male	608	-	-	-	-	-	1	-	-	-	2	5
Female	1,122	-	-	-	-	-	-	-	1	1	4	6
Black, both sexes	105	-	-	-	-	-	1	-	-	1	1	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	1	-	-	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Iowa												
All races, both sexes	1,050	-	-	-	-	-	-	-	-	-	2	4
Male	354	-	-	-	-	-	-	-	-	-	2	2
Female	696	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	1,037	-	-	-	-	-	-	-	-	-	2	2
Male	348	-	-	-	-	-	-	-	-	-	2	2
Female	689	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	838	-	-	-	-	-	-	-	-	1	3	3
Male	297	-	-	-	-	-	-	-	-	1	1	-
Female	541	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	796	-	-	-	-	-	-	-	-	1	3	2
Male	280	-	-	-	-	-	-	-	-	1	1	-
Female	516	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Kentucky												
All races, both sexes	1,247	-	-	-	-	-	1	1	2	5	5	14
Male	481	-	-	-	-	-	1	1	1	3	5	7
Female	766	-	-	-	-	-	-	-	1	2	-	7
White, both sexes	1,162	-	-	-	-	-	1	-	2	3	4	11
Male	442	-	-	-	-	-	1	-	1	2	4	7
Female	720	-	-	-	-	-	-	-	1	1	-	4
Black, both sexes	80	-	-	-	-	-	-	1	-	2	1	3
Male	38	-	-	-	-	-	-	1	-	1	1	-
Female	42	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,243	-	-	-	-	-	1	-	3	4	9	16
Male	489	-	-	-	-	-	-	-	2	3	2	8
Female	754	-	-	-	-	-	1	-	1	1	7	8
White, both sexes	840	-	-	-	-	-	-	-	2	1	1	6
Male	325	-	-	-	-	-	-	-	2	1	1	4
Female	515	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	397	-	-	-	-	-	1	-	1	3	8	10
Male	162	-	-	-	-	-	-	-	-	2	1	4
Female	235	-	-	-	-	-	1	-	1	1	7	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Maine												
All races, both sexes	398	-	1	-	-	-	-	-	-	-	-	2
Male	130	-	1	-	-	-	-	-	-	-	-	1
Female	268	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	398	-	1	-	-	-	-	-	-	-	-	2
Male	130	-	1	-	-	-	-	-	-	-	-	1
Female	268	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,206	1	-	-	1	-	-	1	-	5	3	9
Male	433	1	-	-	-	-	-	1	-	1	1	6
Female	773	-	-	-	1	-	-	-	-	4	2	3
White, both sexes	880	1	-	-	-	-	-	1	-	-	2	3
Male	297	1	-	-	-	-	-	1	-	-	1	3
Female	583	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	299	-	-	-	1	-	-	-	-	5	1	6
Male	126	-	-	-	-	-	-	-	-	1	-	3
Female	173	-	-	-	1	-	-	-	-	4	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Massachusetts												
All races, both sexes	1,446	-	-	-	-	-	-	-	2	-	1	2
Male	487	-	-	-	-	-	-	-	1	-	-	2
Female	959	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	1,361	-	-	-	-	-	-	-	1	-	1	1
Male	457	-	-	-	-	-	-	-	1	-	-	1
Female	904	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	64	-	-	-	-	-	-	-	1	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,225	-	-	-	-	-	1	1	4	1	10	15
Male	820	-	-	-	-	-	1	1	3	-	3	8
Female	1,405	-	-	-	-	-	-	-	1	1	7	7
White, both sexes	1,954	-	-	-	-	-	1	1	2	1	8	14
Male	721	-	-	-	-	-	1	1	2	-	2	8
Female	1,233	-	-	-	-	-	-	-	-	1	6	6
Black, both sexes	260	-	-	-	-	-	-	-	2	-	2	1
Male	94	-	-	-	-	-	-	-	1	-	1	-
Female	166	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Minnesota												
All races, both sexes	1,117	-	-	-	-	-	1	-	1	-	1	4
Male	396	-	-	-	-	-	1	-	-	-	1	1
Female	721	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	1,088	-	-	-	-	-	1	-	1	-	1	2
Male	385	-	-	-	-	-	1	-	-	-	1	1
Female	703	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Mississippi												
All races, both sexes	997	-	-	-	-	1	-	4	1	1	9	27
Male	392	-	-	-	-	1	-	2	1	1	4	19
Female	605	-	-	-	-	-	-	2	-	-	5	8
White, both sexes	640	-	-	-	-	1	-	1	1	-	-	9
Male	243	-	-	-	-	1	-	1	1	-	-	7
Female	397	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	356	-	-	-	-	-	-	3	-	1	9	18
Male	149	-	-	-	-	-	-	1	-	1	4	12
Female	207	-	-	-	-	-	-	2	-	-	5	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Missouri												
All races, both sexes	1,710	-	-	-	-	-	-	-	1	-	8	12
Male	618	-	-	-	-	-	-	-	1	-	6	7
Female	1,092	-	-	-	-	-	-	-	-	-	2	5
White, both sexes	1,554	-	-	-	-	-	-	-	-	-	5	10
Male	552	-	-	-	-	-	-	-	-	-	4	6
Female	1,002	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	151	-	-	-	-	-	-	-	1	-	3	2
Male	64	-	-	-	-	-	-	-	1	-	2	1
Female	87	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	254	-	-	-	-	-	-	-	-	-	1	1
Male	101	-	-	-	-	-	-	-	-	-	1	-
Female	153	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	245	-	-	-	-	-	-	-	-	-	-	1
Male	96	-	-	-	-	-	-	-	-	-	-	-
Female	149	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Nebraska												
All races, both sexes	523	-	-	-	-	-	-	-	-	1	1	2
Male	180	-	-	-	-	-	-	-	-	1	1	2
Female	343	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	508	-	-	-	-	-	-	-	-	1	1	2
Male	175	-	-	-	-	-	-	-	-	1	1	2
Female	333	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	492	-	-	-	-	-	-	-	1	2	1	7
Male	209	-	-	-	-	-	-	-	-	1	1	5
Female	283	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	430	-	-	-	-	-	-	-	1	2	1	6
Male	178	-	-	-	-	-	-	-	-	1	1	4
Female	252	-	-	-	-	-	-	-	1	1	-	2
Black, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
New Hampshire												
All races, both sexes	250	-	-	-	-	-	-	-	1	-	-	-
Male	75	-	-	-	-	-	-	-	-	-	-	-
Female	175	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	249	-	-	-	-	-	-	-	1	-	-	-
Male	74	-	-	-	-	-	-	-	-	-	-	-
Female	175	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,888	-	-	-	-	-	-	2	-	6	7	13
Male	699	-	-	-	-	-	-	1	-	4	4	7
Female	1,189	-	-	-	-	-	-	1	-	2	3	6
White, both sexes	1,593	-	-	-	-	-	-	-	-	3	1	7
Male	567	-	-	-	-	-	-	-	-	2	1	5
Female	1,026	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	250	-	-	-	-	-	-	2	-	3	6	5
Male	112	-	-	-	-	-	-	1	-	2	3	1
Female	138	-	-	-	-	-	-	1	-	1	3	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
New Mexico												
All races, both sexes	356	-	-	-	-	1	-	-	1	-	3	5
Male	131	-	-	-	-	-	-	-	1	-	1	2
Female	225	-	-	-	-	1	-	-	-	-	2	3
White, both sexes	332	-	-	-	-	1	-	-	-	-	2	5
Male	117	-	-	-	-	-	-	-	-	-	-	2
Female	215	-	-	-	-	1	-	-	-	-	2	3
Black, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3,339	-	-	-	-	-	-	-	3	4	8	28
Male	1,239	-	-	-	-	-	-	-	1	1	4	16
Female	2,100	-	-	-	-	-	-	-	2	3	4	12
White, both sexes	2,875	-	-	-	-	-	-	-	2	1	4	19
Male	1,062	-	-	-	-	-	-	-	1	1	2	11
Female	1,813	-	-	-	-	-	-	-	1	-	2	8
Black, both sexes	378	-	-	-	-	-	-	-	1	3	4	9
Male	139	-	-	-	-	-	-	-	-	-	2	5
Female	239	-	-	-	-	-	-	-	1	3	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	86	-	-	-	-	-	-	-	-	-	-	-
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
North Carolina												
All races, both sexes	2,421	-	-	-	-	1	-	-	1	4	10	11
Male	919	-	-	-	-	1	-	-	1	4	5	7
Female	1,502	-	-	-	-	-	-	-	-	-	5	4
White, both sexes	1,859	-	-	-	-	1	-	-	-	1	7	6
Male	681	-	-	-	-	1	-	-	-	1	4	4
Female	1,178	-	-	-	-	-	-	-	-	-	3	2
Black, both sexes	535	-	-	-	-	-	-	-	1	3	3	5
Male	225	-	-	-	-	-	-	-	1	3	1	3
Female	310	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	266	-	-	-	-	-	-	-	1	-	-	2
Male	101	-	-	-	-	-	-	-	-	-	-	1
Female	165	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	262	-	-	-	-	-	-	-	-	-	-	2
Male	99	-	-	-	-	-	-	-	-	-	-	1
Female	163	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Ohio												
All races, both sexes	2,664	-	-	-	-	1	-	1	1	6	10	18
Male	977	-	-	-	-	1	-	-	1	4	6	14
Female	1,687	-	-	-	-	-	-	1	-	2	4	4
White, both sexes	2,373	-	-	-	-	1	-	1	1	5	7	15
Male	842	-	-	-	-	1	-	-	1	3	4	12
Female	1,531	-	-	-	-	-	-	1	-	2	3	3
Black, both sexes	282	-	-	-	-	-	-	-	-	1	3	3
Male	130	-	-	-	-	-	-	-	-	1	2	2
Female	152	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,330	-	-	-	-	1	-	1	1	3	3	13
Male	522	-	-	-	-	-	-	-	1	2	3	5
Female	808	-	-	-	-	1	-	1	-	1	-	8
White, both sexes	1,171	-	-	-	-	1	-	-	-	3	3	10
Male	455	-	-	-	-	-	-	-	-	2	3	3
Female	716	-	-	-	-	1	-	-	-	1	-	7
Black, both sexes	99	-	-	-	-	-	-	1	1	-	-	2
Male	43	-	-	-	-	-	-	-	1	-	-	2
Female	56	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Oregon												
All races, both sexes	1,104	-	-	-	-	-	-	-	1	-	3	3
Male	457	-	-	-	-	-	-	-	-	-	2	2
Female	647	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	1,060	-	-	-	-	-	-	-	1	-	2	3
Male	437	-	-	-	-	-	-	-	-	-	1	2
Female	623	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,802	-	1	-	-	1	-	2	-	4	14	33
Male	1,369	-	-	-	-	-	-	1	-	4	9	24
Female	2,433	-	1	-	-	1	-	1	-	-	5	9
White, both sexes	3,467	-	1	-	-	1	-	1	-	4	10	21
Male	1,240	-	-	-	-	-	-	-	-	4	6	14
Female	2,227	-	1	-	-	1	-	1	-	-	4	7
Black, both sexes	312	-	-	-	-	-	-	1	-	-	4	12
Male	122	-	-	-	-	-	-	1	-	-	3	10
Female	190	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Rhode Island												
All races, both sexes	213	-	-	-	-	-	-	-	-	-	-	2
Male	73	-	-	-	-	-	-	-	-	-	-	2
Female	140	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	208	-	-	-	-	-	-	-	-	-	-	2
Male	72	-	-	-	-	-	-	-	-	-	-	2
Female	136	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,203	1	-	-	-	-	-	-	1	5	6	22
Male	465	1	-	-	-	-	-	-	-	1	5	13
Female	738	-	-	-	-	-	-	-	1	4	1	9
White, both sexes	813	-	-	-	-	-	-	-	1	2	4	12
Male	306	-	-	-	-	-	-	-	-	1	3	7
Female	507	-	-	-	-	-	-	-	1	1	1	5
Black, both sexes	381	1	-	-	-	-	-	-	-	2	2	10
Male	157	1	-	-	-	-	-	-	-	-	2	6
Female	224	-	-	-	-	-	-	-	-	2	-	4
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
South Dakota												
All races, both sexes	267	-	-	-	-	-	-	-	1	-	-	-
Male	112	-	-	-	-	-	-	-	1	-	-	-
Female	155	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	257	-	-	-	-	-	-	-	1	-	-	-
Male	108	-	-	-	-	-	-	-	1	-	-	-
Female	149	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,921	-	-	-	-	-	1	-	3	4	9	17
Male	706	-	-	-	-	-	1	-	3	1	4	12
Female	1,215	-	-	-	-	-	-	-	-	3	5	5
White, both sexes	1,612	-	-	-	-	-	-	-	2	4	6	12
Male	577	-	-	-	-	-	-	-	2	1	2	10
Female	1,035	-	-	-	-	-	-	-	-	3	4	2
Black, both sexes	305	-	-	-	-	-	1	-	1	-	3	5
Male	126	-	-	-	-	-	1	-	1	-	2	2
Female	179	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Texas												
All races, both sexes	4,876	2	-	-	-	1	-	1	4	10	24	58
Male	1,888	1	-	-	-	1	-	-	-	6	13	28
Female	2,988	1	-	-	-	-	-	1	4	4	11	30
White, both sexes	4,185	1	-	-	-	1	-	1	2	9	16	37
Male	1,597	-	-	-	-	1	-	-	-	5	10	18
Female	2,588	1	-	-	-	-	-	1	2	4	6	19
Black, both sexes	636	1	-	-	-	-	-	-	2	1	8	20
Male	268	1	-	-	-	-	-	-	-	1	3	10
Female	368	-	-	-	-	-	-	-	2	-	5	10
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	50	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	369	1	-	-	-	-	1	-	-	1	-	1
Male	147	-	-	-	-	-	-	-	-	1	-	1
Female	222	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	360	1	-	-	-	-	1	-	-	1	-	1
Male	147	-	-	-	-	-	-	-	-	1	-	1
Female	213	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Vermont												
All races, both sexes	136	-	-	-	-	-	-	-	-	1	1	1
Male	52	-	-	-	-	-	-	-	-	1	1	1
Female	84	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	136	-	-	-	-	-	-	-	-	1	1	1
Male	52	-	-	-	-	-	-	-	-	1	1	1
Female	84	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,925	1	-	-	1	-	2	2	1	1	7	13
Male	713	1	-	-	-	-	2	2	-	-	3	8
Female	1,212	-	-	-	1	-	-	-	1	1	4	5
White, both sexes	1,469	1	-	-	-	-	2	1	-	-	3	6
Male	517	1	-	-	-	-	2	1	-	-	1	4
Female	952	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	423	-	-	-	-	-	-	1	1	1	4	7
Male	174	-	-	-	-	-	-	1	-	-	2	4
Female	249	-	-	-	-	-	-	-	1	1	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	1	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Washington												
All races, both sexes	1,422	-	-	-	-	-	-	-	1	2	2	5
Male	504	-	-	-	-	-	-	-	1	2	2	5
Female	918	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1,310	-	-	-	-	-	-	-	1	2	1	3
Male	451	-	-	-	-	-	-	-	1	2	1	3
Female	859	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	39	-	-	-	-	-	-	-	-	-	1	2
Male	15	-	-	-	-	-	-	-	-	-	1	2
Female	24	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	62	-	-	-	-	-	-	-	-	-	-	-
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	610	-	-	-	-	-	-	-	-	-	3	3
Male	227	-	-	-	-	-	-	-	-	-	3	2
Female	383	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	592	-	-	-	-	-	-	-	-	-	3	3
Male	219	-	-	-	-	-	-	-	-	-	3	2
Female	373	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Wisconsin												
All races, both sexes	1,528	1	-	-	-	-	-	-	1	2	2	5
Male	555	1	-	-	-	-	-	-	-	1	1	3
Female	973	-	-	-	-	-	-	-	1	1	1	2
White, both sexes	1,458	1	-	-	-	-	-	-	-	1	2	4
Male	518	1	-	-	-	-	-	-	-	-	1	2
Female	940	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	50	-	-	-	-	-	-	-	1	1	-	1
Male	28	-	-	-	-	-	-	-	-	1	-	1
Female	22	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	150	-	-	-	-	-	-	-	-	-	1	-
Male	56	-	-	-	-	-	-	-	-	-	1	-
Female	94	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	148	-	-	-	-	-	-	-	-	-	1	-
Male	55	-	-	-	-	-	-	-	-	-	1	-
Female	93	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
United States												
All races, both sexes	1,020	1,681	2,423	3,531	5,471	9,356	14,573	16,149	11,785	4,546	811	1
Male	617	1,041	1,375	1,965	2,712	4,224	5,561	5,189	2,982	823	106	-
Female	403	640	1,048	1,566	2,759	5,132	9,012	10,960	8,803	3,723	705	1
White, both sexes	675	1,134	1,700	2,614	4,262	7,866	12,846	14,616	10,759	4,169	719	1
Male	402	703	973	1,463	2,109	3,587	4,925	4,702	2,718	744	93	-
Female	273	431	727	1,151	2,153	4,279	7,921	9,914	8,041	3,425	626	1
Black, both sexes	316	481	634	785	1,024	1,219	1,366	1,173	796	304	88	-
Male	198	299	353	420	511	527	492	332	179	59	13	-
Female	118	182	281	365	513	692	874	841	617	245	75	-
American Indian 1/, both sexes	7	9	16	22	32	34	45	40	34	4	1	-
Male	3	8	13	10	15	12	16	12	7	1	-	-
Female	4	1	3	12	17	22	29	28	27	3	1	-
Asian or Pacific Islander 2/, both sexes	22	57	73	110	153	237	316	320	196	69	3	-
Male	14	31	36	72	77	98	128	143	78	19	-	-
Female	8	26	37	38	76	139	188	177	118	50	3	-
Alabama												
All races, both sexes	38	55	69	101	154	246	319	325	201	67	13	-
Male	22	41	41	58	88	122	112	96	43	11	2	-
Female	16	14	28	43	66	124	207	229	158	56	11	-
White, both sexes	20	23	44	70	95	172	237	259	152	55	10	-
Male	14	16	23	39	53	85	78	80	31	6	2	-
Female	6	7	21	31	42	87	159	179	121	49	8	-
Black, both sexes	18	31	25	31	57	73	82	65	47	12	3	-
Male	8	24	18	19	33	37	34	16	11	5	-	-
Female	10	7	7	12	24	36	48	49	36	7	3	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	-	-	1	2	-	-	-
Male	-	1	-	-	2	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Alaska												
All races, both sexes	2	8	4	4	9	10	16	14	8	3	1	-
Male	1	5	4	4	5	6	6	5	3	-	-	-
Female	1	3	-	-	4	4	10	9	5	3	1	-
White, both sexes	1	6	3	2	5	8	14	10	7	2	1	-
Male	1	4	3	2	4	4	5	4	3	-	-	-
Female	-	2	-	-	1	4	9	6	4	2	1	-
Black, both sexes	-	1	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	1	1	3	2	1	3	1	-	-	-
Male	-	-	1	1	1	2	1	1	-	-	-	-
Female	1	-	-	-	2	-	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Arizona												
All races, both sexes	9	25	42	56	92	152	283	277	207	79	14	-
Male	7	12	23	30	48	71	107	87	53	16	1	-
Female	2	13	19	26	44	81	176	190	154	63	13	-
White, both sexes	8	24	36	52	87	139	273	266	202	77	14	-
Male	6	11	19	29	44	64	102	82	51	14	1	-
Female	2	13	17	23	43	75	171	184	151	63	13	-
Black, both sexes	1	1	3	3	3	7	8	7	3	2	-	-
Male	1	1	2	1	3	5	4	3	2	2	-	-
Female	-	-	1	2	-	2	4	4	1	-	-	-
American Indian 1/, both sexes	-	-	1	1	2	3	2	4	1	-	-	-
Male	-	-	1	-	1	2	1	2	-	-	-	-
Female	-	-	-	1	1	1	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	3	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	3	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Arkansas												
All races, both sexes	22	38	49	70	93	130	217	247	170	63	11	-
Male	13	19	26	36	38	54	83	81	45	9	1	-
Female	9	19	23	34	55	76	134	166	125	54	10	-
White, both sexes	16	24	40	52	65	107	189	221	152	58	7	-
Male	10	10	21	30	25	45	71	69	41	9	-	-
Female	6	14	19	22	40	62	118	152	111	49	7	-
Black, both sexes	5	14	9	18	27	22	26	25	15	5	4	-
Male	3	9	5	6	13	9	12	11	4	-	1	-
Female	2	5	4	12	14	13	14	14	11	5	3	-
American Indian 1/, both sexes	1	-	-	-	-	1	2	1	3	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	2	-	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
California												
All races, both sexes	98	174	229	335	531	900	1,424	1,654	1,122	480	91	-
Male	55	102	122	189	248	392	552	567	339	96	7	-
Female	43	72	107	146	283	508	872	1,087	783	384	84	-
White, both sexes	73	107	162	245	397	727	1,210	1,423	977	422	82	-
Male	44	64	91	135	183	321	472	481	283	82	6	-
Female	29	43	71	110	214	406	738	942	694	340	76	-
Black, both sexes	17	33	30	49	62	62	78	79	54	28	8	-
Male	7	20	14	25	30	28	28	19	15	6	1	-
Female	10	13	16	24	32	34	50	60	39	22	7	-
American Indian 1/, both sexes	-	2	3	2	4	3	2	6	1	-	-	-
Male	-	2	-	1	2	1	1	3	1	-	-	-
Female	-	-	3	1	2	2	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	32	34	39	68	108	134	146	90	30	1	-
Male	4	16	17	28	33	42	51	64	40	8	-	-
Female	4	16	17	11	35	66	83	82	50	22	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Colorado												
All races, both sexes	11	8	18	44	55	96	153	189	142	53	10	-
Male	5	6	6	22	21	34	64	52	34	5	1	-
Female	6	2	12	22	34	62	89	137	108	48	9	-
White, both sexes	10	6	14	39	51	89	147	182	137	51	10	-
Male	4	5	4	19	20	30	61	49	33	5	1	-
Female	6	1	10	20	31	59	86	133	104	46	9	-
Black, both sexes	1	2	3	3	1	5	2	6	4	1	-	-
Male	1	1	1	1	-	3	2	2	1	-	-	-
Female	-	1	2	2	1	2	-	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	3	2	3	1	-	1	-	-
Male	-	-	1	2	1	1	1	1	-	-	-	-
Female	-	-	-	-	2	1	2	-	-	1	-	-
Connecticut												
All races, both sexes	8	14	13	32	41	85	145	224	121	69	24	-
Male	6	8	8	23	23	35	62	78	27	8	-	-
Female	2	6	5	9	18	50	83	146	94	61	24	-
White, both sexes	5	10	12	26	35	79	134	220	111	67	24	-
Male	4	5	7	18	18	33	56	77	25	8	-	-
Female	1	5	5	8	17	46	78	143	86	59	24	-
Black, both sexes	3	3	1	6	5	6	9	4	9	2	-	-
Male	2	2	1	5	4	2	5	1	2	-	-	-
Female	1	1	-	1	1	4	4	3	7	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Delaware												
All races, both sexes	1	2	8	14	20	21	51	53	29	10	1	-
Male	1	-	3	11	8	12	17	12	7	1	-	-
Female	-	2	5	3	12	9	34	41	22	9	1	-
White, both sexes	1	1	6	10	12	20	48	47	26	8	1	-
Male	1	-	1	8	4	12	14	9	6	-	-	-
Female	-	1	5	2	8	8	34	38	20	8	1	-
Black, both sexes	-	1	2	4	8	1	3	6	3	2	-	-
Male	-	-	2	3	4	-	3	3	1	1	-	-
Female	-	1	-	1	4	1	-	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	4	3	4	12	11	23	18	17	8	2	-
Male	2	2	-	3	4	7	7	7	6	2	-	-
Female	1	2	3	1	8	4	16	11	11	6	2	-
White, both sexes	-	-	-	-	2	4	5	6	6	3	1	-
Male	-	-	-	-	1	3	-	4	3	-	-	-
Female	-	-	-	-	1	1	5	2	3	3	1	-
Black, both sexes	3	4	3	4	10	7	18	12	11	4	1	-
Male	2	2	-	3	3	4	7	3	3	1	-	-
Female	1	2	3	1	7	3	11	9	8	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Florida												
All races, both sexes	74	109	167	263	345	595	894	978	693	259	42	-
Male	43	67	93	145	171	271	373	372	217	62	11	-
Female	31	42	74	118	174	324	521	606	476	197	31	-
White, both sexes	51	66	111	191	272	506	803	900	627	237	40	-
Male	32	39	67	111	137	233	337	350	206	55	11	-
Female	19	27	44	80	135	273	466	550	421	182	29	-
Black, both sexes	23	42	54	68	68	82	81	73	61	22	2	-
Male	11	27	26	33	32	32	32	19	10	7	-	-
Female	12	15	28	35	36	50	49	54	51	15	2	-
American Indian 1/, both sexes	-	1	-	-	2	2	1	1	2	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	4	3	5	9	4	3	-	-	-
Male	-	-	-	1	1	5	4	3	1	-	-	-
Female	-	-	2	3	2	-	5	1	2	-	-	-
Georgia												
All races, both sexes	47	73	126	150	212	303	409	426	273	105	14	-
Male	29	49	79	88	105	133	153	124	61	17	2	-
Female	18	24	47	62	107	170	256	302	212	88	12	-
White, both sexes	23	32	69	88	141	200	325	347	223	82	8	-
Male	13	23	41	59	69	88	128	98	53	14	1	-
Female	10	9	28	29	72	112	197	249	170	68	7	-
Black, both sexes	22	38	56	59	70	96	80	76	49	22	6	-
Male	14	24	38	29	35	43	23	23	8	3	1	-
Female	8	14	18	30	35	53	57	53	41	19	5	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	3	1	7	4	3	-	1	-	-
Male	2	1	-	-	1	2	2	3	-	-	-	-
Female	-	1	1	3	-	5	2	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Hawaii												
All races, both sexes	3	8	13	14	24	42	69	76	72	24	2	-
Male	1	5	10	9	16	18	27	32	31	7	-	-
Female	2	3	3	5	8	24	42	44	41	17	2	-
White, both sexes	-	2	3	3	7	11	17	14	18	9	1	-
Male	-	2	2	1	7	6	4	4	8	2	-	-
Female	-	-	1	2	-	5	13	10	10	7	1	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	6	10	10	17	31	52	61	54	15	1	-
Male	1	3	8	7	9	12	23	28	23	5	-	-
Female	2	3	2	3	8	19	29	33	31	10	1	-
Idaho												
All races, both sexes	5	10	11	15	34	57	93	79	71	27	4	-
Male	2	6	8	9	19	28	42	23	29	8	1	-
Female	3	4	3	6	15	29	51	56	42	19	3	-
White, both sexes	5	8	11	14	34	55	92	78	70	27	4	-
Male	2	5	8	9	19	27	42	23	29	8	1	-
Female	3	3	3	5	15	28	50	55	41	19	3	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Illinois												
All races, both sexes	36	50	103	150	236	397	651	726	586	227	43	-
Male	19	28	57	85	121	180	240	236	137	29	6	-
Female	17	22	46	65	115	217	411	490	449	198	37	-
White, both sexes	23	39	73	97	182	341	577	661	547	208	37	-
Male	11	21	42	53	96	160	216	215	129	25	6	-
Female	12	18	31	44	86	181	361	446	418	183	31	-
Black, both sexes	12	10	29	48	50	49	67	53	36	15	6	-
Male	7	6	15	29	22	19	22	15	8	3	-	-
Female	5	4	14	19	28	30	45	38	28	12	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	5	4	7	7	12	3	4	-	-
Male	1	1	-	3	3	1	2	6	-	1	-	-
Female	-	-	1	2	1	6	5	6	3	3	-	-
Indiana												
All races, both sexes	30	39	54	91	127	258	390	397	298	98	33	-
Male	17	24	27	48	75	116	133	130	57	15	4	-
Female	13	15	27	43	52	142	257	267	241	83	29	-
White, both sexes	24	29	45	83	116	240	373	385	288	96	31	-
Male	13	18	22	45	68	107	127	125	56	15	4	-
Female	11	11	23	38	48	133	246	260	232	81	27	-
Black, both sexes	6	9	9	7	11	18	17	11	10	2	2	-
Male	4	5	5	2	7	9	6	5	1	-	-	-
Female	2	4	4	5	4	9	11	6	9	2	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Iowa												
All races, both sexes	10	16	25	28	64	95	188	288	220	89	21	-
Male	6	10	10	12	33	37	80	80	63	15	4	-
Female	4	6	15	16	31	58	108	208	157	74	17	-
White, both sexes	10	15	23	28	59	94	187	287	220	89	21	-
Male	6	10	8	12	31	36	80	79	63	15	4	-
Female	4	5	15	16	28	58	107	208	157	74	17	-
Black, both sexes	-	-	2	-	4	1	1	-	-	-	-	-
Male	-	-	2	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	17	16	31	55	108	176	209	138	63	13	-
Male	3	10	6	16	30	53	62	64	32	16	3	-
Female	2	7	10	15	25	55	114	145	106	47	10	-
White, both sexes	4	13	13	29	49	103	168	203	135	60	13	-
Male	3	8	3	15	27	50	59	62	32	16	3	-
Female	1	5	10	14	22	53	109	141	103	44	10	-
Black, both sexes	1	4	2	2	6	5	6	5	3	2	-	-
Male	-	2	2	1	3	3	2	2	-	-	-	-
Female	1	2	-	1	3	2	4	3	3	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Kentucky												
All races, both sexes	22	32	49	77	98	164	248	240	214	65	10	-
Male	15	19	26	36	47	88	99	70	53	9	1	-
Female	7	13	23	41	51	76	149	170	161	56	9	-
White, both sexes	19	28	44	69	90	153	230	230	206	63	9	-
Male	13	16	23	29	44	81	94	68	50	9	-	-
Female	6	12	21	40	46	72	136	162	156	54	9	-
Black, both sexes	3	4	4	8	8	10	16	9	8	2	1	-
Male	2	3	3	7	3	7	4	2	3	-	1	-
Female	1	1	1	1	5	3	12	7	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
Louisiana												
All races, both sexes	35	41	46	85	135	189	241	236	147	52	3	-
Male	28	21	26	42	73	76	85	80	34	7	2	-
Female	7	20	20	43	62	113	156	156	113	45	1	-
White, both sexes	14	19	26	49	76	133	170	186	116	39	2	-
Male	12	11	13	28	39	57	62	63	26	5	1	-
Female	2	8	13	21	37	76	108	123	90	34	1	-
Black, both sexes	21	21	20	36	57	55	71	48	31	13	1	-
Male	16	10	13	14	33	19	23	16	8	2	1	-
Female	5	11	7	22	24	36	48	32	23	11	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Maine												
All races, both sexes	1	10	10	13	36	55	60	101	78	28	3	-
Male	-	8	5	5	17	27	19	27	13	7	-	-
Female	1	2	5	8	19	28	41	74	65	21	3	-
White, both sexes	1	10	10	13	36	55	60	101	78	28	3	-
Male	-	8	5	5	17	27	19	27	13	7	-	-
Female	1	2	5	8	19	28	41	74	65	21	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	12	36	47	52	88	172	234	294	181	59	11	-
Male	7	24	20	29	44	79	75	87	43	11	4	-
Female	5	12	27	23	44	93	159	207	138	48	7	-
White, both sexes	3	15	18	32	51	115	184	240	153	52	10	-
Male	2	9	8	16	26	54	55	72	36	9	4	-
Female	1	6	10	16	25	61	129	168	117	43	6	-
Black, both sexes	8	20	28	19	34	54	45	49	22	6	1	-
Male	5	14	11	13	16	24	18	14	6	1	-	-
Female	3	6	17	6	18	30	27	35	16	5	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	3	3	5	4	6	1	-	-
Male	-	1	-	-	2	1	2	1	1	1	-	-
Female	1	-	-	1	1	2	3	3	5	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Massachusetts												
All races, both sexes	7	21	36	52	75	161	300	341	318	103	27	-
Male	4	8	22	30	42	83	111	96	67	18	3	-
Female	3	13	14	22	33	78	189	245	251	85	24	-
White, both sexes	7	20	31	46	70	151	286	328	300	94	25	-
Male	4	7	20	29	38	76	107	89	65	17	3	-
Female	3	13	11	17	32	75	179	239	235	77	22	-
Black, both sexes	-	1	5	4	4	7	9	9	15	6	2	-
Male	-	1	2	-	3	5	3	5	2	1	-	-
Female	-	-	3	4	1	2	6	4	13	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	3	4	4	3	3	-	-
Male	-	-	-	1	1	2	1	2	-	-	-	-
Female	-	-	-	1	-	1	3	2	3	3	-	-
Michigan												
All races, both sexes	27	38	70	84	148	303	459	534	359	149	22	-
Male	18	25	35	43	76	133	177	173	91	29	4	-
Female	9	13	35	41	72	170	282	361	268	120	18	-
White, both sexes	18	27	45	66	119	257	408	492	335	141	19	-
Male	14	20	23	36	62	113	157	162	87	29	4	-
Female	4	7	22	30	57	144	251	330	248	112	15	-
Black, both sexes	9	11	24	18	29	44	48	39	22	8	3	-
Male	4	5	11	7	14	18	19	10	4	-	-	-
Female	5	6	13	11	15	26	29	29	18	8	3	-
American Indian 1/, both sexes	-	-	1	-	-	1	2	1	2	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Minnesota												
All races, both sexes	12	18	23	40	66	117	207	270	234	111	12	-
Male	6	15	13	24	29	56	85	90	52	22	1	-
Female	6	3	10	16	37	61	122	180	182	89	11	-
White, both sexes	9	17	19	36	64	113	206	265	233	109	12	-
Male	6	14	11	21	28	55	84	89	52	21	1	-
Female	3	3	8	15	36	58	122	176	181	88	11	-
Black, both sexes	2	-	2	-	1	-	-	1	1	2	-	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	2	-	1	-	-	-	-	1	1	1	-	-
American Indian 1/, both sexes	1	1	1	1	-	3	-	3	-	-	-	-
Male	-	1	1	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	1	1	1	1	-	-	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	1	1	1	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	20	39	62	72	97	153	185	151	123	47	5	-
Male	12	27	36	39	45	64	64	38	32	6	1	-
Female	8	12	26	33	52	89	121	113	91	41	4	-
White, both sexes	9	15	25	42	53	102	134	116	96	32	4	-
Male	4	8	14	27	27	48	48	29	23	4	1	-
Female	5	7	11	15	26	54	86	87	73	28	3	-
Black, both sexes	11	24	37	30	44	51	51	34	27	15	1	-
Male	8	19	22	12	18	16	16	9	9	2	-	-
Female	3	5	15	18	26	35	35	25	18	13	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Missouri												
All races, both sexes	32	37	49	76	146	224	340	367	297	102	19	-
Male	19	27	23	46	72	113	113	100	67	20	4	-
Female	13	10	26	30	74	111	227	267	230	82	15	-
White, both sexes	24	24	38	65	128	203	312	348	283	95	19	-
Male	12	17	21	39	63	102	105	97	64	18	4	-
Female	12	7	17	26	65	101	207	251	219	77	15	-
Black, both sexes	8	13	10	11	18	21	27	18	12	7	-	-
Male	7	10	2	7	9	11	8	3	1	2	-	-
Female	1	3	8	4	9	10	19	15	11	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	1	2	6	15	13	21	53	71	44	22	4	-
Male	1	2	3	9	7	13	24	19	15	7	-	-
Female	-	-	3	6	6	8	29	52	29	15	4	-
White, both sexes	-	2	5	14	12	19	52	71	43	22	4	-
Male	-	2	2	9	7	12	23	19	15	7	-	-
Female	-	-	3	5	5	7	29	52	28	15	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	2	1	-	1	-	-	-
Male	1	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Nebraska												
All races, both sexes	6	8	8	14	24	55	98	142	106	56	2	-
Male	2	6	5	6	11	30	43	37	26	10	-	-
Female	4	2	3	8	13	25	55	105	80	46	2	-
White, both sexes	4	8	6	12	22	55	97	139	103	56	2	-
Male	2	6	4	5	10	30	42	37	25	10	-	-
Female	2	2	2	7	12	25	55	102	78	46	2	-
Black, both sexes	2	-	2	1	2	-	1	2	3	-	-	-
Male	-	-	1	1	1	-	1	-	1	-	-	-
Female	2	-	1	-	1	-	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	10	16	22	39	52	65	107	94	57	19	-	-
Male	5	11	9	22	30	36	39	32	12	6	-	-
Female	5	5	13	17	22	29	68	62	45	13	-	-
White, both sexes	8	14	16	28	41	57	99	85	53	19	-	-
Male	3	10	6	15	21	32	37	30	12	6	-	-
Female	5	4	10	13	20	25	62	55	41	13	-	-
Black, both sexes	1	1	4	8	7	5	5	7	1	-	-	-
Male	1	1	3	6	6	2	-	2	-	-	-	-
Female	-	-	1	2	1	3	5	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	2	4	3	3	2	2	-	-	-
Male	1	-	-	1	3	2	2	-	-	-	-	-
Female	-	1	2	1	1	1	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
New Hampshire												
All races, both sexes	3	4	8	4	15	30	46	62	47	29	1	-
Male	-	3	6	4	8	15	15	13	8	3	-	-
Female	3	1	2	-	7	15	31	49	39	26	1	-
White, both sexes	3	4	8	4	15	30	46	61	47	29	1	-
Male	-	3	6	4	8	15	15	12	8	3	-	-
Female	3	1	2	-	7	15	31	49	39	26	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	22	43	67	74	124	235	386	432	335	118	24	-
Male	14	29	42	44	66	91	148	149	89	9	2	-
Female	8	14	25	30	58	144	238	283	246	109	22	-
White, both sexes	15	31	41	56	91	196	326	396	304	105	21	-
Male	9	22	24	35	49	72	126	133	80	7	2	-
Female	6	9	17	21	42	124	200	263	224	98	19	-
Black, both sexes	7	12	24	13	26	34	50	26	28	11	3	-
Male	5	7	17	6	13	18	17	11	9	2	-	-
Female	2	5	7	7	13	16	33	15	19	9	3	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	5	6	5	10	10	3	2	-	-
Male	-	-	1	3	4	1	5	5	-	-	-	-
Female	-	-	1	2	2	4	5	5	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
New Mexico												
All races, both sexes	2	8	11	17	23	46	71	73	69	21	5	-
Male	1	2	6	9	10	24	26	26	16	5	2	-
Female	1	6	5	8	13	22	45	47	53	16	3	-
White, both sexes	2	7	9	15	22	43	67	71	63	21	4	-
Male	1	1	6	7	9	22	24	26	12	5	2	-
Female	1	6	3	8	13	21	43	45	51	16	2	-
Black, both sexes	-	1	2	1	1	2	1	2	1	-	-	-
Male	-	1	-	1	1	1	-	-	1	-	-	-
Female	-	-	2	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	2	-	5	-	1	-
Male	-	-	-	1	-	1	1	-	3	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	45	83	103	148	255	415	697	748	539	224	39	-
Male	30	52	60	79	131	178	273	243	132	36	3	-
Female	15	31	43	69	124	237	424	505	407	188	36	-
White, both sexes	31	62	73	103	195	361	608	664	505	214	33	-
Male	20	40	43	56	103	158	245	220	126	34	2	-
Female	11	22	30	47	92	203	363	444	379	180	31	-
Black, both sexes	13	19	26	35	51	39	69	64	30	9	6	-
Male	9	10	16	17	23	16	21	13	5	1	1	-
Female	4	9	10	18	28	23	48	51	25	8	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	4	10	9	15	20	20	4	1	-	-
Male	1	2	1	6	5	4	7	10	1	1	-	-
Female	-	-	3	4	4	11	13	10	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
North Carolina												
All races, both sexes	36	65	95	142	184	351	511	524	340	118	28	-
Male	24	39	60	86	99	163	174	157	74	21	4	-
Female	12	26	35	56	85	188	337	367	266	97	24	-
White, both sexes	22	39	53	85	114	255	414	448	294	99	21	-
Male	11	24	34	51	58	122	143	142	65	19	2	-
Female	11	15	19	34	56	133	271	306	229	80	19	-
Black, both sexes	14	26	39	55	65	92	91	73	44	18	6	-
Male	13	15	23	33	38	40	28	14	9	2	2	-
Female	1	11	16	22	27	52	63	59	35	16	4	-
American Indian 1/, both sexes	-	-	2	2	2	4	4	1	-	1	-	-
Male	-	-	2	2	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	3	3	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	3	-	2	2	2	-	1	-
Male	-	-	1	-	2	-	2	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	2	-	1	-
North Dakota												
All races, both sexes	1	2	3	7	11	32	54	60	61	26	6	-
Male	1	2	2	5	5	15	26	25	12	7	-	-
Female	-	-	1	2	6	17	28	35	49	19	6	-
White, both sexes	1	2	3	5	11	32	53	60	61	26	6	-
Male	1	2	2	4	5	15	25	25	12	7	-	-
Female	-	-	1	1	6	17	28	35	49	19	6	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Ohio												
All races, both sexes	34	47	73	125	193	381	563	595	435	155	26	-
Male	23	27	39	74	95	172	213	186	97	23	2	-
Female	11	20	34	51	98	209	350	409	338	132	24	-
White, both sexes	23	35	56	94	158	332	521	557	398	146	23	-
Male	14	21	30	50	72	154	198	172	89	20	1	-
Female	9	14	26	44	86	178	323	385	309	126	22	-
Black, both sexes	10	11	16	29	35	48	40	38	36	9	3	-
Male	9	5	9	22	23	18	13	14	8	3	1	-
Female	1	6	7	7	12	30	27	24	28	6	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	-	1	2	-	1	-	-	-
Male	-	-	-	2	-	-	2	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	1	-	-	-
Oklahoma												
All races, both sexes	27	32	52	78	127	194	249	263	194	76	16	-
Male	13	20	32	46	64	87	95	91	45	13	5	-
Female	14	12	20	32	63	107	154	172	149	63	11	-
White, both sexes	18	19	38	61	112	176	230	235	181	70	14	-
Male	7	11	26	35	57	82	88	83	41	13	4	-
Female	11	8	12	26	55	94	142	152	140	57	10	-
Black, both sexes	7	10	12	11	8	11	9	15	6	4	2	-
Male	4	7	4	8	3	4	3	4	2	-	1	-
Female	3	3	8	3	5	7	6	11	4	4	1	-
American Indian 1/, both sexes	1	1	2	4	6	4	10	13	7	1	-	-
Male	1	-	2	2	4	-	4	4	2	-	-	-
Female	-	1	-	2	2	4	6	9	5	1	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	2	1	3	-	-	-	1	-	-
Male	1	2	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	2	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Oregon												
All races, both sexes	13	25	34	45	70	134	225	259	193	80	19	-
Male	9	22	23	29	36	61	99	90	57	23	4	-
Female	4	3	11	16	34	73	126	169	136	57	15	-
White, both sexes	12	21	31	40	68	128	218	250	189	78	19	-
Male	9	19	22	26	35	57	97	85	57	23	4	-
Female	3	2	9	14	33	71	121	165	132	55	15	-
Black, both sexes	-	4	1	1	1	4	3	2	-	1	-	-
Male	-	3	-	1	1	4	-	2	-	-	-	-
Female	-	1	1	-	-	-	3	-	-	1	-	-
American Indian 1/, both sexes	1	-	1	2	1	-	-	1	2	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	2	1	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	2	4	6	2	1	-	-
Male	-	-	-	2	-	-	2	2	-	-	-	-
Female	-	-	1	-	-	2	2	4	2	1	-	-
Pennsylvania												
All races, both sexes	35	70	91	140	247	467	784	960	665	255	33	-
Male	21	45	62	75	110	197	311	296	168	42	4	-
Female	14	25	29	65	137	270	473	664	497	213	29	-
White, both sexes	28	53	73	113	209	415	732	902	631	242	31	-
Male	18	33	50	65	95	177	297	277	161	39	4	-
Female	10	20	23	48	114	238	435	625	470	203	27	-
Black, both sexes	7	17	18	24	36	50	47	51	32	11	2	-
Male	3	12	12	8	13	19	13	18	7	3	-	-
Female	4	5	6	16	23	31	34	33	25	8	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	2	2	5	7	2	2	-	-
Male	-	-	-	2	2	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	4	6	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Rhode Island												
All races, both sexes	2	4	3	7	10	23	47	59	43	11	2	-
Male	1	3	2	5	7	10	15	17	10	1	-	-
Female	1	1	1	2	3	13	32	42	33	10	2	-
White, both sexes	2	4	3	7	9	23	45	59	41	11	2	-
Male	1	3	2	5	7	10	14	17	10	1	-	-
Female	1	1	1	2	2	13	31	42	31	10	2	-
Black, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
South Carolina												
All races, both sexes	28	37	65	83	102	167	237	211	168	53	17	-
Male	17	20	40	51	50	81	80	62	35	8	1	-
Female	11	17	25	32	52	86	157	149	133	45	16	-
White, both sexes	10	18	35	44	55	124	172	159	123	43	11	-
Male	7	11	22	27	31	65	54	45	27	6	-	-
Female	3	7	13	17	24	59	118	114	96	37	11	-
Black, both sexes	18	19	30	39	47	42	62	51	43	9	6	-
Male	10	9	18	24	19	16	26	17	7	1	1	-
Female	8	10	12	15	28	26	36	34	36	8	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
South Dakota												
All races, both sexes	2	3	2	4	11	24	58	62	59	36	5	-
Male	2	1	2	1	5	12	31	32	17	8	-	-
Female	-	2	-	3	6	12	27	30	42	28	5	-
White, both sexes	1	2	2	4	7	23	57	62	57	36	5	-
Male	1	-	2	1	4	11	31	32	17	8	-	-
Female	-	2	-	3	3	12	26	30	40	28	5	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	1	-	-	4	1	1	-	1	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	3	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	28	41	71	113	165	255	392	407	287	113	15	-
Male	18	24	46	66	69	105	138	137	62	18	2	-
Female	10	17	25	47	96	150	254	270	225	95	13	-
White, both sexes	11	27	52	78	131	213	347	363	259	97	10	-
Male	6	16	33	50	54	88	117	123	57	16	2	-
Female	5	11	19	28	77	125	230	240	202	81	8	-
Black, both sexes	17	14	19	35	33	41	44	43	28	16	5	-
Male	12	8	13	16	14	17	20	13	5	2	-	-
Female	5	6	6	19	19	24	24	30	23	14	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Texas												
All races, both sexes	91	156	180	289	399	654	996	964	737	262	47	1
Male	50	98	107	148	197	283	399	305	193	51	8	-
Female	41	58	73	141	202	371	597	659	544	211	39	1
White, both sexes	54	120	136	216	328	542	887	878	674	241	41	1
Male	28	77	84	110	161	238	352	282	177	47	7	-
Female	26	43	52	106	167	304	535	596	497	194	34	1
Black, both sexes	35	33	41	69	64	98	98	82	58	20	6	-
Male	20	20	21	36	35	36	45	21	14	4	1	-
Female	15	13	20	33	29	62	53	61	44	16	5	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	3	3	7	13	11	4	4	-	-	-
Male	2	1	2	2	1	9	2	2	2	-	-	-
Female	-	2	1	1	6	4	9	2	2	-	-	-
Utah												
All races, both sexes	2	5	14	13	29	59	67	99	57	20	-	-
Male	1	2	7	5	13	31	27	41	15	3	-	-
Female	1	3	7	8	16	28	40	58	42	17	-	-
White, both sexes	2	5	14	13	26	56	65	98	57	20	-	-
Male	1	2	7	5	13	31	27	41	15	3	-	-
Female	1	3	7	8	13	25	38	57	42	17	-	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Vermont												
All races, both sexes	-	-	3	5	8	17	30	32	25	10	3	-
Male	-	-	2	3	4	7	12	11	8	2	-	-
Female	-	-	1	2	4	10	18	21	17	8	3	-
White, both sexes	-	-	3	5	8	17	30	32	25	10	3	-
Male	-	-	2	3	4	7	12	11	8	2	-	-
Female	-	-	1	2	4	10	18	21	17	8	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	25	44	68	90	170	254	382	417	320	108	19	-
Male	15	30	36	48	83	118	142	119	82	22	2	-
Female	10	14	32	42	87	136	240	298	238	86	17	-
White, both sexes	17	24	31	59	111	195	295	338	280	94	12	-
Male	8	17	16	27	54	87	113	96	70	18	2	-
Female	9	7	15	32	57	108	182	242	210	76	10	-
Black, both sexes	7	19	34	28	56	54	78	76	37	13	7	-
Male	6	12	17	18	27	28	23	22	11	3	-	-
Female	1	7	17	10	29	26	55	54	26	10	7	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	3	3	5	8	3	3	1	-	-
Male	1	1	2	3	2	3	5	1	1	1	-	-
Female	-	-	-	-	1	2	3	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Washington												
All races, both sexes	11	24	37	46	84	155	296	354	274	117	14	-
Male	7	11	19	21	33	75	104	120	76	27	1	-
Female	4	13	18	25	51	80	192	234	198	90	13	-
White, both sexes	10	22	34	40	71	134	266	338	263	111	14	-
Male	6	11	17	19	28	63	91	114	69	25	1	-
Female	4	11	17	21	43	71	175	224	194	86	13	-
Black, both sexes	1	2	1	2	5	8	9	2	3	3	-	-
Male	1	-	1	-	1	4	3	-	1	1	-	-
Female	-	2	-	2	4	4	6	2	2	2	-	-
American Indian 1/, both sexes	-	-	-	1	2	2	5	1	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	1	2	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	3	6	11	16	13	8	3	-	-
Male	-	-	1	2	3	8	8	6	6	1	-	-
Female	-	-	1	1	3	3	8	7	2	2	-	-
West Virginia												
All races, both sexes	11	19	21	24	56	88	113	138	90	38	6	-
Male	8	11	12	12	33	41	41	37	22	4	1	-
Female	3	8	9	12	23	47	72	101	68	34	5	-
White, both sexes	11	19	20	24	54	82	109	135	89	38	5	-
Male	8	11	12	12	32	37	39	37	22	4	-	-
Female	3	8	8	12	22	45	70	98	67	34	5	-
Black, both sexes	-	-	1	-	2	6	4	3	1	-	1	-
Male	-	-	-	-	1	4	2	-	-	-	1	-
Female	-	-	1	-	1	2	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Wisconsin												
All races, both sexes	13	19	36	49	94	169	311	379	293	127	27	-
Male	11	12	19	31	43	80	129	129	66	27	2	-
Female	2	7	17	18	51	89	182	250	227	100	25	-
White, both sexes	10	14	29	41	85	160	296	370	293	125	27	-
Male	8	9	16	24	37	74	123	128	66	27	2	-
Female	2	5	13	17	48	86	173	242	227	98	25	-
Black, both sexes	3	5	6	5	6	8	7	6	-	1	-	-
Male	3	3	2	5	4	5	4	-	-	-	-	-
Female	-	2	4	-	2	3	3	6	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	5	-	-	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	4	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	1	1	3	3	-	1	-	-
Male	-	-	1	2	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	-	2	2	-	1	-	-
Wyoming												
All races, both sexes	2	2	8	7	12	21	25	30	28	12	2	-
Male	2	1	5	4	5	11	9	8	9	1	-	-
Female	-	1	3	3	7	10	16	22	19	11	2	-
White, both sexes	2	2	8	6	11	21	25	30	28	12	2	-
Male	2	1	5	3	5	11	9	8	9	1	-	-
Female	-	1	3	3	6	10	16	22	19	11	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
United States												
All races, both sexes	24,544	55	34	23	16	11	18	13	16	49	95	186
Male	9,450	32	15	16	10	7	10	8	8	18	50	91
Female	15,094	23	19	7	6	4	8	5	8	31	45	95
White, both sexes	21,002	28	24	15	14	8	12	9	10	28	61	129
Male	7,984	16	9	11	9	6	8	5	5	11	34	64
Female	13,018	12	15	4	5	2	4	4	5	17	27	65
Black, both sexes	2,874	23	9	7	1	2	5	3	5	18	30	53
Male	1,183	14	5	4	-	1	1	3	2	5	13	24
Female	1,691	9	4	3	1	1	4	-	3	13	17	29
American Indian 1/, both sexes	99	2	1	-	-	1	-	-	1	-	2	1
Male	35	1	1	-	-	-	-	-	1	-	1	1
Female	64	1	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	569	2	-	1	1	-	1	1	-	3	2	3
Male	248	1	-	1	1	-	1	-	-	2	2	2
Female	321	1	-	-	-	-	-	1	-	1	-	1
Alabama												
All races, both sexes	330	1	-	1	-	-	-	-	-	2	-	7
Male	124	-	-	1	-	-	-	-	-	-	-	2
Female	206	1	-	-	-	-	-	-	-	2	-	5
White, both sexes	230	-	-	-	-	-	-	-	-	1	-	5
Male	88	-	-	-	-	-	-	-	-	-	-	1
Female	142	-	-	-	-	-	-	-	-	1	-	4
Black, both sexes	98	-	-	1	-	-	-	-	-	1	-	2
Male	36	-	-	1	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Alaska												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	348	-	-	-	-	-	1	1	-	1	5	-
Male	144	-	-	-	-	-	-	1	-	1	2	-
Female	204	-	-	-	-	-	1	-	-	-	3	-
White, both sexes	329	-	-	-	-	-	1	1	-	1	5	-
Male	134	-	-	-	-	-	-	1	-	1	2	-
Female	195	-	-	-	-	-	1	-	-	-	3	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Arkansas												
All races, both sexes	319	2	-	-	-	1	-	-	1	3	-	1
Male	119	-	-	-	-	-	-	-	-	-	-	1
Female	200	2	-	-	-	1	-	-	1	3	-	-
White, both sexes	258	-	-	-	-	-	-	-	1	2	-	1
Male	102	-	-	-	-	-	-	-	-	-	-	1
Female	156	-	-	-	-	-	-	-	1	2	-	-
Black, both sexes	59	2	-	-	-	-	-	-	-	1	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	42	2	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3,763	9	2	2	2	-	3	3	5	10	13	35
Male	1,460	4	2	1	1	-	2	1	2	3	10	20
Female	2,303	5	-	1	1	-	1	2	3	7	3	15
White, both sexes	3,137	6	1	1	1	-	3	2	3	6	9	24
Male	1,199	2	1	1	-	-	2	1	2	2	7	15
Female	1,938	4	-	-	1	-	1	1	1	4	2	9
Black, both sexes	270	3	1	1	-	-	-	-	2	3	3	8
Male	112	2	1	-	-	-	-	-	-	1	2	3
Female	158	1	-	1	-	-	-	-	2	2	1	5
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	343	-	-	-	1	-	-	1	-	1	1	3
Male	146	-	-	-	1	-	-	-	-	-	1	2
Female	197	-	-	-	-	-	-	1	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Colorado												
All races, both sexes	294	-	-	1	1	-	-	-	-	-	1	1
Male	111	-	-	1	1	-	-	-	-	-	1	-
Female	183	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	282	-	-	1	1	-	-	-	-	-	1	1
Male	104	-	-	1	1	-	-	-	-	-	1	-
Female	178	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	264	-	-	1	-	-	-	-	-	1	-	1
Male	87	-	-	1	-	-	-	-	-	-	-	-
Female	177	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	250	-	-	1	-	-	-	-	-	1	-	1
Male	84	-	-	1	-	-	-	-	-	-	-	-
Female	166	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Delaware												
All races, both sexes	53	-	1	-	-	-	-	-	1	-	-	2
Male	19	-	1	-	-	-	-	-	1	-	-	2
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	1	-	-	-	-	-	1	-	-	1
Male	4	-	1	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Florida												
All races, both sexes	1,743	4	2	-	2	-	1	1	-	4	3	16
Male	695	2	1	-	2	-	-	1	-	1	1	7
Female	1,048	2	1	-	-	-	1	-	-	3	2	9
White, both sexes	1,456	1	2	-	2	-	-	-	-	3	1	11
Male	593	1	1	-	2	-	-	-	-	1	-	5
Female	863	-	1	-	-	-	-	-	-	2	1	6
Black, both sexes	276	3	-	-	-	-	1	1	-	1	2	5
Male	98	1	-	-	-	-	-	1	-	-	1	2
Female	178	2	-	-	-	-	1	-	-	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	526	3	1	-	-	1	1	1	1	2	3	5
Male	212	1	1	-	-	1	1	1	1	1	1	2
Female	314	2	-	-	-	-	-	-	-	1	2	3
White, both sexes	379	1	1	-	-	1	1	-	-	-	1	2
Male	139	-	1	-	-	1	1	-	-	-	1	1
Female	240	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	146	2	-	-	-	-	-	1	1	2	2	3
Male	73	1	-	-	-	-	-	1	1	1	-	1
Female	73	1	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Hawaii												
All races, both sexes	69	-	-	-	-	-	-	-	-	-	-	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	173	-	-	-	-	-	1	-	-	1	1	-
Male	65	-	-	-	-	-	1	-	-	-	1	-
Female	108	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	170	-	-	-	-	-	1	-	-	1	1	-
Male	63	-	-	-	-	-	1	-	-	-	1	-
Female	107	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Illinois												
All races, both sexes	1,047	2	4	-	1	1	-	-	-	-	4	6
Male	427	2	1	-	1	1	-	-	-	-	-	4
Female	620	-	3	-	-	-	-	-	-	-	4	2
White, both sexes	864	1	2	-	1	1	-	-	-	-	2	5
Male	338	1	1	-	1	1	-	-	-	-	-	3
Female	526	-	1	-	-	-	-	-	-	-	2	2
Black, both sexes	169	1	2	-	-	-	-	-	-	-	2	1
Male	83	1	-	-	-	-	-	-	-	-	-	1
Female	86	-	2	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	509	-	-	-	1	-	-	-	-	1	7	4
Male	197	-	-	-	-	-	-	-	-	1	4	1
Female	312	-	-	-	1	-	-	-	-	-	3	3
White, both sexes	447	-	-	-	1	-	-	-	-	1	5	3
Male	173	-	-	-	-	-	-	-	-	1	3	1
Female	274	-	-	-	1	-	-	-	-	-	2	2
Black, both sexes	62	-	-	-	-	-	-	-	-	-	2	1
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	38	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Iowa												
All races, both sexes	237	-	-	-	-	-	-	-	-	-	1	1
Male	97	-	-	-	-	-	-	-	-	-	-	-
Female	140	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	235	-	-	-	-	-	-	-	-	-	1	1
Male	97	-	-	-	-	-	-	-	-	-	-	-
Female	138	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	266	1	1	-	-	-	-	-	1	-	-	-
Male	90	1	-	-	-	-	-	-	-	-	-	-
Female	176	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	255	-	1	-	-	-	-	-	1	-	-	-
Male	85	-	-	-	-	-	-	-	-	-	-	-
Female	170	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	9	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Kentucky												
All races, both sexes	268	-	1	-	1	-	-	-	-	-	-	2
Male	109	-	1	-	-	-	-	-	-	-	-	1
Female	159	-	-	-	1	-	-	-	-	-	-	1
White, both sexes	250	-	1	-	1	-	-	-	-	-	-	2
Male	104	-	1	-	-	-	-	-	-	-	-	1
Female	146	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	370	1	1	-	-	-	-	1	-	2	2	7
Male	136	1	-	-	-	-	-	-	-	-	1	6
Female	234	-	1	-	-	-	-	1	-	2	1	1
White, both sexes	264	-	-	-	-	-	-	1	-	-	-	4
Male	93	-	-	-	-	-	-	-	-	-	-	3
Female	171	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	104	1	1	-	-	-	-	-	-	2	1	3
Male	42	1	-	-	-	-	-	-	-	-	1	3
Female	62	-	1	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Maine												
All races, both sexes	102	-	-	1	-	-	-	-	-	-	-	1
Male	44	-	-	1	-	-	-	-	-	-	-	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	102	-	-	1	-	-	-	-	-	-	-	1
Male	44	-	-	1	-	-	-	-	-	-	-	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	357	1	2	-	-	-	-	-	-	2	1	3
Male	152	-	-	-	-	-	-	-	-	-	-	2
Female	205	1	2	-	-	-	-	-	-	2	1	1
White, both sexes	273	1	1	-	-	-	-	-	-	1	-	2
Male	110	-	-	-	-	-	-	-	-	-	-	1
Female	163	1	1	-	-	-	-	-	-	1	-	1
Black, both sexes	76	-	1	-	-	-	-	-	-	1	1	1
Male	39	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	1	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Massachusetts												
All races, both sexes	586	-	-	1	-	1	1	-	-	-	2	3
Male	226	-	-	1	-	-	-	-	-	-	-	2
Female	360	-	-	-	-	1	1	-	-	-	2	1
White, both sexes	549	-	-	1	-	1	1	-	-	-	1	3
Male	205	-	-	1	-	-	-	-	-	-	-	2
Female	344	-	-	-	-	1	1	-	-	-	1	1
Black, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,059	-	-	1	-	2	2	-	1	-	2	8
Male	413	-	-	-	-	1	2	-	1	-	1	4
Female	646	-	-	1	-	1	-	-	-	-	1	4
White, both sexes	896	-	-	1	-	1	1	-	1	-	1	4
Male	339	-	-	-	-	1	1	-	1	-	1	2
Female	557	-	-	1	-	-	-	-	-	-	-	2
Black, both sexes	153	-	-	-	-	1	-	-	-	-	1	4
Male	72	-	-	-	-	-	-	-	-	-	-	2
Female	81	-	-	-	-	1	-	-	-	-	1	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Minnesota												
All races, both sexes	431	1	-	-	-	1	-	-	-	-	1	1
Male	170	1	-	-	-	1	-	-	-	-	-	1
Female	261	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	411	-	-	-	-	1	-	-	-	-	1	1
Male	159	-	-	-	-	1	-	-	-	-	-	1
Female	252	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	167	1	-	-	-	-	-	-	-	2	3	2
Male	76	1	-	-	-	-	-	-	-	1	-	1
Female	91	-	-	-	-	-	-	-	-	1	3	1
White, both sexes	111	1	-	-	-	-	-	-	-	-	1	1
Male	48	1	-	-	-	-	-	-	-	-	-	1
Female	63	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	56	-	-	-	-	-	-	-	-	2	2	1
Male	28	-	-	-	-	-	-	-	-	1	-	-
Female	28	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Missouri												
All races, both sexes	589	2	1	2	-	-	-	-	-	-	5	3
Male	201	1	-	1	-	-	-	-	-	-	2	-
Female	388	1	1	1	-	-	-	-	-	-	3	3
White, both sexes	522	-	1	2	-	-	-	-	-	-	5	2
Male	180	-	-	1	-	-	-	-	-	-	2	-
Female	342	-	1	1	-	-	-	-	-	-	3	2
Black, both sexes	64	2	-	-	-	-	-	-	-	-	-	1
Male	20	1	-	-	-	-	-	-	-	-	-	-
Female	44	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	1
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	84	-	-	-	-	-	-	-	-	-	-	1
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Nebraska												
All races, both sexes	167	-	-	-	-	-	-	-	-	-	1	2
Male	61	-	-	-	-	-	-	-	-	-	1	1
Female	106	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	160	-	-	-	-	-	-	-	-	-	1	1
Male	57	-	-	-	-	-	-	-	-	-	1	-
Female	103	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	89	-	-	-	-	-	-	-	-	-	-	2
Male	37	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	84	-	-	-	-	-	-	-	-	-	-	2
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Hampshire												
All races, both sexes	85	2	-	-	-	-	-	-	-	-	-	1
Male	37	1	-	-	-	-	-	-	-	-	-	1
Female	48	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	85	2	-	-	-	-	-	-	-	-	-	1
Male	37	1	-	-	-	-	-	-	-	-	-	1
Female	48	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	391	-	-	3	-	1	-	-	-	-	1	1
Male	166	-	-	1	-	1	-	-	-	-	1	1
Female	225	-	-	2	-	-	-	-	-	-	-	-
White, both sexes	335	-	-	2	-	-	-	-	-	-	-	-
Male	145	-	-	1	-	-	-	-	-	-	-	-
Female	190	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	46	-	-	1	-	1	-	-	-	-	1	1
Male	19	-	-	-	-	1	-	-	-	-	1	1
Female	27	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Mexico												
All races, both sexes	200	-	-	-	-	-	1	-	-	-	-	1
Male	70	-	-	-	-	-	1	-	-	-	-	1
Female	130	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	186	-	-	-	-	-	1	-	-	-	-	1
Male	67	-	-	-	-	-	1	-	-	-	-	1
Female	119	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	768	7	2	2	3	-	1	1	1	2	6	9
Male	286	3	1	2	1	-	-	1	1	-	5	5
Female	482	4	1	-	2	-	1	-	-	2	1	4
White, both sexes	676	4	2	1	3	-	-	-	1	-	2	8
Male	246	2	1	1	1	-	-	-	1	-	1	5
Female	430	2	1	-	2	-	-	-	-	-	1	3
Black, both sexes	79	2	-	1	-	-	1	1	-	2	4	1
Male	35	1	-	1	-	-	-	1	-	-	4	-
Female	44	1	-	-	-	-	1	-	-	2	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
North Carolina												
All races, both sexes	783	3	1	-	-	-	2	-	-	2	9	8
Male	304	3	-	-	-	-	1	-	-	1	4	3
Female	479	-	1	-	-	-	1	-	-	1	5	5
White, both sexes	588	2	1	-	-	-	1	-	-	1	5	4
Male	221	2	-	-	-	-	-	-	-	-	3	-
Female	367	-	1	-	-	-	1	-	-	1	2	4
Black, both sexes	188	1	-	-	-	-	1	-	-	1	4	3
Male	79	1	-	-	-	-	1	-	-	1	1	2
Female	109	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	71	-	1	-	-	-	-	-	-	-	-	1
Male	31	-	1	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	-	-	1
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Ohio												
All races, both sexes	1,344	4	4	1	1	-	-	1	2	-	2	10
Male	498	2	1	1	1	-	-	-	-	-	-	2
Female	846	2	3	-	-	-	-	1	2	-	2	8
White, both sexes	1,192	3	3	1	1	-	-	1	1	-	2	9
Male	442	2	-	1	1	-	-	-	-	-	-	2
Female	750	1	3	-	-	-	-	1	1	-	2	7
Black, both sexes	149	1	1	-	-	-	-	-	1	-	-	1
Male	54	-	1	-	-	-	-	-	-	-	-	-
Female	95	1	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	306	1	1	-	-	-	-	-	1	-	1	4
Male	110	1	-	-	-	-	-	-	1	-	1	3
Female	196	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	273	1	1	-	-	-	-	-	1	-	-	4
Male	94	1	-	-	-	-	-	-	1	-	-	3
Female	179	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Oregon												
All races, both sexes	383	-	-	-	1	-	-	1	-	-	-	3
Male	163	-	-	-	1	-	-	-	-	-	-	2
Female	220	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	367	-	-	-	1	-	-	1	-	-	-	2
Male	156	-	-	-	1	-	-	-	-	-	-	1
Female	211	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,111	1	2	2	1	-	1	-	1	4	5	2
Male	367	-	1	1	-	-	1	-	-	3	4	-
Female	744	1	1	1	1	-	-	-	1	1	1	2
White, both sexes	995	1	2	1	-	-	1	-	1	2	3	2
Male	319	-	1	-	-	-	1	-	-	2	3	-
Female	676	1	1	1	-	-	-	-	1	-	-	2
Black, both sexes	112	-	-	1	1	-	-	-	-	2	1	-
Male	46	-	-	1	-	-	-	-	-	1	-	-
Female	66	-	-	-	1	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Rhode Island												
All races, both sexes	72	-	1	-	-	-	-	-	-	-	-	-
Male	27	-	1	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	69	-	1	-	-	-	-	-	-	-	-	-
Male	25	-	1	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	349	-	-	-	-	-	-	-	-	1	2	2
Male	138	-	-	-	-	-	-	-	-	1	1	-
Female	211	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	245	-	-	-	-	-	-	-	-	1	-	-
Male	97	-	-	-	-	-	-	-	-	1	-	-
Female	148	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	101	-	-	-	-	-	-	-	-	-	2	2
Male	40	-	-	-	-	-	-	-	-	-	1	-
Female	61	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
South Dakota												
All races, both sexes	68	1	-	-	-	-	-	-	1	-	-	-
Male	32	1	-	-	-	-	-	-	1	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	1	-	-	-
Male	3	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	479	1	-	1	-	-	-	-	-	-	1	5
Male	205	1	-	1	-	-	-	-	-	-	-	3
Female	274	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	406	-	-	1	-	-	-	-	-	-	1	1
Male	168	-	-	1	-	-	-	-	-	-	-	1
Female	238	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	70	1	-	-	-	-	-	-	-	-	-	4
Male	34	1	-	-	-	-	-	-	-	-	-	2
Female	36	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Texas												
All races, both sexes	1,812	4	5	4	1	-	2	2	-	4	6	12
Male	666	3	2	3	1	-	-	2	-	3	4	3
Female	1,146	1	3	1	-	-	2	-	-	1	2	9
White, both sexes	1,507	3	4	1	1	-	-	2	-	3	5	7
Male	546	2	1	1	1	-	-	2	-	2	3	2
Female	961	1	3	-	-	-	-	-	-	1	2	5
Black, both sexes	285	-	1	2	-	-	2	-	-	-	1	5
Male	107	-	1	1	-	-	-	-	-	-	1	1
Female	178	-	-	1	-	-	2	-	-	-	-	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	1	-	1	-	-	-	-	-	1	-	-
Male	13	1	-	1	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	161	-	-	-	-	1	1	-	-	1	1	-
Male	62	-	-	-	-	1	1	-	-	1	1	-
Female	99	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	157	-	-	-	-	1	1	-	-	1	1	-
Male	62	-	-	-	-	1	1	-	-	1	1	-
Female	95	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Vermont												
All races, both sexes	58	1	-	-	-	-	-	-	-	-	-	-
Male	22	1	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	1	-	-	-	-	-	-	-	-	-	-
Male	22	1	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	589	1	1	-	-	1	-	1	-	1	3	7
Male	211	1	1	-	-	1	-	1	-	-	2	3
Female	378	-	-	-	-	-	-	-	-	1	1	4
White, both sexes	452	-	-	-	-	1	-	1	-	1	3	4
Male	167	-	-	-	-	1	-	1	-	-	2	2
Female	285	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	130	1	1	-	-	-	-	-	-	-	-	3
Male	42	1	1	-	-	-	-	-	-	-	-	1
Female	88	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Washington												
All races, both sexes	498	-	-	-	1	-	-	-	-	2	1	2
Male	199	-	-	-	1	-	-	-	-	-	1	2
Female	299	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	463	-	-	-	1	-	-	-	-	2	1	2
Male	182	-	-	-	1	-	-	-	-	-	1	2
Female	281	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	150	-	-	-	-	-	-	-	-	-	-	1
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	106	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	145	-	-	-	-	-	-	-	-	-	-	1
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	102	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Wisconsin												
All races, both sexes	576	1	-	-	-	1	-	-	-	1	1	2
Male	242	1	-	-	-	-	-	-	-	1	1	1
Female	334	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	555	-	-	-	-	1	-	-	-	-	1	2
Male	233	-	-	-	-	-	-	-	-	-	1	1
Female	322	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	15	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
United States												
All races, both sexes	309	442	679	1,037	1,737	2,989	4,759	5,703	4,237	1,781	355	-
Male	157	258	400	575	877	1,433	1,970	1,995	1,143	336	41	-
Female	152	184	279	462	860	1,556	2,789	3,708	3,094	1,445	314	-
White, both sexes	201	305	490	777	1,378	2,478	4,153	5,115	3,855	1,609	303	-
Male	102	179	284	420	689	1,204	1,739	1,806	1,033	311	39	-
Female	99	126	206	357	689	1,274	2,414	3,309	2,822	1,298	264	-
Black, both sexes	94	117	170	219	289	409	476	459	299	141	45	-
Male	49	67	107	132	152	177	185	137	84	20	1	-
Female	45	50	63	87	137	232	291	322	215	121	44	-
American Indian 1/, both sexes	1	4	8	7	9	12	15	18	11	4	2	-
Male	1	3	3	5	4	5	5	2	2	-	-	-
Female	-	1	5	2	5	7	10	16	9	4	2	-
Asian or Pacific Islander 2/, both sexes	13	16	11	34	61	90	115	111	72	27	5	-
Male	5	9	6	18	32	47	41	50	24	5	1	-
Female	8	7	5	16	29	43	74	61	48	22	4	-
Alabama												
All races, both sexes	11	9	4	22	28	43	60	75	49	13	5	-
Male	3	5	2	10	16	18	30	23	13	1	-	-
Female	8	4	2	12	12	25	30	52	36	12	5	-
White, both sexes	7	6	4	11	16	30	45	59	37	8	1	-
Male	3	4	2	5	10	14	22	18	8	1	-	-
Female	4	2	2	6	6	16	23	41	29	7	1	-
Black, both sexes	4	3	-	11	11	13	15	16	12	5	4	-
Male	-	1	-	5	6	4	8	5	5	-	-	-
Female	4	2	-	6	5	9	7	11	7	5	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Alaska												
All races, both sexes	-	2	-	3	2	3	7	6	2	-	-	-
Male	-	-	-	-	1	1	1	3	1	-	-	-
Female	-	2	-	3	1	2	6	3	1	-	-	-
White, both sexes	-	2	-	2	1	3	6	5	2	-	-	-
Male	-	-	-	-	1	1	-	3	1	-	-	-
Female	-	2	-	2	-	2	6	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	7	7	10	15	34	45	65	89	50	17	1	-
Male	3	6	5	7	17	23	30	31	14	4	-	-
Female	4	1	5	8	17	22	35	58	36	13	1	-
White, both sexes	7	6	10	14	30	41	61	87	48	16	1	-
Male	3	5	5	6	15	21	28	30	13	4	-	-
Female	4	1	5	8	15	20	33	57	35	12	1	-
Black, both sexes	-	1	-	-	4	1	-	1	1	1	-	-
Male	-	1	-	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	2	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	4	1	1	-	-	-
Male	-	-	-	1	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Arkansas												
All races, both sexes	5	9	10	17	26	46	57	73	50	14	4	-
Male	-	6	7	9	10	21	28	20	15	2	-	-
Female	5	3	3	8	16	25	29	53	35	12	4	-
White, both sexes	4	4	8	12	21	36	48	63	43	13	2	-
Male	-	2	6	7	9	20	23	19	13	2	-	-
Female	4	2	2	5	12	16	25	44	30	11	2	-
Black, both sexes	1	5	2	5	5	10	8	10	7	1	2	-
Male	-	4	1	2	1	1	5	1	2	-	-	-
Female	1	1	1	3	4	9	3	9	5	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
California												
All races, both sexes	45	74	90	143	246	474	763	829	662	292	61	-
Male	16	41	49	78	128	226	315	290	194	71	6	-
Female	29	33	41	65	118	248	448	539	468	221	55	-
White, both sexes	31	50	63	103	191	377	653	718	580	258	57	-
Male	11	25	34	57	105	176	273	246	167	67	5	-
Female	20	25	29	46	86	201	380	472	413	191	52	-
Black, both sexes	8	13	17	17	27	36	37	41	36	14	3	-
Male	3	10	10	10	11	20	15	12	11	1	-	-
Female	5	3	7	7	16	16	22	29	25	13	3	-
American Indian 1/, both sexes	-	-	2	-	1	1	3	4	1	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	2	-	1	-	2	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	6	11	8	23	27	60	70	66	45	19	1	-
Male	2	6	5	11	12	29	26	31	16	3	1	-
Female	4	5	3	12	15	31	44	35	29	16	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Colorado												
All races, both sexes	2	3	7	14	27	30	63	68	54	18	4	-
Male	1	1	3	7	13	12	24	26	19	1	1	-
Female	1	2	4	7	14	18	39	42	35	17	3	-
White, both sexes	2	3	6	12	23	28	61	67	54	18	4	-
Male	1	1	2	5	11	11	23	26	19	1	1	-
Female	1	2	4	7	12	17	38	41	35	17	3	-
Black, both sexes	-	-	1	2	3	-	-	-	-	-	-	-
Male	-	-	1	2	1	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Connecticut												
All races, both sexes	2	4	4	3	18	31	47	67	52	29	4	-
Male	-	2	4	2	9	10	21	24	10	4	-	-
Female	2	2	-	1	9	21	26	43	42	25	4	-
White, both sexes	2	3	3	3	16	29	44	65	51	29	2	-
Male	-	2	3	2	7	10	21	24	10	4	-	-
Female	2	1	-	1	9	19	23	41	41	25	2	-
Black, both sexes	-	1	1	-	1	2	3	1	1	-	2	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	3	1	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Delaware												
All races, both sexes	-	-	-	3	4	2	6	16	12	6	-	-
Male	-	-	-	2	2	1	1	5	4	-	-	-
Female	-	-	-	1	2	1	5	11	8	6	-	-
White, both sexes	-	-	-	3	4	2	4	14	10	6	-	-
Male	-	-	-	2	2	1	1	5	3	-	-	-
Female	-	-	-	1	2	1	3	9	7	6	-	-
Black, both sexes	-	-	-	-	-	-	2	2	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	2	1	1	3	8	3	5	2	-	-
Male	-	-	1	1	1	1	3	2	-	-	-	-
Female	-	1	1	-	-	2	5	1	5	2	-	-
White, both sexes	-	-	1	-	1	1	3	2	4	1	-	-
Male	-	-	-	-	1	1	1	2	-	-	-	-
Female	-	-	1	-	-	-	2	-	4	1	-	-
Black, both sexes	-	1	1	1	-	2	5	1	1	1	-	-
Male	-	-	1	1	-	-	2	-	-	-	-	-
Female	-	1	-	-	-	2	3	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Florida												
All races, both sexes	25	22	43	62	129	217	334	419	303	125	31	-
Male	15	12	28	31	66	108	147	157	90	19	7	-
Female	10	10	15	31	63	109	187	262	213	106	24	-
White, both sexes	10	18	27	40	92	183	284	372	279	108	23	-
Male	6	11	17	21	50	98	132	141	82	19	6	-
Female	4	7	10	19	42	85	152	231	197	89	17	-
Black, both sexes	15	4	16	21	36	32	46	45	23	17	8	-
Male	9	1	11	10	15	9	14	15	8	-	1	-
Female	6	3	5	11	21	23	32	30	15	17	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	4	2	1	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	3	1	1	-	-	-
Georgia												
All races, both sexes	12	8	28	37	33	74	98	103	76	33	6	-
Male	7	6	18	16	20	38	40	31	17	9	-	-
Female	5	2	10	21	13	36	58	72	59	24	6	-
White, both sexes	6	3	14	25	21	49	75	86	60	28	5	-
Male	3	3	8	8	8	24	32	26	15	7	-	-
Female	3	-	6	17	13	25	43	60	45	21	5	-
Black, both sexes	6	5	14	12	12	24	23	17	16	5	1	-
Male	4	3	10	8	12	14	8	5	2	2	-	-
Female	2	2	4	4	-	10	15	12	14	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Hawaii												
All races, both sexes	2	-	2	2	10	6	9	19	14	3	2	-
Male	1	-	2	2	6	3	5	9	4	1	-	-
Female	1	-	-	-	4	3	4	10	10	2	2	-
White, both sexes	-	-	2	1	1	2	2	4	-	-	-	-
Male	-	-	2	1	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	9	4	7	15	14	3	2	-
Male	1	-	-	1	5	2	4	8	4	1	-	-
Female	1	-	-	-	4	2	3	7	10	2	2	-
Idaho												
All races, both sexes	2	2	3	10	7	30	31	47	26	12	-	-
Male	1	1	2	5	5	12	9	20	6	2	-	-
Female	1	1	1	5	2	18	22	27	20	10	-	-
White, both sexes	2	2	3	9	7	29	30	47	26	12	-	-
Male	1	1	2	4	5	12	8	20	6	2	-	-
Female	1	1	1	5	2	17	22	27	20	10	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Illinois												
All races, both sexes	11	16	29	52	67	138	182	251	190	77	16	-
Male	7	8	19	31	42	70	75	90	60	13	3	-
Female	4	8	10	21	25	68	107	161	130	64	13	-
White, both sexes	5	8	15	32	47	113	150	230	170	69	13	-
Male	3	4	9	18	30	55	61	80	56	12	3	-
Female	2	4	6	14	17	58	89	150	114	57	10	-
Black, both sexes	6	8	14	19	19	23	29	17	17	8	3	-
Male	4	4	10	13	11	14	13	7	4	1	-	-
Female	2	4	4	6	8	9	16	10	13	7	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	3	4	3	-	-	-
Male	-	-	-	-	1	1	1	3	-	-	-	-
Female	-	-	-	1	-	1	2	1	3	-	-	-
Indiana												
All races, both sexes	8	11	21	34	37	57	84	113	79	44	8	-
Male	3	10	12	21	17	23	37	35	27	5	1	-
Female	5	1	9	13	20	34	47	78	52	39	7	-
White, both sexes	7	7	15	24	33	52	77	102	72	42	6	-
Male	2	7	9	16	14	22	33	33	26	5	1	-
Female	5	-	6	8	19	30	44	69	46	37	5	-
Black, both sexes	1	4	6	10	4	5	7	11	7	2	2	-
Male	1	3	3	5	3	1	4	2	1	-	-	-
Female	-	1	3	5	1	4	3	9	6	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Iowa												
All races, both sexes	2	4	4	7	12	30	47	44	57	22	6	-
Male	2	2	2	6	6	17	24	19	12	6	1	-
Female	-	2	2	1	6	13	23	25	45	16	5	-
White, both sexes	2	3	4	7	12	30	46	44	57	22	6	-
Male	2	2	2	6	6	17	24	19	12	6	1	-
Female	-	1	2	1	6	13	22	25	45	16	5	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	4	7	15	29	50	68	58	27	4	-
Male	-	1	2	3	9	16	22	23	11	2	-	-
Female	-	-	2	4	6	13	28	45	47	25	4	-
White, both sexes	-	1	4	7	14	24	49	65	58	27	4	-
Male	-	1	2	3	8	14	22	22	11	2	-	-
Female	-	-	2	4	6	10	27	43	47	25	4	-
Black, both sexes	-	-	-	-	1	3	1	3	-	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Kentucky												
All races, both sexes	4	6	12	16	16	36	55	58	45	12	4	-
Male	4	1	10	12	10	17	23	12	13	4	1	-
Female	-	5	2	4	6	19	32	46	32	8	3	-
White, both sexes	4	5	12	15	15	33	51	53	42	12	4	-
Male	4	1	10	11	10	16	22	11	12	4	1	-
Female	-	4	2	4	5	17	29	42	30	8	3	-
Black, both sexes	-	1	-	1	1	3	4	5	3	-	-	-
Male	-	-	-	1	-	1	1	1	1	-	-	-
Female	-	1	-	-	1	2	3	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	11	12	13	22	25	50	67	85	49	19	3	-
Male	7	6	8	14	13	21	24	20	10	5	-	-
Female	4	6	5	8	12	29	43	65	39	14	3	-
White, both sexes	3	7	8	13	16	38	53	64	43	14	-	-
Male	2	4	5	6	8	16	20	18	8	3	-	-
Female	1	3	3	7	8	22	33	46	35	11	-	-
Black, both sexes	8	4	5	9	9	12	14	21	6	5	3	-
Male	5	1	3	8	5	5	4	2	2	2	-	-
Female	3	3	2	1	4	7	10	19	4	3	3	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Maine												
All races, both sexes	1	1	3	4	7	10	13	24	24	13	-	-
Male	1	1	2	3	2	4	8	9	8	4	-	-
Female	-	-	1	1	5	6	5	15	16	9	-	-
White, both sexes	1	1	3	4	7	10	13	24	24	13	-	-
Male	1	1	2	3	2	4	8	9	8	4	-	-
Female	-	-	1	1	5	6	5	15	16	9	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	11	13	16	29	40	62	84	50	36	4	-
Male	3	5	9	11	13	21	28	31	18	11	-	-
Female	-	6	4	5	16	19	34	53	32	25	4	-
White, both sexes	2	6	4	9	18	31	47	73	45	29	4	-
Male	2	1	2	6	8	13	21	28	17	11	-	-
Female	-	5	2	3	10	18	26	45	28	18	4	-
Black, both sexes	1	5	9	7	11	7	13	9	4	6	-	-
Male	1	4	7	5	5	6	6	3	1	-	-	-
Female	-	1	2	2	6	1	7	6	3	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	2	2	1	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Massachusetts												
All races, both sexes	2	8	8	22	36	64	126	146	119	32	15	-
Male	1	5	6	14	21	39	50	53	28	4	2	-
Female	1	3	2	8	15	25	76	93	91	28	13	-
White, both sexes	2	8	7	17	32	54	117	142	118	31	14	-
Male	1	5	5	11	18	31	46	51	28	4	2	-
Female	1	3	2	6	14	23	71	91	90	27	12	-
Black, both sexes	-	-	1	4	3	8	3	3	1	-	-	-
Male	-	-	1	3	2	6	2	2	-	-	-	-
Female	-	-	-	1	1	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	6	1	-	1	1	-
Male	-	-	-	-	1	2	2	-	-	-	-	-
Female	-	-	-	1	-	-	4	1	-	1	1	-
Michigan												
All races, both sexes	19	22	31	44	68	132	207	254	182	66	18	-
Male	9	15	21	29	28	68	91	88	42	10	3	-
Female	10	7	10	15	40	64	116	166	140	56	15	-
White, both sexes	11	15	21	34	57	107	177	219	172	59	15	-
Male	5	11	13	21	24	59	79	72	39	7	3	-
Female	6	4	8	13	33	48	98	147	133	52	12	-
Black, both sexes	8	7	10	10	9	23	30	33	8	7	2	-
Male	4	4	8	8	4	8	12	16	3	3	-	-
Female	4	3	2	2	5	15	18	17	5	4	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	2	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Minnesota												
All races, both sexes	2	6	6	11	22	49	74	98	101	49	9	-
Male	1	5	3	9	15	23	34	44	22	10	1	-
Female	1	1	3	2	7	26	40	54	79	39	8	-
White, both sexes	1	4	4	9	19	47	71	97	100	47	9	-
Male	1	3	3	7	12	23	32	43	22	10	1	-
Female	-	1	1	2	7	24	39	54	78	37	8	-
Black, both sexes	-	1	-	-	-	-	1	-	1	1	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	1	1	-	1	2	-	-	-	1	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	2	-	2	1	-	-	-	-
Male	-	-	-	2	2	-	1	1	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	3	7	9	11	16	28	19	37	20	9	-	-
Male	3	7	7	8	9	15	6	13	3	2	-	-
Female	-	-	2	3	7	13	13	24	17	7	-	-
White, both sexes	-	3	5	8	10	24	15	27	11	5	-	-
Male	-	3	4	6	5	13	6	8	1	-	-	-
Female	-	-	1	2	5	11	9	19	10	5	-	-
Black, both sexes	3	4	4	3	6	4	4	10	9	4	-	-
Male	3	4	3	2	4	2	-	5	2	2	-	-
Female	-	-	1	1	2	2	4	5	7	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Missouri												
All races, both sexes	5	12	12	30	36	64	117	143	100	43	14	-
Male	4	9	4	17	16	22	39	47	26	10	3	-
Female	1	3	8	13	20	42	78	96	74	33	11	-
White, both sexes	4	7	7	22	33	56	105	131	94	40	13	-
Male	4	6	3	12	15	22	35	45	22	10	3	-
Female	-	1	4	10	18	34	70	86	72	30	10	-
Black, both sexes	1	5	5	8	3	8	12	10	5	3	1	-
Male	-	3	1	5	1	-	4	2	3	-	-	-
Female	1	2	4	3	2	8	8	8	2	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	3	2	-	3	10	6	14	23	15	7	1	-
Male	1	1	-	1	6	1	8	14	5	-	-	-
Female	2	1	-	2	4	5	6	9	10	7	1	-
White, both sexes	3	2	-	3	10	6	14	22	15	7	1	-
Male	1	1	-	1	6	1	8	14	5	-	-	-
Female	2	1	-	2	4	5	6	8	10	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Nebraska												
All races, both sexes	5	4	3	5	7	20	35	39	22	22	2	-
Male	3	3	2	2	3	11	12	10	9	4	-	-
Female	2	1	1	3	4	9	23	29	13	18	2	-
White, both sexes	5	3	3	5	7	18	33	38	22	22	2	-
Male	3	2	2	2	3	10	11	10	9	4	-	-
Female	2	1	1	3	4	8	22	28	13	18	2	-
Black, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Nevada												
All races, both sexes	3	-	5	5	10	6	21	19	14	4	-	-
Male	2	-	4	3	3	3	6	12	3	-	-	-
Female	1	-	1	2	7	3	15	7	11	4	-	-
White, both sexes	2	-	5	5	8	6	19	19	14	4	-	-
Male	1	-	4	3	2	3	5	12	3	-	-	-
Female	1	-	1	2	6	3	14	7	11	4	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Hampshire												
All races, both sexes	-	-	2	3	2	13	18	24	11	8	1	-
Male	-	-	1	2	1	8	10	7	3	2	1	-
Female	-	-	1	1	1	5	8	17	8	6	-	-
White, both sexes	-	-	2	3	2	13	18	24	11	8	1	-
Male	-	-	1	2	1	8	10	7	3	2	1	-
Female	-	-	1	1	1	5	8	17	8	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	7	6	12	25	52	83	100	62	29	5	-
Male	2	4	-	7	18	28	34	43	23	3	-	-
Female	2	3	6	5	7	24	49	57	39	26	5	-
White, both sexes	3	6	6	11	18	42	76	86	54	26	5	-
Male	2	3	-	6	14	23	31	41	21	3	-	-
Female	1	3	6	5	4	19	45	45	33	23	5	-
Black, both sexes	1	1	-	1	6	9	5	12	5	2	-	-
Male	-	1	-	1	4	4	3	2	1	-	-	-
Female	1	-	-	-	2	5	2	10	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	2	2	2	1	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	2	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Mexico												
All races, both sexes	6	1	7	10	17	24	45	45	28	12	3	-
Male	3	1	3	5	8	11	22	8	6	1	-	-
Female	3	-	4	5	9	13	23	37	22	11	3	-
White, both sexes	6	1	7	10	15	23	42	38	27	12	3	-
Male	3	1	3	5	7	10	21	8	6	1	-	-
Female	3	-	4	5	8	13	21	30	21	11	3	-
Black, both sexes	-	-	-	-	1	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	6	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	6	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5	8	21	28	59	79	151	185	128	51	19	-
Male	3	6	13	12	27	32	65	65	36	8	-	-
Female	2	2	8	16	32	47	86	120	92	43	19	-
White, both sexes	-	8	15	22	48	68	136	169	122	50	17	-
Male	-	6	9	9	20	29	59	59	35	8	-	-
Female	-	2	6	13	28	39	77	110	87	42	17	-
Black, both sexes	4	-	5	5	9	10	12	14	6	-	2	-
Male	3	-	4	3	6	3	4	4	1	-	-	-
Female	1	-	1	2	3	7	8	10	5	-	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	2	1	2	2	-	1	-	-
Male	-	-	-	-	1	-	2	2	-	-	-	-
Female	1	-	1	-	1	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
North Carolina												
All races, both sexes	14	13	39	40	53	89	149	170	131	53	7	-
Male	7	11	22	25	26	43	59	60	28	10	1	-
Female	7	2	17	15	27	46	90	110	103	43	6	-
White, both sexes	7	8	22	22	35	64	112	148	109	42	5	-
Male	4	7	12	16	14	33	45	53	23	8	1	-
Female	3	1	10	6	21	31	67	95	86	34	4	-
Black, both sexes	6	4	15	18	18	24	37	22	22	10	2	-
Male	2	3	9	9	12	10	14	7	5	2	-	-
Female	4	1	6	9	6	14	23	15	17	8	2	-
American Indian 1/, both sexes	-	1	2	-	-	1	-	-	-	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	5	4	16	21	13	7	1	-
Male	-	-	1	-	3	2	10	8	5	-	-	-
Female	-	-	-	1	2	2	6	13	8	7	1	-
White, both sexes	-	-	1	1	4	4	15	21	13	7	1	-
Male	-	-	1	-	2	2	10	8	5	-	-	-
Female	-	-	-	1	2	2	5	13	8	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Ohio												
All races, both sexes	19	37	37	55	91	148	285	335	221	79	12	-
Male	11	20	22	28	39	73	120	103	58	16	1	-
Female	8	17	15	27	52	75	165	232	163	63	11	-
White, both sexes	17	26	32	48	71	122	257	306	209	73	10	-
Male	10	13	19	23	31	62	107	98	57	15	1	-
Female	7	13	13	25	40	60	150	208	152	58	9	-
Black, both sexes	2	10	5	7	19	26	28	28	12	6	2	-
Male	1	6	3	5	8	11	13	4	1	1	-	-
Female	1	4	2	2	11	15	15	24	11	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	4	14	16	22	35	57	66	50	23	6	-
Male	2	2	6	9	9	19	21	18	13	5	-	-
Female	3	2	8	7	13	16	36	48	37	18	6	-
White, both sexes	3	3	12	11	22	30	53	61	44	23	4	-
Male	1	2	4	6	9	15	20	16	11	5	-	-
Female	2	1	8	5	13	15	33	45	33	18	4	-
Black, both sexes	1	1	1	4	-	5	3	2	3	-	-	-
Male	-	-	1	2	-	4	1	1	2	-	-	-
Female	1	1	-	2	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	1	3	3	-	1	-
Male	1	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	3	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Oregon												
All races, both sexes	4	3	6	7	23	42	84	101	69	35	4	-
Male	2	2	4	4	15	21	38	36	25	13	-	-
Female	2	1	2	3	8	21	46	65	44	22	4	-
White, both sexes	4	2	4	6	19	40	80	101	68	35	4	-
Male	2	2	3	4	12	20	38	36	24	13	-	-
Female	2	-	1	2	7	20	42	65	44	22	4	-
Black, both sexes	-	-	-	1	1	2	1	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	3	-	2	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	2	-	-	-	-	-
Pennsylvania												
All races, both sexes	8	17	36	41	62	119	210	275	204	103	17	-
Male	5	11	22	21	29	49	70	87	47	15	1	-
Female	3	6	14	20	33	70	140	188	157	88	16	-
White, both sexes	5	13	33	35	54	101	190	255	189	92	15	-
Male	3	8	22	17	24	42	62	78	42	13	1	-
Female	2	5	11	18	30	59	128	177	147	79	14	-
Black, both sexes	3	4	3	6	8	18	19	19	14	11	2	-
Male	2	3	-	4	5	7	7	9	5	2	-	-
Female	1	1	3	2	3	11	12	10	9	9	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Rhode Island												
All races, both sexes	2	-	1	3	3	9	22	17	8	4	2	-
Male	2	-	1	2	1	4	9	5	1	-	1	-
Female	-	-	-	1	2	5	13	12	7	4	1	-
White, both sexes	2	-	1	2	3	9	21	17	8	3	2	-
Male	2	-	1	1	1	4	8	5	1	-	1	-
Female	-	-	-	1	2	5	13	12	7	3	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	9	18	16	29	48	78	73	46	21	4	-
Male	2	4	9	12	11	28	34	23	9	4	-	-
Female	-	5	9	4	18	20	44	50	37	17	4	-
White, both sexes	-	5	12	9	23	31	56	51	37	18	2	-
Male	-	4	7	8	8	19	24	16	7	3	-	-
Female	-	1	5	1	15	12	32	35	30	15	2	-
Black, both sexes	2	4	6	6	6	17	20	22	9	3	2	-
Male	2	-	2	3	3	9	10	7	2	1	-	-
Female	-	4	4	3	3	8	10	15	7	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
South Dakota												
All races, both sexes	-	-	-	1	4	11	9	17	16	5	3	-
Male	-	-	-	1	3	6	3	11	3	2	1	-
Female	-	-	-	-	1	5	6	6	13	3	2	-
White, both sexes	-	-	-	1	4	10	9	17	16	5	3	-
Male	-	-	-	1	3	5	3	11	3	2	1	-
Female	-	-	-	-	1	5	6	6	13	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	13	17	23	38	74	88	96	76	38	5	-
Male	1	9	12	10	19	43	36	40	22	7	1	-
Female	2	4	5	13	19	31	52	56	54	31	4	-
White, both sexes	2	9	10	17	33	58	81	89	65	34	5	-
Male	-	5	7	7	17	33	36	37	17	6	1	-
Female	2	4	3	10	16	25	45	52	48	28	4	-
Black, both sexes	1	3	7	6	5	15	7	7	10	4	-	-
Male	1	3	5	3	2	9	-	3	4	1	-	-
Female	-	-	2	3	3	6	7	4	6	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Texas												
All races, both sexes	22	39	64	79	171	247	344	389	290	111	16	-
Male	8	18	31	41	87	117	130	125	68	17	3	-
Female	14	21	33	38	84	130	214	264	222	94	13	-
White, both sexes	17	25	47	62	134	201	291	342	253	94	15	-
Male	6	12	22	31	67	95	112	114	56	14	3	-
Female	11	13	25	31	67	106	179	228	197	80	12	-
Black, both sexes	5	12	17	15	35	44	50	44	35	16	1	-
Male	2	5	9	8	18	20	18	10	11	2	-	-
Female	3	7	8	7	17	24	32	34	24	14	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	2	2	3	3	2	1	-	-
Male	-	1	-	2	2	2	-	1	1	1	-	-
Female	-	-	-	-	-	-	3	2	1	-	-	-
Utah												
All races, both sexes	1	3	4	5	15	14	43	36	26	10	-	-
Male	1	1	2	2	7	6	15	10	10	4	-	-
Female	-	2	2	3	8	8	28	26	16	6	-	-
White, both sexes	1	2	4	5	15	13	43	35	25	10	-	-
Male	1	1	2	2	7	6	15	10	10	4	-	-
Female	-	1	2	3	8	7	28	25	15	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Vermont												
All races, both sexes	-	-	1	1	4	7	8	18	13	3	2	-
Male	-	-	1	-	2	2	4	5	5	2	-	-
Female	-	-	-	1	2	5	4	13	8	1	2	-
White, both sexes	-	-	1	1	4	7	8	18	13	3	2	-
Male	-	-	1	-	2	2	4	5	5	2	-	-
Female	-	-	-	1	2	5	4	13	8	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8	11	8	21	47	79	99	136	110	47	8	-
Male	3	6	5	12	19	32	43	56	25	1	-	-
Female	5	5	3	9	28	47	56	80	85	46	8	-
White, both sexes	4	7	6	16	31	58	76	108	92	39	5	-
Male	2	5	4	8	13	26	34	47	21	1	-	-
Female	2	2	2	8	18	32	42	61	71	38	5	-
Black, both sexes	4	4	2	5	12	18	23	28	18	8	3	-
Male	1	1	1	4	5	5	9	9	4	-	-	-
Female	3	3	1	1	7	13	14	19	14	8	3	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	3	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	2	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Washington												
All races, both sexes	6	5	7	20	35	50	91	129	104	37	8	-
Male	5	1	6	12	14	26	39	51	30	10	1	-
Female	1	4	1	8	21	24	52	78	74	27	7	-
White, both sexes	4	5	4	17	29	47	83	123	100	37	8	-
Male	4	1	3	10	11	23	37	49	29	10	1	-
Female	-	4	1	7	18	24	46	74	71	27	7	-
Black, both sexes	1	-	2	2	2	-	2	1	2	-	-	-
Male	1	-	2	1	1	-	-	-	1	-	-	-
Female	-	-	-	1	1	-	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	1	2	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	3	2	5	4	-	-	-	-
Male	-	-	1	1	2	2	1	2	-	-	-	-
Female	1	-	-	-	1	-	4	2	-	-	-	-
West Virginia												
All races, both sexes	2	4	5	11	10	19	35	29	14	15	5	-
Male	1	1	2	5	6	7	9	8	1	3	1	-
Female	1	3	3	6	4	12	26	21	13	12	4	-
White, both sexes	2	4	4	10	10	19	33	28	14	15	5	-
Male	1	1	2	4	6	7	9	8	1	3	1	-
Female	1	3	2	6	4	12	24	20	13	12	4	-
Black, both sexes	-	-	1	1	-	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Wisconsin												
All races, both sexes	3	2	9	21	41	60	116	138	132	43	5	-
Male	1	1	5	11	24	29	55	64	39	9	-	-
Female	2	1	4	10	17	31	61	74	93	34	5	-
White, both sexes	1	2	9	17	39	56	114	136	130	42	5	-
Male	1	1	5	10	22	29	53	63	38	9	-	-
Female	-	1	4	7	17	27	61	73	92	33	5	-
Black, both sexes	2	-	-	2	1	4	2	1	1	1	-	-
Male	-	-	-	-	1	-	2	1	-	-	-	-
Female	2	-	-	2	-	4	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	-	2	-	2	3	3	7	3	5	1	-	-
Male	-	1	-	-	2	2	3	1	2	-	-	-
Female	-	1	-	2	1	1	4	2	3	1	-	-
White, both sexes	-	2	-	2	3	3	6	3	5	1	-	-
Male	-	1	-	-	2	2	3	1	2	-	-	-
Female	-	1	-	2	1	1	3	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
United States												
All races, both sexes	31,136	20	2	3	9	27	42	71	84	176	299	559
Male	15,195	13	-	2	7	16	29	52	59	123	217	388
Female	15,941	7	2	1	2	11	13	19	25	53	82	171
White, both sexes	27,317	15	1	2	7	23	33	50	60	127	205	405
Male	13,365	10	-	2	7	12	24	37	43	85	151	291
Female	13,952	5	1	-	-	11	9	13	17	42	54	114
Black, both sexes	3,191	5	1	-	2	4	7	18	21	38	87	136
Male	1,496	3	-	-	-	4	4	14	14	28	60	88
Female	1,695	2	1	-	2	-	3	4	7	10	27	48
American Indian 1/, both sexes	106	-	-	-	-	-	1	-	-	1	1	7
Male	40	-	-	-	-	-	-	-	-	1	1	2
Female	66	-	-	-	-	-	1	-	-	-	-	5
Asian or Pacific Islander 2/, both sexes	522	-	-	1	-	-	1	3	3	10	6	11
Male	294	-	-	-	-	-	1	1	2	9	5	7
Female	228	-	-	1	-	-	-	2	1	1	1	4
Alabama												
All races, both sexes	527	-	-	-	-	-	2	1	2	5	12	20
Male	270	-	-	-	-	-	2	1	-	5	9	13
Female	257	-	-	-	-	-	-	-	2	-	3	7
White, both sexes	378	-	-	-	-	-	1	-	1	1	6	8
Male	192	-	-	-	-	-	1	-	-	1	4	7
Female	186	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	148	-	-	-	-	-	1	1	1	4	6	12
Male	78	-	-	-	-	-	1	1	-	4	5	6
Female	70	-	-	-	-	-	-	-	1	-	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Alaska												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	563	-	-	-	-	1	1	2	1	1	3	10
Male	299	-	-	-	-	-	1	2	1	-	2	6
Female	264	-	-	-	-	1	-	-	-	1	1	4
White, both sexes	541	-	-	-	-	1	1	1	-	1	3	8
Male	286	-	-	-	-	-	1	1	-	-	2	6
Female	255	-	-	-	-	1	-	-	-	1	1	2
Black, both sexes	11	-	-	-	-	-	-	1	1	-	-	-
Male	9	-	-	-	-	-	-	1	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Arkansas												
All races, both sexes	330	-	-	-	1	-	-	1	2	2	5	10
Male	182	-	-	-	1	-	-	1	2	1	3	8
Female	148	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	290	-	-	-	1	-	-	-	2	-	4	9
Male	168	-	-	-	1	-	-	-	2	-	2	7
Female	122	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	39	-	-	-	-	-	-	1	-	2	1	1
Male	14	-	-	-	-	-	-	1	-	1	1	1
Female	25	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,949	1	-	-	-	4	5	5	11	24	44	65
Male	1,502	1	-	-	-	2	3	4	4	16	35	47
Female	1,447	-	-	-	-	2	2	1	7	8	9	18
White, both sexes	2,503	1	-	-	-	3	3	4	7	16	37	49
Male	1,269	1	-	-	-	1	2	4	3	9	31	36
Female	1,234	-	-	-	-	2	1	-	4	7	6	13
Black, both sexes	204	-	-	-	-	1	1	-	2	3	5	10
Male	92	-	-	-	-	1	-	-	-	2	3	7
Female	112	-	-	-	-	-	1	-	2	1	2	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	238	-	-	-	-	-	1	1	2	5	2	5
Male	139	-	-	-	-	-	1	-	1	5	1	4
Female	99	-	-	-	-	-	-	1	1	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Colorado												
All races, both sexes	650	-	1	1	-	-	1	2	-	5	3	11
Male	268	-	-	1	-	-	1	1	-	2	3	8
Female	382	-	1	-	-	-	-	1	-	3	-	3
White, both sexes	614	-	1	1	-	-	1	2	-	5	3	10
Male	249	-	-	1	-	-	1	1	-	2	3	8
Female	365	-	1	-	-	-	-	1	-	3	-	2
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	414	-	-	-	-	-	-	3	2	2	1	4
Male	189	-	-	-	-	-	-	3	1	1	1	2
Female	225	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	380	-	-	-	-	-	-	2	2	2	1	4
Male	170	-	-	-	-	-	-	2	1	1	1	2
Female	210	-	-	-	-	-	-	-	1	1	-	2
Black, both sexes	30	-	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	1	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Delaware												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	1	-
Male	32	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	61	-	-	-	-	-	-	-	-	1	-	2
Male	28	-	-	-	-	-	-	-	-	1	-	2
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	45	-	-	-	-	-	-	-	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Florida												
All races, both sexes	2,218	1	-	2	1	2	-	3	2	9	12	36
Male	1,192	1	-	1	1	1	-	2	2	7	8	30
Female	1,026	-	-	1	-	1	-	1	-	2	4	6
White, both sexes	2,000	-	-	1	1	1	-	3	1	3	6	20
Male	1,080	-	-	1	1	-	-	2	1	3	4	18
Female	920	-	-	-	-	1	-	1	-	-	2	2
Black, both sexes	209	1	-	-	-	1	-	-	1	6	5	15
Male	107	1	-	-	-	1	-	-	1	4	3	11
Female	102	-	-	-	-	-	-	-	-	2	2	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	1	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	741	1	-	-	-	-	2	2	3	9	12	22
Male	357	-	-	-	-	-	-	1	2	8	10	12
Female	384	1	-	-	-	-	2	1	1	1	2	10
White, both sexes	549	1	-	-	-	-	1	-	1	6	2	15
Male	253	-	-	-	-	-	-	-	-	5	2	8
Female	296	1	-	-	-	-	1	-	1	1	-	7
Black, both sexes	184	-	-	-	-	-	1	2	2	3	10	7
Male	100	-	-	-	-	-	-	1	2	3	8	4
Female	84	-	-	-	-	-	1	1	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Hawaii												
All races, both sexes	115	-	-	-	-	-	-	-	-	-	1	4
Male	58	-	-	-	-	-	-	-	-	-	1	3
Female	57	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	3
Male	22	-	-	-	-	-	-	-	-	-	-	3
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	84	-	-	-	-	-	-	-	-	-	1	1
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	135	-	-	-	1	-	-	1	-	2	-	3
Male	80	-	-	-	1	-	-	1	-	1	-	3
Female	55	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	130	-	-	-	1	-	-	1	-	2	-	2
Male	77	-	-	-	1	-	-	1	-	1	-	2
Female	53	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Illinois												
All races, both sexes	1,309	-	-	-	-	2	1	3	3	6	10	24
Male	582	-	-	-	-	1	1	3	3	4	8	18
Female	727	-	-	-	-	1	-	-	-	2	2	6
White, both sexes	1,136	-	-	-	-	1	-	2	3	3	9	17
Male	508	-	-	-	-	-	-	2	3	1	8	14
Female	628	-	-	-	-	1	-	-	-	2	1	3
Black, both sexes	160	-	-	-	-	1	1	1	-	2	1	6
Male	66	-	-	-	-	1	1	1	-	2	-	3
Female	94	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	850	2	-	-	-	-	-	1	3	6	7	17
Male	430	1	-	-	-	-	-	1	3	3	6	13
Female	420	1	-	-	-	-	-	-	-	3	1	4
White, both sexes	793	1	-	-	-	-	-	-	2	5	6	15
Male	403	-	-	-	-	-	-	-	2	3	5	12
Female	390	1	-	-	-	-	-	-	-	2	1	3
Black, both sexes	55	1	-	-	-	-	-	1	1	1	1	2
Male	26	1	-	-	-	-	-	1	1	-	1	1
Female	29	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Iowa												
All races, both sexes	472	-	-	-	-	-	-	-	1	2	3	5
Male	224	-	-	-	-	-	-	-	-	2	1	4
Female	248	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	463	-	-	-	-	-	-	-	1	2	3	5
Male	219	-	-	-	-	-	-	-	-	2	1	4
Female	244	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	559	-	-	-	-	-	-	3	-	1	7	6
Male	228	-	-	-	-	-	-	2	-	-	5	4
Female	331	-	-	-	-	-	-	1	-	1	2	2
White, both sexes	536	-	-	-	-	-	-	2	-	1	7	6
Male	218	-	-	-	-	-	-	1	-	-	5	4
Female	318	-	-	-	-	-	-	1	-	1	2	2
Black, both sexes	21	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Kentucky												
All races, both sexes	500	-	-	-	-	-	1	4	-	2	6	6
Male	246	-	-	-	-	-	1	2	-	1	6	3
Female	254	-	-	-	-	-	-	2	-	1	-	3
White, both sexes	463	-	-	-	-	-	1	4	-	2	4	4
Male	224	-	-	-	-	-	1	2	-	1	4	2
Female	239	-	-	-	-	-	-	2	-	1	-	2
Black, both sexes	35	-	-	-	-	-	-	-	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	2	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	461	-	-	-	-	-	-	2	2	3	8	15
Male	211	-	-	-	-	-	-	1	2	3	5	7
Female	250	-	-	-	-	-	-	1	-	-	3	8
White, both sexes	325	-	-	-	-	-	-	-	1	3	4	5
Male	152	-	-	-	-	-	-	-	1	3	3	3
Female	173	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	132	-	-	-	-	-	-	2	1	-	4	10
Male	57	-	-	-	-	-	-	1	1	-	2	4
Female	75	-	-	-	-	-	-	1	-	-	2	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Maine												
All races, both sexes	155	-	-	-	-	-	1	-	-	-	2	6
Male	80	-	-	-	-	-	1	-	-	-	1	5
Female	75	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	154	-	-	-	-	-	1	-	-	-	2	6
Male	79	-	-	-	-	-	1	-	-	-	1	5
Female	75	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	492	-	-	-	-	1	2	1	2	5	6	11
Male	210	-	-	-	-	1	2	1	1	3	5	8
Female	282	-	-	-	-	-	-	-	1	2	1	3
White, both sexes	351	-	-	-	-	1	1	-	-	3	3	5
Male	152	-	-	-	-	1	1	-	-	1	3	2
Female	199	-	-	-	-	-	-	-	-	2	-	3
Black, both sexes	134	-	-	-	-	-	1	1	2	2	3	6
Male	54	-	-	-	-	-	1	1	1	2	2	6
Female	80	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Massachusetts												
All races, both sexes	745	-	-	-	-	3	1	-	1	4	3	5
Male	348	-	-	-	-	3	1	-	1	4	2	5
Female	397	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	721	-	-	-	-	3	1	-	1	3	3	5
Male	335	-	-	-	-	3	1	-	1	3	2	5
Female	386	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,288	2	-	-	-	-	1	2	4	4	14	23
Male	610	2	-	-	-	-	1	2	4	3	11	16
Female	678	-	-	-	-	-	-	-	-	1	3	7
White, both sexes	1,096	2	-	-	-	-	1	2	4	3	9	18
Male	521	2	-	-	-	-	1	2	4	2	7	12
Female	575	-	-	-	-	-	-	-	-	1	2	6
Black, both sexes	179	-	-	-	-	-	-	-	-	1	5	4
Male	85	-	-	-	-	-	-	-	-	1	4	4
Female	94	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Minnesota												
All races, both sexes	479	-	-	-	-	-	-	-	-	2	1	4
Male	236	-	-	-	-	-	-	-	-	1	1	2
Female	243	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	470	-	-	-	-	-	-	-	-	2	1	4
Male	230	-	-	-	-	-	-	-	-	1	1	2
Female	240	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	231	-	-	-	-	-	1	1	-	2	5	4
Male	128	-	-	-	-	-	1	-	-	1	4	2
Female	103	-	-	-	-	-	-	1	-	1	1	2
White, both sexes	165	-	-	-	-	-	1	1	-	2	1	1
Male	93	-	-	-	-	-	1	-	-	1	-	-
Female	72	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	65	-	-	-	-	-	-	-	-	-	4	3
Male	34	-	-	-	-	-	-	-	-	-	4	2
Female	31	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Missouri												
All races, both sexes	768	2	-	-	-	-	-	-	3	-	6	12
Male	350	1	-	-	-	-	-	-	2	-	1	11
Female	418	1	-	-	-	-	-	-	1	-	5	1
White, both sexes	669	2	-	-	-	-	-	-	2	-	3	10
Male	316	1	-	-	-	-	-	-	2	-	1	9
Female	353	1	-	-	-	-	-	-	-	-	2	1
Black, both sexes	95	-	-	-	-	-	-	-	1	-	3	2
Male	33	-	-	-	-	-	-	-	-	-	-	2
Female	62	-	-	-	-	-	-	-	1	-	3	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	96	-	-	-	-	-	-	-	-	1	-	1
Male	53	-	-	-	-	-	-	-	-	1	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	93	-	-	-	-	-	-	-	-	1	-	1
Male	51	-	-	-	-	-	-	-	-	1	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Nebraska												
All races, both sexes	223	-	-	-	-	-	1	-	1	-	1	1
Male	95	-	-	-	-	-	1	-	1	-	1	1
Female	128	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	215	-	-	-	-	-	1	-	1	-	1	1
Male	91	-	-	-	-	-	1	-	1	-	1	1
Female	124	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	296	-	-	-	-	1	-	3	1	2	1	5
Male	136	-	-	-	-	1	-	1	1	2	1	1
Female	160	-	-	-	-	-	-	2	-	-	-	4
White, both sexes	273	-	-	-	-	1	-	1	1	1	-	4
Male	124	-	-	-	-	1	-	1	1	1	-	1
Female	149	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	16	-	-	-	-	-	-	1	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	8	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Hampshire												
All races, both sexes	133	-	-	-	-	-	-	-	-	2	1	4
Male	73	-	-	-	-	-	-	-	-	1	1	4
Female	60	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	132	-	-	-	-	-	-	-	-	2	1	4
Male	72	-	-	-	-	-	-	-	-	1	1	4
Female	60	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	902	-	-	-	1	-	2	2	-	3	2	8
Male	461	-	-	-	1	-	2	2	-	3	1	3
Female	441	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	783	-	-	-	1	-	2	-	-	2	1	6
Male	404	-	-	-	1	-	2	-	-	2	1	2
Female	379	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	100	-	-	-	-	-	-	2	-	-	1	2
Male	46	-	-	-	-	-	-	2	-	-	-	1
Female	54	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Mexico												
All races, both sexes	163	-	-	-	-	-	1	-	-	1	1	1
Male	86	-	-	-	-	-	1	-	-	1	1	1
Female	77	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	152	-	-	-	-	-	1	-	-	1	1	1
Male	80	-	-	-	-	-	1	-	-	1	1	1
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,671	1	1	-	2	-	-	3	1	3	15	34
Male	812	1	-	-	2	-	-	3	-	3	11	25
Female	859	-	1	-	-	-	-	-	1	-	4	9
White, both sexes	1,451	1	-	-	2	-	-	3	1	3	9	24
Male	709	1	-	-	2	-	-	3	-	3	9	19
Female	742	-	-	-	-	-	-	-	1	-	-	5
Black, both sexes	197	-	1	-	-	-	-	-	-	-	6	10
Male	90	-	-	-	-	-	-	-	-	-	2	6
Female	107	-	1	-	-	-	-	-	-	-	4	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
North Carolina												
All races, both sexes	956	-	-	-	-	1	1	1	2	6	10	25
Male	484	-	-	-	-	-	1	1	2	6	9	15
Female	472	-	-	-	-	1	-	-	-	-	1	10
White, both sexes	753	-	-	-	-	1	1	-	-	4	4	18
Male	388	-	-	-	-	-	1	-	-	4	4	11
Female	365	-	-	-	-	1	-	-	-	-	-	7
Black, both sexes	197	-	-	-	-	-	-	-	2	2	6	7
Male	93	-	-	-	-	-	-	-	2	2	5	4
Female	104	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	76	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	75	-	-	-	-	-	-	-	-	-	-	1
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Ohio												
All races, both sexes	1,455	2	-	-	-	1	4	1	5	6	14	26
Male	695	1	-	-	-	1	2	1	4	4	9	18
Female	760	1	-	-	-	-	2	-	1	2	5	8
White, both sexes	1,297	1	-	-	-	1	4	1	4	4	7	18
Male	627	1	-	-	-	1	2	1	4	2	4	12
Female	670	-	-	-	-	-	2	-	-	2	3	6
Black, both sexes	154	1	-	-	-	-	-	-	1	2	7	8
Male	67	-	-	-	-	-	-	-	-	2	5	6
Female	87	1	-	-	-	-	-	-	1	-	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	393	-	-	-	-	-	1	-	1	3	3	10
Male	189	-	-	-	-	-	1	-	-	2	2	9
Female	204	-	-	-	-	-	-	-	1	1	1	1
White, both sexes	340	-	-	-	-	-	1	-	1	3	2	10
Male	172	-	-	-	-	-	1	-	-	2	1	9
Female	168	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Oregon												
All races, both sexes	431	-	-	-	-	1	-	1	-	2	1	4
Male	217	-	-	-	-	-	-	-	-	-	1	3
Female	214	-	-	-	-	1	-	1	-	2	-	1
White, both sexes	422	-	-	-	-	1	-	1	-	2	1	4
Male	213	-	-	-	-	-	-	-	-	-	1	3
Female	209	-	-	-	-	1	-	1	-	2	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,680	2	-	-	-	2	-	1	3	7	18	12
Male	775	1	-	-	-	1	-	1	3	3	15	8
Female	905	1	-	-	-	1	-	-	-	4	3	4
White, both sexes	1,553	2	-	-	-	2	-	1	3	7	16	9
Male	724	1	-	-	-	1	-	1	3	3	13	6
Female	829	1	-	-	-	1	-	-	-	4	3	3
Black, both sexes	123	-	-	-	-	-	-	-	-	-	1	2
Male	48	-	-	-	-	-	-	-	-	-	1	1
Female	75	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Rhode Island												
All races, both sexes	143	-	-	-	-	-	-	-	-	1	-	-
Male	62	-	-	-	-	-	-	-	-	1	-	-
Female	81	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	140	-	-	-	-	-	-	-	-	1	-	-
Male	62	-	-	-	-	-	-	-	-	1	-	-
Female	78	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	401	-	-	-	-	1	3	1	2	2	3	8
Male	202	-	-	-	-	1	3	1	2	1	1	5
Female	199	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	310	-	-	-	-	1	2	1	1	1	1	5
Male	164	-	-	-	-	1	2	1	1	1	-	3
Female	146	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	89	-	-	-	-	-	1	-	1	1	2	3
Male	37	-	-	-	-	-	1	-	1	-	1	2
Female	52	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
South Dakota												
All races, both sexes	99	-	-	-	-	-	-	-	-	1	-	1
Male	42	-	-	-	-	-	-	-	-	1	-	1
Female	57	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	96	-	-	-	-	-	-	-	-	1	-	-
Male	40	-	-	-	-	-	-	-	-	1	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	670	-	-	-	-	2	2	4	4	4	11	17
Male	308	-	-	-	-	1	1	4	4	3	7	10
Female	362	-	-	-	-	1	1	-	-	1	4	7
White, both sexes	578	-	-	-	-	2	1	3	1	4	9	12
Male	265	-	-	-	-	1	1	3	1	3	5	7
Female	313	-	-	-	-	1	-	-	-	1	4	5
Black, both sexes	91	-	-	-	-	-	1	1	3	-	2	5
Male	43	-	-	-	-	-	-	1	3	-	2	3
Female	48	-	-	-	-	-	1	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Texas												
All races, both sexes	1,665	4	-	-	2	3	5	9	12	17	26	40
Male	833	2	-	-	-	2	1	4	9	14	12	26
Female	832	2	-	-	2	1	4	5	3	3	14	14
White, both sexes	1,444	3	-	-	-	3	5	8	12	12	21	30
Male	716	2	-	-	-	2	1	3	9	10	8	20
Female	728	1	-	-	-	1	4	5	3	2	13	10
Black, both sexes	203	1	-	-	2	-	-	1	-	4	5	10
Male	101	-	-	-	-	-	-	1	-	3	4	6
Female	102	1	-	-	2	-	-	-	-	1	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	184	-	-	-	-	-	-	1	1	-	3	3
Male	108	-	-	-	-	-	-	1	-	-	3	2
Female	76	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	177	-	-	-	-	-	-	1	1	-	1	2
Male	103	-	-	-	-	-	-	1	-	-	1	2
Female	74	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Vermont												
All races, both sexes	69	-	-	-	-	-	-	1	-	-	1	-
Male	30	-	-	-	-	-	-	1	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	69	-	-	-	-	-	-	1	-	-	1	-
Male	30	-	-	-	-	-	-	1	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	679	1	-	-	1	1	-	4	5	3	3	14
Male	331	1	-	-	1	1	-	3	2	1	2	12
Female	348	-	-	-	-	-	-	1	3	2	1	2
White, both sexes	534	-	-	-	1	-	-	3	3	2	1	8
Male	254	-	-	-	1	-	-	3	1	1	1	7
Female	280	-	-	-	-	-	-	-	2	1	-	1
Black, both sexes	138	1	-	-	-	1	-	1	2	1	2	6
Male	72	1	-	-	-	1	-	-	1	-	1	5
Female	66	-	-	-	-	-	-	1	1	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Washington												
All races, both sexes	684	-	-	-	-	1	1	-	1	8	6	2
Male	346	-	-	-	-	-	-	-	1	7	5	-
Female	338	-	-	-	-	1	1	-	-	1	1	2
White, both sexes	645	-	-	-	-	1	-	-	-	5	5	2
Male	326	-	-	-	-	-	-	-	-	5	5	-
Female	319	-	-	-	-	1	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	1	2	-	-
Male	13	-	-	-	-	-	-	-	1	1	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	284	-	-	-	-	-	-	1	-	1	1	6
Male	147	-	-	-	-	-	-	-	-	1	1	5
Female	137	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	280	-	-	-	-	-	-	1	-	1	1	6
Male	143	-	-	-	-	-	-	-	-	1	1	5
Female	137	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Wisconsin												
All races, both sexes	629	1	-	-	-	-	2	1	3	6	6	8
Male	284	1	-	-	-	-	1	1	2	1	4	5
Female	345	-	-	-	-	-	1	-	1	5	2	3
White, both sexes	607	1	-	-	-	-	2	1	3	4	5	8
Male	275	1	-	-	-	-	1	1	2	1	3	5
Female	332	-	-	-	-	-	1	-	1	3	2	3
Black, both sexes	16	-	-	-	-	-	-	-	-	2	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
United States												
All races, both sexes	743	1,112	1,429	2,025	2,928	4,407	5,631	5,529	4,036	1,632	372	-
Male	517	776	932	1,280	1,696	2,373	2,659	2,323	1,280	392	61	-
Female	226	336	497	745	1,232	2,034	2,972	3,206	2,756	1,240	311	-
White, both sexes	532	866	1,187	1,699	2,550	3,922	5,027	5,077	3,710	1,489	330	-
Male	383	607	791	1,086	1,503	2,157	2,415	2,155	1,196	356	54	-
Female	149	259	396	613	1,047	1,765	2,612	2,922	2,514	1,133	276	-
Black, both sexes	187	222	216	278	314	391	478	363	266	123	34	-
Male	118	153	125	171	155	167	177	122	59	29	5	-
Female	69	69	91	107	159	224	301	241	207	94	29	-
American Indian 1/, both sexes	3	9	3	15	12	15	19	9	7	4	-	-
Male	3	5	1	5	6	7	5	4	-	-	-	-
Female	-	4	2	10	6	8	14	5	7	4	-	-
Asian or Pacific Islander 2/, both sexes	21	15	23	33	52	79	107	80	53	16	8	-
Male	13	11	15	18	32	42	62	42	25	7	2	-
Female	8	4	8	15	20	37	45	38	28	9	6	-
Alabama												
All races, both sexes	14	28	29	52	57	62	93	83	37	26	4	-
Male	10	19	19	30	35	35	48	33	4	7	-	-
Female	4	9	10	22	22	27	45	50	33	19	4	-
White, both sexes	8	18	22	38	44	48	73	63	25	19	3	-
Male	7	11	16	21	26	25	41	26	1	5	-	-
Female	1	7	6	17	18	23	32	37	24	14	3	-
Black, both sexes	6	10	7	14	13	14	20	20	12	6	1	-
Male	3	8	3	9	9	10	7	7	3	2	-	-
Female	3	2	4	5	4	4	13	13	9	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Alaska												
All races, both sexes	2	2	4	2	9	5	3	4	-	1	-	-
Male	1	2	3	1	6	4	1	1	-	-	-	-
Female	1	-	1	1	3	1	2	3	-	1	-	-
White, both sexes	2	2	4	1	8	4	3	3	-	1	-	-
Male	1	2	3	1	6	3	1	1	-	-	-	-
Female	1	-	1	-	2	1	2	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	21	12	29	50	55	64	105	116	67	23	2	-
Male	14	8	18	32	34	33	55	51	33	9	-	-
Female	7	4	11	18	21	31	50	65	34	14	2	-
White, both sexes	20	11	28	48	49	62	102	116	65	23	2	-
Male	14	7	17	31	31	31	53	51	32	9	-	-
Female	6	4	11	17	18	31	49	65	33	14	2	-
Black, both sexes	-	1	1	1	4	1	-	-	1	-	-	-
Male	-	1	1	1	2	1	-	-	1	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	-	3	-	1	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Arkansas												
All races, both sexes	15	16	25	28	37	55	45	48	28	10	2	-
Male	9	13	16	21	27	33	21	16	9	1	-	-
Female	6	3	9	7	10	22	24	32	19	9	2	-
White, both sexes	12	12	22	27	33	52	40	43	22	9	2	-
Male	8	11	14	20	25	33	20	16	8	1	-	-
Female	4	1	8	7	8	19	20	27	14	8	2	-
Black, both sexes	2	4	3	1	4	3	5	5	6	1	-	-
Male	1	2	2	1	2	-	1	-	1	-	-	-
Female	1	2	1	-	2	3	4	5	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	81	101	119	180	271	392	533	528	377	162	46	-
Male	57	70	74	114	159	206	275	241	148	35	11	-
Female	24	31	45	66	112	186	258	287	229	127	35	-
White, both sexes	55	82	95	150	224	327	463	463	341	143	40	-
Male	41	58	61	96	132	174	237	209	135	28	11	-
Female	14	24	34	54	92	153	226	254	206	115	29	-
Black, both sexes	15	11	16	20	21	26	23	22	15	9	4	-
Male	6	8	9	14	9	10	9	9	4	1	-	-
Female	9	3	7	6	12	16	14	13	11	8	4	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	8	7	10	25	39	47	43	21	10	2	-
Male	9	4	3	4	18	22	29	23	9	6	-	-
Female	1	4	4	6	7	17	18	20	12	4	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Colorado												
All races, both sexes	8	10	24	34	41	89	123	139	90	52	16	-
Male	5	9	17	22	19	49	38	56	24	10	3	-
Female	3	1	7	12	22	40	85	83	66	42	13	-
White, both sexes	7	8	22	31	40	84	117	131	88	49	14	-
Male	5	7	15	20	19	46	35	52	23	9	2	-
Female	2	1	7	11	21	38	82	79	65	40	12	-
Black, both sexes	-	2	2	2	1	5	4	5	1	3	-	-
Male	-	2	2	2	-	3	2	1	-	1	-	-
Female	-	-	-	-	1	2	2	4	1	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	2	1	-	2	-
Male	-	-	-	-	-	-	1	2	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	4	10	13	15	33	53	81	70	83	35	5	-
Male	3	8	9	12	20	27	35	30	31	5	1	-
Female	1	2	4	3	13	26	46	40	52	30	4	-
White, both sexes	2	6	10	14	27	49	75	67	79	35	5	-
Male	2	6	8	12	15	25	32	29	28	5	1	-
Female	-	-	2	2	12	24	43	38	51	30	4	-
Black, both sexes	1	4	3	1	6	3	6	3	2	-	-	-
Male	-	2	1	-	5	1	3	1	1	-	-	-
Female	1	2	2	1	1	2	3	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	2	-	-	-
Male	1	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Delaware												
All races, both sexes	1	-	2	4	9	9	8	13	4	5	-	-
Male	1	-	1	2	8	6	4	8	-	1	-	-
Female	-	-	1	2	1	3	4	5	4	4	-	-
White, both sexes	-	-	1	4	5	9	8	13	3	5	-	-
Male	-	-	1	2	5	6	4	8	-	1	-	-
Female	-	-	-	2	-	3	4	5	3	4	-	-
Black, both sexes	1	-	1	-	4	-	-	-	1	-	-	-
Male	1	-	-	-	3	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	4	3	5	6	8	14	9	6	1	-	-
Male	2	2	1	4	5	4	4	2	1	-	-	-
Female	-	2	2	1	1	4	10	7	5	1	-	-
White, both sexes	-	-	2	1	1	2	2	5	1	-	-	-
Male	-	-	-	1	1	1	-	1	-	-	-	-
Female	-	-	2	-	-	1	2	4	1	-	-	-
Black, both sexes	2	4	1	4	5	6	11	4	5	1	-	-
Male	2	2	1	3	4	3	3	1	1	-	-	-
Female	-	2	-	1	1	3	8	3	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Florida												
All races, both sexes	60	79	112	153	243	332	361	383	289	115	23	-
Male	43	57	79	105	147	194	200	164	115	30	5	-
Female	17	22	33	48	96	138	161	219	174	85	18	-
White, both sexes	50	67	96	134	217	312	332	357	275	104	20	-
Male	36	47	70	95	133	183	184	155	113	29	5	-
Female	14	20	26	39	84	129	148	202	162	75	15	-
Black, both sexes	10	12	15	19	25	19	26	26	14	11	3	-
Male	7	10	9	10	13	10	15	9	2	1	-	-
Female	3	2	6	9	12	9	11	17	12	10	3	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	24	47	54	55	69	89	114	115	76	38	9	-
Male	14	33	40	36	40	39	54	42	18	7	1	-
Female	10	14	14	19	29	50	60	73	58	31	8	-
White, both sexes	10	32	33	35	57	72	94	95	62	27	6	-
Male	7	21	23	22	33	30	47	35	14	5	1	-
Female	3	11	10	13	24	42	47	60	48	22	5	-
Black, both sexes	13	15	20	19	10	17	20	19	13	11	2	-
Male	7	12	16	13	6	9	7	6	4	2	-	-
Female	6	3	4	6	4	8	13	13	9	9	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	2	-	-	1	1	-	1	-
Male	-	-	1	1	1	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Hawaii												
All races, both sexes	1	5	2	9	12	17	33	17	8	4	2	-
Male	-	4	2	4	7	7	17	8	3	1	1	-
Female	1	1	-	5	5	10	16	9	5	3	1	-
White, both sexes	-	3	1	-	4	2	8	5	1	2	-	-
Male	-	2	1	-	3	1	6	4	1	1	-	-
Female	-	1	-	-	1	1	2	1	-	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	8	8	14	25	12	7	2	2	-
Male	-	2	1	3	4	6	11	4	2	-	1	-
Female	1	-	-	5	4	8	14	8	5	2	1	-
Idaho												
All races, both sexes	2	3	5	9	12	23	30	24	12	7	1	-
Male	2	1	4	6	7	15	14	14	7	3	1	-
Female	-	2	1	3	5	8	16	10	5	4	-	-
White, both sexes	2	3	5	9	12	23	28	22	12	7	1	-
Male	2	1	4	6	7	15	13	13	7	3	1	-
Female	-	2	1	3	5	8	15	9	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Illinois												
All races, both sexes	29	40	53	71	111	160	242	267	192	72	23	-
Male	13	28	39	39	57	85	106	102	57	16	2	-
Female	16	12	14	32	54	75	136	165	135	56	21	-
White, both sexes	19	31	45	56	95	141	203	244	180	67	20	-
Male	9	24	31	30	46	79	92	96	56	16	1	-
Female	10	7	14	26	49	62	111	148	124	51	19	-
Black, both sexes	10	9	8	14	16	18	35	21	10	4	3	-
Male	4	4	8	8	11	5	12	5	-	-	1	-
Female	6	5	-	6	5	13	23	16	10	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	3	2	2	1	-	-
Male	-	-	-	1	-	1	2	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
Indiana												
All races, both sexes	12	30	33	53	71	128	162	131	123	57	14	-
Male	10	22	21	41	42	78	74	59	39	15	2	-
Female	2	8	12	12	29	50	88	72	84	42	12	-
White, both sexes	12	28	31	51	64	122	150	125	117	52	12	-
Male	10	21	20	39	39	76	68	59	34	13	2	-
Female	2	7	11	12	25	46	82	66	83	39	10	-
Black, both sexes	-	2	2	2	6	6	11	6	6	5	2	-
Male	-	1	1	2	2	2	6	-	5	2	-	-
Female	-	1	1	-	4	4	5	6	1	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Iowa												
All races, both sexes	7	13	15	19	29	69	93	88	85	33	10	-
Male	6	11	11	9	20	40	45	36	31	8	-	-
Female	1	2	4	10	9	29	48	52	54	25	10	-
White, both sexes	7	13	13	19	27	67	92	88	85	31	10	-
Male	6	11	10	9	20	38	44	36	31	7	-	-
Female	1	2	3	10	7	29	48	52	54	24	10	-
Black, both sexes	-	-	1	-	2	2	-	-	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	2	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	15	15	13	37	83	98	113	105	40	17	-
Male	5	11	10	9	23	37	48	47	19	6	2	-
Female	1	4	5	4	14	46	50	66	86	34	15	-
White, both sexes	5	12	15	12	36	77	94	111	103	38	17	-
Male	5	8	10	9	22	34	47	47	19	5	2	-
Female	-	4	5	3	14	43	47	64	84	33	15	-
Black, both sexes	1	2	-	1	1	5	4	2	2	2	-	-
Male	-	2	-	-	1	2	1	-	-	1	-	-
Female	1	-	-	1	-	3	3	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Kentucky												
All races, both sexes	8	20	29	45	46	71	102	82	61	13	4	-
Male	4	13	21	27	30	38	46	35	16	3	-	-
Female	4	7	8	18	16	33	56	47	45	10	4	-
White, both sexes	4	19	27	37	43	64	101	80	57	13	3	-
Male	1	12	19	24	28	35	46	33	13	3	-	-
Female	3	7	8	13	15	29	55	47	44	10	3	-
Black, both sexes	4	1	1	7	3	7	1	2	4	-	1	-
Male	3	1	1	2	2	3	-	2	3	-	-	-
Female	1	-	-	5	1	4	1	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	12	19	26	33	60	49	85	72	52	19	4	-
Male	6	10	10	23	42	25	34	21	18	4	-	-
Female	6	9	16	10	18	24	51	51	34	15	4	-
White, both sexes	6	10	15	18	48	39	63	57	39	14	3	-
Male	4	6	7	12	36	19	24	15	15	4	-	-
Female	2	4	8	6	12	20	39	42	24	10	3	-
Black, both sexes	6	9	10	14	11	10	21	15	13	5	1	-
Male	2	4	2	10	6	6	10	6	3	-	-	-
Female	4	5	8	4	5	4	11	9	10	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Maine												
All races, both sexes	5	1	10	10	13	23	32	22	19	9	2	-
Male	3	1	7	3	8	12	17	11	7	4	-	-
Female	2	-	3	7	5	11	15	11	12	5	2	-
White, both sexes	4	1	10	10	13	23	32	22	19	9	2	-
Male	2	1	7	3	8	12	17	11	7	4	-	-
Female	2	-	3	7	5	11	15	11	12	5	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	10	21	21	27	43	80	83	87	64	22	6	-
Male	8	12	10	18	17	36	36	31	16	4	1	-
Female	2	9	11	9	26	44	47	56	48	18	5	-
White, both sexes	7	9	8	19	29	54	62	74	54	18	4	-
Male	6	5	5	12	13	28	27	30	13	4	1	-
Female	1	4	3	7	16	26	35	44	41	14	3	-
Black, both sexes	3	11	12	8	14	25	20	12	9	4	1	-
Male	2	6	4	6	4	8	8	1	2	-	-	-
Female	1	5	8	2	10	17	12	11	7	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	1	1	1	-	1	-
Male	-	1	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Massachusetts												
All races, both sexes	11	18	32	28	67	93	145	166	103	56	9	-
Male	9	15	20	20	35	48	73	68	31	12	1	-
Female	2	3	12	8	32	45	72	98	72	44	8	-
White, both sexes	11	16	31	27	64	92	140	161	99	55	9	-
Male	9	13	19	19	33	48	69	67	31	11	1	-
Female	2	3	12	8	31	44	71	94	68	44	8	-
Black, both sexes	-	2	1	-	3	-	4	4	3	1	-	-
Male	-	2	1	-	2	-	3	1	-	1	-	-
Female	-	-	-	-	1	-	1	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	1	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Michigan												
All races, both sexes	34	54	57	100	107	192	236	211	168	74	5	-
Male	27	36	34	66	59	91	103	90	50	14	1	-
Female	7	18	23	34	48	101	133	121	118	60	4	-
White, both sexes	19	39	46	83	91	168	200	193	147	66	5	-
Male	17	26	28	53	49	86	89	82	47	13	1	-
Female	2	13	18	30	42	82	111	111	100	53	4	-
Black, both sexes	15	15	10	16	15	22	33	18	17	8	-	-
Male	10	10	6	12	9	5	13	8	2	1	-	-
Female	5	5	4	4	6	17	20	10	15	7	-	-
American Indian 1/, both sexes	-	-	-	1	1	1	2	-	2	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Minnesota												
All races, both sexes	6	19	17	30	45	70	98	93	73	16	5	-
Male	5	16	12	18	23	32	53	41	26	5	1	-
Female	1	3	5	12	22	38	45	52	47	11	4	-
White, both sexes	6	18	16	29	44	68	97	91	73	16	5	-
Male	5	15	11	17	22	32	52	40	26	5	1	-
Female	1	3	5	12	22	36	45	51	47	11	4	-
Black, both sexes	-	-	1	-	1	1	1	1	-	-	-	-
Male	-	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	15	16	20	27	22	34	30	33	17	3	1	-
Male	11	10	15	20	14	16	13	16	4	1	-	-
Female	4	6	5	7	8	18	17	17	13	2	1	-
White, both sexes	8	10	11	19	17	30	22	26	13	2	1	-
Male	7	7	7	14	12	16	10	13	4	1	-	-
Female	1	3	4	5	5	14	12	13	9	1	1	-
Black, both sexes	7	6	8	8	5	4	8	7	4	1	-	-
Male	4	3	7	6	2	-	3	3	-	-	-	-
Female	3	3	1	2	3	4	5	4	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Missouri												
All races, both sexes	17	25	29	60	91	96	127	134	114	44	8	-
Male	10	20	18	33	50	53	57	53	29	11	1	-
Female	7	5	11	27	41	43	70	81	85	33	7	-
White, both sexes	14	19	23	51	80	86	111	120	99	41	8	-
Male	9	17	14	30	46	48	53	47	28	10	1	-
Female	5	2	9	21	34	38	58	73	71	31	7	-
Black, both sexes	3	6	6	9	11	9	15	13	14	3	-	-
Male	1	3	4	3	4	4	4	6	1	1	-	-
Female	2	3	2	6	7	5	11	7	13	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	5	3	5	2	9	13	15	23	13	6	-	-
Male	3	2	3	2	5	10	10	8	5	3	-	-
Female	2	1	2	-	4	3	5	15	8	3	-	-
White, both sexes	4	3	5	2	9	13	15	22	12	6	-	-
Male	2	2	3	2	5	10	10	7	5	3	-	-
Female	2	1	2	-	4	3	5	15	7	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Nebraska												
All races, both sexes	5	6	10	9	15	31	39	43	33	25	3	-
Male	4	2	9	6	7	16	17	16	8	6	-	-
Female	1	4	1	3	8	15	22	27	25	19	3	-
White, both sexes	3	6	9	9	14	31	38	42	31	25	3	-
Male	2	2	8	6	7	16	17	16	7	6	-	-
Female	1	4	1	3	7	15	21	26	24	19	3	-
Black, both sexes	2	-	1	-	1	-	1	1	2	-	-	-
Male	2	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	7	13	14	19	24	37	65	50	36	17	1	-
Male	4	6	8	12	16	21	29	21	11	1	-	-
Female	3	7	6	7	8	16	36	29	25	16	1	-
White, both sexes	7	12	13	16	23	29	63	49	36	16	1	-
Male	4	5	7	10	15	16	29	21	11	1	-	-
Female	3	7	6	6	8	13	34	28	25	15	1	-
Black, both sexes	-	-	-	3	1	6	2	-	-	-	-	-
Male	-	-	-	2	1	3	-	-	-	-	-	-
Female	-	-	-	1	-	3	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	2	-	1	-	-	-	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Hampshire												
All races, both sexes	3	5	8	7	18	20	17	21	17	10	-	-
Male	3	3	6	2	10	13	9	12	5	4	-	-
Female	-	2	2	5	8	7	8	9	12	6	-	-
White, both sexes	3	4	8	7	18	20	17	21	17	10	-	-
Male	3	2	6	2	10	13	9	12	5	4	-	-
Female	-	2	2	5	8	7	8	9	12	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	21	32	41	52	76	133	187	170	124	36	12	-
Male	14	28	27	37	40	66	101	74	45	13	4	-
Female	7	4	14	15	36	67	86	96	79	23	8	-
White, both sexes	16	22	33	38	62	113	167	158	117	34	11	-
Male	12	19	21	29	37	58	92	69	42	13	4	-
Female	4	3	12	9	25	55	75	89	75	21	7	-
Black, both sexes	4	10	7	12	13	16	16	11	3	2	1	-
Male	2	9	5	7	2	7	6	5	-	-	-	-
Female	2	1	2	5	11	9	10	6	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	1	3	4	1	4	-	-	-
Male	-	-	1	1	1	-	3	-	3	-	-	-
Female	1	-	-	1	-	3	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Mexico												
All races, both sexes	5	6	5	14	13	24	25	29	25	12	1	-
Male	3	4	3	13	10	17	12	12	7	1	-	-
Female	2	2	2	1	3	7	13	17	18	11	1	-
White, both sexes	4	4	5	10	13	23	25	28	25	10	1	-
Male	2	2	3	10	10	17	12	12	7	1	-	-
Female	2	2	2	-	3	6	13	16	18	9	1	-
Black, both sexes	1	1	-	2	-	-	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	2	-	1	-	1	-	2	-	-
Male	-	1	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	45	59	86	120	147	253	276	281	226	94	24	-
Male	36	44	56	75	82	133	123	114	78	23	3	-
Female	9	15	30	45	65	120	153	167	148	71	21	-
White, both sexes	29	45	70	101	128	217	242	257	208	87	24	-
Male	22	32	49	65	73	116	109	107	76	20	3	-
Female	7	13	21	36	55	101	133	150	132	67	21	-
Black, both sexes	16	13	14	17	17	32	29	20	15	7	-	-
Male	14	11	7	9	7	14	10	5	2	3	-	-
Female	2	2	7	8	10	18	19	15	13	4	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	2	3	5	4	3	-	-	-
Male	-	1	-	1	2	3	4	2	-	-	-	-
Female	-	-	2	-	-	-	1	2	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
North Carolina												
All races, both sexes	28	36	62	56	99	124	184	165	114	33	9	-
Male	20	27	37	27	57	72	84	76	39	10	1	-
Female	8	9	25	29	42	52	100	89	75	23	8	-
White, both sexes	20	24	48	41	76	100	153	135	97	26	5	-
Male	13	19	31	20	48	59	73	64	34	6	1	-
Female	7	5	17	21	28	41	80	71	63	20	4	-
Black, both sexes	8	12	14	15	22	23	29	30	16	7	4	-
Male	7	8	6	7	9	12	11	12	4	4	-	-
Female	1	4	8	8	13	11	18	18	12	3	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	4	7	2	17	6	18	12	4	-	-
Male	3	-	3	6	1	10	2	5	7	2	-	-
Female	2	-	1	1	1	7	4	13	5	2	-	-
White, both sexes	5	-	4	6	2	17	6	18	12	4	-	-
Male	3	-	3	5	1	10	2	5	7	2	-	-
Female	2	-	1	1	1	7	4	13	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Ohio												
All races, both sexes	31	48	64	99	125	238	268	264	178	67	14	-
Male	21	36	42	60	61	127	124	109	56	16	3	-
Female	10	12	22	39	64	111	144	155	122	51	11	-
White, both sexes	22	40	56	88	113	208	242	250	160	65	13	-
Male	16	29	39	57	58	114	115	104	50	16	2	-
Female	6	11	17	31	55	94	127	146	110	49	11	-
Black, both sexes	9	8	8	10	11	30	25	14	17	2	1	-
Male	5	7	3	3	3	13	9	5	5	-	1	-
Female	4	1	5	7	8	17	16	9	12	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	11	13	22	26	57	53	55	59	58	19	2	-
Male	7	9	12	15	25	32	28	27	13	7	-	-
Female	4	4	10	11	32	21	27	32	45	12	2	-
White, both sexes	10	12	19	22	43	46	45	55	52	17	2	-
Male	6	8	11	14	21	29	24	26	13	7	-	-
Female	4	4	8	8	22	17	21	29	39	10	2	-
Black, both sexes	-	-	3	1	6	5	3	2	3	1	-	-
Male	-	-	1	1	3	2	2	-	-	-	-	-
Female	-	-	2	-	3	3	1	2	3	1	-	-
American Indian 1/, both sexes	1	1	-	3	6	2	7	2	3	1	-	-
Male	1	1	-	-	1	1	2	1	-	-	-	-
Female	-	-	-	3	5	1	5	1	3	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Oregon												
All races, both sexes	11	18	17	26	45	61	88	95	43	14	4	-
Male	7	14	12	16	30	36	35	41	17	3	2	-
Female	4	4	5	10	15	25	53	54	26	11	2	-
White, both sexes	11	16	17	25	43	59	88	94	42	14	4	-
Male	7	13	12	16	28	36	35	41	16	3	2	-
Female	4	3	5	9	15	23	53	53	26	11	2	-
Black, both sexes	-	2	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	28	45	59	71	148	285	339	310	239	97	14	-
Male	17	30	34	51	75	153	173	126	56	27	1	-
Female	11	15	25	20	73	132	166	184	183	70	13	-
White, both sexes	15	40	48	63	137	264	317	299	224	92	14	-
Male	11	26	28	48	73	145	163	123	54	24	1	-
Female	4	14	20	15	64	119	154	176	170	68	13	-
Black, both sexes	12	5	11	7	11	21	22	11	15	5	-	-
Male	6	4	6	2	2	8	10	3	2	3	-	-
Female	6	1	5	5	9	13	12	8	13	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Rhode Island												
All races, both sexes	1	6	5	6	13	18	31	29	20	9	4	-
Male	1	4	3	4	9	12	11	10	6	1	-	-
Female	-	2	2	2	4	6	20	19	14	8	4	-
White, both sexes	1	5	5	6	13	18	30	29	20	8	4	-
Male	1	4	3	4	9	12	11	10	6	1	-	-
Female	-	1	2	2	4	6	19	19	14	7	4	-
Black, both sexes	-	1	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	10	18	24	31	56	69	64	60	33	16	-	-
Male	8	8	14	21	31	37	27	27	11	4	-	-
Female	2	10	10	10	25	32	37	33	22	12	-	-
White, both sexes	6	8	17	24	44	61	50	51	28	9	-	-
Male	5	4	11	17	24	36	22	23	11	2	-	-
Female	1	4	6	7	20	25	28	28	17	7	-	-
Black, both sexes	4	9	7	7	11	8	14	9	5	7	-	-
Male	3	4	3	4	6	1	5	4	-	2	-	-
Female	1	5	4	3	5	7	9	5	5	5	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
South Dakota												
All races, both sexes	-	3	3	5	10	9	18	22	14	10	3	-
Male	-	-	2	4	5	6	8	7	4	2	2	-
Female	-	3	1	1	5	3	10	15	10	8	1	-
White, both sexes	-	2	3	5	10	8	18	22	14	10	3	-
Male	-	-	2	4	5	5	8	7	4	2	2	-
Female	-	2	1	1	5	3	10	15	10	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	17	24	36	58	73	87	113	109	75	25	9	-
Male	13	13	26	27	47	40	43	45	18	3	3	-
Female	4	11	10	31	26	47	70	64	57	22	6	-
White, both sexes	10	20	33	47	65	76	99	93	71	24	8	-
Male	6	11	23	22	43	36	39	40	18	3	3	-
Female	4	9	10	25	22	40	60	53	53	21	5	-
Black, both sexes	7	4	3	10	8	11	14	16	4	1	1	-
Male	7	2	3	5	4	4	4	5	-	-	-	-
Female	-	2	-	5	4	7	10	11	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Texas												
All races, both sexes	57	69	78	124	161	223	290	280	167	80	18	-
Male	45	43	50	78	101	121	124	126	54	17	4	-
Female	12	26	28	46	60	102	166	154	113	63	14	-
White, both sexes	38	48	69	107	143	203	257	255	148	69	13	-
Male	30	27	44	64	88	113	118	113	48	14	2	-
Female	8	21	25	43	55	90	139	142	100	55	11	-
Black, both sexes	17	21	8	16	14	18	32	20	18	11	5	-
Male	13	16	5	13	9	7	6	8	5	3	2	-
Female	4	5	3	3	5	11	26	12	13	8	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	1	4	2	1	5	1	-	-	-
Male	2	-	1	1	4	1	-	5	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Utah												
All races, both sexes	2	8	11	15	19	24	45	30	18	4	-	-
Male	1	6	8	7	16	16	25	16	6	1	-	-
Female	1	2	3	8	3	8	20	14	12	3	-	-
White, both sexes	2	7	11	15	19	22	44	30	18	4	-	-
Male	1	5	8	7	16	15	24	16	6	1	-	-
Female	1	2	3	8	3	7	20	14	12	3	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Vermont												
All races, both sexes	-	2	-	5	5	10	16	13	11	3	2	-
Male	-	-	-	1	3	6	6	7	3	1	1	-
Female	-	2	-	4	2	4	10	6	8	2	1	-
White, both sexes	-	2	-	5	5	10	16	13	11	3	2	-
Male	-	-	-	1	3	6	6	7	3	1	1	-
Female	-	2	-	4	2	4	10	6	8	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	14	27	32	56	65	98	113	114	95	26	7	-
Male	10	16	26	37	41	60	43	43	29	2	1	-
Female	4	11	6	19	24	38	70	71	66	24	6	-
White, both sexes	9	21	24	40	51	83	92	93	78	22	3	-
Male	6	13	20	24	32	50	37	36	21	1	-	-
Female	3	8	4	16	19	33	55	57	57	21	3	-
Black, both sexes	5	6	8	14	13	14	20	21	16	3	4	-
Male	4	3	6	11	9	9	5	7	7	1	1	-
Female	1	3	2	3	4	5	15	14	9	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	1	1	-	1	1	-	-
Male	-	-	-	2	-	1	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Washington												
All races, both sexes	12	29	26	37	55	97	131	128	98	41	11	-
Male	10	23	18	22	36	55	66	59	24	20	-	-
Female	2	6	8	15	19	42	65	69	74	21	11	-
White, both sexes	11	28	22	34	54	91	122	122	96	41	11	-
Male	9	23	16	21	35	52	61	55	24	20	-	-
Female	2	5	6	13	19	39	61	67	72	21	11	-
Black, both sexes	-	-	-	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	2	-	5	6	4	2	-	-	-
Male	1	-	2	-	-	2	3	3	-	-	-	-
Female	-	-	1	2	-	3	3	1	2	-	-	-
West Virginia												
All races, both sexes	3	10	12	26	29	37	58	49	36	12	3	-
Male	2	6	7	17	19	21	29	23	13	3	-	-
Female	1	4	5	9	10	16	29	26	23	9	3	-
White, both sexes	3	9	12	25	28	37	58	48	36	12	3	-
Male	2	5	7	16	18	21	29	22	13	3	-	-
Female	1	4	5	9	10	16	29	26	23	9	3	-
Black, both sexes	-	1	-	1	1	-	-	1	-	-	-	-
Male	-	1	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Wisconsin												
All races, both sexes	13	21	24	38	65	88	104	107	96	33	13	-
Male	7	19	13	22	33	42	54	39	29	9	2	-
Female	6	2	11	16	32	46	50	68	67	24	11	-
White, both sexes	12	17	21	36	62	84	103	106	96	33	13	-
Male	7	15	13	21	32	40	54	39	29	9	2	-
Female	5	2	8	15	30	44	49	67	67	24	11	-
Black, both sexes	1	2	3	1	3	3	1	-	-	-	-	-
Male	-	2	-	1	1	2	-	-	-	-	-	-
Female	1	-	3	-	2	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	1	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	3	3	4	3	12	8	12	2	2	-	-
Male	-	2	2	3	2	8	5	4	-	1	-	-
Female	2	1	1	1	1	4	3	8	2	1	-	-
White, both sexes	1	2	3	4	3	12	8	12	2	2	-	-
Male	-	2	2	3	2	8	5	4	-	1	-	-
Female	1	-	1	1	1	4	3	8	2	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
United States												
All races, both sexes	8,652	2	1	-	-	-	1	5	2	8	23	51
Male	3,359	1	-	-	-	-	1	5	-	6	15	36
Female	5,293	1	1	-	-	-	-	-	2	2	8	15
White, both sexes	7,820	2	-	-	-	-	1	3	2	6	15	35
Male	2,992	1	-	-	-	-	1	3	-	5	8	26
Female	4,828	1	-	-	-	-	-	-	2	1	7	9
Black, both sexes	709	-	1	-	-	-	-	2	-	2	7	12
Male	312	-	-	-	-	-	-	2	-	1	6	8
Female	397	-	1	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	96	-	-	-	-	-	-	-	-	-	1	1
Male	49	-	-	-	-	-	-	-	-	-	1	1
Female	47	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	174	-	-	-	-	-	-	-	1	2	5	4
Male	81	-	-	-	-	-	-	-	-	2	4	4
Female	93	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	123	-	-	-	-	-	-	-	1	1	2	1
Male	48	-	-	-	-	-	-	-	-	1	1	1
Female	75	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	51	-	-	-	-	-	-	-	-	1	3	3
Male	33	-	-	-	-	-	-	-	-	1	3	3
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Alaska												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	187	-	-	-	-	-	-	-	-	-	1	1
Male	94	-	-	-	-	-	-	-	-	-	1	-
Female	93	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	183	-	-	-	-	-	-	-	-	-	1	-
Male	92	-	-	-	-	-	-	-	-	-	1	-
Female	91	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Arkansas												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	964	-	-	-	-	-	-	-	1	-	2	6
Male	410	-	-	-	-	-	-	-	-	-	1	5
Female	554	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	866	-	-	-	-	-	-	-	1	-	2	3
Male	361	-	-	-	-	-	-	-	-	-	1	3
Female	505	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	48	-	-	-	-	-	-	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Colorado												
All races, both sexes	341	-	-	-	-	-	-	1	-	-	-	3
Male	103	-	-	-	-	-	-	1	-	-	-	2
Female	238	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	323	-	-	-	-	-	-	1	-	-	-	2
Male	97	-	-	-	-	-	-	1	-	-	-	2
Female	226	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	105	-	-	-	-	-	-	-	-	2	-	-
Male	34	-	-	-	-	-	-	-	-	1	-	-
Female	71	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	100	-	-	-	-	-	-	-	-	2	-	-
Male	31	-	-	-	-	-	-	-	-	1	-	-
Female	69	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Delaware												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Florida												
All races, both sexes	651	-	-	-	-	-	-	-	-	1	3	3
Male	279	-	-	-	-	-	-	-	-	-	2	2
Female	372	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	602	-	-	-	-	-	-	-	-	-	1	1
Male	256	-	-	-	-	-	-	-	-	-	1	1
Female	346	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	49	-	-	-	-	-	-	-	-	1	2	2
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	26	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	176	-	-	-	-	-	-	1	-	1	1	3
Male	65	-	-	-	-	-	-	1	-	1	1	1
Female	111	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	138	-	-	-	-	-	-	-	-	1	-	3
Male	48	-	-	-	-	-	-	-	-	1	-	1
Female	90	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	35	-	-	-	-	-	-	1	-	-	1	-
Male	16	-	-	-	-	-	-	1	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Hawaii												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Illinois												
All races, both sexes	371	-	-	-	-	-	-	-	-	-	-	-
Male	124	-	-	-	-	-	-	-	-	-	-	-
Female	247	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	335	-	-	-	-	-	-	-	-	-	-	-
Male	109	-	-	-	-	-	-	-	-	-	-	-
Female	226	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	316	-	-	-	-	-	-	1	-	-	1	4
Male	118	-	-	-	-	-	-	1	-	-	1	4
Female	198	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	291	-	-	-	-	-	-	-	-	-	1	4
Male	107	-	-	-	-	-	-	-	-	-	1	4
Female	184	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	24	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Iowa												
All races, both sexes	185	-	-	-	-	-	-	-	-	-	2	1
Male	67	-	-	-	-	-	-	-	-	-	-	-
Female	118	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	181	-	-	-	-	-	-	-	-	-	2	1
Male	64	-	-	-	-	-	-	-	-	-	-	-
Female	117	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	321	-	-	-	-	-	-	-	-	-	-	2
Male	107	-	-	-	-	-	-	-	-	-	-	1
Female	214	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	310	-	-	-	-	-	-	-	-	-	-	2
Male	105	-	-	-	-	-	-	-	-	-	-	1
Female	205	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Kentucky												
All races, both sexes	122	-	-	-	-	-	-	1	-	-	-	1
Male	48	-	-	-	-	-	-	1	-	-	-	-
Female	74	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	115	-	-	-	-	-	-	1	-	-	-	-
Male	45	-	-	-	-	-	-	1	-	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	127	-	-	-	-	-	-	-	-	-	1	-
Male	48	-	-	-	-	-	-	-	-	-	1	-
Female	79	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	100	-	-	-	-	-	-	-	-	-	1	-
Male	37	-	-	-	-	-	-	-	-	-	1	-
Female	63	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Maine												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	98	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	66	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Massachusetts												
All races, both sexes	216	-	-	-	-	-	-	-	-	-	-	1
Male	78	-	-	-	-	-	-	-	-	-	-	1
Female	138	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	211	-	-	-	-	-	-	-	-	-	-	1
Male	75	-	-	-	-	-	-	-	-	-	-	1
Female	136	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	382	-	-	-	-	-	-	-	-	-	1	-
Male	136	-	-	-	-	-	-	-	-	-	-	-
Female	246	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	312	-	-	-	-	-	-	-	-	-	1	-
Male	114	-	-	-	-	-	-	-	-	-	-	-
Female	198	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Minnesota												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	71	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Missouri												
All races, both sexes	259	-	-	-	-	-	-	-	-	-	-	-
Male	94	-	-	-	-	-	-	-	-	-	-	-
Female	165	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	225	-	-	-	-	-	-	-	-	-	-	-
Male	83	-	-	-	-	-	-	-	-	-	-	-
Female	142	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Nebraska												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	141	-	-	-	-	-	-	-	-	-	-	1
Male	55	-	-	-	-	-	-	-	-	-	-	-
Female	86	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	137	-	-	-	-	-	-	-	-	-	-	1
Male	54	-	-	-	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
New Hampshire												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	207	-	-	-	-	-	-	-	-	-	-	1
Male	89	-	-	-	-	-	-	-	-	-	-	-
Female	118	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	198	-	-	-	-	-	-	-	-	-	-	1
Male	86	-	-	-	-	-	-	-	-	-	-	-
Female	112	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
New Mexico												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	434	-	1	-	-	-	-	-	-	-	-	1
Male	171	-	-	-	-	-	-	-	-	-	-	1
Female	263	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	381	-	-	-	-	-	-	-	-	-	-	-
Male	148	-	-	-	-	-	-	-	-	-	-	-
Female	233	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	46	-	1	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
North Carolina												
All races, both sexes	222	-	-	-	-	-	-	-	-	-	-	3
Male	94	-	-	-	-	-	-	-	-	-	-	1
Female	128	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	178	-	-	-	-	-	-	-	-	-	-	2
Male	71	-	-	-	-	-	-	-	-	-	-	-
Female	107	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Ohio												
All races, both sexes	339	-	-	-	-	-	-	-	-	-	-	2
Male	125	-	-	-	-	-	-	-	-	-	-	2
Female	214	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	313	-	-	-	-	-	-	-	-	-	-	1
Male	109	-	-	-	-	-	-	-	-	-	-	1
Female	204	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	124	-	-	-	-	-	-	-	-	-	-	1
Male	54	-	-	-	-	-	-	-	-	-	-	1
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	107	-	-	-	-	-	-	-	-	-	-	1
Male	51	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Oregon												
All races, both sexes	116	-	-	-	-	-	-	-	-	-	-	1
Male	52	-	-	-	-	-	-	-	-	-	-	1
Female	64	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	115	-	-	-	-	-	-	-	-	-	-	1
Male	51	-	-	-	-	-	-	-	-	-	-	1
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	400	1	-	-	-	-	-	-	-	-	1	1
Male	144	-	-	-	-	-	-	-	-	-	1	1
Female	256	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	386	1	-	-	-	-	-	-	-	-	1	1
Male	138	-	-	-	-	-	-	-	-	-	1	1
Female	248	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Rhode Island												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
South Dakota												
All races, both sexes	18	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	166	-	-	-	-	-	1	-	-	-	-	1
Male	60	-	-	-	-	-	1	-	-	-	-	1
Female	106	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	143	-	-	-	-	-	1	-	-	-	-	1
Male	48	-	-	-	-	-	1	-	-	-	-	1
Female	95	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Texas												
All races, both sexes	372	-	-	-	-	-	-	-	-	-	1	1
Male	144	-	-	-	-	-	-	-	-	-	1	-
Female	228	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	324	-	-	-	-	-	-	-	-	-	-	-
Male	119	-	-	-	-	-	-	-	-	-	-	-
Female	205	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	42	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	1	-
Male	30	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Vermont												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	185	-	-	-	-	-	-	1	-	-	-	2
Male	64	-	-	-	-	-	-	1	-	-	-	2
Female	121	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	155	-	-	-	-	-	-	1	-	-	-	2
Male	50	-	-	-	-	-	-	1	-	-	-	2
Female	105	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Washington												
All races, both sexes	189	-	-	-	-	-	-	-	-	1	-	1
Male	71	-	-	-	-	-	-	-	-	1	-	-
Female	118	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	182	-	-	-	-	-	-	-	-	1	-	1
Male	68	-	-	-	-	-	-	-	-	1	-	-
Female	114	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	74	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Wisconsin												
All races, both sexes	115	1	-	-	-	-	-	-	-	-	1	1
Male	42	1	-	-	-	-	-	-	-	-	1	1
Female	73	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	113	1	-	-	-	-	-	-	-	-	1	1
Male	41	1	-	-	-	-	-	-	-	-	1	1
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
United States												
All races, both sexes	89	155	233	328	521	929	1,540	1,852	1,774	881	257	-
Male	63	104	140	183	280	442	643	708	498	199	35	-
Female	26	51	93	145	241	487	897	1,144	1,276	682	222	-
White, both sexes	69	124	191	276	444	830	1,387	1,712	1,663	822	238	-
Male	52	80	119	152	237	397	574	655	468	182	32	-
Female	17	44	72	124	207	433	813	1,057	1,195	640	206	-
Black, both sexes	17	28	40	45	70	85	124	122	89	51	14	-
Male	9	22	21	28	37	37	57	43	25	14	2	-
Female	8	6	19	17	33	48	67	79	64	37	12	-
American Indian 1/, both sexes	1	1	-	3	1	1	6	3	5	3	-	-
Male	1	1	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	2	-	1	6	2	5	3	-	-
Asian or Pacific Islander 2/, both sexes	2	2	2	4	6	13	23	15	17	5	5	-
Male	1	1	-	2	5	8	12	9	5	3	1	-
Female	1	1	2	2	1	5	11	6	12	2	4	-
Alabama												
All races, both sexes	5	13	11	17	17	15	24	29	16	14	1	-
Male	4	10	7	11	11	4	11	8	2	3	-	-
Female	1	3	4	6	6	11	13	21	14	11	1	-
White, both sexes	3	7	8	11	11	13	21	20	11	12	1	-
Male	3	4	6	7	6	3	9	5	-	2	-	-
Female	-	3	2	4	5	10	12	15	11	10	1	-
Black, both sexes	2	6	3	6	6	2	3	9	5	2	-	-
Male	1	6	1	4	5	1	2	3	2	1	-	-
Female	1	-	2	2	1	1	1	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Alaska												
All races, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	2	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	2	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	1	6	6	20	13	37	54	25	17	2	-
Male	4	-	3	3	13	8	18	26	13	5	-	-
Female	-	1	3	3	7	5	19	28	12	12	2	-
White, both sexes	4	1	6	6	18	13	37	54	24	17	2	-
Male	4	-	3	3	11	8	18	26	13	5	-	-
Female	-	1	3	3	7	5	19	28	11	12	2	-
Black, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Arkansas												
All races, both sexes	2	-	3	2	5	4	3	6	5	2	-	-
Male	-	-	2	2	1	-	2	2	1	1	-	-
Female	2	-	1	-	4	4	1	4	4	1	-	-
White, both sexes	1	-	3	2	4	4	3	4	4	2	-	-
Male	-	-	2	2	1	-	2	2	1	1	-	-
Female	1	-	1	-	3	4	1	2	3	1	-	-
Black, both sexes	1	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13	10	20	33	52	115	163	226	188	99	36	-
Male	9	7	12	20	25	58	81	101	64	18	9	-
Female	4	3	8	13	27	57	82	125	124	81	27	-
White, both sexes	11	9	13	28	45	99	146	211	175	91	32	-
Male	7	7	8	16	21	48	71	94	60	16	9	-
Female	4	2	5	12	24	51	75	117	115	75	23	-
Black, both sexes	-	-	6	4	4	7	5	7	5	5	3	-
Male	-	-	4	3	2	4	3	4	3	-	-	-
Female	-	-	2	1	2	3	2	3	2	5	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	3	9	12	8	8	3	1	-
Male	1	-	-	1	2	6	7	3	1	2	-	-
Female	-	1	1	-	1	3	5	5	7	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Colorado												
All races, both sexes	-	1	4	11	16	34	70	78	66	41	16	-
Male	-	-	3	7	6	20	20	24	12	5	3	-
Female	-	1	1	4	10	14	50	54	54	36	13	-
White, both sexes	-	1	4	10	15	33	67	73	65	38	14	-
Male	-	-	3	7	6	19	19	22	12	4	2	-
Female	-	1	1	3	9	14	48	51	53	34	12	-
Black, both sexes	-	-	-	-	1	1	2	4	1	3	-	-
Male	-	-	-	-	-	1	1	1	-	1	-	-
Female	-	-	-	-	1	-	1	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	2	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	-	2	3	-	-	10	15	15	31	24	3	-
Male	-	2	2	-	-	3	12	4	6	3	1	-
Female	-	-	1	-	-	7	3	11	25	21	2	-
White, both sexes	-	1	2	-	-	9	15	15	29	24	3	-
Male	-	1	1	-	-	3	12	4	5	3	1	-
Female	-	-	1	-	-	6	3	11	24	21	2	-
Black, both sexes	-	1	1	-	-	1	-	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Delaware												
All races, both sexes	-	-	-	1	-	-	-	2	-	2	-	-
Male	-	-	-	-	-	-	-	2	-	1	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	-	2	-	2	-	-
Male	-	-	-	-	-	-	-	2	-	1	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	2	1	5	2	3	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	4	2	3	-	-	-
White, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	2	-	3	-	2	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Florida												
All races, both sexes	4	10	22	26	41	85	101	131	138	67	19	-
Male	4	7	13	14	22	39	48	51	53	19	5	-
Female	-	3	9	12	19	46	53	80	85	48	14	-
White, both sexes	4	7	18	23	39	79	92	123	135	62	18	-
Male	4	5	11	13	20	36	42	49	51	18	5	-
Female	-	2	7	10	19	43	50	74	84	44	13	-
Black, both sexes	-	3	4	3	2	6	9	8	3	5	1	-
Male	-	2	2	1	2	3	6	2	2	1	-	-
Female	-	1	2	2	-	3	3	6	1	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	6	9	12	11	20	30	28	29	19	5	-
Male	-	4	7	6	3	7	13	10	7	4	-	-
Female	1	2	2	6	8	13	17	18	22	15	5	-
White, both sexes	-	5	8	8	9	16	26	19	25	15	3	-
Male	-	3	6	3	2	4	12	7	6	3	-	-
Female	-	2	2	5	7	12	14	12	19	12	3	-
Black, both sexes	-	1	1	4	2	4	4	8	4	4	1	-
Male	-	1	1	3	1	3	1	2	1	1	-	-
Female	-	-	-	1	1	1	3	6	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Hawaii												
All races, both sexes	-	2	-	2	1	-	8	2	1	-	-	-
Male	-	1	-	-	1	-	4	1	1	-	-	-
Female	-	1	-	2	-	-	4	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	1	-	5	2	-	-	-	-
Male	-	1	-	-	1	-	3	1	-	-	-	-
Female	-	-	-	2	-	-	2	1	-	-	-	-
Idaho												
All races, both sexes	-	1	-	2	2	2	5	2	3	1	1	-
Male	-	-	-	2	2	2	4	1	1	1	1	-
Female	-	1	-	-	-	-	1	1	2	-	-	-
White, both sexes	-	1	-	2	2	2	5	2	3	1	1	-
Male	-	-	-	2	2	2	4	1	1	1	1	-
Female	-	1	-	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Illinois												
All races, both sexes	4	4	7	10	25	32	66	83	76	47	17	-
Male	2	-	4	6	18	15	19	30	16	12	2	-
Female	2	4	3	4	7	17	47	53	60	35	15	-
White, both sexes	3	2	6	7	19	30	55	79	74	45	15	-
Male	1	-	3	4	14	15	15	28	16	12	1	-
Female	2	2	3	3	5	15	40	51	58	33	14	-
Black, both sexes	1	2	1	3	6	2	11	3	2	1	2	-
Male	1	-	1	2	4	-	4	1	-	-	1	-
Female	-	2	-	1	2	2	7	2	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Indiana												
All races, both sexes	2	9	4	5	16	31	53	63	77	39	11	-
Male	2	7	3	4	8	14	21	20	22	10	1	-
Female	-	2	1	1	8	17	32	43	55	29	10	-
White, both sexes	2	8	4	4	14	29	46	59	73	37	10	-
Male	2	6	3	3	7	14	18	20	18	10	1	-
Female	-	2	1	1	7	15	28	39	55	27	9	-
Black, both sexes	-	1	-	1	2	2	6	4	4	2	1	-
Male	-	1	-	1	1	-	3	-	4	-	-	-
Female	-	-	-	-	1	2	3	4	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Iowa												
All races, both sexes	3	5	2	3	7	21	33	34	41	27	6	-
Male	3	5	2	-	4	11	12	12	13	5	-	-
Female	-	-	-	3	3	10	21	22	28	22	6	-
White, both sexes	3	5	2	3	6	19	33	34	41	26	6	-
Male	3	5	2	-	4	9	12	12	13	4	-	-
Female	-	-	-	3	2	10	21	22	28	22	6	-
Black, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	5	4	6	12	42	59	70	77	30	13	-
Male	-	4	3	4	8	16	27	27	13	3	1	-
Female	1	1	1	2	4	26	32	43	64	27	12	-
White, both sexes	-	5	4	6	12	39	56	68	76	29	13	-
Male	-	4	3	4	8	15	26	27	13	3	1	-
Female	-	1	1	2	4	24	30	41	63	26	12	-
Black, both sexes	1	-	-	-	-	3	3	2	1	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	2	2	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Kentucky												
All races, both sexes	1	2	4	7	3	14	26	29	29	4	1	-
Male	1	2	1	5	2	9	7	12	6	2	-	-
Female	-	-	3	2	1	5	19	17	23	2	1	-
White, both sexes	1	2	4	6	3	13	25	28	27	4	1	-
Male	1	2	1	4	2	9	7	11	5	2	-	-
Female	-	-	3	2	1	4	18	17	22	2	1	-
Black, both sexes	-	-	-	1	-	1	1	1	2	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	3	2	6	13	10	28	30	18	11	4	-
Male	-	2	2	3	7	5	8	10	6	4	-	-
Female	1	1	-	3	6	5	20	20	12	7	4	-
White, both sexes	-	2	2	3	9	8	23	26	14	9	3	-
Male	-	1	2	1	6	3	6	8	5	4	-	-
Female	-	1	-	2	3	5	17	18	9	5	3	-
Black, both sexes	1	1	-	3	4	2	4	4	4	2	1	-
Male	-	1	-	2	1	2	2	2	1	-	-	-
Female	1	-	-	1	3	-	2	2	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Maine												
All races, both sexes	1	-	1	-	-	1	6	5	4	3	-	-
Male	1	-	1	-	-	1	3	3	1	2	-	-
Female	-	-	-	-	-	-	3	2	3	1	-	-
White, both sexes	1	-	1	-	-	1	6	5	4	3	-	-
Male	1	-	1	-	-	1	3	3	1	2	-	-
Female	-	-	-	-	-	-	3	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	2	1	6	8	12	17	18	22	7	2	-
Male	2	1	-	4	2	5	7	3	6	2	-	-
Female	1	1	1	2	6	7	10	15	16	5	2	-
White, both sexes	2	1	-	5	6	7	10	15	19	7	1	-
Male	1	-	-	3	2	4	4	3	5	2	-	-
Female	1	1	-	2	4	3	6	12	14	5	1	-
Black, both sexes	1	1	1	1	2	5	7	3	3	-	-	-
Male	1	1	-	1	-	1	3	-	1	-	-	-
Female	-	-	1	-	2	4	4	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Massachusetts												
All races, both sexes	4	5	6	4	15	21	32	45	49	27	7	-
Male	4	5	2	2	9	10	8	19	13	5	-	-
Female	-	-	4	2	6	11	24	26	36	22	7	-
White, both sexes	4	4	6	4	15	21	31	44	48	26	7	-
Male	4	4	2	2	9	10	7	19	13	4	-	-
Female	-	-	4	2	6	11	24	25	35	22	7	-
Black, both sexes	-	1	-	-	-	-	1	1	1	1	-	-
Male	-	1	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5	10	11	16	19	47	67	85	73	45	3	-
Male	3	7	4	11	5	20	26	36	17	6	1	-
Female	2	3	7	5	14	27	41	49	56	39	2	-
White, both sexes	3	7	6	12	16	39	52	76	59	38	3	-
Male	2	4	3	9	4	19	20	31	16	5	1	-
Female	1	3	3	3	12	20	32	45	43	33	2	-
Black, both sexes	2	3	4	3	3	8	15	9	11	7	-	-
Male	1	3	1	1	1	1	6	5	1	1	-	-
Female	1	-	3	2	2	7	9	4	10	6	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Minnesota												
All races, both sexes	-	4	6	4	4	10	13	11	14	6	1	-
Male	-	3	5	2	2	5	8	4	5	2	-	-
Female	-	1	1	2	2	5	5	7	9	4	1	-
White, both sexes	-	4	5	4	4	10	12	11	14	6	1	-
Male	-	3	4	2	2	5	7	4	5	2	-	-
Female	-	1	1	2	2	5	5	7	9	4	1	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	-	2	1	4	6	5	-	1	-
Male	-	-	-	-	2	-	3	3	1	-	-	-
Female	-	-	1	-	-	1	1	3	4	-	1	-
White, both sexes	-	-	1	-	2	1	3	4	4	-	1	-
Male	-	-	-	-	2	-	2	3	1	-	-	-
Female	-	-	1	-	-	1	1	1	3	-	1	-
Black, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Missouri												
All races, both sexes	5	6	7	12	16	25	39	57	62	22	8	-
Male	1	4	5	4	9	12	18	23	11	6	1	-
Female	4	2	2	8	7	13	21	34	51	16	7	-
White, both sexes	4	5	6	9	15	22	35	48	54	19	8	-
Male	1	4	4	3	8	10	17	19	11	5	1	-
Female	3	1	2	6	7	12	18	29	43	14	7	-
Black, both sexes	1	1	1	3	1	3	3	8	7	3	-	-
Male	-	-	1	1	1	2	1	4	-	1	-	-
Female	1	1	-	2	-	1	2	4	7	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	2	-	2	-	1	2	8	6	5	5	-	-
Male	1	-	1	-	1	2	4	1	1	2	-	-
Female	1	-	1	-	-	-	4	5	4	3	-	-
White, both sexes	2	-	2	-	1	2	8	6	5	5	-	-
Male	1	-	1	-	1	2	4	1	1	2	-	-
Female	1	-	1	-	-	-	4	5	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Nebraska												
All races, both sexes	-	2	1	1	2	5	12	18	16	14	3	-
Male	-	1	1	-	-	4	6	4	4	3	-	-
Female	-	1	-	1	2	1	6	14	12	11	3	-
White, both sexes	-	2	1	1	1	5	11	17	15	14	3	-
Male	-	1	1	-	-	4	6	4	4	3	-	-
Female	-	1	-	1	1	1	5	13	11	11	3	-
Black, both sexes	-	-	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	4	4	5	8	12	33	35	28	11	-	-
Male	-	1	2	4	5	5	13	16	8	1	-	-
Female	-	3	2	1	3	7	20	19	20	10	-	-
White, both sexes	-	4	4	5	7	11	32	35	28	10	-	-
Male	-	1	2	4	4	5	13	16	8	1	-	-
Female	-	3	2	1	3	6	19	19	20	9	-	-
Black, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
New Hampshire												
All races, both sexes	1	1	1	5	3	4	2	5	5	4	-	-
Male	1	1	1	2	1	3	1	4	2	2	-	-
Female	-	-	-	3	2	1	1	1	3	2	-	-
White, both sexes	1	1	1	5	3	4	2	5	5	4	-	-
Male	1	1	1	2	1	3	1	4	2	2	-	-
Female	-	-	-	3	2	1	1	1	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	2	3	7	9	25	42	45	52	15	3	-
Male	2	1	3	7	3	16	19	15	18	4	1	-
Female	1	1	-	-	6	9	23	30	34	11	2	-
White, both sexes	2	2	3	7	8	24	38	45	50	15	3	-
Male	2	1	3	7	3	16	17	15	17	4	1	-
Female	-	1	-	-	5	8	21	30	33	11	2	-
Black, both sexes	1	-	-	-	1	1	3	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	1	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
New Mexico												
All races, both sexes	-	-	1	1	2	3	3	9	11	5	1	-
Male	-	-	1	1	1	2	1	4	2	-	-	-
Female	-	-	-	-	1	1	2	5	9	5	1	-
White, both sexes	-	-	1	1	2	3	3	9	11	4	1	-
Male	-	-	1	1	1	2	1	4	2	-	-	-
Female	-	-	-	-	1	1	2	5	9	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	4	16	17	30	54	78	82	94	35	19	-
Male	3	3	8	10	16	22	39	31	30	7	1	-
Female	-	1	8	7	14	32	39	51	64	28	18	-
White, both sexes	2	3	13	13	26	43	71	73	86	32	19	-
Male	2	2	7	8	14	16	35	29	28	6	1	-
Female	-	1	6	5	12	27	36	44	58	26	18	-
Black, both sexes	1	1	3	3	3	10	6	9	5	3	-	-
Male	1	1	1	2	1	5	4	2	2	1	-	-
Female	-	-	2	1	2	5	2	7	3	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	3	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
North Carolina												
All races, both sexes	3	3	7	8	14	17	51	55	36	19	6	-
Male	3	2	3	2	8	10	24	23	13	4	1	-
Female	-	1	4	6	6	7	27	32	23	15	5	-
White, both sexes	1	3	5	6	6	15	41	43	34	17	5	-
Male	1	2	2	1	4	9	18	20	11	2	1	-
Female	-	1	3	5	2	6	23	23	23	15	4	-
Black, both sexes	2	-	2	2	8	2	10	12	1	2	1	-
Male	2	-	1	1	4	1	6	3	1	2	-	-
Female	-	-	1	1	4	1	4	9	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	4	2	4	1	3	-	-
Male	1	-	-	-	-	2	1	1	-	1	-	-
Female	-	-	-	-	-	2	1	3	1	2	-	-
White, both sexes	1	-	-	-	-	4	2	4	1	3	-	-
Male	1	-	-	-	-	2	1	1	-	1	-	-
Female	-	-	-	-	-	2	1	3	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Ohio												
All races, both sexes	1	5	12	16	20	41	51	63	82	34	12	-
Male	-	4	6	5	7	21	22	22	26	8	2	-
Female	1	1	6	11	13	20	29	41	56	26	10	-
White, both sexes	1	3	10	15	19	36	43	61	78	34	12	-
Male	-	2	5	5	6	18	18	20	24	8	2	-
Female	1	1	5	10	13	18	25	41	54	26	10	-
Black, both sexes	-	2	2	1	1	5	7	2	4	-	-	-
Male	-	2	1	-	1	3	4	2	2	-	-	-
Female	-	-	1	1	-	2	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	3	2	3	4	15	16	15	24	26	13	2	-
Male	2	1	2	3	10	9	5	10	6	5	-	-
Female	1	1	1	1	5	7	10	14	20	8	2	-
White, both sexes	3	2	2	3	11	13	12	22	24	12	2	-
Male	2	1	2	2	8	9	5	10	6	5	-	-
Female	1	1	-	1	3	4	7	12	18	7	2	-
Black, both sexes	-	-	1	1	3	2	-	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	2	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	3	1	2	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Oregon												
All races, both sexes	3	3	4	4	6	4	25	34	21	9	2	-
Male	1	3	3	1	6	1	11	14	8	2	1	-
Female	2	-	1	3	-	3	14	20	13	7	1	-
White, both sexes	3	3	4	4	5	4	25	34	21	9	2	-
Male	1	3	3	1	5	1	11	14	8	2	1	-
Female	2	-	1	3	-	3	14	20	13	7	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	5	8	8	25	45	80	84	88	44	7	-
Male	2	3	5	5	12	21	34	25	22	13	-	-
Female	1	2	3	3	13	24	46	59	66	31	7	-
White, both sexes	2	5	7	8	25	42	76	83	85	43	7	-
Male	2	3	5	5	12	19	33	24	21	12	-	-
Female	-	2	2	3	13	23	43	59	64	31	7	-
Black, both sexes	1	-	1	-	-	3	4	1	3	1	-	-
Male	-	-	-	-	-	2	1	1	1	1	-	-
Female	1	-	1	-	-	1	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Rhode Island												
All races, both sexes	-	1	-	-	3	8	9	11	6	6	4	-
Male	-	1	-	-	2	5	2	5	2	-	-	-
Female	-	-	-	-	1	3	7	6	4	6	4	-
White, both sexes	-	1	-	-	3	8	8	11	6	6	4	-
Male	-	1	-	-	2	5	2	5	2	-	-	-
Female	-	-	-	-	1	3	6	6	4	6	4	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	6	-	3	6	9	8	6	3	-	-
Male	-	-	2	-	1	2	3	4	2	1	-	-
Female	-	-	4	-	2	4	6	4	4	2	-	-
White, both sexes	-	-	4	-	2	4	8	7	4	3	-	-
Male	-	-	2	-	1	2	3	4	2	1	-	-
Female	-	-	2	-	1	2	5	3	2	2	-	-
Black, both sexes	-	-	2	-	1	2	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	1	2	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
South Dakota												
All races, both sexes	-	-	-	-	1	1	3	6	3	3	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	2	5	3	3	-	-
White, both sexes	-	-	-	-	1	1	3	6	3	3	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	2	5	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	4	6	7	8	19	29	43	33	9	5	-
Male	1	2	4	4	7	7	9	16	5	2	1	-
Female	-	2	2	3	1	12	20	27	28	7	4	-
White, both sexes	1	2	4	5	5	16	28	35	32	8	5	-
Male	1	1	2	3	5	5	9	12	5	2	1	-
Female	-	1	2	2	-	11	19	23	27	6	4	-
Black, both sexes	-	2	2	2	3	3	1	8	1	1	-	-
Male	-	1	2	1	2	2	-	4	-	-	-	-
Female	-	1	-	1	1	1	1	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Texas												
All races, both sexes	2	6	7	24	29	38	65	80	72	34	13	-
Male	2	1	6	13	16	18	25	31	19	10	2	-
Female	-	5	1	11	13	20	40	49	53	24	11	-
White, both sexes	1	4	5	20	24	33	60	72	66	29	10	-
Male	1	-	4	10	14	16	22	26	17	8	1	-
Female	-	4	1	10	10	17	38	46	49	21	9	-
Black, both sexes	1	2	2	3	4	4	5	6	5	5	3	-
Male	1	1	2	2	1	1	3	3	1	2	1	-
Female	-	1	-	1	3	3	2	3	4	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	2	1	-	-	-
Male	-	-	-	1	1	1	-	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	2	2	7	7	10	14	10	2	-	-
Male	-	1	-	1	6	5	5	6	5	-	-	-
Female	-	-	2	1	1	2	5	8	5	2	-	-
White, both sexes	-	1	2	2	7	7	10	14	10	2	-	-
Male	-	1	-	1	6	5	5	6	5	-	-	-
Female	-	-	2	1	1	2	5	8	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Vermont												
All races, both sexes	-	-	-	-	-	1	1	6	1	-	-	-
Male	-	-	-	-	-	1	-	3	-	-	-	-
Female	-	-	-	-	-	-	1	3	1	-	-	-
White, both sexes	-	-	-	-	-	1	1	6	1	-	-	-
Male	-	-	-	-	-	1	-	3	-	-	-	-
Female	-	-	-	-	-	-	1	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	5	2	16	12	12	32	31	49	18	3	-
Male	2	3	2	8	9	8	7	11	10	1	-	-
Female	-	2	-	8	3	4	25	20	39	17	3	-
White, both sexes	1	5	1	15	8	10	28	26	41	15	2	-
Male	1	3	1	7	6	7	6	8	8	-	-	-
Female	-	2	-	8	2	3	22	18	33	15	2	-
Black, both sexes	1	-	1	1	4	2	4	5	8	3	1	-
Male	1	-	1	1	3	1	1	3	2	1	-	-
Female	-	-	-	-	1	1	3	2	6	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Washington												
All races, both sexes	1	2	7	6	4	18	33	40	40	27	9	-
Male	1	2	5	4	2	6	11	17	11	11	-	-
Female	-	-	2	2	2	12	22	23	29	16	9	-
White, both sexes	1	2	7	6	4	15	31	38	40	27	9	-
Male	1	2	5	4	2	5	10	16	11	11	-	-
Female	-	-	2	2	2	10	21	22	29	16	9	-
Black, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	2	3	3	3	6	20	20	9	5	3	-
Male	-	1	1	1	2	2	7	6	5	2	-	-
Female	-	1	2	2	1	4	13	14	4	3	3	-
White, both sexes	-	2	3	3	3	6	20	20	9	5	3	-
Male	-	1	1	1	2	2	7	6	5	2	-	-
Female	-	1	2	2	1	4	13	14	4	3	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Wisconsin												
All races, both sexes	1	2	2	3	7	14	22	16	28	9	8	-
Male	1	2	1	2	1	5	11	5	9	1	1	-
Female	-	-	1	1	6	9	11	11	19	8	7	-
White, both sexes	1	1	1	3	7	14	22	16	28	9	8	-
Male	1	1	1	2	1	5	11	5	9	1	1	-
Female	-	-	-	1	6	9	11	11	19	8	7	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
United States												
All races, both sexes	13,238	2	-	1	6	16	27	49	58	124	216	356
Male	7,732	1	-	1	6	11	21	41	46	99	170	269
Female	5,506	1	-	-	-	5	6	8	12	25	46	87
White, both sexes	11,613	-	-	1	6	13	23	36	38	93	147	258
Male	6,836	-	-	1	6	8	19	30	32	71	123	203
Female	4,777	-	-	-	-	5	4	6	6	22	24	55
Black, both sexes	1,267	2	-	-	-	3	3	11	17	21	64	87
Male	678	1	-	-	-	3	2	10	12	19	43	59
Female	589	1	-	-	-	-	1	1	5	2	21	28
American Indian 1/, both sexes	38	-	-	-	-	-	1	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	320	-	-	-	-	-	-	2	3	10	5	9
Male	198	-	-	-	-	-	-	1	2	9	4	6
Female	122	-	-	-	-	-	-	1	1	1	1	3
Alabama												
All races, both sexes	184	-	-	-	-	-	2	1	1	3	4	11
Male	106	-	-	-	-	-	2	1	-	3	2	5
Female	78	-	-	-	-	-	-	-	1	-	2	6
White, both sexes	140	-	-	-	-	-	1	-	-	-	2	3
Male	84	-	-	-	-	-	1	-	-	-	1	3
Female	56	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	44	-	-	-	-	-	1	1	1	3	2	8
Male	22	-	-	-	-	-	1	1	-	3	1	2
Female	22	-	-	-	-	-	-	-	1	-	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Alaska												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	240	-	-	-	-	1	1	2	1	1	2	7
Male	139	-	-	-	-	-	1	2	1	-	1	6
Female	101	-	-	-	-	1	-	-	-	1	1	1
White, both sexes	229	-	-	-	-	1	1	1	-	1	2	7
Male	131	-	-	-	-	-	1	1	-	-	1	6
Female	98	-	-	-	-	1	-	-	-	1	1	1
Black, both sexes	7	-	-	-	-	-	-	1	1	-	-	-
Male	6	-	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Arkansas												
All races, both sexes	166	-	-	-	1	-	-	1	1	1	2	4
Male	115	-	-	-	1	-	-	1	1	1	2	3
Female	51	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	147	-	-	-	1	-	-	-	1	-	1	3
Male	104	-	-	-	1	-	-	-	1	-	1	2
Female	43	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	18	-	-	-	-	-	-	1	-	1	1	1
Male	11	-	-	-	-	-	-	1	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,219	-	-	-	-	2	2	4	6	15	33	45
Male	759	-	-	-	-	2	2	4	4	13	27	36
Female	460	-	-	-	-	-	-	-	2	2	6	9
White, both sexes	988	-	-	-	-	1	2	4	4	9	26	36
Male	620	-	-	-	-	1	2	4	3	7	23	28
Female	368	-	-	-	-	-	-	-	1	2	3	8
Black, both sexes	89	-	-	-	-	1	-	-	-	1	5	5
Male	45	-	-	-	-	1	-	-	-	1	3	5
Female	44	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	142	-	-	-	-	-	-	-	2	5	2	4
Male	94	-	-	-	-	-	-	-	1	5	1	3
Female	48	-	-	-	-	-	-	-	1	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Colorado												
All races, both sexes	201	-	-	1	-	-	-	-	-	5	3	6
Male	119	-	-	1	-	-	-	-	-	2	3	6
Female	82	-	-	-	-	-	-	-	-	3	-	-
White, both sexes	187	-	-	1	-	-	-	-	-	5	3	6
Male	109	-	-	1	-	-	-	-	-	2	3	6
Female	78	-	-	-	-	-	-	-	-	3	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	170	-	-	-	-	-	-	2	1	-	1	3
Male	95	-	-	-	-	-	-	2	-	-	1	2
Female	75	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	162	-	-	-	-	-	-	2	1	-	1	3
Male	90	-	-	-	-	-	-	2	-	-	1	2
Female	72	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Delaware												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	26	-	-	-	-	-	-	-	-	1	-	2
Male	16	-	-	-	-	-	-	-	-	1	-	2
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Florida												
All races, both sexes	952	1	-	-	1	1	-	2	1	5	7	25
Male	602	1	-	-	1	-	-	2	1	4	5	24
Female	350	-	-	-	-	1	-	-	-	1	2	1
White, both sexes	862	-	-	-	1	1	-	2	-	2	3	16
Male	547	-	-	-	1	-	-	2	-	2	2	15
Female	315	-	-	-	-	1	-	-	-	-	1	1
Black, both sexes	84	1	-	-	-	-	-	-	1	3	3	8
Male	51	1	-	-	-	-	-	-	1	2	2	8
Female	33	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	329	-	-	-	-	-	1	-	2	6	9	15
Male	186	-	-	-	-	-	-	-	2	6	7	10
Female	143	-	-	-	-	-	1	-	-	-	2	5
White, both sexes	260	-	-	-	-	-	1	-	-	4	1	10
Male	141	-	-	-	-	-	-	-	-	4	1	6
Female	119	-	-	-	-	-	1	-	-	-	-	4
Black, both sexes	65	-	-	-	-	-	-	-	2	2	8	5
Male	42	-	-	-	-	-	-	-	2	2	6	4
Female	23	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Hawaii												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	1	4
Male	35	-	-	-	-	-	-	-	-	-	1	3
Female	33	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	3
Male	13	-	-	-	-	-	-	-	-	-	-	3
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	71	-	-	-	1	-	-	1	-	-	-	1
Male	39	-	-	-	1	-	-	1	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	1	-	-	1	-	-	-	1
Male	37	-	-	-	1	-	-	1	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Illinois												
All races, both sexes	528	-	-	-	-	1	-	2	3	5	9	17
Male	295	-	-	-	-	1	-	2	3	4	7	14
Female	233	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	455	-	-	-	-	-	-	1	3	2	8	13
Male	257	-	-	-	-	-	-	1	3	1	7	11
Female	198	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	64	-	-	-	-	1	-	1	-	2	1	3
Male	32	-	-	-	-	1	-	1	-	2	-	2
Female	32	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	335	-	-	-	-	-	-	-	2	2	5	12
Male	206	-	-	-	-	-	-	-	2	2	4	8
Female	129	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	319	-	-	-	-	-	-	-	1	2	4	10
Male	198	-	-	-	-	-	-	-	1	2	3	7
Female	121	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	15	-	-	-	-	-	-	-	1	-	1	2
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Iowa												
All races, both sexes	187	-	-	-	-	-	-	-	1	2	1	3
Male	115	-	-	-	-	-	-	-	-	2	1	3
Female	72	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	185	-	-	-	-	-	-	-	1	2	1	3
Male	114	-	-	-	-	-	-	-	-	2	1	3
Female	71	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	142	-	-	-	-	-	-	2	-	1	4	4
Male	76	-	-	-	-	-	-	1	-	-	3	3
Female	66	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	137	-	-	-	-	-	-	2	-	1	4	4
Male	74	-	-	-	-	-	-	1	-	-	3	3
Female	63	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Kentucky												
All races, both sexes	200	-	-	-	-	-	1	2	-	2	5	3
Male	129	-	-	-	-	-	1	1	-	1	5	2
Female	71	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	182	-	-	-	-	-	1	2	-	2	4	2
Male	117	-	-	-	-	-	1	1	-	1	4	1
Female	65	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	175	-	-	-	-	-	-	1	1	2	5	7
Male	93	-	-	-	-	-	-	1	1	2	3	4
Female	82	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	123	-	-	-	-	-	-	-	1	2	2	3
Male	70	-	-	-	-	-	-	-	1	2	2	2
Female	53	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	50	-	-	-	-	-	-	1	-	-	3	4
Male	21	-	-	-	-	-	-	1	-	-	1	2
Female	29	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Maine												
All races, both sexes	79	-	-	-	-	-	1	-	-	-	2	2
Male	40	-	-	-	-	-	1	-	-	-	1	2
Female	39	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	78	-	-	-	-	-	1	-	-	-	2	2
Male	39	-	-	-	-	-	1	-	-	-	1	2
Female	39	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	211	-	-	-	-	1	1	1	2	3	5	7
Male	106	-	-	-	-	1	1	1	1	2	4	6
Female	105	-	-	-	-	-	-	-	1	1	1	1
White, both sexes	149	-	-	-	-	1	1	-	-	2	3	3
Male	77	-	-	-	-	1	1	-	-	1	3	2
Female	72	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	56	-	-	-	-	-	-	1	2	1	2	4
Male	25	-	-	-	-	-	-	1	1	1	1	4
Female	31	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Massachusetts												
All races, both sexes	302	-	-	-	-	3	1	-	1	4	2	2
Male	175	-	-	-	-	3	1	-	1	4	2	2
Female	127	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	289	-	-	-	-	3	1	-	1	3	2	2
Male	168	-	-	-	-	3	1	-	1	3	2	2
Female	121	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	575	-	-	-	-	-	1	2	4	3	10	19
Male	338	-	-	-	-	-	1	2	4	3	10	14
Female	237	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	512	-	-	-	-	-	1	2	4	2	7	15
Male	301	-	-	-	-	-	1	2	4	2	7	11
Female	211	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	59	-	-	-	-	-	-	-	-	1	3	3
Male	36	-	-	-	-	-	-	-	-	1	3	3
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Minnesota												
All races, both sexes	251	-	-	-	-	-	-	-	-	2	1	2
Male	133	-	-	-	-	-	-	-	-	1	1	2
Female	118	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	245	-	-	-	-	-	-	-	-	2	1	2
Male	129	-	-	-	-	-	-	-	-	1	1	2
Female	116	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	134	-	-	-	-	-	1	-	-	2	4	2
Male	76	-	-	-	-	-	1	-	-	1	4	2
Female	58	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	103	-	-	-	-	-	1	-	-	2	-	-
Male	60	-	-	-	-	-	1	-	-	1	-	-
Female	43	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	-	4	2
Male	16	-	-	-	-	-	-	-	-	-	4	2
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Missouri												
All races, both sexes	305	-	-	-	-	-	-	-	3	-	6	10
Male	175	-	-	-	-	-	-	-	2	-	1	9
Female	130	-	-	-	-	-	-	-	1	-	5	1
White, both sexes	269	-	-	-	-	-	-	-	2	-	3	8
Male	160	-	-	-	-	-	-	-	2	-	1	7
Female	109	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	35	-	-	-	-	-	-	-	1	-	3	2
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	1	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Nebraska												
All races, both sexes	83	-	-	-	-	-	1	-	1	-	1	1
Male	49	-	-	-	-	-	1	-	1	-	1	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	80	-	-	-	-	-	1	-	1	-	1	1
Male	46	-	-	-	-	-	1	-	1	-	1	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	89	-	-	-	-	-	-	3	1	2	1	1
Male	49	-	-	-	-	-	-	1	1	2	1	-
Female	40	-	-	-	-	-	-	2	-	-	-	1
White, both sexes	74	-	-	-	-	-	-	1	1	1	-	-
Male	41	-	-	-	-	-	-	1	1	1	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	1	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	6	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
New Hampshire												
All races, both sexes	60	-	-	-	-	-	-	-	-	1	1	2
Male	34	-	-	-	-	-	-	-	-	1	1	2
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	1	1	2
Male	33	-	-	-	-	-	-	-	-	1	1	2
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	403	-	-	-	1	-	2	2	-	3	1	4
Male	241	-	-	-	1	-	2	2	-	3	1	2
Female	162	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	339	-	-	-	1	-	2	-	-	2	1	3
Male	205	-	-	-	1	-	2	-	-	2	1	2
Female	134	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	50	-	-	-	-	-	-	2	-	-	-	1
Male	27	-	-	-	-	-	-	2	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
New Mexico												
All races, both sexes	77	-	-	-	-	-	-	-	-	1	1	1
Male	53	-	-	-	-	-	-	-	-	1	1	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	71	-	-	-	-	-	-	-	-	1	1	1
Male	48	-	-	-	-	-	-	-	-	1	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	806	-	-	-	1	-	-	3	1	2	14	24
Male	449	-	-	-	1	-	-	3	-	2	11	17
Female	357	-	-	-	-	-	-	-	1	-	3	7
White, both sexes	701	-	-	-	1	-	-	3	1	2	9	18
Male	399	-	-	-	1	-	-	3	-	2	9	15
Female	302	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	93	-	-	-	-	-	-	-	-	-	5	6
Male	41	-	-	-	-	-	-	-	-	-	2	2
Female	52	-	-	-	-	-	-	-	-	-	3	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
North Carolina												
All races, both sexes	425	-	-	-	-	1	-	1	2	6	8	17
Male	251	-	-	-	-	-	-	1	2	6	7	10
Female	174	-	-	-	-	1	-	-	-	-	1	7
White, both sexes	356	-	-	-	-	1	-	-	-	4	3	11
Male	214	-	-	-	-	-	-	-	-	4	3	7
Female	142	-	-	-	-	1	-	-	-	-	-	4
Black, both sexes	67	-	-	-	-	-	-	-	2	2	5	6
Male	35	-	-	-	-	-	-	-	2	2	4	3
Female	32	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Ohio												
All races, both sexes	618	1	-	-	-	-	2	-	3	4	12	17
Male	345	-	-	-	-	-	2	-	2	3	8	11
Female	273	1	-	-	-	-	-	-	1	1	4	6
White, both sexes	545	-	-	-	-	-	2	-	2	3	6	11
Male	307	-	-	-	-	-	2	-	2	2	4	7
Female	238	-	-	-	-	-	-	-	-	1	2	4
Black, both sexes	72	1	-	-	-	-	-	-	1	1	6	6
Male	37	-	-	-	-	-	-	-	-	1	4	4
Female	35	1	-	-	-	-	-	-	1	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	161	-	-	-	-	-	1	-	-	3	3	5
Male	97	-	-	-	-	-	1	-	-	2	2	5
Female	64	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	142	-	-	-	-	-	1	-	-	3	2	5
Male	90	-	-	-	-	-	1	-	-	2	1	5
Female	52	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Oregon												
All races, both sexes	210	-	-	-	-	1	-	1	-	2	1	2
Male	117	-	-	-	-	-	-	-	-	-	1	2
Female	93	-	-	-	-	1	-	1	-	2	-	-
White, both sexes	207	-	-	-	-	1	-	1	-	2	1	2
Male	116	-	-	-	-	-	-	-	-	-	1	2
Female	91	-	-	-	-	1	-	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	741	-	-	-	-	-	-	1	2	5	13	8
Male	393	-	-	-	-	-	-	1	2	2	12	6
Female	348	-	-	-	-	-	-	-	-	3	1	2
White, both sexes	679	-	-	-	-	-	-	1	2	5	11	6
Male	362	-	-	-	-	-	-	1	2	2	10	5
Female	317	-	-	-	-	-	-	-	-	3	1	1
Black, both sexes	58	-	-	-	-	-	-	-	-	-	1	1
Male	28	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Rhode Island												
All races, both sexes	62	-	-	-	-	-	-	-	-	1	-	-
Male	31	-	-	-	-	-	-	-	-	1	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	1	-	-
Male	31	-	-	-	-	-	-	-	-	1	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	199	-	-	-	-	-	3	-	2	1	2	5
Male	116	-	-	-	-	-	3	-	2	1	1	4
Female	83	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	153	-	-	-	-	-	2	-	1	1	-	2
Male	92	-	-	-	-	-	2	-	1	1	-	2
Female	61	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	45	-	-	-	-	-	1	-	1	-	2	3
Male	23	-	-	-	-	-	1	-	1	-	1	2
Female	22	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
South Dakota												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	272	-	-	-	-	1	1	3	2	3	7	12
Male	156	-	-	-	-	1	-	3	2	3	5	6
Female	116	-	-	-	-	-	1	-	-	-	2	6
White, both sexes	238	-	-	-	-	1	-	2	-	3	6	8
Male	140	-	-	-	-	1	-	2	-	3	4	4
Female	98	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	33	-	-	-	-	-	1	1	2	-	1	4
Male	16	-	-	-	-	-	-	1	2	-	1	2
Female	17	-	-	-	-	-	1	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Texas												
All races, both sexes	661	-	-	-	-	2	3	6	7	12	16	25
Male	412	-	-	-	-	2	-	4	6	11	9	19
Female	249	-	-	-	-	-	3	2	1	1	7	6
White, both sexes	581	-	-	-	-	2	3	5	7	10	14	18
Male	358	-	-	-	-	2	-	3	6	10	8	14
Female	223	-	-	-	-	-	3	2	1	-	6	4
Black, both sexes	70	-	-	-	-	-	-	1	-	2	2	7
Male	46	-	-	-	-	-	-	1	-	1	1	5
Female	24	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	79	-	-	-	-	-	-	1	-	-	2	1
Male	49	-	-	-	-	-	-	1	-	-	2	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	75	-	-	-	-	-	-	1	-	-	1	1
Male	46	-	-	-	-	-	-	1	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Vermont												
All races, both sexes	36	-	-	-	-	-	-	1	-	-	1	-
Male	16	-	-	-	-	-	-	1	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	1	-	-	1	-
Male	16	-	-	-	-	-	-	1	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	289	-	-	-	1	1	-	2	4	2	2	6
Male	170	-	-	-	1	1	-	2	2	1	2	6
Female	119	-	-	-	-	-	-	-	2	1	-	-
White, both sexes	240	-	-	-	1	-	-	2	2	2	1	4
Male	141	-	-	-	1	-	-	2	1	1	1	4
Female	99	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	44	-	-	-	-	1	-	-	2	-	1	2
Male	26	-	-	-	-	1	-	-	1	-	1	2
Female	18	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Washington												
All races, both sexes	306	-	-	-	-	1	1	-	1	7	5	1
Male	196	-	-	-	-	-	-	-	1	6	5	-
Female	110	-	-	-	-	1	1	-	-	1	-	1
White, both sexes	282	-	-	-	-	1	-	-	-	4	5	1
Male	183	-	-	-	-	-	-	-	-	4	5	-
Female	99	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	1	2	-	-
Male	8	-	-	-	-	-	-	-	1	1	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	132	-	-	-	-	-	-	1	-	1	1	4
Male	71	-	-	-	-	-	-	-	-	1	1	3
Female	61	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	130	-	-	-	-	-	-	1	-	1	1	4
Male	69	-	-	-	-	-	-	-	-	1	1	3
Female	61	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Wisconsin												
All races, both sexes	276	-	-	-	-	-	1	1	2	3	2	4
Male	151	-	-	-	-	-	1	1	2	1	2	2
Female	125	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	266	-	-	-	-	-	1	1	2	3	2	4
Male	146	-	-	-	-	-	1	1	2	1	2	2
Female	120	-	-	-	-	-	-	-	-	2	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
United States												
All races, both sexes	439	615	751	1,096	1,539	2,177	2,445	1,994	1,039	254	34	-
Male	331	467	537	735	952	1,281	1,285	947	427	96	9	-
Female	108	148	214	361	587	896	1,160	1,047	612	158	25	-
White, both sexes	303	483	639	935	1,367	1,981	2,232	1,844	953	230	31	-
Male	240	375	468	637	862	1,191	1,195	880	400	87	8	-
Female	63	108	171	298	505	790	1,037	964	553	143	23	-
Black, both sexes	120	116	97	131	128	141	144	99	60	21	2	-
Male	81	80	58	79	64	59	51	37	12	7	1	-
Female	39	36	39	52	64	82	93	62	48	14	1	-
American Indian 1/, both sexes	1	4	-	5	6	7	6	5	1	-	-	-
Male	1	3	-	3	2	4	4	2	-	-	-	-
Female	-	1	-	2	4	3	2	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	15	12	15	25	38	48	63	46	25	3	1	-
Male	9	9	11	16	24	27	35	28	15	2	-	-
Female	6	3	4	9	14	21	28	18	10	1	1	-
Alabama												
All races, both sexes	2	8	12	26	22	21	37	23	6	4	1	-
Male	2	6	9	12	13	15	20	13	1	2	-	-
Female	-	2	3	14	9	6	17	10	5	2	1	-
White, both sexes	2	7	10	21	21	16	31	18	5	2	1	-
Male	2	5	7	10	13	11	19	10	1	1	-	-
Female	-	2	3	11	8	5	12	8	4	1	1	-
Black, both sexes	-	1	2	5	1	5	6	5	1	2	-	-
Male	-	1	2	2	-	4	1	3	-	1	-	-
Female	-	-	-	3	1	1	5	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Alaska												
All races, both sexes	1	2	1	-	6	4	2	2	-	-	-	-
Male	-	2	1	-	4	3	1	1	-	-	-	-
Female	1	-	-	-	2	1	1	1	-	-	-	-
White, both sexes	1	2	1	-	5	3	2	1	-	-	-	-
Male	-	2	1	-	4	2	1	1	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	12	6	15	30	25	33	48	34	19	3	-	-
Male	7	4	11	20	14	16	28	16	10	2	-	-
Female	5	2	4	10	11	17	20	18	9	1	-	-
White, both sexes	12	5	14	30	23	32	45	34	18	3	-	-
Male	7	3	10	20	14	15	26	16	9	2	-	-
Female	5	2	4	10	9	17	19	18	9	1	-	-
Black, both sexes	-	1	1	-	1	1	-	-	1	-	-	-
Male	-	1	1	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Arkansas												
All races, both sexes	7	9	13	15	19	32	25	26	6	4	-	-
Male	5	8	12	11	17	23	14	11	5	-	-	-
Female	2	1	1	4	2	9	11	15	1	4	-	-
White, both sexes	5	7	12	14	18	31	22	25	4	3	-	-
Male	4	7	11	10	16	23	13	11	4	-	-	-
Female	1	-	1	4	2	8	9	14	-	3	-	-
Black, both sexes	1	2	1	1	1	1	3	1	2	1	-	-
Male	1	1	1	1	1	-	1	-	1	-	-	-
Female	-	1	-	-	-	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	49	58	71	97	149	183	233	166	86	19	1	-
Male	38	41	45	66	95	107	131	90	49	9	-	-
Female	11	17	26	31	54	76	102	76	37	10	1	-
White, both sexes	31	44	59	80	119	149	198	137	73	16	-	-
Male	26	32	39	57	77	89	111	72	42	7	-	-
Female	5	12	20	23	42	60	87	65	31	9	-	-
Black, both sexes	11	7	6	11	9	12	10	5	5	1	-	-
Male	6	5	3	6	3	5	5	1	1	-	-	-
Female	5	2	3	5	6	7	5	4	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	7	6	6	21	22	25	24	8	2	1	-
Male	6	4	3	3	15	13	15	17	6	2	-	-
Female	1	3	3	3	6	9	10	7	2	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Colorado												
All races, both sexes	5	8	13	17	18	34	34	39	12	6	-	-
Male	2	8	10	12	10	19	14	21	8	3	-	-
Female	3	-	3	5	8	15	20	18	4	3	-	-
White, both sexes	4	6	12	15	18	33	31	36	11	6	-	-
Male	2	6	9	10	10	19	12	19	7	3	-	-
Female	2	-	3	5	8	14	19	17	4	3	-	-
Black, both sexes	-	2	1	2	-	1	2	1	-	-	-	-
Male	-	2	1	2	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	4	6	9	18	27	38	30	23	3	1	-
Male	3	3	5	6	10	18	14	19	10	2	-	-
Female	1	1	1	3	8	9	24	11	13	1	1	-
White, both sexes	2	3	6	8	16	26	38	29	23	3	1	-
Male	2	3	5	6	8	17	14	18	10	2	-	-
Female	-	-	1	2	8	9	24	11	13	1	1	-
Black, both sexes	1	1	-	1	2	-	-	1	-	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Delaware												
All races, both sexes	1	-	1	1	6	7	4	6	1	-	-	-
Male	1	-	1	1	5	5	3	3	-	-	-	-
Female	-	-	-	-	1	2	1	3	1	-	-	-
White, both sexes	-	-	1	1	4	7	4	6	1	-	-	-
Male	-	-	1	1	4	5	3	3	-	-	-	-
Female	-	-	-	-	-	2	1	3	1	-	-	-
Black, both sexes	1	-	-	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	3	2	2	1	5	4	3	1	-	-	-
Male	2	2	-	2	1	3	1	1	1	-	-	-
Female	-	1	2	-	-	2	3	2	-	-	-	-
White, both sexes	-	-	2	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Black, both sexes	2	3	-	1	-	4	4	3	1	-	-	-
Male	2	2	-	1	-	2	1	1	1	-	-	-
Female	-	1	-	-	-	2	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Florida												
All races, both sexes	38	45	58	86	130	149	172	144	71	14	2	-
Male	30	35	45	64	81	101	103	64	37	4	-	-
Female	8	10	13	22	49	48	69	80	34	10	2	-
White, both sexes	30	40	50	76	119	142	160	140	66	12	2	-
Male	24	31	40	58	75	97	98	61	37	4	-	-
Female	6	9	10	18	44	45	62	79	29	8	2	-
Black, both sexes	8	5	8	10	11	6	9	4	5	2	-	-
Male	6	4	5	6	6	3	4	3	-	-	-	-
Female	2	1	3	4	5	3	5	1	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	15	26	29	24	38	44	49	45	21	3	2	-
Male	10	17	21	16	23	21	26	19	7	-	1	-
Female	5	9	8	8	15	23	23	26	14	3	1	-
White, both sexes	8	18	18	16	34	39	43	44	19	3	2	-
Male	6	12	12	11	20	20	24	18	6	-	1	-
Female	2	6	6	5	14	19	19	26	13	3	1	-
Black, both sexes	7	8	10	7	2	5	6	1	2	-	-	-
Male	4	5	8	4	2	1	2	1	1	-	-	-
Female	3	3	2	3	-	4	4	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	-	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Hawaii												
All races, both sexes	1	3	1	7	8	10	17	10	6	-	-	-
Male	-	3	1	4	5	4	8	5	1	-	-	-
Female	1	-	-	3	3	6	9	5	5	-	-	-
White, both sexes	-	2	-	-	3	-	3	3	-	-	-	-
Male	-	2	-	-	3	-	3	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	6	5	10	14	7	6	-	-	-
Male	-	1	1	3	2	4	5	3	1	-	-	-
Female	1	-	-	3	3	6	9	4	5	-	-	-
Idaho												
All races, both sexes	2	2	2	6	5	17	18	9	4	3	-	-
Male	2	1	1	4	3	10	7	5	3	-	-	-
Female	-	1	1	2	2	7	11	4	1	3	-	-
White, both sexes	2	2	2	6	5	17	17	7	4	3	-	-
Male	2	1	1	4	3	10	6	4	3	-	-	-
Female	-	1	1	2	2	7	11	3	1	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Illinois												
All races, both sexes	15	25	32	40	55	69	94	102	50	8	1	-
Male	9	20	26	23	29	43	49	42	20	3	-	-
Female	6	5	6	17	26	26	45	60	30	5	1	-
White, both sexes	9	21	26	33	47	62	85	92	45	7	1	-
Male	7	17	20	19	23	39	46	41	19	3	-	-
Female	2	4	6	14	24	23	39	51	26	4	1	-
Black, both sexes	6	4	6	6	8	6	7	9	3	1	-	-
Male	2	3	6	3	6	3	2	1	-	-	-	-
Female	4	1	-	3	2	3	5	8	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	2	1	2	-	-	-
Male	-	-	-	1	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Indiana												
All races, both sexes	6	16	22	35	42	62	64	39	22	6	-	-
Male	5	11	15	25	26	42	35	23	7	1	-	-
Female	1	5	7	10	16	20	29	16	15	5	-	-
White, both sexes	6	15	20	34	37	60	63	39	22	6	-	-
Male	5	11	14	24	24	41	35	23	7	1	-	-
Female	1	4	6	10	13	19	28	16	15	5	-	-
Black, both sexes	-	1	2	1	4	2	1	-	-	-	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	1	1	-	3	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Iowa												
All races, both sexes	3	7	5	13	12	34	47	29	26	4	-	-
Male	2	5	5	8	10	21	27	15	14	2	-	-
Female	1	2	-	5	2	13	20	14	12	2	-	-
White, both sexes	3	7	5	13	12	34	46	29	26	3	-	-
Male	2	5	5	8	10	21	26	15	14	2	-	-
Female	1	2	-	5	2	13	20	14	12	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	5	7	4	18	23	23	23	19	4	1	-
Male	4	5	5	3	12	10	14	10	6	-	-	-
Female	-	-	2	1	6	13	9	13	13	4	1	-
White, both sexes	4	3	7	3	18	22	23	23	18	4	1	-
Male	4	3	5	3	12	10	14	10	6	-	-	-
Female	-	-	2	-	6	12	9	13	12	4	1	-
Black, both sexes	-	2	-	1	-	1	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Kentucky												
All races, both sexes	5	10	19	20	26	33	38	27	8	1	-	-
Male	3	6	17	15	19	21	23	13	2	-	-	-
Female	2	4	2	5	7	12	15	14	6	1	-	-
White, both sexes	1	10	18	15	24	30	38	26	8	1	-	-
Male	-	6	16	13	18	19	23	12	2	-	-	-
Female	1	4	2	2	6	11	15	14	6	1	-	-
Black, both sexes	4	-	1	4	2	3	-	1	-	-	-	-
Male	3	-	1	1	1	2	-	1	-	-	-	-
Female	1	-	-	3	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	7	11	10	13	23	28	31	20	15	1	-	-
Male	3	7	4	8	19	14	14	5	8	-	-	-
Female	4	4	6	5	4	14	17	15	7	1	-	-
White, both sexes	3	5	6	9	20	22	24	16	10	-	-	-
Male	2	4	3	6	17	12	9	4	6	-	-	-
Female	1	1	3	3	3	10	15	12	4	-	-	-
Black, both sexes	4	6	3	3	3	6	7	4	5	1	-	-
Male	1	3	-	1	2	2	5	1	2	-	-	-
Female	3	3	3	2	1	4	2	3	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Maine												
All races, both sexes	1	-	5	4	8	19	9	13	9	4	2	-
Male	1	-	4	1	5	10	4	7	3	1	-	-
Female	-	-	1	3	3	9	5	6	6	3	2	-
White, both sexes	-	-	5	4	8	19	9	13	9	4	2	-
Male	-	-	4	1	5	10	4	7	3	1	-	-
Female	-	-	1	3	3	9	5	6	6	3	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	7	10	9	21	36	42	33	19	6	2	-
Male	5	6	3	5	8	16	23	17	5	2	-	-
Female	1	1	7	4	13	20	19	16	14	4	2	-
White, both sexes	4	3	3	5	13	27	34	29	15	4	2	-
Male	4	3	1	2	5	14	18	17	3	2	-	-
Female	-	-	2	3	8	13	16	12	12	2	2	-
Black, both sexes	2	3	6	4	8	8	7	3	3	2	-	-
Male	1	2	1	3	3	2	4	-	1	-	-	-
Female	1	1	5	1	5	6	3	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	1	1	1	-	-	-
Male	-	1	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Massachusetts												
All races, both sexes	5	8	17	13	31	50	60	63	27	13	2	-
Male	3	6	14	11	19	27	36	31	10	4	1	-
Female	2	2	3	2	12	23	24	32	17	9	1	-
White, both sexes	5	7	16	12	29	49	58	62	24	13	2	-
Male	3	5	13	10	18	27	35	30	10	4	1	-
Female	2	2	3	2	11	22	23	32	14	9	1	-
Black, both sexes	-	1	1	-	2	-	1	1	2	-	-	-
Male	-	1	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Michigan												
All races, both sexes	23	27	33	50	57	97	112	74	53	8	2	-
Male	18	21	25	34	37	53	57	34	22	3	-	-
Female	5	6	8	16	20	44	55	40	31	5	2	-
White, both sexes	12	20	29	45	52	89	105	69	50	8	2	-
Male	11	17	22	29	33	50	56	32	21	3	-	-
Female	1	3	7	16	19	39	49	37	29	5	2	-
Black, both sexes	11	7	4	5	5	7	6	5	2	-	-	-
Male	7	4	3	5	4	3	1	2	-	-	-	-
Female	4	3	1	-	1	4	5	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Minnesota												
All races, both sexes	5	10	7	24	25	43	55	40	30	6	1	-
Male	4	10	5	15	14	21	30	17	9	3	1	-
Female	1	-	2	9	11	22	25	23	21	3	-	-
White, both sexes	5	9	7	23	24	42	55	38	30	6	1	-
Male	4	9	5	14	13	21	30	16	9	3	1	-
Female	1	-	2	9	11	21	25	22	21	3	-	-
Black, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	8	10	19	14	23	19	18	6	2	-	-
Male	3	4	8	13	10	11	6	10	2	1	-	-
Female	3	4	2	6	4	12	13	8	4	1	-	-
White, both sexes	4	4	7	16	11	22	14	16	5	1	-	-
Male	3	2	5	11	8	11	5	10	2	1	-	-
Female	1	2	2	5	3	11	9	6	3	-	-	-
Black, both sexes	2	4	3	3	3	1	5	2	1	1	-	-
Male	-	2	3	2	2	-	1	-	-	-	-	-
Female	2	2	-	1	1	1	4	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Missouri												
All races, both sexes	7	13	13	27	53	47	50	42	25	9	-	-
Male	6	13	8	18	33	28	23	18	11	5	-	-
Female	1	-	5	9	20	19	27	24	14	4	-	-
White, both sexes	7	11	10	25	46	44	45	38	21	9	-	-
Male	6	11	7	17	30	27	21	16	10	5	-	-
Female	1	-	3	8	16	17	24	22	11	4	-	-
Black, both sexes	-	2	3	2	7	2	5	4	4	-	-	-
Male	-	2	1	1	3	-	2	2	1	-	-	-
Female	-	-	2	1	4	2	3	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	2	2	1	4	8	4	12	1	1	-	-
Male	2	1	1	1	2	6	3	5	1	1	-	-
Female	-	1	1	-	2	2	1	7	-	-	-	-
White, both sexes	1	2	2	1	4	8	4	11	1	1	-	-
Male	1	1	1	1	2	6	3	4	1	1	-	-
Female	-	1	1	-	2	2	1	7	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Nebraska												
All races, both sexes	4	4	7	7	9	18	11	13	2	4	-	-
Male	3	1	7	5	6	8	4	9	1	1	-	-
Female	1	3	-	2	3	10	7	4	1	3	-	-
White, both sexes	2	4	6	7	9	18	11	13	2	4	-	-
Male	1	1	6	5	6	8	4	9	1	1	-	-
Female	1	3	-	2	3	10	7	4	1	3	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	5	7	8	8	14	20	7	7	2	-	-
Male	1	4	5	4	6	10	9	3	2	-	-	-
Female	2	1	2	4	2	4	11	4	5	2	-	-
White, both sexes	3	4	6	6	8	9	19	7	7	2	-	-
Male	1	3	4	3	6	7	9	3	2	-	-	-
Female	2	1	2	3	2	2	10	4	5	2	-	-
Black, both sexes	-	-	-	2	-	4	1	-	-	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
New Hampshire												
All races, both sexes	1	2	5	1	10	10	7	11	8	1	-	-
Male	1	2	3	-	7	6	5	3	3	-	-	-
Female	-	-	2	1	3	4	2	8	5	1	-	-
White, both sexes	1	1	5	1	10	10	7	11	8	1	-	-
Male	1	1	3	-	7	6	5	3	3	-	-	-
Female	-	-	2	1	3	4	2	8	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	17	22	24	33	44	68	81	60	35	4	2	-
Male	12	19	18	24	26	33	49	31	15	2	1	-
Female	5	3	6	9	18	35	32	29	20	2	1	-
White, both sexes	13	15	16	21	34	62	74	57	33	3	2	-
Male	10	13	12	16	24	31	44	31	13	2	1	-
Female	3	2	4	5	10	31	30	26	20	1	1	-
Black, both sexes	3	7	7	11	9	3	4	2	-	1	-	-
Male	2	6	5	7	1	1	3	-	-	-	-	-
Female	1	1	2	4	8	2	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	2	3	1	2	-	-	-
Male	-	-	1	1	1	-	2	-	2	-	-	-
Female	1	-	-	-	-	2	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
New Mexico												
All races, both sexes	3	4	3	8	9	17	13	9	6	2	-	-
Male	1	4	2	8	7	13	7	3	4	1	-	-
Female	2	-	1	-	2	4	6	6	2	1	-	-
White, both sexes	2	2	3	6	9	17	13	8	6	2	-	-
Male	-	2	2	6	7	13	7	3	4	1	-	-
Female	2	-	1	-	2	4	6	5	2	1	-	-
Black, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	27	42	53	75	85	135	121	129	66	26	2	-
Male	24	34	34	47	50	82	56	51	28	8	1	-
Female	3	8	19	28	35	53	65	78	38	18	1	-
White, both sexes	17	31	43	64	75	123	106	121	61	24	2	-
Male	16	25	30	41	46	77	50	48	28	7	1	-
Female	1	6	13	23	29	46	56	73	33	17	1	-
Black, both sexes	10	10	9	10	9	9	12	6	5	2	-	-
Male	8	8	4	5	3	3	3	2	-	1	-	-
Female	2	2	5	5	6	6	9	4	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	1	2	3	2	-	-	-	-
Male	-	1	-	1	1	2	3	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
North Carolina												
All races, both sexes	13	18	26	32	56	61	85	60	33	6	-	-
Male	10	14	20	17	35	37	46	29	14	3	-	-
Female	3	4	6	15	21	24	39	31	19	3	-	-
White, both sexes	11	12	23	26	50	51	77	52	30	5	-	-
Male	8	10	18	15	34	32	43	25	13	2	-	-
Female	3	2	5	11	16	19	34	27	17	3	-	-
Black, both sexes	2	6	3	6	6	9	8	8	3	1	-	-
Male	2	4	2	2	1	4	3	4	1	1	-	-
Female	-	2	1	4	5	5	5	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	4	3	2	9	2	11	5	-	-	-
Male	1	-	3	3	1	5	1	3	4	-	-	-
Female	2	-	1	-	1	4	1	8	1	-	-	-
White, both sexes	3	-	4	2	2	9	2	11	5	-	-	-
Male	1	-	3	2	1	5	1	3	4	-	-	-
Female	2	-	1	-	1	4	1	8	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Ohio												
All races, both sexes	18	24	38	48	62	104	122	118	37	6	2	-
Male	14	18	27	37	33	60	60	53	12	4	1	-
Female	4	6	11	11	29	44	62	65	25	2	1	-
White, both sexes	11	18	32	46	58	91	114	111	33	6	1	-
Male	9	13	25	36	32	54	57	50	10	4	-	-
Female	2	5	7	10	26	37	57	61	23	2	1	-
Black, both sexes	7	6	6	2	4	13	8	7	3	-	1	-
Male	5	5	2	1	1	6	3	3	1	-	1	-
Female	2	1	4	1	3	7	5	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	7	12	12	25	22	29	21	15	2	-	-
Male	3	6	8	9	12	15	20	11	3	-	-	-
Female	1	1	4	3	13	7	9	10	12	2	-	-
White, both sexes	4	7	11	12	17	20	26	21	12	1	-	-
Male	3	6	8	9	10	13	18	11	3	-	-	-
Female	1	1	3	3	7	7	8	10	9	1	-	-
Black, both sexes	-	-	1	-	3	2	-	-	2	1	-	-
Male	-	-	-	-	2	2	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	4	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	4	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Oregon												
All races, both sexes	7	9	8	14	25	34	46	41	15	2	2	-
Male	6	8	6	9	15	22	20	19	7	1	1	-
Female	1	1	2	5	10	12	26	22	8	1	1	-
White, both sexes	7	9	8	13	25	34	46	40	14	2	2	-
Male	6	8	6	9	15	22	20	19	6	1	1	-
Female	1	1	2	4	10	12	26	21	8	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	20	27	23	43	75	161	156	114	70	21	2	-
Male	14	21	14	33	38	87	80	56	19	8	-	-
Female	6	6	9	10	37	74	76	58	51	13	2	-
White, both sexes	9	24	18	39	70	149	148	111	64	20	2	-
Male	8	18	11	31	36	82	76	54	19	7	-	-
Female	1	6	7	8	34	67	72	57	45	13	2	-
Black, both sexes	10	3	5	3	5	12	8	3	6	1	-	-
Male	6	3	3	1	2	5	4	2	-	1	-	-
Female	4	-	2	2	3	7	4	1	6	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Rhode Island												
All races, both sexes	1	5	3	4	9	5	17	7	10	-	-	-
Male	1	3	2	2	6	5	7	2	2	-	-	-
Female	-	2	1	2	3	-	10	5	8	-	-	-
White, both sexes	1	4	3	4	9	5	17	7	10	-	-	-
Male	1	3	2	2	6	5	7	2	2	-	-	-
Female	-	1	1	2	3	-	10	5	8	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	9	11	7	19	31	38	30	25	12	4	-	-
Male	7	4	5	13	19	25	10	13	7	2	-	-
Female	2	7	2	6	12	13	20	12	5	2	-	-
White, both sexes	6	4	5	14	25	36	24	20	11	2	-	-
Male	5	1	4	10	15	24	9	11	7	-	-	-
Female	1	3	1	4	10	12	15	9	4	2	-	-
Black, both sexes	3	7	2	5	5	2	6	5	1	2	-	-
Male	2	3	1	3	3	1	1	2	-	2	-	-
Female	1	4	1	2	2	1	5	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
South Dakota												
All races, both sexes	-	1	2	3	6	5	12	10	8	-	1	-
Male	-	-	2	3	2	4	6	5	1	-	-	-
Female	-	1	-	-	4	1	6	5	7	-	1	-
White, both sexes	-	1	2	3	6	4	12	10	8	-	1	-
Male	-	-	2	3	2	3	6	5	1	-	-	-
Female	-	1	-	-	4	1	6	5	7	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	11	11	17	32	42	37	39	32	17	5	-	-
Male	10	8	13	14	28	18	17	19	8	1	-	-
Female	1	3	4	18	14	19	22	13	9	4	-	-
White, both sexes	5	10	17	27	41	34	35	28	16	5	-	-
Male	4	8	13	13	27	18	16	18	8	1	-	-
Female	1	2	4	14	14	16	19	10	8	4	-	-
Black, both sexes	6	1	-	4	1	3	4	4	1	-	-	-
Male	6	-	-	1	1	-	1	1	-	-	-	-
Female	-	1	-	3	-	3	3	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Texas												
All races, both sexes	38	34	36	55	78	96	114	94	28	15	2	-
Male	33	27	21	42	55	61	55	45	16	5	1	-
Female	5	7	15	13	23	35	59	49	12	10	1	-
White, both sexes	26	25	32	45	68	91	106	86	27	14	2	-
Male	22	20	20	33	46	59	54	41	15	4	1	-
Female	4	5	12	12	22	32	52	45	12	10	1	-
Black, both sexes	10	9	3	10	7	4	7	6	1	1	-	-
Male	9	7	-	9	6	2	1	2	1	1	-	-
Female	1	2	3	1	1	2	6	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	3	1	1	2	-	-	-	-
Male	2	-	1	-	3	-	-	2	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Utah												
All races, both sexes	2	5	7	9	10	11	19	6	6	-	-	-
Male	1	3	6	4	8	8	11	3	1	-	-	-
Female	1	2	1	5	2	3	8	3	5	-	-	-
White, both sexes	2	4	7	9	10	10	18	6	6	-	-	-
Male	1	2	6	4	8	8	10	3	1	-	-	-
Female	1	2	1	5	2	2	8	3	5	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Vermont												
All races, both sexes	-	1	-	4	3	6	8	5	5	1	1	-
Male	-	-	-	1	2	3	3	3	1	1	-	-
Female	-	1	-	3	1	3	5	2	4	-	1	-
White, both sexes	-	1	-	4	3	6	8	5	5	1	1	-
Male	-	-	-	1	2	3	3	3	1	1	-	-
Female	-	1	-	3	1	3	5	2	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	6	15	17	29	34	58	47	42	19	3	1	-
Male	6	8	14	23	20	36	23	19	6	-	-	-
Female	-	7	3	6	14	22	24	23	13	3	1	-
White, both sexes	3	13	15	20	27	51	41	37	19	2	-	-
Male	3	8	12	15	16	32	21	18	6	-	-	-
Female	-	5	3	5	11	19	20	19	13	2	-	-
Black, both sexes	3	2	2	7	6	6	6	5	-	-	1	-
Male	3	-	2	6	4	3	2	1	-	-	-	-
Female	-	2	-	1	2	3	4	4	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	1	-	-	-	1	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Washington												
All races, both sexes	8	22	11	22	30	57	57	51	27	5	-	-
Male	7	18	8	13	24	36	37	28	8	5	-	-
Female	1	4	3	9	6	21	20	23	19	-	-	-
White, both sexes	8	21	10	19	29	54	52	47	26	5	-	-
Male	7	18	7	12	23	34	35	25	8	5	-	-
Female	1	3	3	7	6	20	17	22	18	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	3	4	3	1	-	-	-
Male	-	-	1	-	-	2	1	2	-	-	-	-
Female	-	-	-	2	-	1	3	1	1	-	-	-
West Virginia												
All races, both sexes	3	5	8	14	16	23	23	16	14	3	-	-
Male	2	4	5	8	8	12	12	10	4	1	-	-
Female	1	1	3	6	8	11	11	6	10	2	-	-
White, both sexes	3	4	8	14	16	23	23	15	14	3	-	-
Male	2	3	5	8	8	12	12	9	4	1	-	-
Female	1	1	3	6	8	11	11	6	10	2	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Wisconsin												
All races, both sexes	6	16	14	26	35	42	52	35	27	9	1	-
Male	4	14	10	16	19	21	31	12	9	5	1	-
Female	2	2	4	10	16	21	21	23	18	4	-	-
White, both sexes	5	14	14	24	34	40	51	34	27	9	1	-
Male	4	12	10	15	18	20	31	12	9	5	1	-
Female	1	2	4	9	16	20	20	22	18	4	-	-
Black, both sexes	1	1	-	1	1	2	1	-	-	-	-	-
Male	-	1	-	1	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	2	3	3	1	8	5	5	1	1	-	-
Male	-	2	2	2	1	6	2	2	-	-	-	-
Female	1	-	1	1	-	2	3	3	1	1	-	-
White, both sexes	-	2	3	3	1	8	5	5	1	1	-	-
Male	-	2	2	2	1	6	2	2	-	-	-	-
Female	-	-	1	1	-	2	3	3	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
United States												
All races, both sexes	9,246	16	1	2	3	11	14	17	24	44	60	152
Male	4,104	11	-	1	1	5	7	6	13	18	32	83
Female	5,142	5	1	1	2	6	7	11	11	26	28	69
White, both sexes	7,884	13	1	1	1	10	9	11	20	28	43	112
Male	3,537	9	-	1	1	4	4	4	11	9	20	62
Female	4,347	4	1	-	-	6	5	7	9	19	23	50
Black, both sexes	1,215	3	-	-	2	1	4	5	4	15	16	37
Male	506	2	-	-	-	1	2	2	2	8	11	21
Female	709	1	-	-	2	-	2	3	2	7	5	16
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-	1	1	2
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	106	-	-	1	-	-	1	1	-	-	-	1
Male	47	-	-	-	-	-	1	-	-	-	-	-
Female	59	-	-	1	-	-	-	1	-	-	-	1
Alabama												
All races, both sexes	169	-	-	-	-	-	-	-	-	-	3	5
Male	83	-	-	-	-	-	-	-	-	-	3	4
Female	86	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	115	-	-	-	-	-	-	-	-	-	2	4
Male	60	-	-	-	-	-	-	-	-	-	2	3
Female	55	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	53	-	-	-	-	-	-	-	-	-	1	1
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Alaska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	136	-	-	-	-	-	-	-	-	-	-	2
Male	66	-	-	-	-	-	-	-	-	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	129	-	-	-	-	-	-	-	-	-	-	1
Male	63	-	-	-	-	-	-	-	-	-	-	-
Female	66	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Arkansas												
All races, both sexes	131	-	-	-	-	-	-	-	1	1	2	6
Male	56	-	-	-	-	-	-	-	1	-	1	5
Female	75	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	115	-	-	-	-	-	-	-	1	-	2	6
Male	53	-	-	-	-	-	-	-	1	-	1	5
Female	62	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	766	1	-	-	-	2	3	1	4	9	9	14
Male	333	1	-	-	-	-	1	-	-	3	7	6
Female	433	-	-	-	-	2	2	1	4	6	2	8
White, both sexes	649	1	-	-	-	2	1	-	2	7	9	10
Male	288	1	-	-	-	-	-	-	-	2	7	5
Female	361	-	-	-	-	2	1	-	2	5	2	5
Black, both sexes	67	-	-	-	-	-	1	-	2	2	-	3
Male	23	-	-	-	-	-	-	-	-	1	-	1
Female	44	-	-	-	-	-	1	-	2	1	-	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	1	1	-	-	-	-
Male	21	-	-	-	-	-	1	-	-	-	-	-
Female	26	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Colorado												
All races, both sexes	108	-	1	-	-	-	1	1	-	-	-	2
Male	46	-	-	-	-	-	1	-	-	-	-	-
Female	62	-	1	-	-	-	-	1	-	-	-	2
White, both sexes	104	-	1	-	-	-	1	1	-	-	-	2
Male	43	-	-	-	-	-	1	-	-	-	-	-
Female	61	-	1	-	-	-	-	1	-	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	139	-	-	-	-	-	-	1	1	-	-	1
Male	60	-	-	-	-	-	-	1	1	-	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	118	-	-	-	-	-	-	-	1	-	-	1
Male	49	-	-	-	-	-	-	-	1	-	-	-
Female	69	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	20	-	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Delaware												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Florida												
All races, both sexes	615	-	-	2	-	1	-	1	1	3	2	8
Male	311	-	-	1	-	1	-	-	1	3	1	4
Female	304	-	-	1	-	-	-	1	-	-	1	4
White, both sexes	536	-	-	1	-	-	-	1	1	1	2	3
Male	277	-	-	1	-	-	-	-	1	1	1	2
Female	259	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	76	-	-	-	-	1	-	-	-	2	-	5
Male	33	-	-	-	-	1	-	-	-	2	-	2
Female	43	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	236	1	-	-	-	-	1	1	1	2	2	4
Male	106	-	-	-	-	-	-	-	-	1	2	1
Female	130	1	-	-	-	-	1	1	1	1	-	3
White, both sexes	151	1	-	-	-	-	-	-	1	1	1	2
Male	64	-	-	-	-	-	-	-	-	-	1	1
Female	87	1	-	-	-	-	-	-	1	1	-	1
Black, both sexes	84	-	-	-	-	-	1	1	-	1	1	2
Male	42	-	-	-	-	-	-	-	-	1	1	-
Female	42	-	-	-	-	-	1	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hawaii												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	44	-	-	-	-	-	-	-	-	2	-	1
Male	26	-	-	-	-	-	-	-	-	1	-	1
Female	18	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	2	-	1
Male	26	-	-	-	-	-	-	-	-	1	-	1
Female	17	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Illinois												
All races, both sexes	410	-	-	-	-	1	1	1	-	1	1	7
Male	163	-	-	-	-	-	1	1	-	-	1	4
Female	247	-	-	-	-	1	-	-	-	1	-	3
White, both sexes	346	-	-	-	-	1	-	1	-	1	1	4
Male	142	-	-	-	-	-	-	1	-	-	1	3
Female	204	-	-	-	-	1	-	-	-	1	-	1
Black, both sexes	62	-	-	-	-	-	1	-	-	-	-	3
Male	20	-	-	-	-	-	1	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	199	2	-	-	-	-	-	-	1	4	1	1
Male	106	1	-	-	-	-	-	-	1	1	1	1
Female	93	1	-	-	-	-	-	-	-	3	-	-
White, both sexes	183	1	-	-	-	-	-	-	1	3	1	1
Male	98	-	-	-	-	-	-	-	1	1	1	1
Female	85	1	-	-	-	-	-	-	-	2	-	-
Black, both sexes	16	1	-	-	-	-	-	-	-	1	-	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Iowa												
All races, both sexes	100	-	-	-	-	-	-	-	-	-	-	1
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	97	-	-	-	-	-	-	-	-	-	-	1
Male	41	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	96	-	-	-	-	-	-	1	-	-	3	-
Male	45	-	-	-	-	-	-	1	-	-	2	-
Female	51	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	89	-	-	-	-	-	-	-	-	-	3	-
Male	39	-	-	-	-	-	-	-	-	-	2	-
Female	50	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Kentucky												
All races, both sexes	178	-	-	-	-	-	-	1	-	-	1	2
Male	69	-	-	-	-	-	-	-	-	-	1	1
Female	109	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	166	-	-	-	-	-	-	1	-	-	-	2
Male	62	-	-	-	-	-	-	-	-	-	-	1
Female	104	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	159	-	-	-	-	-	-	1	1	1	2	8
Male	70	-	-	-	-	-	-	-	1	1	1	3
Female	89	-	-	-	-	-	-	1	-	-	1	5
White, both sexes	102	-	-	-	-	-	-	-	-	1	1	2
Male	45	-	-	-	-	-	-	-	-	1	-	1
Female	57	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	56	-	-	-	-	-	-	1	1	-	1	6
Male	25	-	-	-	-	-	-	-	1	-	1	2
Female	31	-	-	-	-	-	-	1	-	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Maine												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	183	-	-	-	-	-	1	-	-	2	1	4
Male	72	-	-	-	-	-	1	-	-	1	1	2
Female	111	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	129	-	-	-	-	-	-	-	-	1	-	2
Male	51	-	-	-	-	-	-	-	-	-	-	-
Female	78	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	54	-	-	-	-	-	1	-	-	1	1	2
Male	21	-	-	-	-	-	1	-	-	1	1	2
Female	33	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Massachusetts												
All races, both sexes	227	-	-	-	-	-	-	-	-	-	1	2
Male	95	-	-	-	-	-	-	-	-	-	-	2
Female	132	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	221	-	-	-	-	-	-	-	-	-	1	2
Male	92	-	-	-	-	-	-	-	-	-	-	2
Female	129	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	331	2	-	-	-	-	-	-	-	1	3	4
Male	136	2	-	-	-	-	-	-	-	-	1	2
Female	195	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	272	2	-	-	-	-	-	-	-	1	1	3
Male	106	2	-	-	-	-	-	-	-	-	-	1
Female	166	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	55	-	-	-	-	-	-	-	-	-	2	1
Male	28	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Minnesota												
All races, both sexes	155	-	-	-	-	-	-	-	-	-	-	2
Male	67	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	154	-	-	-	-	-	-	-	-	-	-	2
Male	67	-	-	-	-	-	-	-	-	-	-	-
Female	87	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	77	-	-	-	-	-	-	1	-	-	1	2
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	46	-	-	-	-	-	-	1	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Missouri												
All races, both sexes	204	2	-	-	-	-	-	-	-	-	-	2
Male	81	1	-	-	-	-	-	-	-	-	-	2
Female	123	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	175	2	-	-	-	-	-	-	-	-	-	2
Male	73	1	-	-	-	-	-	-	-	-	-	2
Female	102	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	27	-	-	-	-	-	-	-	-	1	-	-
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	1	-	-
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Nebraska												
All races, both sexes	66	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	66	-	-	-	-	1	-	-	-	-	-	3
Male	32	-	-	-	-	1	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	62	-	-	-	-	1	-	-	-	-	-	3
Male	29	-	-	-	-	1	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Hampshire												
All races, both sexes	40	-	-	-	-	-	-	-	-	1	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	1	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	292	-	-	-	-	-	-	-	-	-	1	3
Male	131	-	-	-	-	-	-	-	-	-	-	1
Female	161	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	246	-	-	-	-	-	-	-	-	-	-	2
Male	113	-	-	-	-	-	-	-	-	-	-	-
Female	133	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	43	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Mexico												
All races, both sexes	50	-	-	-	-	-	1	-	-	-	-	-
Male	21	-	-	-	-	-	1	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	1	-	-	-	-	-
Male	20	-	-	-	-	-	1	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	431	1	-	-	1	-	-	-	-	1	1	9
Male	192	1	-	-	1	-	-	-	-	1	-	7
Female	239	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	369	1	-	-	1	-	-	-	-	1	-	6
Male	162	1	-	-	1	-	-	-	-	1	-	4
Female	207	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	58	-	-	-	-	-	-	-	-	-	1	3
Male	28	-	-	-	-	-	-	-	-	-	-	3
Female	30	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
North Carolina												
All races, both sexes	309	-	-	-	-	-	1	-	-	-	2	5
Male	139	-	-	-	-	-	1	-	-	-	2	4
Female	170	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	219	-	-	-	-	-	1	-	-	-	1	5
Male	103	-	-	-	-	-	1	-	-	-	1	4
Female	116	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	87	-	-	-	-	-	-	-	-	-	1	-
Male	36	-	-	-	-	-	-	-	-	-	1	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Ohio												
All races, both sexes	498	1	-	-	-	1	2	1	2	2	2	7
Male	225	1	-	-	-	1	-	1	2	1	1	5
Female	273	-	-	-	-	-	2	-	-	1	1	2
White, both sexes	439	1	-	-	-	1	2	1	2	1	1	6
Male	211	1	-	-	-	1	-	1	2	-	-	4
Female	228	-	-	-	-	-	2	-	-	1	1	2
Black, both sexes	57	-	-	-	-	-	-	-	-	1	1	1
Male	14	-	-	-	-	-	-	-	-	1	1	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	108	-	-	-	-	-	-	-	1	-	-	4
Male	38	-	-	-	-	-	-	-	-	-	-	3
Female	70	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	91	-	-	-	-	-	-	-	1	-	-	4
Male	31	-	-	-	-	-	-	-	-	-	-	3
Female	60	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Oregon												
All races, both sexes	105	-	-	-	-	-	-	-	-	-	-	1
Male	48	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	100	-	-	-	-	-	-	-	-	-	-	1
Male	46	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	539	1	-	-	-	2	-	-	1	2	4	3
Male	238	1	-	-	-	1	-	-	1	1	2	1
Female	301	-	-	-	-	1	-	-	-	1	2	2
White, both sexes	488	1	-	-	-	2	-	-	1	2	4	2
Male	224	1	-	-	-	1	-	-	1	1	2	-
Female	264	-	-	-	-	1	-	-	-	1	2	2
Black, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Rhode Island												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	160	-	-	-	-	1	-	1	-	1	-	3
Male	71	-	-	-	-	1	-	1	-	-	-	1
Female	89	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	124	-	-	-	-	1	-	1	-	-	-	3
Male	57	-	-	-	-	1	-	1	-	-	-	1
Female	67	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	35	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
South Dakota												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	232	-	-	-	-	1	-	1	2	1	4	4
Male	92	-	-	-	-	-	-	1	2	-	2	3
Female	140	-	-	-	-	1	-	-	-	1	2	1
White, both sexes	197	-	-	-	-	1	-	1	1	1	3	3
Male	77	-	-	-	-	-	-	1	1	-	1	2
Female	120	-	-	-	-	1	-	-	-	1	2	1
Black, both sexes	35	-	-	-	-	-	-	-	1	-	1	1
Male	15	-	-	-	-	-	-	-	1	-	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Texas												
All races, both sexes	632	4	-	-	2	1	2	3	5	5	9	14
Male	277	2	-	-	-	-	1	-	3	3	2	7
Female	355	2	-	-	2	1	1	3	2	2	7	7
White, both sexes	539	3	-	-	-	1	2	3	5	2	7	12
Male	239	2	-	-	-	-	1	-	3	-	-	6
Female	300	1	-	-	-	1	1	3	2	2	7	6
Black, both sexes	91	1	-	-	2	-	-	-	-	2	2	2
Male	36	-	-	-	-	-	-	-	-	2	2	1
Female	55	1	-	-	2	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	49	-	-	-	-	-	-	-	1	-	-	2
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	47	-	-	-	-	-	-	-	1	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Vermont												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	205	1	-	-	-	-	-	1	1	1	1	6
Male	97	1	-	-	-	-	-	-	-	-	-	4
Female	108	-	-	-	-	-	-	1	1	1	1	2
White, both sexes	139	-	-	-	-	-	-	-	1	-	-	2
Male	63	-	-	-	-	-	-	-	-	-	-	1
Female	76	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	64	1	-	-	-	-	-	1	-	1	1	4
Male	32	1	-	-	-	-	-	-	-	-	-	3
Female	32	-	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Washington												
All races, both sexes	189	-	-	-	-	-	-	-	-	-	1	-
Male	79	-	-	-	-	-	-	-	-	-	-	-
Female	110	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	181	-	-	-	-	-	-	-	-	-	-	-
Male	75	-	-	-	-	-	-	-	-	-	-	-
Female	106	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	78	-	-	-	-	-	-	-	-	-	-	2
Male	49	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	76	-	-	-	-	-	-	-	-	-	-	2
Male	47	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Wisconsin												
All races, both sexes	238	-	-	-	-	-	1	-	1	3	3	3
Male	91	-	-	-	-	-	-	-	-	-	1	2
Female	147	-	-	-	-	-	1	-	1	3	2	1
White, both sexes	228	-	-	-	-	-	1	-	1	1	2	3
Male	88	-	-	-	-	-	-	-	-	-	-	2
Female	140	-	-	-	-	-	1	-	1	1	2	1
Black, both sexes	8	-	-	-	-	-	-	-	-	2	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
United States												
All races, both sexes	215	342	445	601	868	1,301	1,646	1,683	1,223	497	81	-
Male	123	205	255	362	464	650	731	668	355	97	17	-
Female	92	137	190	239	404	651	915	1,015	868	400	64	-
White, both sexes	160	259	357	488	739	1,111	1,408	1,521	1,094	437	61	-
Male	91	152	204	297	404	569	646	620	328	87	14	-
Female	69	107	153	191	335	542	762	901	766	350	47	-
Black, both sexes	50	78	79	102	116	165	210	142	117	51	18	-
Male	28	51	46	64	54	71	69	42	22	8	2	-
Female	22	27	33	38	62	94	141	100	95	43	16	-
American Indian 1/, both sexes	1	4	3	7	5	7	7	1	1	1	-	-
Male	1	1	1	1	3	3	1	1	-	-	-	-
Female	-	3	2	6	2	4	6	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	4	1	6	4	8	18	21	19	11	8	2	-
Male	3	1	4	-	3	7	15	5	5	2	1	-
Female	1	-	2	4	5	11	6	14	6	6	1	-
Alabama												
All races, both sexes	7	7	6	9	18	26	32	31	15	8	2	-
Male	4	3	3	7	11	16	17	12	1	2	-	-
Female	3	4	3	2	7	10	15	19	14	6	2	-
White, both sexes	3	4	4	6	12	19	21	25	9	5	1	-
Male	2	2	3	4	7	11	13	11	-	2	-	-
Female	1	2	1	2	5	8	8	14	9	3	1	-
Black, both sexes	4	3	2	3	6	7	11	6	6	2	1	-
Male	2	1	-	3	4	5	4	1	1	-	-	-
Female	2	2	2	-	2	2	7	5	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Alaska												
All races, both sexes	1	-	1	2	2	-	1	2	-	1	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	1	2	-	1	-	-
White, both sexes	1	-	1	1	2	-	1	2	-	1	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	5	8	14	10	18	20	28	23	3	-	-
Male	3	4	4	9	7	9	9	9	10	2	-	-
Female	2	1	4	5	3	9	11	19	13	1	-	-
White, both sexes	4	5	8	12	8	17	20	28	23	3	-	-
Male	3	4	4	8	6	8	9	9	10	2	-	-
Female	1	1	4	4	2	9	11	19	13	1	-	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Arkansas												
All races, both sexes	6	7	9	11	13	19	17	16	17	4	2	-
Male	4	5	2	8	9	10	5	3	3	-	-	-
Female	2	2	7	3	4	9	12	13	14	4	2	-
White, both sexes	6	5	7	11	11	17	15	14	14	4	2	-
Male	4	4	1	8	8	10	5	3	3	-	-	-
Female	2	1	6	3	3	7	10	11	11	4	2	-
Black, both sexes	-	2	2	-	2	2	2	2	3	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	1	2	2	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	19	33	28	50	70	94	137	136	103	44	9	-
Male	10	22	17	28	39	41	63	50	35	8	2	-
Female	9	11	11	22	31	53	74	86	68	36	7	-
White, both sexes	13	29	23	42	60	79	119	115	93	36	8	-
Male	8	19	14	23	34	37	55	43	33	5	2	-
Female	5	10	9	19	26	42	64	72	60	31	6	-
Black, both sexes	4	4	4	5	8	7	8	10	5	3	1	-
Male	-	3	2	5	4	1	1	4	-	1	-	-
Female	4	1	2	-	4	6	7	6	5	2	1	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	3	1	8	10	11	5	5	-	-
Male	2	-	-	-	1	3	7	3	2	2	-	-
Female	-	-	-	3	-	5	3	8	3	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Colorado												
All races, both sexes	3	1	7	6	7	21	19	22	12	5	-	-
Male	3	1	4	3	3	10	4	11	4	2	-	-
Female	-	-	3	3	4	11	15	11	8	3	-	-
White, both sexes	3	1	6	6	7	18	19	22	12	5	-	-
Male	3	1	3	3	3	8	4	11	4	2	-	-
Female	-	-	3	3	4	10	15	11	8	3	-	-
Black, both sexes	-	-	1	-	-	3	-	-	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	4	4	6	15	16	28	25	29	8	1	-
Male	-	3	2	6	10	6	9	7	15	-	-	-
Female	-	1	2	-	5	10	19	18	14	8	1	-
White, both sexes	-	2	2	6	11	14	22	23	27	8	1	-
Male	-	2	2	6	7	5	6	7	13	-	-	-
Female	-	-	-	-	4	9	16	16	14	8	1	-
Black, both sexes	-	2	2	-	4	2	6	2	1	-	-	-
Male	-	1	-	-	3	1	3	-	1	-	-	-
Female	-	1	2	-	1	1	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Delaware												
All races, both sexes	-	-	1	2	3	2	4	5	3	3	-	-
Male	-	-	-	1	3	1	1	3	-	-	-	-
Female	-	-	1	1	-	1	3	2	3	3	-	-
White, both sexes	-	-	-	2	1	2	4	5	2	3	-	-
Male	-	-	-	1	1	1	1	3	-	-	-	-
Female	-	-	-	1	-	1	3	2	2	3	-	-
Black, both sexes	-	-	1	-	2	-	-	-	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	3	3	2	5	4	2	1	-	-
Male	-	-	1	2	2	1	2	1	-	-	-	-
Female	-	1	-	1	1	1	3	3	2	1	-	-
White, both sexes	-	-	-	-	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	1	1	3	3	2	4	1	2	1	-	-
Male	-	-	1	2	2	1	2	-	-	-	-	-
Female	-	1	-	1	1	1	2	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Florida												
All races, both sexes	18	24	32	41	72	98	88	108	80	34	2	-
Male	9	15	21	27	44	54	49	49	25	7	-	-
Female	9	9	11	14	28	44	39	59	55	27	2	-
White, both sexes	16	20	28	35	59	91	80	94	74	30	-	-
Male	8	11	19	24	38	50	44	45	25	7	-	-
Female	8	9	9	11	21	41	36	49	49	23	-	-
Black, both sexes	2	4	3	6	12	7	8	14	6	4	2	-
Male	1	4	2	3	5	4	5	4	-	-	-	-
Female	1	-	1	3	7	3	3	10	6	4	2	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	8	15	16	19	20	25	35	42	26	16	2	-
Male	4	12	12	14	14	11	15	13	4	3	-	-
Female	4	3	4	5	6	14	20	29	22	13	2	-
White, both sexes	2	9	7	11	14	17	25	32	18	9	1	-
Male	1	6	5	8	11	6	11	10	2	2	-	-
Female	1	3	2	3	3	11	14	22	16	7	1	-
Black, both sexes	6	6	9	8	6	8	10	10	7	7	1	-
Male	3	6	7	6	3	5	4	3	2	1	-	-
Female	3	-	2	2	3	3	6	7	5	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hawaii												
All races, both sexes	-	-	1	-	3	7	8	5	1	4	2	-
Male	-	-	1	-	1	3	5	2	1	1	1	-
Female	-	-	-	-	2	4	3	3	-	3	1	-
White, both sexes	-	-	1	-	1	2	2	2	-	2	-	-
Male	-	-	1	-	-	1	2	2	-	1	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	4	6	3	1	2	2	-
Male	-	-	-	-	1	2	3	-	1	-	1	-
Female	-	-	-	-	1	2	3	3	-	2	1	-
Idaho												
All races, both sexes	-	-	3	1	5	4	7	13	5	3	-	-
Male	-	-	3	-	2	3	3	8	3	2	-	-
Female	-	-	-	1	3	1	4	5	2	1	-	-
White, both sexes	-	-	3	1	5	4	6	13	5	3	-	-
Male	-	-	3	-	2	3	3	8	3	2	-	-
Female	-	-	-	1	3	1	3	5	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Illinois												
All races, both sexes	10	11	14	21	31	59	82	82	66	17	5	-
Male	2	8	9	10	10	27	38	30	21	1	-	-
Female	8	3	5	11	21	32	44	52	45	16	5	-
White, both sexes	7	8	13	16	29	49	63	73	61	15	4	-
Male	1	7	8	7	9	25	31	27	21	1	-	-
Female	6	1	5	9	20	24	32	46	40	14	4	-
Black, both sexes	3	3	1	5	2	10	17	9	5	2	1	-
Male	1	1	1	3	1	2	6	3	-	-	-	-
Female	2	2	-	2	1	8	11	6	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	4	5	7	13	13	35	45	29	24	12	3	-
Male	3	4	3	12	8	22	18	16	10	4	1	-
Female	1	1	4	1	5	13	27	13	14	8	2	-
White, both sexes	4	5	7	13	13	33	41	27	22	9	2	-
Male	3	4	3	12	8	21	15	16	9	2	1	-
Female	1	1	4	1	5	12	26	11	13	7	1	-
Black, both sexes	-	-	-	-	-	2	4	2	2	3	1	-
Male	-	-	-	-	-	1	3	-	1	2	-	-
Female	-	-	-	-	-	1	1	2	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Iowa												
All races, both sexes	1	1	8	3	10	14	13	25	18	2	4	-
Male	1	1	4	1	6	8	6	9	4	1	-	-
Female	-	-	4	2	4	6	7	16	14	1	4	-
White, both sexes	1	1	6	3	9	14	13	25	18	2	4	-
Male	1	1	3	1	6	8	6	9	4	1	-	-
Female	-	-	3	2	3	6	7	16	14	1	4	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	5	4	3	7	18	16	20	9	6	3	-
Male	1	2	2	2	3	11	7	10	-	3	1	-
Female	-	3	2	1	4	7	9	10	9	3	2	-
White, both sexes	1	4	4	3	6	16	15	20	9	5	3	-
Male	1	1	2	2	2	9	7	10	-	2	1	-
Female	-	3	2	1	4	7	8	10	9	3	2	-
Black, both sexes	-	-	-	-	1	1	1	-	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Kentucky												
All races, both sexes	2	8	6	18	17	24	38	26	24	8	3	-
Male	-	5	3	7	9	8	16	10	8	1	-	-
Female	2	3	3	11	8	16	22	16	16	7	3	-
White, both sexes	2	7	5	16	16	21	38	26	22	8	2	-
Male	-	4	2	7	8	7	16	10	6	1	-	-
Female	2	3	3	9	8	14	22	16	16	7	2	-
Black, both sexes	-	1	-	2	1	3	-	-	2	-	1	-
Male	-	1	-	-	1	1	-	-	2	-	-	-
Female	-	-	-	2	-	2	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	5	14	14	24	11	26	22	19	7	-	-
Male	3	1	4	12	16	6	12	6	4	-	-	-
Female	1	4	10	2	8	5	14	16	15	7	-	-
White, both sexes	3	3	7	6	19	9	16	15	15	5	-	-
Male	2	1	2	5	13	4	9	3	4	-	-	-
Female	1	2	5	1	6	5	7	12	11	5	-	-
Black, both sexes	1	2	7	8	4	2	10	7	4	2	-	-
Male	1	-	2	7	3	2	3	3	-	-	-	-
Female	-	2	5	1	1	-	7	4	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Maine												
All races, both sexes	3	1	4	6	5	3	17	4	6	2	-	-
Male	1	1	2	2	3	1	10	1	3	1	-	-
Female	2	-	2	4	2	2	7	3	3	1	-	-
White, both sexes	3	1	4	6	5	3	17	4	6	2	-	-
Male	1	1	2	2	3	1	10	1	3	1	-	-
Female	2	-	2	4	2	2	7	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	12	10	12	14	32	24	36	23	9	2	-
Male	1	5	7	9	7	15	6	11	5	-	1	-
Female	-	7	3	3	7	17	18	25	18	9	1	-
White, both sexes	1	5	5	9	10	20	18	30	20	7	1	-
Male	1	2	4	7	6	10	5	10	5	-	1	-
Female	-	3	1	2	4	10	13	20	15	7	-	-
Black, both sexes	-	7	5	3	4	12	6	6	3	2	1	-
Male	-	3	3	2	1	5	1	1	-	-	-	-
Female	-	4	2	1	3	7	5	5	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Massachusetts												
All races, both sexes	2	5	9	11	21	22	53	58	27	16	-	-
Male	2	4	4	7	7	11	29	18	8	3	-	-
Female	-	1	5	4	14	11	24	40	19	13	-	-
White, both sexes	2	5	9	11	20	22	51	55	27	16	-	-
Male	2	4	4	7	6	11	27	18	8	3	-	-
Female	-	1	5	4	14	11	24	37	19	13	-	-
Black, both sexes	-	-	-	-	1	-	2	2	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	6	17	13	34	31	48	57	52	42	21	-	-
Male	6	8	5	21	17	18	20	20	11	5	-	-
Female	-	9	8	13	14	30	37	32	31	16	-	-
White, both sexes	4	12	11	26	23	40	43	48	38	20	-	-
Male	4	5	3	15	12	17	13	19	10	5	-	-
Female	-	7	8	11	11	23	30	29	28	15	-	-
Black, both sexes	2	5	2	8	7	7	12	4	4	1	-	-
Male	2	3	2	6	4	1	6	1	1	-	-	-
Female	-	2	-	2	3	6	6	3	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Minnesota												
All races, both sexes	1	5	4	2	16	17	30	42	29	4	3	-
Male	1	3	2	1	7	6	15	20	12	-	-	-
Female	-	2	2	1	9	11	15	22	17	4	3	-
White, both sexes	1	5	4	2	16	16	30	42	29	4	3	-
Male	1	3	2	1	7	6	15	20	12	-	-	-
Female	-	2	2	1	9	10	15	22	17	4	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	9	8	9	8	6	10	7	9	6	1	-	-
Male	8	6	7	7	2	5	4	3	1	-	-	-
Female	1	2	2	1	4	5	3	6	5	1	-	-
White, both sexes	4	6	3	3	4	7	5	6	4	1	-	-
Male	4	5	2	3	2	5	3	-	1	-	-	-
Female	-	1	1	-	2	2	2	6	3	1	-	-
Black, both sexes	5	2	5	5	2	3	2	3	2	-	-	-
Male	4	1	4	4	-	-	1	3	-	-	-	-
Female	1	1	1	1	2	3	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Missouri												
All races, both sexes	5	6	9	21	22	24	38	35	27	13	-	-
Male	3	3	5	11	8	13	16	12	7	-	-	-
Female	2	3	4	10	14	11	22	23	20	13	-	-
White, both sexes	3	3	7	17	19	20	31	34	24	13	-	-
Male	2	2	3	10	8	11	15	12	7	-	-	-
Female	1	1	4	7	11	9	16	22	17	13	-	-
Black, both sexes	2	3	2	4	3	4	7	1	3	-	-	-
Male	1	1	2	1	-	2	1	-	-	-	-	-
Female	1	2	-	3	3	2	6	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	1	1	1	4	3	3	5	7	-	-	-
Male	-	1	1	1	2	2	3	2	3	-	-	-
Female	1	-	-	-	2	1	-	3	4	-	-	-
White, both sexes	1	1	1	1	4	3	3	5	6	-	-	-
Male	-	1	1	1	2	2	3	2	3	-	-	-
Female	1	-	-	-	2	1	-	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Nebraska												
All races, both sexes	1	-	2	1	4	8	16	12	15	7	-	-
Male	1	-	1	1	1	4	7	3	3	2	-	-
Female	-	-	1	-	3	4	9	9	12	5	-	-
White, both sexes	1	-	2	1	4	8	16	12	14	7	-	-
Male	1	-	1	1	1	4	7	3	2	2	-	-
Female	-	-	1	-	3	4	9	9	12	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	4	3	6	8	11	12	8	1	4	1	-
Male	3	1	1	4	5	6	7	2	1	-	-	-
Female	1	3	2	2	3	5	5	6	-	4	1	-
White, both sexes	4	4	3	5	8	9	12	7	1	4	1	-
Male	3	1	1	3	5	4	7	2	1	-	-	-
Female	1	3	2	2	3	5	5	5	-	4	1	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Hampshire												
All races, both sexes	1	2	2	1	5	6	8	5	4	5	-	-
Male	1	-	2	-	2	4	3	5	-	2	-	-
Female	-	2	-	1	3	2	5	-	4	3	-	-
White, both sexes	1	2	2	1	5	6	8	5	4	5	-	-
Male	1	-	2	-	2	4	3	5	-	2	-	-
Female	-	2	-	1	3	2	5	-	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	8	14	12	23	40	64	65	37	17	7	-
Male	-	8	6	6	11	17	33	28	12	7	2	-
Female	1	-	8	6	12	23	31	37	25	10	5	-
White, both sexes	1	5	14	10	20	27	55	56	34	16	6	-
Male	-	5	6	6	10	11	31	23	12	7	2	-
Female	1	-	8	4	10	16	24	33	22	9	4	-
Black, both sexes	-	3	-	1	3	12	9	9	2	1	1	-
Male	-	3	-	-	1	6	2	5	-	-	-	-
Female	-	-	-	1	2	6	7	4	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Mexico												
All races, both sexes	2	2	1	5	2	4	9	11	8	5	-	-
Male	2	-	-	4	2	2	4	5	1	-	-	-
Female	-	2	1	1	-	2	5	6	7	5	-	-
White, both sexes	2	2	1	3	2	3	9	11	8	4	-	-
Male	2	-	-	3	2	2	4	5	1	-	-	-
Female	-	2	1	-	-	1	5	6	7	4	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	15	13	17	28	32	64	77	70	66	33	3	-
Male	9	7	14	18	16	29	28	32	20	8	1	-
Female	6	6	3	10	16	35	49	38	46	25	2	-
White, both sexes	10	11	14	24	27	51	65	63	61	31	3	-
Male	4	5	12	16	13	23	24	30	20	7	1	-
Female	6	6	2	8	14	28	41	33	41	24	2	-
Black, both sexes	5	2	2	4	5	13	11	5	5	2	-	-
Male	5	2	2	2	3	6	3	1	-	1	-	-
Female	-	-	-	2	2	7	8	4	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
North Carolina												
All races, both sexes	12	15	29	16	29	46	48	50	45	8	3	-
Male	7	11	14	8	14	25	14	24	12	3	-	-
Female	5	4	15	8	15	21	34	26	33	5	3	-
White, both sexes	8	9	20	9	20	34	35	40	33	4	-	-
Male	4	7	11	4	10	18	12	19	10	2	-	-
Female	4	2	9	5	10	16	23	21	23	2	-	-
Black, both sexes	4	6	9	7	8	12	11	10	12	4	3	-
Male	3	4	3	4	4	7	2	5	2	1	-	-
Female	1	2	6	3	4	5	9	5	10	3	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	4	-	4	2	3	6	1	-	-
Male	1	-	-	3	-	3	-	1	3	1	-	-
Female	-	-	-	1	-	1	2	2	3	-	-	-
White, both sexes	1	-	-	4	-	4	2	3	6	1	-	-
Male	1	-	-	3	-	3	-	1	3	1	-	-
Female	-	-	-	1	-	1	2	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Ohio												
All races, both sexes	12	19	14	35	43	93	95	83	59	27	-	-
Male	7	14	9	18	21	46	42	34	18	4	-	-
Female	5	5	5	17	22	47	53	49	41	23	-	-
White, both sexes	10	19	14	27	36	81	85	78	49	25	-	-
Male	7	14	9	16	20	42	40	34	16	4	-	-
Female	3	5	5	11	16	39	45	44	33	21	-	-
Black, both sexes	2	-	-	7	6	12	10	5	10	2	-	-
Male	-	-	-	2	1	4	2	-	2	-	-	-
Female	2	-	-	5	5	8	8	5	8	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	4	7	10	17	15	11	14	17	4	-	-
Male	2	2	2	3	3	8	3	6	4	2	-	-
Female	2	2	5	7	14	7	8	8	13	2	-	-
White, both sexes	3	3	6	7	15	13	7	12	16	4	-	-
Male	1	1	1	3	3	7	1	5	4	2	-	-
Female	2	2	5	4	12	6	6	7	12	2	-	-
Black, both sexes	-	-	1	-	-	1	3	1	1	-	-	-
Male	-	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
American Indian 1/, both sexes	1	1	-	3	1	1	1	1	-	-	-	-
Male	1	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	3	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Oregon												
All races, both sexes	1	6	5	8	14	23	17	20	7	3	-	-
Male	-	3	3	6	9	13	4	8	2	-	-	-
Female	1	3	2	2	5	10	13	12	5	3	-	-
White, both sexes	1	4	5	8	13	21	17	20	7	3	-	-
Male	-	2	3	6	8	13	4	8	2	-	-	-
Female	1	2	2	2	5	8	13	12	5	3	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	13	28	20	48	79	103	112	81	32	5	-
Male	1	6	15	13	25	45	59	45	15	6	1	-
Female	4	7	13	7	23	34	44	67	66	26	4	-
White, both sexes	4	11	23	16	42	73	93	105	75	29	5	-
Male	1	5	12	12	25	44	54	45	14	5	1	-
Female	3	6	11	4	17	29	39	60	61	24	4	-
Black, both sexes	1	2	5	4	6	6	10	7	6	3	-	-
Male	-	1	3	1	-	1	5	-	1	1	-	-
Female	1	1	2	3	6	5	5	7	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Rhode Island												
All races, both sexes	-	-	2	2	1	5	5	11	4	3	-	-
Male	-	-	1	2	1	2	2	3	2	1	-	-
Female	-	-	1	-	-	3	3	8	2	2	-	-
White, both sexes	-	-	2	2	1	5	5	11	4	2	-	-
Male	-	-	1	2	1	2	2	3	2	1	-	-
Female	-	-	1	-	-	3	3	8	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	7	11	12	22	25	25	27	15	9	-	-
Male	1	4	7	8	11	10	14	10	2	1	-	-
Female	-	3	4	4	11	15	11	17	13	8	-	-
White, both sexes	-	4	8	10	17	21	18	24	13	4	-	-
Male	-	3	5	7	8	10	10	8	2	1	-	-
Female	-	1	3	3	9	11	8	16	11	3	-	-
Black, both sexes	1	2	3	2	5	4	7	3	2	5	-	-
Male	1	1	2	1	3	-	4	2	-	-	-	-
Female	-	1	1	1	2	4	3	1	2	5	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
South Dakota												
All races, both sexes	-	2	1	2	3	3	3	6	3	7	2	-
Male	-	-	-	1	3	2	1	1	3	2	2	-
Female	-	2	1	1	-	1	2	5	-	5	-	-
White, both sexes	-	1	1	2	3	3	3	6	3	7	2	-
Male	-	-	-	1	3	2	1	1	3	2	2	-
Female	-	1	1	1	-	1	2	5	-	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	9	13	19	23	31	45	34	25	11	4	-
Male	2	3	9	9	12	15	17	10	5	-	2	-
Female	3	6	4	10	11	16	28	24	20	11	2	-
White, both sexes	4	8	12	15	19	26	36	30	23	11	3	-
Male	1	2	8	6	11	13	14	10	5	-	2	-
Female	3	6	4	9	8	13	22	20	18	11	1	-
Black, both sexes	1	1	1	4	4	5	9	4	2	-	1	-
Male	1	1	1	3	1	2	3	-	-	-	-	-
Female	-	-	-	1	3	3	6	4	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Texas												
All races, both sexes	17	29	35	45	54	89	111	106	67	31	3	-
Male	10	15	23	23	30	42	44	50	19	2	1	-
Female	7	14	12	22	24	47	67	56	48	29	2	-
White, both sexes	11	19	32	42	51	79	91	97	55	26	1	-
Male	7	7	20	21	28	38	42	46	16	2	-	-
Female	4	12	12	21	23	41	49	51	39	24	1	-
Black, both sexes	6	10	3	3	3	10	20	8	12	5	2	-
Male	3	8	3	2	2	4	2	3	3	-	1	-
Female	3	2	-	1	1	6	18	5	9	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	2	2	4	2	6	16	10	2	2	-	-
Male	-	2	2	2	2	3	9	7	-	1	-	-
Female	-	-	-	2	-	3	7	3	2	1	-	-
White, both sexes	-	2	2	4	2	5	16	10	2	2	-	-
Male	-	2	2	2	2	2	9	7	-	1	-	-
Female	-	-	-	2	-	3	7	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Vermont												
All races, both sexes	-	1	-	1	2	3	7	2	5	2	1	-
Male	-	-	-	-	1	2	3	1	2	-	1	-
Female	-	1	-	1	1	1	4	1	3	2	-	-
White, both sexes	-	1	-	1	2	3	7	2	5	2	1	-
Male	-	-	-	-	1	2	3	1	2	-	1	-
Female	-	1	-	1	1	1	4	1	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	6	7	13	11	19	28	34	41	27	5	3	-
Male	2	5	10	6	12	16	13	13	13	1	1	-
Female	4	2	3	5	7	12	21	28	14	4	2	-
White, both sexes	5	3	8	5	16	22	23	30	18	5	1	-
Male	2	2	7	2	10	11	10	10	7	1	-	-
Female	3	1	1	3	6	11	13	20	11	4	1	-
Black, both sexes	1	4	5	6	3	6	10	11	8	-	2	-
Male	-	3	3	4	2	5	2	3	5	-	1	-
Female	1	1	2	2	1	1	8	8	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Washington												
All races, both sexes	3	5	8	9	21	22	41	37	31	9	2	-
Male	2	3	5	5	10	13	18	14	5	4	-	-
Female	1	2	3	4	11	9	23	23	26	5	2	-
White, both sexes	2	5	5	9	21	22	39	37	30	9	2	-
Male	1	3	4	5	10	13	16	14	5	4	-	-
Female	1	2	1	4	11	9	23	23	25	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	-	-	2	-	1	-	-	-
Male	1	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	-	3	1	9	10	8	15	13	13	4	-	-
Male	-	1	1	8	9	7	10	7	4	-	-	-
Female	-	2	-	1	1	1	5	6	9	4	-	-
White, both sexes	-	3	1	8	9	8	15	13	13	4	-	-
Male	-	1	1	7	8	7	10	7	4	-	-	-
Female	-	2	-	1	1	1	5	6	9	4	-	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Wisconsin												
All races, both sexes	6	3	8	9	23	32	30	56	41	15	4	-
Male	2	3	2	4	13	16	12	22	11	3	-	-
Female	4	-	6	5	10	16	18	34	30	12	4	-
White, both sexes	6	2	6	9	21	30	30	56	41	15	4	-
Male	2	2	2	4	13	15	12	22	11	3	-	-
Female	4	-	4	5	8	15	18	34	30	12	4	-
Black, both sexes	-	1	2	-	2	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	-	1	1	4	2	5	1	1	-	-
Male	-	-	-	1	-	2	2	1	-	1	-	-
Female	1	1	-	-	1	2	-	4	1	-	-	-
White, both sexes	1	-	-	1	1	4	2	5	1	1	-	-
Male	-	-	-	1	-	2	2	1	-	1	-	-
Female	1	-	-	-	1	2	-	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
United States												
All races, both sexes	3,995	35	3	3	2	14	32	35	79	116	162	260
Male	1,772	18	1	3	-	7	12	16	38	64	85	141
Female	2,223	17	2	-	2	7	20	19	41	52	77	119
White, both sexes	3,213	29	3	2	1	9	19	20	52	78	101	178
Male	1,414	14	1	2	-	3	6	7	26	48	56	104
Female	1,799	15	2	-	1	6	13	13	26	30	45	74
Black, both sexes	713	4	-	1	1	5	13	15	26	37	55	76
Male	324	2	-	1	-	4	6	9	11	15	25	34
Female	389	2	-	-	1	1	7	6	15	22	30	42
American Indian 1/, both sexes	22	1	-	-	-	-	-	-	1	-	3	5
Male	8	1	-	-	-	-	-	-	1	-	2	2
Female	14	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	47	1	-	-	-	-	-	-	-	1	3	1
Male	26	1	-	-	-	-	-	-	-	1	2	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	72	-	-	-	-	-	-	-	-	1	2	2
Male	27	-	-	-	-	-	-	-	-	1	1	1
Female	45	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	56	-	-	-	-	-	-	-	-	1	1	2
Male	20	-	-	-	-	-	-	-	-	1	1	1
Female	36	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	84	-	-	-	-	2	1	1	5	-	2	4
Male	35	-	-	-	-	1	-	1	2	-	1	2
Female	49	-	-	-	-	1	1	-	3	-	1	2
White, both sexes	77	-	-	-	-	2	1	1	5	-	1	4
Male	33	-	-	-	-	1	-	1	2	-	1	2
Female	44	-	-	-	-	1	1	-	3	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Arkansas												
All races, both sexes	50	-	-	-	-	-	1	1	-	-	2	4
Male	22	-	-	-	-	-	1	-	-	-	2	1
Female	28	-	-	-	-	-	-	1	-	-	-	3
White, both sexes	43	-	-	-	-	-	-	-	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	2	1
Female	23	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	1	1	-	-	-	2
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	463	4	-	-	-	-	3	3	8	14	23	33
Male	220	2	-	-	-	-	1	-	7	9	15	21
Female	243	2	-	-	-	-	2	3	1	5	8	12
White, both sexes	393	4	-	-	-	-	3	3	6	10	14	26
Male	185	2	-	-	-	-	1	-	6	6	9	18
Female	208	2	-	-	-	-	2	3	-	4	5	8
Black, both sexes	54	-	-	-	-	-	-	-	2	4	8	6
Male	26	-	-	-	-	-	-	-	1	3	5	2
Female	28	-	-	-	-	-	-	-	1	1	3	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Colorado												
All races, both sexes	62	2	-	-	-	1	-	-	1	4	1	4
Male	33	2	-	-	-	-	-	-	1	3	1	4
Female	29	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	60	2	-	-	-	1	-	-	1	4	1	4
Male	32	2	-	-	-	-	-	-	1	3	1	4
Female	28	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	1	3
Male	17	-	-	-	-	-	-	-	-	-	1	2
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Delaware												
All races, both sexes	12	1	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	12	1	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	17	-	-	-	-	-	-	-	1	3	2	2
Male	7	-	-	-	-	-	-	-	1	-	2	-
Female	10	-	-	-	-	-	-	-	-	3	-	2
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	1	3	2	2
Male	7	-	-	-	-	-	-	-	1	-	2	-
Female	10	-	-	-	-	-	-	-	-	3	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Florida												
All races, both sexes	255	3	-	-	-	-	-	3	2	5	7	18
Male	114	1	-	-	-	-	-	2	1	4	3	9
Female	141	2	-	-	-	-	-	1	1	1	4	9
White, both sexes	201	3	-	-	-	-	-	1	2	3	3	12
Male	82	1	-	-	-	-	-	1	1	3	1	4
Female	119	2	-	-	-	-	-	-	1	-	2	8
Black, both sexes	53	-	-	-	-	-	-	2	-	2	4	6
Male	31	-	-	-	-	-	-	1	-	1	2	5
Female	22	-	-	-	-	-	-	1	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	125	-	-	-	-	3	2	1	3	3	3	6
Male	52	-	-	-	-	1	1	1	-	2	-	2
Female	73	-	-	-	-	2	1	-	3	1	3	4
White, both sexes	81	-	-	-	-	1	1	-	1	2	1	3
Male	36	-	-	-	-	-	1	-	-	2	-	1
Female	45	-	-	-	-	1	-	-	1	-	1	2
Black, both sexes	42	-	-	-	-	2	1	1	2	1	2	3
Male	16	-	-	-	-	1	-	1	-	-	-	1
Female	26	-	-	-	-	1	1	-	2	1	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Hawaii												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	17	1	-	1	-	-	-	-	-	-	-	3
Male	7	-	-	1	-	-	-	-	-	-	-	2
Female	10	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	1	-	-	-	-	-	-	-	-	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	10	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Illinois												
All races, both sexes	171	3	-	-	-	-	4	7	2	3	10	6
Male	75	3	-	-	-	-	2	4	-	3	5	3
Female	96	-	-	-	-	-	2	3	2	-	5	3
White, both sexes	128	1	-	-	-	-	2	3	-	2	5	5
Male	51	1	-	-	-	-	1	-	-	2	3	2
Female	77	-	-	-	-	-	1	3	-	-	2	3
Black, both sexes	40	1	-	-	-	-	2	4	2	1	5	1
Male	22	1	-	-	-	-	1	4	-	1	2	1
Female	18	-	-	-	-	-	1	-	2	-	3	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	71	2	-	-	-	-	-	-	-	1	4	5
Male	34	2	-	-	-	-	-	-	-	-	2	4
Female	37	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	65	2	-	-	-	-	-	-	-	1	3	5
Male	33	2	-	-	-	-	-	-	-	-	1	4
Female	32	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Iowa												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	4	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	25	-	-	-	-	-	-	-	-	-	4	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	35	-	-	-	-	-	-	-	2	1	1	3
Male	14	-	-	-	-	-	-	-	-	1	-	2
Female	21	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	32	-	-	-	-	-	-	-	2	1	1	2
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	21	-	-	-	-	-	-	-	2	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Kentucky												
All races, both sexes	95	-	-	-	-	-	1	1	1	2	4	-
Male	36	-	-	-	-	-	1	-	-	1	2	-
Female	59	-	-	-	-	-	-	1	1	1	2	-
White, both sexes	89	-	-	-	-	-	1	1	1	2	2	-
Male	34	-	-	-	-	-	1	-	-	1	1	-
Female	55	-	-	-	-	-	-	1	1	1	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	56	-	-	-	-	1	-	1	1	1	3	4
Male	23	-	-	-	-	1	-	-	-	-	1	3
Female	33	-	-	-	-	-	-	1	1	1	2	1
White, both sexes	34	-	-	-	-	-	-	1	1	1	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	20	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	22	-	-	-	-	1	-	-	-	-	2	3
Male	9	-	-	-	-	1	-	-	-	-	-	2
Female	13	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Maine												
All races, both sexes	23	-	-	-	-	-	-	1	1	2	-	2
Male	11	-	-	-	-	-	-	-	1	-	-	2
Female	12	-	-	-	-	-	-	1	-	2	-	-
White, both sexes	23	-	-	-	-	-	-	1	1	2	-	2
Male	11	-	-	-	-	-	-	-	1	-	-	2
Female	12	-	-	-	-	-	-	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	65	-	-	-	-	1	-	2	2	7	6	3
Male	28	-	-	-	-	-	-	1	2	2	1	1
Female	37	-	-	-	-	1	-	1	-	5	5	2
White, both sexes	36	-	-	-	-	1	-	1	-	4	1	1
Male	17	-	-	-	-	-	-	1	-	2	-	1
Female	19	-	-	-	-	1	-	-	-	2	1	-
Black, both sexes	29	-	-	-	-	-	-	1	2	3	5	2
Male	11	-	-	-	-	-	-	-	2	-	1	-
Female	18	-	-	-	-	-	-	1	-	3	4	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Massachusetts												
All races, both sexes	91	1	-	-	1	-	-	-	4	1	2	5
Male	39	1	-	-	-	-	-	-	3	1	-	5
Female	52	-	-	-	1	-	-	-	1	-	2	-
White, both sexes	84	-	-	-	1	-	-	-	4	1	2	5
Male	33	-	-	-	-	-	-	-	3	1	-	5
Female	51	-	-	-	1	-	-	-	1	-	2	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	178	-	1	-	-	-	2	1	4	11	13	18
Male	90	-	-	-	-	-	-	1	1	7	6	8
Female	88	-	1	-	-	-	2	-	3	4	7	10
White, both sexes	141	-	1	-	-	-	-	-	3	9	12	10
Male	74	-	-	-	-	-	-	-	1	6	5	5
Female	67	-	1	-	-	-	-	-	2	3	7	5
Black, both sexes	35	-	-	-	-	-	2	1	1	2	1	8
Male	15	-	-	-	-	-	-	1	-	1	1	3
Female	20	-	-	-	-	-	2	-	1	1	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Minnesota												
All races, both sexes	43	-	-	-	-	-	1	1	2	2	1	1
Male	17	-	-	-	-	-	-	-	1	2	1	-
Female	26	-	-	-	-	-	1	1	1	-	-	1
White, both sexes	42	-	-	-	-	-	1	1	2	2	1	1
Male	17	-	-	-	-	-	-	-	1	2	1	-
Female	25	-	-	-	-	-	1	1	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	50	-	-	1	-	-	1	-	-	3	2	3
Male	18	-	-	1	-	-	1	-	-	1	1	2
Female	32	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	28	-	-	1	-	-	-	-	-	1	1	-
Male	9	-	-	1	-	-	-	-	-	1	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	1	-	-	2	1	3
Male	9	-	-	-	-	-	1	-	-	-	-	2
Female	13	-	-	-	-	-	-	-	-	2	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Missouri												
All races, both sexes	87	2	1	-	-	1	2	-	2	6	-	4
Male	37	1	-	-	-	1	-	-	1	3	-	1
Female	50	1	1	-	-	-	2	-	1	3	-	3
White, both sexes	75	2	1	-	-	1	-	-	2	4	-	3
Male	32	1	-	-	-	1	-	-	1	2	-	1
Female	43	1	1	-	-	-	-	-	1	2	-	2
Black, both sexes	12	-	-	-	-	-	2	-	-	2	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	2	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Nebraska												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	31	-	-	-	-	-	-	1	-	-	1	2
Male	19	-	-	-	-	-	-	1	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	27	-	-	-	-	-	-	1	-	-	1	1
Male	17	-	-	-	-	-	-	1	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
New Hampshire												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	133	-	-	-	-	-	1	-	3	6	3	9
Male	57	-	-	-	-	-	-	-	1	3	-	6
Female	76	-	-	-	-	-	1	-	2	3	3	3
White, both sexes	100	-	-	-	-	-	1	-	-	3	-	4
Male	45	-	-	-	-	-	-	-	-	2	-	3
Female	55	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	30	-	-	-	-	-	-	-	3	3	3	4
Male	10	-	-	-	-	-	-	-	1	1	-	2
Female	20	-	-	-	-	-	-	-	2	2	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
New Mexico												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	269	2	-	1	1	1	3	3	8	7	13	15
Male	127	1	-	1	-	1	1	-	3	4	7	13
Female	142	1	-	-	1	-	2	3	5	3	6	2
White, both sexes	193	2	-	1	-	-	1	1	5	2	7	13
Male	92	1	-	1	-	-	-	-	2	-	4	11
Female	101	1	-	-	-	-	1	1	3	2	3	2
Black, both sexes	75	-	-	-	1	1	2	2	3	5	6	2
Male	34	-	-	-	-	1	1	-	1	4	3	2
Female	41	-	-	-	1	-	1	2	2	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
North Carolina												
All races, both sexes	111	3	-	-	-	-	-	-	-	3	3	9
Male	47	1	-	-	-	-	-	-	-	1	1	7
Female	64	2	-	-	-	-	-	-	-	2	2	2
White, both sexes	64	1	-	-	-	-	-	-	-	2	1	-
Male	25	-	-	-	-	-	-	-	-	1	-	-
Female	39	1	-	-	-	-	-	-	-	1	1	-
Black, both sexes	44	2	-	-	-	-	-	-	-	1	1	9
Male	21	1	-	-	-	-	-	-	-	-	-	7
Female	23	1	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Ohio												
All races, both sexes	195	1	-	-	-	1	-	3	3	6	8	17
Male	87	-	-	-	-	1	-	2	2	3	4	8
Female	108	1	-	-	-	-	-	1	1	3	4	9
White, both sexes	160	1	-	-	-	-	-	1	2	5	6	13
Male	73	-	-	-	-	-	-	1	1	2	3	7
Female	87	1	-	-	-	-	-	-	1	3	3	6
Black, both sexes	34	-	-	-	-	1	-	2	1	1	1	4
Male	14	-	-	-	-	1	-	1	1	1	1	1
Female	20	-	-	-	-	-	-	1	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	59	2	-	-	-	-	1	-	2	4	5	6
Male	23	1	-	-	-	-	-	-	1	1	2	2
Female	36	1	-	-	-	-	1	-	1	3	3	4
White, both sexes	48	2	-	-	-	-	1	-	2	2	3	4
Male	19	1	-	-	-	-	-	-	1	1	1	1
Female	29	1	-	-	-	-	1	-	1	1	2	3
Black, both sexes	7	-	-	-	-	-	-	-	-	2	1	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	2	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Oregon												
All races, both sexes	45	-	-	-	-	-	1	1	-	-	2	4
Male	22	-	-	-	-	-	-	1	-	-	1	1
Female	23	-	-	-	-	-	1	-	-	-	1	3
White, both sexes	43	-	-	-	-	-	1	1	-	-	2	2
Male	22	-	-	-	-	-	-	1	-	-	1	1
Female	21	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	162	-	1	-	-	1	3	1	3	1	5	8
Male	71	-	1	-	-	-	3	1	1	1	4	4
Female	91	-	-	-	-	1	-	-	2	-	1	4
White, both sexes	139	-	1	-	-	1	2	-	2	1	4	3
Male	60	-	1	-	-	-	2	-	1	1	4	1
Female	79	-	-	-	-	1	-	-	1	-	-	2
Black, both sexes	22	-	-	-	-	-	1	1	1	-	1	5
Male	11	-	-	-	-	-	1	1	-	-	-	3
Female	11	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Rhode Island												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	52	1	-	-	-	-	1	-	2	-	-	2
Male	20	-	-	-	-	-	-	-	2	-	-	1
Female	32	1	-	-	-	-	1	-	-	-	-	1
White, both sexes	34	1	-	-	-	-	1	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	22	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	2	-	-	1
Male	8	-	-	-	-	-	-	-	2	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
South Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	100	-	-	-	-	-	-	2	5	4	5	4
Male	39	-	-	-	-	-	-	1	3	3	3	-
Female	61	-	-	-	-	-	-	1	2	1	2	4
White, both sexes	87	-	-	-	-	-	-	2	4	4	3	3
Male	32	-	-	-	-	-	-	1	2	3	1	-
Female	55	-	-	-	-	-	-	1	2	1	2	3
Black, both sexes	13	-	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	1	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Texas												
All races, both sexes	224	2	-	-	-	-	2	1	4	4	14	20
Male	102	-	-	-	-	-	1	-	1	2	10	7
Female	122	2	-	-	-	-	1	1	3	2	4	13
White, both sexes	178	2	-	-	-	-	1	1	3	2	9	13
Male	81	-	-	-	-	-	-	-	1	1	7	6
Female	97	2	-	-	-	-	1	1	2	1	2	7
Black, both sexes	44	-	-	-	-	-	1	-	1	2	5	7
Male	21	-	-	-	-	-	1	-	-	1	3	1
Female	23	-	-	-	-	-	-	-	1	1	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Vermont												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	99	4	-	-	-	-	-	-	3	3	3	7
Male	41	2	-	-	-	-	-	-	2	2	1	3
Female	58	2	-	-	-	-	-	-	1	1	2	4
White, both sexes	64	3	-	-	-	-	-	-	1	1	2	5
Male	25	2	-	-	-	-	-	-	1	1	-	2
Female	39	1	-	-	-	-	-	-	-	-	2	3
Black, both sexes	34	1	-	-	-	-	-	-	2	2	1	2
Male	15	-	-	-	-	-	-	-	1	1	1	1
Female	19	1	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Washington												
All races, both sexes	81	1	-	-	-	-	1	-	1	3	2	6
Male	34	1	-	-	-	-	-	-	-	2	1	2
Female	47	-	-	-	-	-	1	-	1	1	1	4
White, both sexes	77	1	-	-	-	-	1	-	1	2	2	6
Male	32	1	-	-	-	-	-	-	-	1	1	2
Female	45	-	-	-	-	-	1	-	1	1	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	38	-	-	-	-	-	1	-	1	1	1	2
Male	17	-	-	-	-	-	-	-	-	1	-	2
Female	21	-	-	-	-	-	1	-	1	-	1	-
White, both sexes	36	-	-	-	-	-	1	-	-	1	1	2
Male	16	-	-	-	-	-	-	-	-	1	-	2
Female	20	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Wisconsin												
All races, both sexes	64	-	-	-	-	1	-	-	2	3	2	6
Male	35	-	-	-	-	1	-	-	-	1	2	3
Female	29	-	-	-	-	-	-	-	2	2	-	3
White, both sexes	58	-	-	-	-	1	-	-	1	2	2	5
Male	34	-	-	-	-	1	-	-	-	1	2	3
Female	24	-	-	-	-	-	-	-	1	1	-	2
Black, both sexes	6	-	-	-	-	-	-	-	1	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
United States												
All races, both sexes	277	285	255	251	302	413	495	489	322	139	26	-
Male	182	170	144	115	133	189	180	157	80	31	6	-
Female	95	115	111	136	169	224	315	332	242	108	20	-
White, both sexes	199	229	190	202	254	360	424	430	290	122	21	-
Male	129	144	106	94	115	167	148	136	77	26	5	-
Female	70	85	84	108	139	193	276	294	213	96	16	-
Black, both sexes	72	53	57	42	46	48	63	50	28	16	5	-
Male	51	25	35	20	18	18	25	16	3	5	1	-
Female	21	28	22	22	28	30	38	34	25	11	4	-
American Indian 1/, both sexes	1	-	3	2	1	-	1	2	2	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	3	2	1	-	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	3	5	5	1	5	7	7	2	1	-	-
Male	1	1	3	1	-	4	7	4	-	-	-	-
Female	4	2	2	4	1	1	-	3	2	1	-	-
Alabama												
All races, both sexes	5	5	4	6	9	11	16	7	3	1	-	-
Male	4	2	2	2	3	6	4	1	-	-	-	-
Female	1	3	2	4	6	5	12	6	3	1	-	-
White, both sexes	2	2	3	4	7	10	16	4	3	1	-	-
Male	1	-	2	2	2	6	4	-	-	-	-	-
Female	1	2	1	2	5	4	12	4	3	1	-	-
Black, both sexes	3	3	1	2	2	1	-	3	-	-	-	-
Male	3	2	-	-	1	-	-	1	-	-	-	-
Female	-	1	1	2	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	7	6	4	7	11	8	7	10	7	1	1	-
Male	3	3	1	4	5	4	4	2	2	-	-	-
Female	4	3	3	3	6	4	3	8	5	1	1	-
White, both sexes	7	6	2	5	10	7	7	10	7	1	1	-
Male	3	3	-	4	4	4	4	2	2	-	-	-
Female	4	3	2	1	6	3	3	8	5	1	1	-
Black, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Arkansas												
All races, both sexes	4	1	2	5	5	5	5	6	5	2	2	-
Male	1	-	1	2	3	4	3	1	1	1	1	-
Female	3	1	1	3	2	1	2	5	4	1	1	-
White, both sexes	2	1	2	5	5	5	5	6	4	2	2	-
Male	-	-	1	2	3	4	3	1	1	1	1	-
Female	2	1	1	3	2	1	2	5	3	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	38	47	25	30	36	41	49	62	35	10	2	-
Male	19	24	10	13	18	18	23	26	11	3	-	-
Female	19	23	15	17	18	23	26	36	24	7	2	-
White, both sexes	30	43	21	23	31	38	44	53	33	9	2	-
Male	14	22	9	12	14	16	20	22	11	3	-	-
Female	16	21	12	11	17	22	24	31	22	6	2	-
Black, both sexes	6	3	3	5	5	1	3	6	1	1	-	-
Male	4	2	1	1	4	-	1	2	-	-	-	-
Female	2	1	2	4	1	1	2	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	2	-	2	2	3	1	-	-	-
Male	1	-	-	-	-	2	2	2	-	-	-	-
Female	1	1	1	2	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Colorado												
All races, both sexes	6	3	5	2	3	6	7	9	4	4	-	-
Male	4	1	3	2	-	2	3	3	3	1	-	-
Female	2	2	2	-	3	4	4	6	1	3	-	-
White, both sexes	6	3	5	2	2	6	6	9	4	4	-	-
Male	4	1	3	2	-	2	2	3	3	1	-	-
Female	2	2	2	-	2	4	4	6	1	3	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	-	3	4	2	1	5	6	4	2	-	-
Male	4	-	2	2	1	1	1	1	1	1	-	-
Female	-	-	1	2	1	-	4	5	3	1	-	-
White, both sexes	3	-	2	3	2	1	4	6	3	2	-	-
Male	3	-	1	1	1	1	1	1	1	1	-	-
Female	-	-	1	2	1	-	3	5	2	1	-	-
Black, both sexes	1	-	1	1	-	-	1	-	1	-	-	-
Male	1	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Delaware												
All races, both sexes	-	3	1	2	-	-	2	1	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	3	-	1	-	-	2	1	1	-	-	-
White, both sexes	-	3	1	2	-	-	2	1	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	3	-	1	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	1	-	1	-	2	1	1	1	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	1	-	2	1	1	1	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Florida												
All races, both sexes	15	28	22	18	20	27	34	25	22	6	-	-
Male	11	16	10	9	9	12	14	12	1	-	-	-
Female	4	12	12	9	11	15	20	13	21	6	-	-
White, both sexes	11	17	15	16	18	20	29	25	20	6	-	-
Male	7	11	6	7	8	9	11	12	-	-	-	-
Female	4	6	9	9	10	11	18	13	20	6	-	-
Black, both sexes	4	11	7	2	2	7	4	-	2	-	-	-
Male	4	5	4	2	1	3	2	-	1	-	-	-
Female	-	6	3	-	1	4	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	11	8	10	9	10	9	14	21	6	5	1	-
Male	6	5	7	5	3	4	4	10	-	1	-	-
Female	5	3	3	4	7	5	10	11	6	4	1	-
White, both sexes	3	5	8	5	8	7	12	15	4	4	1	-
Male	2	4	6	3	2	4	3	7	-	1	-	-
Female	1	1	2	2	6	3	9	8	4	3	1	-
Black, both sexes	8	3	1	4	2	2	2	6	1	1	-	-
Male	4	1	1	2	1	-	1	3	-	-	-	-
Female	4	2	-	2	1	2	1	3	1	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Hawaii												
All races, both sexes	1	-	-	-	1	-	4	3	-	1	-	-
Male	-	-	-	-	-	-	3	1	-	-	-	-
Female	1	-	-	-	1	-	1	2	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	3	1	-	1	-	-
Male	-	-	-	-	-	-	3	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	1	-	-
Idaho												
All races, both sexes	-	2	-	1	2	1	1	-	5	-	-	-
Male	-	1	-	-	1	-	1	-	1	-	-	-
Female	-	1	-	1	1	1	-	-	4	-	-	-
White, both sexes	-	2	-	1	2	1	1	-	5	-	-	-
Male	-	1	-	-	1	-	1	-	1	-	-	-
Female	-	1	-	1	1	1	-	-	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Illinois												
All races, both sexes	9	10	7	6	16	18	28	21	15	5	1	-
Male	7	8	5	1	6	8	9	5	4	2	-	-
Female	2	2	2	5	10	10	19	16	11	3	1	-
White, both sexes	6	7	4	4	14	14	25	17	14	4	1	-
Male	5	5	3	-	6	6	8	4	4	1	-	-
Female	1	2	1	4	8	8	17	13	10	3	1	-
Black, both sexes	2	2	3	2	2	4	3	4	1	1	-	-
Male	2	2	2	1	-	2	1	1	-	1	-	-
Female	-	-	1	1	2	2	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	7	4	5	7	4	6	11	7	5	3	-	-
Male	6	3	3	1	3	2	3	5	-	-	-	-
Female	1	1	2	6	1	4	8	2	5	3	-	-
White, both sexes	6	3	5	6	4	5	11	7	4	3	-	-
Male	6	3	3	1	3	2	3	5	-	-	-	-
Female	-	-	2	5	1	3	8	2	4	3	-	-
Black, both sexes	1	1	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Iowa												
All races, both sexes	1	2	1	2	1	3	2	6	1	1	-	-
Male	1	1	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	2	1	2	2	6	1	1	-	-
White, both sexes	1	2	1	2	1	3	2	6	1	1	-	-
Male	1	1	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	2	1	2	2	6	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	2	1	2	5	2	2	5	7	1	-	-
Male	1	1	-	1	2	2	-	1	3	-	-	-
Female	-	1	1	1	3	-	2	4	4	1	-	-
White, both sexes	1	2	1	1	5	1	2	5	7	1	-	-
Male	1	1	-	-	2	1	-	1	3	-	-	-
Female	-	1	1	1	3	-	2	4	4	1	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Kentucky												
All races, both sexes	2	7	3	6	7	8	14	21	11	7	-	-
Male	2	4	2	3	4	3	5	8	1	-	-	-
Female	-	3	1	3	3	5	9	13	10	7	-	-
White, both sexes	2	6	3	5	7	8	13	21	10	7	-	-
Male	2	4	2	2	4	3	5	8	1	-	-	-
Female	-	2	1	3	3	5	8	13	9	7	-	-
Black, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	3	5	3	7	7	8	7	2	2	-	-
Male	1	1	4	2	3	-	4	2	-	1	-	-
Female	-	2	1	1	4	7	4	5	2	1	-	-
White, both sexes	1	1	4	2	3	5	7	5	1	-	-	-
Male	1	1	3	2	1	-	3	1	-	-	-	-
Female	-	-	1	-	2	5	4	4	1	-	-	-
Black, both sexes	-	2	1	1	4	2	1	2	1	2	-	-
Male	-	-	1	-	2	-	1	1	-	1	-	-
Female	-	2	-	1	2	2	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Maine												
All races, both sexes	1	1	-	-	3	2	2	5	3	-	-	-
Male	1	1	-	-	2	1	1	2	-	-	-	-
Female	-	-	-	-	1	1	1	3	3	-	-	-
White, both sexes	1	1	-	-	3	2	2	5	3	-	-	-
Male	1	1	-	-	2	1	1	2	-	-	-	-
Female	-	-	-	-	1	1	1	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	5	5	3	3	9	5	4	4	1	-	-
Male	2	3	5	1	1	5	1	2	1	-	-	-
Female	3	2	-	2	2	4	4	2	3	1	-	-
White, both sexes	4	4	2	3	2	8	4	1	-	-	-	-
Male	2	2	2	1	1	4	-	1	-	-	-	-
Female	2	2	-	2	1	4	4	-	-	-	-	-
Black, both sexes	1	1	3	-	1	1	1	3	4	1	-	-
Male	-	1	3	-	-	1	1	1	1	-	-	-
Female	1	-	-	-	1	-	-	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Massachusetts												
All races, both sexes	6	5	9	9	4	7	15	10	7	5	-	-
Male	2	3	6	3	1	3	5	4	1	1	-	-
Female	4	2	3	6	3	4	10	6	6	4	-	-
White, both sexes	6	4	6	9	3	7	15	9	7	5	-	-
Male	2	2	3	3	1	3	5	3	1	1	-	-
Female	4	2	3	6	2	4	10	6	6	4	-	-
Black, both sexes	-	1	3	-	1	-	-	-	-	-	-	-
Male	-	1	3	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	14	21	10	9	7	18	18	11	15	4	1	-
Male	12	15	5	5	4	8	8	4	6	-	-	-
Female	2	6	5	4	3	10	10	7	9	4	1	-
White, both sexes	12	16	5	8	6	15	14	11	14	4	1	-
Male	12	11	3	4	4	7	6	4	6	-	-	-
Female	-	5	2	4	2	8	8	7	8	4	1	-
Black, both sexes	2	5	5	1	1	1	4	-	1	-	-	-
Male	-	4	2	1	-	-	2	-	-	-	-	-
Female	2	1	3	-	1	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Minnesota												
All races, both sexes	2	1	1	2	-	11	5	5	3	3	2	-
Male	1	1	1	-	-	6	2	-	-	1	1	-
Female	1	-	-	2	-	5	3	5	3	2	1	-
White, both sexes	2	1	1	2	-	11	5	5	3	3	1	-
Male	1	1	1	-	-	6	2	-	-	1	1	-
Female	1	-	-	2	-	5	3	5	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	4	3	4	4	6	6	3	2	3	1	-
Male	2	2	2	1	1	2	-	1	-	1	-	-
Female	2	2	1	3	3	4	6	2	2	2	1	-
White, both sexes	2	2	1	2	2	6	4	1	2	2	1	-
Male	1	1	-	-	1	2	-	-	-	1	-	-
Female	1	1	1	2	1	4	4	1	2	1	1	-
Black, both sexes	2	2	2	2	2	-	2	2	-	1	-	-
Male	1	1	2	1	-	-	-	1	-	-	-	-
Female	1	1	-	1	2	-	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Missouri												
All races, both sexes	7	1	6	5	6	8	15	8	10	3	-	-
Male	4	-	1	5	1	3	9	3	4	-	-	-
Female	3	1	5	-	5	5	6	5	6	3	-	-
White, both sexes	5	1	5	5	6	8	11	8	10	3	-	-
Male	2	-	1	5	1	3	7	3	4	-	-	-
Female	3	1	4	-	5	5	4	5	6	3	-	-
Black, both sexes	2	-	1	-	-	-	4	-	-	-	-	-
Male	2	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	1	-	1	1	-	1	1	3	2	-	-
Male	1	1	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	1	1	-	1	1	2	2	-	-
White, both sexes	2	1	-	1	1	-	1	1	3	2	-	-
Male	1	1	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	1	1	-	1	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Nebraska												
All races, both sexes	2	1	1	2	1	4	8	3	5	1	-	-
Male	2	1	-	-	1	1	2	-	1	-	-	-
Female	-	-	1	2	-	3	6	3	4	1	-	-
White, both sexes	1	1	1	2	1	4	8	3	5	1	-	-
Male	1	1	-	-	1	1	2	-	1	-	-	-
Female	-	-	1	2	-	3	6	3	4	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	3	2	3	4	-	4	5	1	-	-	-
Male	3	3	1	2	3	-	3	2	-	-	-	-
Female	2	-	1	1	1	-	1	3	1	-	-	-
White, both sexes	3	2	2	3	4	-	4	5	1	-	-	-
Male	2	2	1	2	3	-	3	2	-	-	-	-
Female	1	-	1	1	1	-	1	3	1	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
New Hampshire												
All races, both sexes	2	1	2	-	1	4	1	2	2	2	-	-
Male	2	1	1	-	-	3	-	1	-	1	-	-
Female	-	-	1	-	1	1	1	1	2	1	-	-
White, both sexes	2	1	2	-	1	4	1	2	2	2	-	-
Male	2	1	1	-	-	3	-	1	-	1	-	-
Female	-	-	1	-	1	1	1	1	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	10	9	4	11	15	19	19	9	2	2	-
Male	9	6	5	1	4	8	2	8	2	1	1	-
Female	2	4	4	3	7	7	17	11	7	1	1	-
White, both sexes	6	10	5	4	9	13	15	18	8	2	2	-
Male	5	6	4	1	4	6	2	8	2	1	1	-
Female	1	4	1	3	5	7	13	10	6	1	1	-
Black, both sexes	5	-	4	-	2	1	4	-	1	-	-	-
Male	4	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	3	-	2	-	4	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
New Mexico												
All races, both sexes	3	3	3	-	-	1	3	4	3	-	-	-
Male	1	2	2	-	-	-	1	2	-	-	-	-
Female	2	1	1	-	-	1	2	2	3	-	-	-
White, both sexes	3	3	3	-	-	1	2	3	2	-	-	-
Male	1	2	2	-	-	-	1	1	-	-	-	-
Female	2	1	1	-	-	1	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	26	22	17	20	21	28	26	28	18	4	5	-
Male	20	15	12	7	13	14	6	4	3	2	-	-
Female	6	7	5	13	8	14	20	24	15	2	5	-
White, both sexes	16	18	11	9	16	21	20	25	17	4	4	-
Male	14	15	8	2	10	13	3	3	3	2	-	-
Female	2	3	3	7	6	8	17	22	14	2	4	-
Black, both sexes	10	4	5	11	5	7	6	3	1	-	1	-
Male	6	-	3	5	3	1	3	1	-	-	-	-
Female	4	4	2	6	2	6	3	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
North Carolina												
All races, both sexes	11	8	8	7	16	11	9	13	5	5	-	-
Male	8	3	3	4	5	4	4	3	2	1	-	-
Female	3	5	5	3	11	7	5	10	3	4	-	-
White, both sexes	3	4	4	6	12	6	6	10	5	4	-	-
Male	2	2	2	3	4	2	3	3	2	1	-	-
Female	1	2	2	3	8	4	3	7	3	3	-	-
Black, both sexes	8	4	3	1	3	5	3	3	-	1	-	-
Male	6	1	1	1	1	2	1	-	-	-	-	-
Female	2	3	2	-	2	3	2	3	-	1	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Ohio												
All races, both sexes	10	7	14	13	16	30	19	28	12	6	1	-
Male	8	3	8	7	8	13	7	8	2	3	-	-
Female	2	4	6	6	8	17	12	20	10	3	1	-
White, both sexes	8	6	10	11	13	28	17	23	12	4	-	-
Male	7	3	6	6	8	12	5	7	2	3	-	-
Female	1	3	4	5	5	16	12	16	10	1	-	-
Black, both sexes	2	1	4	2	3	2	2	5	-	2	1	-
Male	1	-	2	1	-	1	2	1	-	-	-	-
Female	1	1	2	1	3	1	-	4	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	2	5	6	2	5	5	4	4	2	-	-
Male	2	2	2	3	-	2	2	-	2	1	-	-
Female	2	-	3	3	2	3	3	4	2	1	-	-
White, both sexes	4	2	4	5	2	5	5	3	3	1	-	-
Male	2	2	2	2	-	2	2	-	2	-	-	-
Female	2	-	2	3	2	3	3	3	1	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Oregon												
All races, both sexes	1	-	5	2	6	9	1	6	5	2	-	-
Male	1	-	2	1	2	6	-	3	3	1	-	-
Female	-	-	3	1	4	3	1	3	2	1	-	-
White, both sexes	1	-	5	2	6	9	1	6	5	2	-	-
Male	1	-	2	1	2	6	-	3	3	1	-	-
Female	-	-	3	1	4	3	1	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	7	13	5	7	10	17	27	28	15	9	1	-
Male	5	9	3	4	3	5	11	9	4	2	1	-
Female	2	4	2	3	7	12	16	19	11	7	-	-
White, both sexes	6	11	5	7	7	15	25	26	13	9	1	-
Male	4	8	3	4	2	4	9	9	4	2	1	-
Female	2	3	2	3	5	11	16	17	9	7	-	-
Black, both sexes	1	1	-	-	3	2	2	2	2	-	-	-
Male	1	1	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	2	1	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Rhode Island												
All races, both sexes	2	-	-	-	1	1	2	4	-	1	-	-
Male	2	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	1	1	-	3	-	1	-	-
White, both sexes	-	-	-	-	1	1	2	4	-	1	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	1	1	-	3	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	5	5	3	2	5	4	10	7	4	-	-
Male	1	1	5	3	1	3	-	1	-	2	-	-
Female	-	4	-	-	1	2	4	9	7	2	-	-
White, both sexes	-	3	4	3	1	3	1	9	5	2	-	-
Male	-	1	4	3	1	1	-	1	-	-	-	-
Female	-	2	-	-	-	2	1	8	5	2	-	-
Black, both sexes	1	2	1	-	1	2	3	1	2	2	-	-
Male	1	-	1	-	-	2	-	-	-	2	-	-
Female	-	2	-	-	1	-	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
South Dakota												
All races, both sexes	-	-	1	-	1	1	1	2	3	1	1	-
Male	-	-	1	-	-	1	1	-	3	-	-	-
Female	-	-	-	-	1	-	-	2	-	1	1	-
White, both sexes	-	-	1	-	1	1	1	2	3	1	1	-
Male	-	-	1	-	-	1	1	-	3	-	-	-
Female	-	-	-	-	1	-	-	2	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	7	7	8	3	6	11	14	10	8	4	2	-
Male	1	5	4	1	2	6	6	1	2	-	1	-
Female	6	2	4	2	4	5	8	9	6	4	1	-
White, both sexes	6	7	5	3	6	11	13	8	7	3	2	-
Male	1	5	1	1	2	6	5	1	2	-	1	-
Female	5	2	4	2	4	5	8	7	5	3	1	-
Black, both sexes	1	-	3	-	-	-	1	2	1	1	-	-
Male	-	-	3	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Texas												
All races, both sexes	16	12	12	14	11	24	30	25	20	12	1	-
Male	13	8	8	6	6	12	12	7	8	1	-	-
Female	3	4	4	8	5	12	18	18	12	11	1	-
White, both sexes	11	11	8	12	10	21	25	20	18	11	-	-
Male	8	7	6	5	5	11	11	5	7	1	-	-
Female	3	4	2	7	5	10	14	15	11	10	-	-
Black, both sexes	5	1	4	1	1	3	5	4	2	1	1	-
Male	5	1	2	1	1	1	1	2	1	-	-	-
Female	-	-	2	-	-	2	4	2	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	1	-	2	3	-	1	4	1	-	1	-	-
Male	1	-	2	1	-	1	4	-	-	1	-	-
Female	-	-	-	2	-	-	-	1	-	-	-	-
White, both sexes	1	-	2	3	-	1	4	1	-	1	-	-
Male	1	-	2	1	-	1	4	-	-	1	-	-
Female	-	-	-	2	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Vermont												
All races, both sexes	-	-	1	2	1	2	1	-	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	2	1	2	1	-	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	7	6	3	7	13	16	13	7	3	1	-
Male	2	4	5	1	4	3	5	6	-	-	1	-
Female	1	3	1	2	3	10	11	7	7	3	-	-
White, both sexes	3	4	4	1	4	10	9	10	4	3	-	-
Male	2	3	3	-	2	2	3	4	-	-	-	-
Female	1	1	1	1	2	8	6	6	4	3	-	-
Black, both sexes	-	3	1	2	3	3	7	3	3	-	1	-
Male	-	1	1	1	2	1	2	2	-	-	1	-
Female	-	2	-	1	1	2	5	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Washington												
All races, both sexes	9	8	7	6	3	2	11	8	10	2	1	-
Male	4	7	3	4	1	-	2	1	5	1	-	-
Female	5	1	4	2	2	2	9	7	5	1	1	-
White, both sexes	9	8	7	4	3	2	10	8	10	2	1	-
Male	4	7	3	4	1	-	1	1	5	1	-	-
Female	5	1	4	-	2	2	9	7	5	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	3	5	2	5	5	6	1	2	2	-	-
Male	-	2	2	1	3	4	2	-	-	-	-	-
Female	1	1	3	1	2	1	4	1	2	2	-	-
White, both sexes	1	2	5	2	5	5	6	1	2	2	-	-
Male	-	1	2	1	3	4	2	-	-	-	-	-
Female	1	1	3	1	2	1	4	1	2	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Wisconsin												
All races, both sexes	-	3	3	7	7	8	6	8	5	3	-	-
Male	-	2	2	5	4	7	-	5	2	1	-	-
Female	-	1	1	2	3	1	6	3	3	2	-	-
White, both sexes	-	3	3	6	7	7	5	8	5	3	-	-
Male	-	2	2	5	4	6	-	5	2	1	-	-
Female	-	1	1	1	3	1	5	3	3	2	-	-
Black, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
United States												
All races, both sexes	2,849	6	1	1	2	11	32	33	73	107	136	206
Male	1,225	2	-	1	-	7	12	15	35	58	66	103
Female	1,624	4	1	-	2	4	20	18	38	49	70	103
White, both sexes	2,245	5	1	-	1	6	19	19	48	71	79	139
Male	954	2	-	-	-	3	6	7	24	42	40	75
Female	1,291	3	1	-	1	3	13	12	24	29	39	64
Black, both sexes	564	1	-	1	1	5	13	14	25	35	54	64
Male	253	-	-	1	-	4	6	8	11	15	24	27
Female	311	1	-	-	1	1	7	6	14	20	30	37
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	3
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	1	2	-
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	2	1
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	35	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	66	-	-	-	-	2	1	1	4	-	1	2
Male	27	-	-	-	-	1	-	1	2	-	1	1
Female	39	-	-	-	-	1	1	-	2	-	-	1
White, both sexes	60	-	-	-	-	2	1	1	4	-	1	2
Male	25	-	-	-	-	1	-	1	2	-	1	1
Female	35	-	-	-	-	1	1	-	2	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Arkansas												
All races, both sexes	33	-	-	-	-	-	1	1	-	-	1	4
Male	13	-	-	-	-	-	1	-	-	-	1	1
Female	20	-	-	-	-	-	-	1	-	-	-	3
White, both sexes	26	-	-	-	-	-	-	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	1	1	-	-	-	2
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	368	1	-	-	-	-	3	3	7	14	15	28
Male	175	1	-	-	-	-	1	-	6	9	9	17
Female	193	-	-	-	-	-	2	3	1	5	6	11
White, both sexes	305	1	-	-	-	-	3	3	5	10	7	22
Male	143	1	-	-	-	-	1	-	5	6	4	14
Female	162	-	-	-	-	-	2	3	-	4	3	8
Black, both sexes	49	-	-	-	-	-	-	-	2	4	8	5
Male	24	-	-	-	-	-	-	-	1	3	5	2
Female	25	-	-	-	-	-	-	-	1	1	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Colorado												
All races, both sexes	45	-	-	-	-	1	-	-	1	4	-	3
Male	23	-	-	-	-	-	-	-	1	3	-	3
Female	22	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	44	-	-	-	-	1	-	-	1	4	-	3
Male	23	-	-	-	-	-	-	-	1	3	-	3
Female	21	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Delaware												
All races, both sexes	10	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	15	-	-	-	-	-	-	-	1	3	2	2
Male	7	-	-	-	-	-	-	-	1	-	2	-
Female	8	-	-	-	-	-	-	-	-	3	-	2
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	1	3	2	2
Male	7	-	-	-	-	-	-	-	1	-	2	-
Female	8	-	-	-	-	-	-	-	-	3	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Florida												
All races, both sexes	203	1	-	-	-	-	-	3	2	5	7	15
Male	89	-	-	-	-	-	-	2	1	4	3	7
Female	114	1	-	-	-	-	-	1	1	1	4	8
White, both sexes	153	1	-	-	-	-	-	1	2	3	3	9
Male	59	-	-	-	-	-	-	1	1	3	1	2
Female	94	1	-	-	-	-	-	-	1	-	2	7
Black, both sexes	49	-	-	-	-	-	-	2	-	2	4	6
Male	29	-	-	-	-	-	-	1	-	1	2	5
Female	20	-	-	-	-	-	-	1	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	86	-	-	-	-	2	2	1	3	2	3	5
Male	34	-	-	-	-	1	1	1	-	2	-	1
Female	52	-	-	-	-	1	1	-	3	-	3	4
White, both sexes	51	-	-	-	-	-	1	-	1	2	1	2
Male	21	-	-	-	-	-	1	-	-	2	-	-
Female	30	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	34	-	-	-	-	2	1	1	2	-	2	3
Male	13	-	-	-	-	1	-	1	-	-	-	1
Female	21	-	-	-	-	1	1	-	2	-	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Hawaii												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	9	-	-	1	-	-	-	-	-	-	-	2
Male	5	-	-	1	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Illinois												
All races, both sexes	114	-	-	-	-	-	4	7	2	2	8	5
Male	53	-	-	-	-	-	2	4	-	2	3	3
Female	61	-	-	-	-	-	2	3	2	-	5	2
White, both sexes	81	-	-	-	-	-	2	3	-	1	3	4
Male	36	-	-	-	-	-	1	-	-	1	1	2
Female	45	-	-	-	-	-	1	3	-	-	2	2
Black, both sexes	32	-	-	-	-	-	2	4	2	1	5	1
Male	17	-	-	-	-	-	1	4	-	1	2	1
Female	15	-	-	-	-	-	1	-	2	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	46	-	-	-	-	-	-	-	-	1	3	5
Male	21	-	-	-	-	-	-	-	-	-	1	4
Female	25	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	44	-	-	-	-	-	-	-	-	1	3	5
Male	21	-	-	-	-	-	-	-	-	-	1	4
Female	23	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Iowa												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	4	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	4	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	23	-	-	-	-	-	-	-	2	1	1	2
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	21	-	-	-	-	-	-	-	2	1	1	2
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	2	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Kentucky												
All races, both sexes	40	-	-	-	-	-	1	1	1	2	3	-
Male	16	-	-	-	-	-	1	-	-	1	2	-
Female	24	-	-	-	-	-	-	1	1	1	1	-
White, both sexes	36	-	-	-	-	-	1	1	1	2	1	-
Male	15	-	-	-	-	-	1	-	-	1	1	-
Female	21	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	34	-	-	-	-	1	-	-	1	1	3	1
Male	11	-	-	-	-	1	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	1	1	2	1
White, both sexes	20	-	-	-	-	-	-	-	1	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	14	-	-	-	-	1	-	-	-	-	2	1
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Maine												
All races, both sexes	14	-	-	-	-	-	-	1	1	2	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	2
Female	5	-	-	-	-	-	-	1	-	2	-	-
White, both sexes	14	-	-	-	-	-	-	1	1	2	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	2
Female	5	-	-	-	-	-	-	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	50	-	-	-	-	-	-	2	2	7	6	3
Male	19	-	-	-	-	-	-	1	2	2	1	1
Female	31	-	-	-	-	-	-	1	-	5	5	2
White, both sexes	29	-	-	-	-	-	-	1	-	4	1	1
Male	13	-	-	-	-	-	-	1	-	2	-	1
Female	16	-	-	-	-	-	-	-	-	2	1	-
Black, both sexes	21	-	-	-	-	-	-	1	2	3	5	2
Male	6	-	-	-	-	-	-	-	2	-	1	-
Female	15	-	-	-	-	-	-	1	-	3	4	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Massachusetts												
All races, both sexes	57	-	-	-	1	-	-	-	3	1	2	4
Male	23	-	-	-	-	-	-	-	3	1	-	4
Female	34	-	-	-	1	-	-	-	-	-	2	-
White, both sexes	54	-	-	-	1	-	-	-	3	1	2	4
Male	20	-	-	-	-	-	-	-	3	1	-	4
Female	34	-	-	-	1	-	-	-	-	-	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	142	-	1	-	-	-	2	-	4	11	13	15
Male	74	-	-	-	-	-	-	-	1	7	6	5
Female	68	-	1	-	-	-	2	-	3	4	7	10
White, both sexes	111	-	1	-	-	-	-	-	3	9	12	7
Male	61	-	-	-	-	-	-	-	1	6	5	2
Female	50	-	1	-	-	-	-	-	2	3	7	5
Black, both sexes	30	-	-	-	-	-	2	-	1	2	1	8
Male	12	-	-	-	-	-	-	-	-	1	1	3
Female	18	-	-	-	-	-	2	-	1	1	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Minnesota												
All races, both sexes	25	-	-	-	-	-	1	1	2	2	1	-
Male	10	-	-	-	-	-	-	-	1	2	1	-
Female	15	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	24	-	-	-	-	-	1	1	2	2	1	-
Male	10	-	-	-	-	-	-	-	1	2	1	-
Female	14	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	32	-	-	-	-	-	1	-	-	3	1	3
Male	8	-	-	-	-	-	1	-	-	1	-	2
Female	24	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	17	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	1	-	-	2	1	3
Male	5	-	-	-	-	-	1	-	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	2	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Missouri												
All races, both sexes	49	-	-	-	-	1	2	-	2	5	-	2
Male	17	-	-	-	-	1	-	-	1	2	-	-
Female	32	-	-	-	-	-	2	-	1	3	-	2
White, both sexes	40	-	-	-	-	1	-	-	2	3	-	1
Male	13	-	-	-	-	1	-	-	1	1	-	-
Female	27	-	-	-	-	-	-	-	1	2	-	1
Black, both sexes	9	-	-	-	-	-	2	-	-	2	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	2	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Nebraska												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	20	-	-	-	-	-	-	1	-	-	-	2
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	19	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Hampshire												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	102	-	-	-	-	-	1	-	3	6	3	8
Male	40	-	-	-	-	-	-	-	1	3	-	5
Female	62	-	-	-	-	-	1	-	2	3	3	3
White, both sexes	76	-	-	-	-	-	1	-	-	3	-	4
Male	31	-	-	-	-	-	-	-	-	2	-	3
Female	45	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	26	-	-	-	-	-	-	-	3	3	3	4
Male	9	-	-	-	-	-	-	-	1	1	-	2
Female	17	-	-	-	-	-	-	-	2	2	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Mexico												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	212	-	-	-	1	1	3	3	7	7	12	14
Male	102	-	-	-	-	1	1	-	3	4	6	12
Female	110	-	-	-	1	-	2	3	4	3	6	2
White, both sexes	148	-	-	-	-	-	1	1	5	2	6	13
Male	73	-	-	-	-	-	-	-	2	-	3	11
Female	75	-	-	-	-	-	1	1	3	2	3	2
Black, both sexes	64	-	-	-	1	1	2	2	2	5	6	1
Male	29	-	-	-	-	1	1	-	1	4	3	1
Female	35	-	-	-	1	-	1	2	1	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
North Carolina												
All races, both sexes	74	2	-	-	-	-	-	-	-	3	2	5
Male	30	-	-	-	-	-	-	-	-	1	1	5
Female	44	2	-	-	-	-	-	-	-	2	1	-
White, both sexes	42	1	-	-	-	-	-	-	-	2	-	-
Male	15	-	-	-	-	-	-	-	-	1	-	-
Female	27	1	-	-	-	-	-	-	-	1	-	-
Black, both sexes	29	1	-	-	-	-	-	-	-	1	1	5
Male	14	-	-	-	-	-	-	-	-	-	-	5
Female	15	1	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Ohio												
All races, both sexes	149	-	-	-	-	1	-	3	3	4	8	13
Male	62	-	-	-	-	1	-	2	2	2	4	5
Female	87	-	-	-	-	-	-	1	1	2	4	8
White, both sexes	117	-	-	-	-	-	-	1	2	3	6	9
Male	48	-	-	-	-	-	-	1	1	1	3	4
Female	69	-	-	-	-	-	-	-	1	2	3	5
Black, both sexes	31	-	-	-	-	1	-	2	1	1	1	4
Male	14	-	-	-	-	1	-	1	1	1	1	1
Female	17	-	-	-	-	-	-	1	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	44	-	-	-	-	-	1	-	2	4	4	6
Male	15	-	-	-	-	-	-	-	1	1	1	2
Female	29	-	-	-	-	-	1	-	1	3	3	4
White, both sexes	34	-	-	-	-	-	1	-	2	2	3	4
Male	12	-	-	-	-	-	-	-	1	1	1	1
Female	22	-	-	-	-	-	1	-	1	1	2	3
Black, both sexes	7	-	-	-	-	-	-	-	-	2	1	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	2	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Oregon												
All races, both sexes	33	-	-	-	-	-	1	1	-	-	2	3
Male	14	-	-	-	-	-	-	1	-	-	1	1
Female	19	-	-	-	-	-	1	-	-	-	1	2
White, both sexes	32	-	-	-	-	-	1	1	-	-	2	2
Male	14	-	-	-	-	-	-	1	-	-	1	1
Female	18	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	107	-	-	-	-	-	3	1	2	-	4	4
Male	43	-	-	-	-	-	3	1	-	-	3	1
Female	64	-	-	-	-	-	-	-	2	-	1	3
White, both sexes	94	-	-	-	-	-	2	-	1	-	3	2
Male	37	-	-	-	-	-	2	-	-	-	3	-
Female	57	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	13	-	-	-	-	-	1	1	1	-	1	2
Male	6	-	-	-	-	-	1	1	-	-	-	1
Female	7	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	36	-	-	-	-	-	1	-	2	-	-	2
Male	12	-	-	-	-	-	-	-	2	-	-	1
Female	24	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	1	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	2	-	-	1
Male	4	-	-	-	-	-	-	-	2	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	76	-	-	-	-	-	-	2	5	4	4	4
Male	26	-	-	-	-	-	-	1	3	3	2	-
Female	50	-	-	-	-	-	-	1	2	1	2	4
White, both sexes	64	-	-	-	-	-	-	2	4	4	2	3
Male	19	-	-	-	-	-	-	1	2	3	-	-
Female	45	-	-	-	-	-	-	1	2	1	2	3
Black, both sexes	12	-	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	1	-	2	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Texas												
All races, both sexes	152	-	-	-	-	-	2	1	4	3	12	16
Male	70	-	-	-	-	-	1	-	1	1	9	6
Female	82	-	-	-	-	-	1	1	3	2	3	10
White, both sexes	120	-	-	-	-	-	1	1	3	1	7	9
Male	54	-	-	-	-	-	-	-	1	-	6	5
Female	66	-	-	-	-	-	1	1	2	1	1	4
Black, both sexes	32	-	-	-	-	-	1	-	1	2	5	7
Male	16	-	-	-	-	-	1	-	-	1	3	1
Female	16	-	-	-	-	-	-	-	1	1	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	72	2	-	-	-	-	-	-	3	3	3	5
Male	24	1	-	-	-	-	-	-	2	2	1	1
Female	48	1	-	-	-	-	-	-	1	1	2	4
White, both sexes	48	2	-	-	-	-	-	-	1	1	2	3
Male	13	1	-	-	-	-	-	-	1	1	-	-
Female	35	1	-	-	-	-	-	-	-	-	2	3
Black, both sexes	24	-	-	-	-	-	-	-	2	2	1	2
Male	11	-	-	-	-	-	-	-	1	1	1	1
Female	13	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Washington												
All races, both sexes	62	-	-	-	-	-	1	-	1	3	1	6
Male	25	-	-	-	-	-	-	-	-	2	1	2
Female	37	-	-	-	-	-	1	-	1	1	-	4
White, both sexes	60	-	-	-	-	-	1	-	1	2	1	6
Male	24	-	-	-	-	-	-	-	-	1	1	2
Female	36	-	-	-	-	-	1	-	1	1	-	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	24	-	-	-	-	-	1	-	1	1	1	2
Male	11	-	-	-	-	-	-	-	-	1	-	2
Female	13	-	-	-	-	-	1	-	1	-	1	-
White, both sexes	22	-	-	-	-	-	1	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	1	-	2
Female	12	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Wisconsin												
All races, both sexes	36	-	-	-	-	1	-	-	2	2	2	3
Male	18	-	-	-	-	1	-	-	-	1	2	1
Female	18	-	-	-	-	-	-	-	2	1	-	2
White, both sexes	32	-	-	-	-	1	-	-	1	2	2	2
Male	18	-	-	-	-	1	-	-	-	1	2	1
Female	14	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
United States												
All races, both sexes	204	204	183	178	207	286	345	332	201	88	13	-
Male	131	125	95	75	87	121	124	103	44	18	3	-
Female	73	79	88	103	120	165	221	229	157	70	10	-
White, both sexes	138	168	136	138	174	250	291	289	182	80	11	-
Male	87	111	68	58	75	108	98	88	42	17	3	-
Female	51	57	68	80	99	142	193	201	140	63	8	-
Black, both sexes	62	36	41	36	31	33	49	37	16	8	2	-
Male	42	14	26	17	12	10	22	11	2	1	-	-
Female	20	22	15	19	19	23	27	26	14	7	2	-
American Indian 1/, both sexes	1	-	3	1	1	-	1	2	2	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	3	1	1	-	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	3	3	1	3	4	4	1	-	-	-
Male	1	-	1	-	-	3	4	3	-	-	-	-
Female	2	-	2	3	1	-	-	1	1	-	-	-
Alabama												
All races, both sexes	2	4	4	5	6	8	10	3	1	-	-	-
Male	2	1	2	2	2	5	2	1	-	-	-	-
Female	-	3	2	3	4	3	8	2	1	-	-	-
White, both sexes	-	2	3	4	5	7	10	1	1	-	-	-
Male	-	-	2	2	2	5	2	-	-	-	-	-
Female	-	2	1	2	3	2	8	1	1	-	-	-
Black, both sexes	2	2	1	1	1	1	-	2	-	-	-	-
Male	2	1	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	1	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	5	4	6	10	7	6	7	3	1	1	-
Male	2	2	1	3	5	4	3	1	-	-	-	-
Female	3	3	3	3	5	3	3	6	3	1	1	-
White, both sexes	5	5	2	4	9	6	6	7	3	1	1	-
Male	2	2	-	3	4	4	3	1	-	-	-	-
Female	3	3	2	1	5	2	3	6	3	1	1	-
Black, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Arkansas												
All races, both sexes	2	-	-	2	4	3	3	5	4	2	1	-
Male	1	-	-	1	2	2	1	1	-	1	1	-
Female	1	-	-	1	2	1	2	4	4	1	-	-
White, both sexes	-	-	-	2	4	3	3	5	3	2	1	-
Male	-	-	-	1	2	2	1	1	-	1	1	-
Female	-	-	-	1	2	1	2	4	3	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	32	38	23	24	24	33	35	52	26	9	1	-
Male	16	23	10	10	10	12	18	23	7	3	-	-
Female	16	15	13	14	14	21	17	29	19	6	1	-
White, both sexes	24	35	20	19	20	30	30	43	24	8	1	-
Male	11	21	9	10	7	10	15	19	7	3	-	-
Female	13	14	11	9	13	20	15	24	17	5	1	-
Black, both sexes	6	3	2	3	4	1	3	6	1	1	-	-
Male	4	2	1	-	3	-	1	2	-	-	-	-
Female	2	1	1	3	1	1	2	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	2	-	2	2	3	1	-	-	-
Male	1	-	-	-	-	2	2	2	-	-	-	-
Female	1	-	1	2	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Colorado												
All races, both sexes	4	3	4	2	3	4	5	4	4	3	-	-
Male	2	1	2	2	-	2	1	2	3	1	-	-
Female	2	2	2	-	3	2	4	2	1	2	-	-
White, both sexes	4	3	4	2	2	4	5	4	4	3	-	-
Male	2	1	2	2	-	2	1	2	3	1	-	-
Female	2	2	2	-	2	2	4	2	1	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	-	3	2	1	-	5	2	3	2	-	-
Male	2	-	2	1	1	-	1	-	-	1	-	-
Female	-	-	1	1	-	-	4	2	3	1	-	-
White, both sexes	2	-	2	1	1	-	4	2	2	2	-	-
Male	2	-	1	-	1	-	1	-	-	1	-	-
Female	-	-	1	1	-	-	3	2	2	1	-	-
Black, both sexes	-	-	1	1	-	-	1	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Delaware												
All races, both sexes	-	3	1	2	-	-	2	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	3	-	1	-	-	2	-	1	-	-	-
White, both sexes	-	3	1	2	-	-	2	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	3	-	1	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	1	-	1	-	2	-	-	1	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	1	-	2	-	-	1	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Florida												
All races, both sexes	11	22	19	14	16	21	27	17	20	3	-	-
Male	7	13	9	7	7	8	12	8	1	-	-	-
Female	4	9	10	7	9	13	15	9	19	3	-	-
White, both sexes	7	12	12	12	15	15	23	17	18	3	-	-
Male	3	9	5	5	6	6	9	8	-	-	-	-
Female	4	3	7	7	9	9	14	9	18	3	-	-
Black, both sexes	4	10	7	2	1	6	3	-	2	-	-	-
Male	4	4	4	2	1	2	2	-	1	-	-	-
Female	-	6	3	-	-	4	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	10	3	5	7	7	7	11	11	4	3	-	-
Male	5	1	3	4	2	4	4	5	-	-	-	-
Female	5	2	2	3	5	3	7	6	4	3	-	-
White, both sexes	2	1	3	3	6	5	9	8	4	3	-	-
Male	1	1	2	2	2	4	3	3	-	-	-	-
Female	1	-	1	1	4	1	6	5	4	3	-	-
Black, both sexes	8	2	1	4	1	2	2	3	-	-	-	-
Male	4	-	1	2	-	-	1	2	-	-	-	-
Female	4	2	-	2	1	2	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Hawaii												
All races, both sexes	-	-	-	-	1	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	1	-	-	2	1	1	-	1	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
White, both sexes	-	1	-	-	2	1	1	-	1	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Illinois												
All races, both sexes	8	7	6	2	11	14	16	14	7	1	-	-
Male	7	7	4	-	4	7	4	3	2	1	-	-
Female	1	-	2	2	7	7	12	11	5	-	-	-
White, both sexes	5	5	3	2	10	11	14	11	6	1	-	-
Male	5	5	2	-	4	5	4	3	2	1	-	-
Female	-	-	1	2	6	6	10	8	4	-	-	-
Black, both sexes	2	2	3	-	1	3	2	3	1	-	-	-
Male	2	2	2	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	1	1	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	5	2	2	5	3	3	8	4	2	3	-	-
Male	5	2	1	-	2	1	2	3	-	-	-	-
Female	-	-	1	5	1	2	6	1	2	3	-	-
White, both sexes	5	2	2	4	3	3	8	4	1	3	-	-
Male	5	2	1	-	2	1	2	3	-	-	-	-
Female	-	-	1	4	1	2	6	1	1	3	-	-
Black, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Iowa												
All races, both sexes	-	2	1	2	-	1	1	5	1	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	-	1	1	5	1	1	-	-
White, both sexes	-	2	1	2	-	1	1	5	1	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	-	1	1	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	2	3	2	1	2	6	-	-	-
Male	-	-	-	1	2	2	-	1	2	-	-	-
Female	-	1	-	1	1	-	1	1	4	-	-	-
White, both sexes	-	1	-	1	3	1	1	2	6	-	-	-
Male	-	-	-	-	2	1	-	1	2	-	-	-
Female	-	1	-	1	1	-	1	1	4	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Kentucky												
All races, both sexes	2	3	-	2	3	3	7	7	2	3	-	-
Male	2	2	-	-	1	1	3	2	1	-	-	-
Female	-	1	-	2	2	2	4	5	1	3	-	-
White, both sexes	2	2	-	2	3	3	6	7	2	3	-	-
Male	2	2	-	-	1	1	3	2	1	-	-	-
Female	-	-	-	2	2	2	3	5	1	3	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	2	2	1	4	6	6	5	-	1	-	-
Male	-	1	1	-	3	-	3	1	-	-	-	-
Female	-	1	1	1	1	6	3	4	-	1	-	-
White, both sexes	-	1	1	-	2	5	5	3	-	-	-	-
Male	-	1	-	-	1	-	2	-	-	-	-	-
Female	-	-	1	-	1	5	3	3	-	-	-	-
Black, both sexes	-	1	1	1	2	1	1	2	-	1	-	-
Male	-	-	1	-	2	-	1	1	-	-	-	-
Female	-	1	-	1	-	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Maine												
All races, both sexes	1	1	-	-	1	2	1	2	-	-	-	-
Male	1	1	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	1	1	-	-	1	2	1	2	-	-	-	-
Male	1	1	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	4	2	2	3	8	3	3	-	1	-	-
Male	1	2	2	-	1	4	1	1	-	-	-	-
Female	3	2	-	2	2	4	2	2	-	1	-	-
White, both sexes	3	3	1	2	2	8	2	1	-	-	-	-
Male	1	1	1	-	1	4	-	1	-	-	-	-
Female	2	2	-	2	1	4	2	-	-	-	-	-
Black, both sexes	1	1	1	-	1	-	1	2	-	1	-	-
Male	-	1	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	1	-	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Massachusetts												
All races, both sexes	5	3	5	6	1	4	11	3	4	4	-	-
Male	1	2	3	1	1	1	4	1	-	1	-	-
Female	4	1	2	5	-	3	7	2	4	3	-	-
White, both sexes	5	3	3	6	1	4	11	2	4	4	-	-
Male	1	2	1	1	1	1	4	-	-	1	-	-
Female	4	1	2	5	-	3	7	2	4	3	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	11	18	7	6	4	14	12	10	11	3	-	-
Male	9	13	5	4	3	5	7	4	5	-	-	-
Female	2	5	2	2	1	9	5	6	6	3	-	-
White, both sexes	9	15	3	5	3	12	9	10	10	3	-	-
Male	9	11	3	3	3	4	5	4	5	-	-	-
Female	-	4	-	2	-	8	4	6	5	3	-	-
Black, both sexes	2	3	4	1	1	1	3	-	1	-	-	-
Male	-	2	2	1	-	-	2	-	-	-	-	-
Female	2	1	2	-	1	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Minnesota												
All races, both sexes	-	1	-	2	-	6	2	1	2	3	1	-
Male	-	1	-	-	-	4	-	-	-	1	-	-
Female	-	-	-	2	-	2	2	1	2	2	1	-
White, both sexes	-	1	-	2	-	6	2	1	2	3	-	-
Male	-	1	-	-	-	4	-	-	-	1	-	-
Female	-	-	-	2	-	2	2	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	2	1	3	4	3	3	2	2	1	1	-
Male	-	1	-	1	1	-	-	1	-	-	-	-
Female	2	1	1	2	3	3	3	1	2	1	1	-
White, both sexes	1	1	1	1	2	3	2	1	2	1	1	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	1	1	1	3	2	1	2	1	1	-
Black, both sexes	1	1	-	2	2	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	1	-	1	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Missouri												
All races, both sexes	4	1	5	-	6	3	10	2	4	2	-	-
Male	2	-	1	-	1	1	5	2	1	-	-	-
Female	2	1	4	-	5	2	5	-	3	2	-	-
White, both sexes	3	1	5	-	6	3	7	2	4	2	-	-
Male	1	-	1	-	1	1	3	2	1	-	-	-
Female	2	1	4	-	5	2	4	-	3	2	-	-
Black, both sexes	1	-	-	-	-	-	3	-	-	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	1	1	-	-	-	1	2	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	1	1	-	-	-	-	2	-	-
White, both sexes	2	-	-	1	1	-	-	-	1	2	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	1	1	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Nebraska												
All races, both sexes	2	1	1	2	1	2	4	2	1	1	-	-
Male	2	1	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	2	-	1	3	2	1	1	-	-
White, both sexes	1	1	1	2	1	2	4	2	1	1	-	-
Male	1	1	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	2	-	1	3	2	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	2	2	1	4	-	3	3	-	-	-	-
Male	1	2	1	1	3	-	2	1	-	-	-	-
Female	1	-	1	-	1	-	1	2	-	-	-	-
White, both sexes	2	2	2	1	4	-	3	3	-	-	-	-
Male	1	2	1	1	3	-	2	1	-	-	-	-
Female	1	-	1	-	1	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Hampshire												
All races, both sexes	2	1	1	-	1	3	1	2	1	1	-	-
Male	2	1	1	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	1	1	1	-	-
White, both sexes	2	1	1	-	1	3	1	2	1	1	-	-
Male	2	1	1	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	10	8	4	3	7	11	15	14	7	1	1	-
Male	8	5	1	1	1	5	2	5	2	-	1	-
Female	2	3	3	2	6	6	13	9	5	1	-	-
White, both sexes	5	8	2	3	5	10	13	14	6	1	1	-
Male	4	5	1	1	1	4	2	5	2	-	1	-
Female	1	3	1	2	4	6	11	9	4	1	-	-
Black, both sexes	5	-	2	-	2	1	2	-	1	-	-	-
Male	4	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	2	-	2	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Mexico												
All races, both sexes	1	3	2	-	-	1	2	4	3	-	-	-
Male	-	2	1	-	-	-	-	2	-	-	-	-
Female	1	1	1	-	-	1	2	2	3	-	-	-
White, both sexes	1	3	2	-	-	1	1	3	2	-	-	-
Male	-	2	1	-	-	-	-	1	-	-	-	-
Female	1	1	1	-	-	1	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	22	15	13	17	17	21	17	23	14	2	3	-
Male	17	10	8	6	9	11	5	4	3	2	-	-
Female	5	5	5	11	8	10	12	19	11	-	3	-
White, both sexes	13	12	9	7	13	14	12	21	14	2	3	-
Male	12	10	6	2	7	10	2	3	3	2	-	-
Female	1	2	3	5	6	4	10	18	11	-	3	-
Black, both sexes	9	3	4	10	4	7	5	2	-	-	-	-
Male	5	-	2	4	2	1	3	1	-	-	-	-
Female	4	3	2	6	2	6	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
North Carolina												
All races, both sexes	9	4	8	4	8	6	7	10	2	4	-	-
Male	6	1	3	3	3	1	3	2	-	1	-	-
Female	3	3	5	1	5	5	4	8	2	3	-	-
White, both sexes	2	3	4	3	5	4	5	8	2	3	-	-
Male	1	1	2	2	2	1	2	2	-	1	-	-
Female	1	2	2	1	3	3	3	6	2	2	-	-
Black, both sexes	7	1	3	1	2	2	2	2	-	1	-	-
Male	5	-	1	1	1	-	1	-	-	-	-	-
Female	2	1	2	-	1	2	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Ohio												
All races, both sexes	8	6	8	10	12	24	15	24	7	2	1	-
Male	6	2	4	5	6	11	5	7	-	-	-	-
Female	2	4	4	5	6	13	10	17	7	2	1	-
White, both sexes	6	5	5	8	10	22	13	19	7	1	-	-
Male	5	2	2	4	6	10	3	6	-	-	-	-
Female	1	3	3	4	4	12	10	13	7	1	-	-
Black, both sexes	2	1	3	2	2	2	2	5	-	1	1	-
Male	1	-	2	1	-	1	2	1	-	-	-	-
Female	1	1	1	1	2	1	-	4	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	1	5	4	1	4	3	3	3	1	-	-
Male	1	1	2	1	-	2	1	-	1	1	-	-
Female	1	-	3	3	1	2	2	3	2	-	-	-
White, both sexes	2	1	4	3	1	4	3	2	2	-	-	-
Male	1	1	2	-	-	2	1	-	1	-	-	-
Female	1	-	2	3	1	2	2	2	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Oregon												
All races, both sexes	1	-	5	2	5	4	-	5	3	1	-	-
Male	1	-	2	1	2	2	-	2	1	-	-	-
Female	-	-	3	1	3	2	-	3	2	1	-	-
White, both sexes	1	-	5	2	5	4	-	5	3	1	-	-
Male	1	-	2	1	2	2	-	2	1	-	-	-
Female	-	-	3	1	3	2	-	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	7	2	6	5	12	20	22	11	4	-	-
Male	3	4	2	4	2	3	7	5	3	2	-	-
Female	1	3	-	2	3	9	13	17	8	2	-	-
White, both sexes	3	7	2	6	5	11	18	20	10	4	-	-
Male	2	4	2	4	2	3	5	5	3	2	-	-
Female	1	3	-	2	3	8	13	15	7	2	-	-
Black, both sexes	1	-	-	-	-	1	2	2	1	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Rhode Island												
All races, both sexes	1	-	-	-	1	1	1	4	-	1	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	3	-	1	-	-
White, both sexes	-	-	-	-	1	1	1	4	-	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	2	3	2	1	4	2	8	7	2	-	-
Male	-	-	3	2	-	3	-	1	-	-	-	-
Female	-	2	-	-	1	1	2	7	7	2	-	-
White, both sexes	-	-	3	2	-	2	-	7	5	2	-	-
Male	-	-	3	2	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	6	5	2	-	-
Black, both sexes	-	2	-	-	1	2	2	1	2	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	2	-	-	1	-	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
South Dakota												
All races, both sexes	-	-	1	-	1	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	1	-	1	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	3	7	2	6	5	11	7	6	3	2	-
Male	-	3	3	-	2	3	3	-	2	-	1	-
Female	5	-	4	2	4	2	8	7	4	3	1	-
White, both sexes	4	3	4	2	6	5	10	5	6	2	2	-
Male	-	3	-	-	2	3	2	-	2	-	1	-
Female	4	-	4	2	4	2	8	5	4	2	1	-
Black, both sexes	1	-	3	-	-	-	1	2	-	1	-	-
Male	-	-	3	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Texas												
All races, both sexes	11	10	7	9	5	13	22	16	11	10	-	-
Male	10	7	5	4	2	6	9	5	3	1	-	-
Female	1	3	2	5	3	7	13	11	8	9	-	-
White, both sexes	6	10	6	8	4	13	18	14	9	10	-	-
Male	5	7	4	3	1	6	9	4	2	1	-	-
Female	1	3	2	5	3	7	9	10	7	9	-	-
Black, both sexes	5	-	1	1	1	-	4	2	2	-	-	-
Male	5	-	1	1	1	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	2	1	-	1	4	-	-	1	-	-
Male	1	-	2	-	-	1	4	-	-	1	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	2	1	-	1	4	-	-	1	-	-
Male	1	-	2	-	-	1	4	-	-	1	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Vermont												
All races, both sexes	-	-	1	2	1	1	1	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	2	1	1	1	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	5	4	3	5	9	11	9	5	3	-	-
Male	1	3	3	1	2	-	3	4	-	-	-	-
Female	1	2	1	2	3	9	8	5	5	3	-	-
White, both sexes	2	3	3	1	3	8	6	7	3	3	-	-
Male	1	2	2	-	1	-	1	3	-	-	-	-
Female	1	1	1	1	2	8	5	4	3	3	-	-
Black, both sexes	-	2	1	2	2	1	5	2	2	-	-	-
Male	-	1	1	1	1	-	2	1	-	-	-	-
Female	-	1	-	1	1	1	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Washington												
All races, both sexes	7	7	4	4	-	2	8	7	8	2	1	-
Male	3	6	1	3	-	-	1	1	4	1	-	-
Female	4	1	3	1	-	2	7	6	4	1	1	-
White, both sexes	7	7	4	3	-	2	8	7	8	2	1	-
Male	3	6	1	3	-	-	1	1	4	1	-	-
Female	4	1	3	-	-	2	7	6	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	4	2	3	4	3	1	-	-	-	-
Male	-	1	1	1	1	3	1	-	-	-	-	-
Female	-	-	3	1	2	1	2	1	-	-	-	-
White, both sexes	-	-	4	2	3	4	3	1	-	-	-	-
Male	-	-	1	1	1	3	1	-	-	-	-	-
Female	-	-	3	1	2	1	2	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Wisconsin												
All races, both sexes	-	2	3	5	3	3	4	4	2	-	-	-
Male	-	1	2	3	2	3	-	1	1	-	-	-
Female	-	1	1	2	1	-	4	3	1	-	-	-
White, both sexes	-	2	3	4	3	3	3	4	2	-	-	-
Male	-	1	2	3	2	3	-	1	1	-	-	-
Female	-	1	1	1	1	-	3	3	1	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
United States												
All races, both sexes	1,146	29	2	2	-	3	-	2	6	9	26	54
Male	547	16	1	2	-	-	-	1	3	6	19	38
Female	599	13	1	-	-	3	-	1	3	3	7	16
White, both sexes	968	24	2	2	-	3	-	1	4	7	22	39
Male	460	12	1	2	-	-	-	-	2	6	16	29
Female	508	12	1	-	-	3	-	1	2	1	6	10
Black, both sexes	149	3	-	-	-	-	-	1	1	2	1	12
Male	71	2	-	-	-	-	-	1	-	-	1	7
Female	78	1	-	-	-	-	-	-	1	2	-	5
American Indian 1/, both sexes	7	1	-	-	-	-	-	-	1	-	2	2
Male	4	1	-	-	-	-	-	-	1	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	22	1	-	-	-	-	-	-	-	-	1	1
Male	12	1	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	26	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	18	-	-	-	-	-	-	-	1	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	17	-	-	-	-	-	-	-	1	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Arkansas												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	95	3	-	-	-	-	-	-	1	-	8	5
Male	45	1	-	-	-	-	-	-	1	-	6	4
Female	50	2	-	-	-	-	-	-	-	-	2	1
White, both sexes	88	3	-	-	-	-	-	-	1	-	7	4
Male	42	1	-	-	-	-	-	-	1	-	5	4
Female	46	2	-	-	-	-	-	-	-	-	2	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Colorado												
All races, both sexes	17	2	-	-	-	-	-	-	-	-	1	1
Male	10	2	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	2	-	-	-	-	-	-	-	-	1	1
Male	9	2	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	3
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Delaware												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Florida												
All races, both sexes	52	2	-	-	-	-	-	-	-	-	-	3
Male	25	1	-	-	-	-	-	-	-	-	-	2
Female	27	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	48	2	-	-	-	-	-	-	-	-	-	3
Male	23	1	-	-	-	-	-	-	-	-	-	2
Female	25	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	39	-	-	-	-	1	-	-	-	1	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	30	-	-	-	-	1	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Hawaii												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	8	1	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	1	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Illinois												
All races, both sexes	57	3	-	-	-	-	-	-	-	1	2	1
Male	22	3	-	-	-	-	-	-	-	1	2	-
Female	35	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	47	1	-	-	-	-	-	-	-	1	2	1
Male	15	1	-	-	-	-	-	-	-	1	2	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	25	2	-	-	-	-	-	-	-	-	1	-
Male	13	2	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	2	-	-	-	-	-	-	-	-	-	-
Male	12	2	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Iowa												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Kentucky												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	53	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	22	-	-	-	-	-	-	1	-	-	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	3
Female	10	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Maine												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	15	-	-	-	-	1	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Massachusetts												
All races, both sexes	34	1	-	-	-	-	-	-	1	-	-	1
Male	16	1	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	1	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	36	-	-	-	-	-	-	1	-	-	-	3
Male	16	-	-	-	-	-	-	1	-	-	-	3
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	3
Male	13	-	-	-	-	-	-	-	-	-	-	3
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Minnesota												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	18	-	-	1	-	-	-	-	-	-	1	-
Male	10	-	-	1	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	1	-	-	-	-	-	-	1	-
Male	6	-	-	1	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Missouri												
All races, both sexes	38	2	1	-	-	-	-	-	-	1	-	2
Male	20	1	-	-	-	-	-	-	-	1	-	1
Female	18	1	1	-	-	-	-	-	-	-	-	1
White, both sexes	35	2	1	-	-	-	-	-	-	1	-	2
Male	19	1	-	-	-	-	-	-	-	1	-	1
Female	16	1	1	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Nebraska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
New Hampshire												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
New Mexico												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	57	2	-	1	-	-	-	-	1	-	1	1
Male	25	1	-	1	-	-	-	-	-	-	1	1
Female	32	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	45	2	-	1	-	-	-	-	-	-	1	-
Male	19	1	-	1	-	-	-	-	-	-	1	-
Female	26	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
North Carolina												
All races, both sexes	37	1	-	-	-	-	-	-	-	-	1	4
Male	17	1	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	15	1	-	-	-	-	-	-	-	-	-	4
Male	7	1	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Ohio												
All races, both sexes	46	1	-	-	-	-	-	-	-	2	-	4
Male	25	-	-	-	-	-	-	-	-	1	-	3
Female	21	1	-	-	-	-	-	-	-	1	-	1
White, both sexes	43	1	-	-	-	-	-	-	-	2	-	4
Male	25	-	-	-	-	-	-	-	-	1	-	3
Female	18	1	-	-	-	-	-	-	-	1	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	15	2	-	-	-	-	-	-	-	-	1	-
Male	8	1	-	-	-	-	-	-	-	-	1	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	2	-	-	-	-	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Oregon												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	55	-	1	-	-	1	-	-	1	1	1	4
Male	28	-	1	-	-	-	-	-	1	1	1	3
Female	27	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	45	-	1	-	-	1	-	-	1	1	1	1
Male	23	-	1	-	-	-	-	-	1	1	1	1
Female	22	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	3
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Rhode Island												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
South Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Texas												
All races, both sexes	72	2	-	-	-	-	-	-	-	1	2	4
Male	32	-	-	-	-	-	-	-	-	1	1	1
Female	40	2	-	-	-	-	-	-	-	-	1	3
White, both sexes	58	2	-	-	-	-	-	-	-	1	2	4
Male	27	-	-	-	-	-	-	-	-	1	1	1
Female	31	2	-	-	-	-	-	-	-	-	1	3
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	27	2	-	-	-	-	-	-	-	-	-	2
Male	17	1	-	-	-	-	-	-	-	-	-	2
Female	10	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	1	-	-	-	-	-	-	-	-	-	2
Male	12	1	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Washington												
All races, both sexes	19	1	-	-	-	-	-	-	-	-	1	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	1	-	-	-	-	-	-	-	-	1	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Wisconsin												
All races, both sexes	28	-	-	-	-	-	-	-	-	1	-	3
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	3
Male	16	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
United States												
All races, both sexes	73	81	72	73	95	127	150	157	121	51	13	-
Male	51	45	49	40	46	68	56	54	36	13	3	-
Female	22	36	23	33	49	59	94	103	85	38	10	-
White, both sexes	61	61	54	64	80	110	133	141	108	42	10	-
Male	42	33	38	36	40	59	50	48	35	9	2	-
Female	19	28	16	28	40	51	83	93	73	33	8	-
Black, both sexes	10	17	16	6	15	15	14	13	12	8	3	-
Male	9	11	9	3	6	8	3	5	1	4	1	-
Female	1	6	7	3	9	7	11	8	11	4	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	2	2	-	2	3	3	1	1	-	-
Male	-	1	2	1	-	1	3	1	-	-	-	-
Female	2	2	-	1	-	1	-	2	1	1	-	-
Alabama												
All races, both sexes	3	1	-	1	3	3	6	4	2	1	-	-
Male	2	1	-	-	1	1	2	-	-	-	-	-
Female	1	-	-	1	2	2	4	4	2	1	-	-
White, both sexes	2	-	-	-	2	3	6	3	2	1	-	-
Male	1	-	-	-	-	1	2	-	-	-	-	-
Female	1	-	-	-	2	2	4	3	2	1	-	-
Black, both sexes	1	1	-	1	1	-	-	1	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	1	-	1	1	1	1	3	4	-	-	-
Male	1	1	-	1	-	-	1	1	2	-	-	-
Female	1	-	-	-	1	1	-	2	2	-	-	-
White, both sexes	2	1	-	1	1	1	1	3	4	-	-	-
Male	1	1	-	1	-	-	1	1	2	-	-	-
Female	1	-	-	-	1	1	-	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Arkansas												
All races, both sexes	2	1	2	3	1	2	2	1	1	-	1	-
Male	-	-	1	1	1	2	2	-	1	-	-	-
Female	2	1	1	2	-	-	-	1	-	-	1	-
White, both sexes	2	1	2	3	1	2	2	1	1	-	1	-
Male	-	-	1	1	1	2	2	-	1	-	-	-
Female	2	1	1	2	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	9	2	6	12	8	14	10	9	1	1	-
Male	3	1	-	3	8	6	5	3	4	-	-	-
Female	3	8	2	3	4	2	9	7	5	1	1	-
White, both sexes	6	8	1	4	11	8	14	10	9	1	1	-
Male	3	1	-	2	7	6	5	3	4	-	-	-
Female	3	7	1	2	4	2	9	7	5	1	1	-
Black, both sexes	-	-	1	2	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Colorado												
All races, both sexes	2	-	1	-	-	2	2	5	-	1	-	-
Male	2	-	1	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	2	-	4	-	1	-	-
White, both sexes	2	-	1	-	-	2	1	5	-	1	-	-
Male	2	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	2	-	4	-	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	-	-	2	1	1	-	4	1	-	-	-
Male	2	-	-	1	-	1	-	1	1	-	-	-
Female	-	-	-	1	1	-	-	3	-	-	-	-
White, both sexes	1	-	-	2	1	1	-	4	1	-	-	-
Male	1	-	-	1	-	1	-	1	1	-	-	-
Female	-	-	-	1	1	-	-	3	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Florida												
All races, both sexes	4	6	3	4	4	6	7	8	2	3	-	-
Male	4	3	1	2	2	4	2	4	-	-	-	-
Female	-	3	2	2	2	2	5	4	2	3	-	-
White, both sexes	4	5	3	4	3	5	6	8	2	3	-	-
Male	4	2	1	2	2	3	2	4	-	-	-	-
Female	-	3	2	2	1	2	4	4	2	3	-	-
Black, both sexes	-	1	-	-	1	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	5	5	2	3	2	3	10	2	2	1	-
Male	1	4	4	1	1	-	-	5	-	1	-	-
Female	-	1	1	1	2	2	3	5	2	1	1	-
White, both sexes	1	4	5	2	2	2	3	7	-	1	1	-
Male	1	3	4	1	-	-	-	4	-	1	-	-
Female	-	1	1	1	2	2	3	3	-	-	1	-
Black, both sexes	-	1	-	-	1	-	-	3	1	1	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	2	2	-	1	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	2	1	-	1	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	-	1	-	1	-	-	-	-	4	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	1	-	-	-	-	3	-	-	-
White, both sexes	-	1	-	1	-	-	-	-	4	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	1	-	-	-	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Illinois												
All races, both sexes	1	3	1	4	5	4	12	7	8	4	1	-
Male	-	1	1	1	2	1	5	2	2	1	-	-
Female	1	2	-	3	3	3	7	5	6	3	1	-
White, both sexes	1	2	1	2	4	3	11	6	8	3	1	-
Male	-	-	1	-	2	1	4	1	2	-	-	-
Female	1	2	-	2	2	2	7	5	6	3	1	-
Black, both sexes	-	-	-	2	1	1	1	1	-	1	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	2	3	2	1	3	3	3	3	-	-	-
Male	1	1	2	1	1	1	1	2	-	-	-	-
Female	1	1	1	1	-	2	2	1	3	-	-	-
White, both sexes	1	1	3	2	1	2	3	3	3	-	-	-
Male	1	1	2	1	1	1	1	2	-	-	-	-
Female	-	-	1	1	-	1	2	1	3	-	-	-
Black, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Iowa												
All races, both sexes	1	-	-	-	1	2	1	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-
White, both sexes	1	-	-	-	1	2	1	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	1	-	2	-	1	3	1	1	-	-
Male	1	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	2	-	1	3	-	1	-	-
White, both sexes	1	1	1	-	2	-	1	3	1	1	-	-
Male	1	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	2	-	1	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Kentucky												
All races, both sexes	-	4	3	4	4	5	7	14	9	4	-	-
Male	-	2	2	3	3	2	2	6	-	-	-	-
Female	-	2	1	1	1	3	5	8	9	4	-	-
White, both sexes	-	4	3	3	4	5	7	14	8	4	-	-
Male	-	2	2	2	3	2	2	6	-	-	-	-
Female	-	2	1	1	1	3	5	8	8	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	3	2	3	1	2	2	2	1	-	-
Male	1	-	3	2	-	-	1	1	-	1	-	-
Female	-	1	-	-	3	1	1	1	2	-	-	-
White, both sexes	1	-	3	2	1	-	2	2	1	-	-	-
Male	1	-	3	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
Black, both sexes	-	1	-	-	2	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	2	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Maine												
All races, both sexes	-	-	-	-	2	-	1	3	3	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	2	3	-	-	-
White, both sexes	-	-	-	-	2	-	1	3	3	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	3	1	-	1	2	1	4	-	-	-
Male	1	1	3	1	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	2	-	3	-	-	-
White, both sexes	1	1	1	1	-	-	2	-	-	-	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	2	-	-	1	-	1	4	-	-	-
Male	-	-	2	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Massachusetts												
All races, both sexes	1	2	4	3	3	3	4	7	3	1	-	-
Male	1	1	3	2	-	2	1	3	1	-	-	-
Female	-	1	1	1	3	1	3	4	2	1	-	-
White, both sexes	1	1	3	3	2	3	4	7	3	1	-	-
Male	1	-	2	2	-	2	1	3	1	-	-	-
Female	-	1	1	1	2	1	3	4	2	1	-	-
Black, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	3	3	3	3	4	6	1	4	1	1	-
Male	3	2	-	1	1	3	1	-	1	-	-	-
Female	-	1	3	2	2	1	5	1	3	1	1	-
White, both sexes	3	1	2	3	3	3	5	1	4	1	1	-
Male	3	-	-	1	1	3	1	-	1	-	-	-
Female	-	1	2	2	2	-	4	1	3	1	1	-
Black, both sexes	-	2	1	-	-	-	1	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Minnesota												
All races, both sexes	2	-	1	-	-	5	3	4	1	-	1	-
Male	1	-	1	-	-	2	2	-	-	-	1	-
Female	1	-	-	-	-	3	1	4	1	-	-	-
White, both sexes	2	-	1	-	-	5	3	4	1	-	1	-
Male	1	-	1	-	-	2	2	-	-	-	1	-
Female	1	-	-	-	-	3	1	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	2	2	1	-	3	3	1	-	2	-	-
Male	2	1	2	-	-	2	-	-	-	1	-	-
Female	-	1	-	1	-	1	3	1	-	1	-	-
White, both sexes	1	1	-	1	-	3	2	-	-	1	-	-
Male	1	-	-	-	-	2	-	-	-	1	-	-
Female	-	1	-	1	-	1	2	-	-	-	-	-
Black, both sexes	1	1	2	-	-	-	1	1	-	1	-	-
Male	1	1	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Missouri												
All races, both sexes	3	-	1	5	-	5	5	6	6	1	-	-
Male	2	-	-	5	-	2	4	1	3	-	-	-
Female	1	-	1	-	-	3	1	5	3	1	-	-
White, both sexes	2	-	-	5	-	5	4	6	6	1	-	-
Male	1	-	-	5	-	2	4	1	3	-	-	-
Female	1	-	-	-	-	3	-	5	3	1	-	-
Black, both sexes	1	-	1	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	-	-	-	-	1	1	2	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
White, both sexes	-	1	-	-	-	-	1	1	2	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Nebraska												
All races, both sexes	-	-	-	-	-	2	4	1	4	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	2	3	1	3	-	-	-
White, both sexes	-	-	-	-	-	2	4	1	4	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	2	3	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	1	-	2	-	-	1	2	1	-	-	-
Male	2	1	-	1	-	-	1	1	-	-	-	-
Female	1	-	-	1	-	-	-	1	1	-	-	-
White, both sexes	1	-	-	2	-	-	1	2	1	-	-	-
Male	1	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
New Hampshire												
All races, both sexes	-	-	1	-	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	5	1	4	4	4	5	2	1	1	-
Male	1	1	4	-	3	3	-	3	-	1	-	-
Female	-	1	1	1	1	1	4	2	2	-	1	-
White, both sexes	1	2	3	1	4	3	2	4	2	1	1	-
Male	1	1	3	-	3	2	-	3	-	1	-	-
Female	-	1	-	1	1	1	2	1	2	-	1	-
Black, both sexes	-	-	2	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
New Mexico												
All races, both sexes	2	-	1	-	-	-	1	-	-	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	1	-	-	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	7	4	3	4	7	9	5	4	2	2	-
Male	3	5	4	1	4	3	1	-	-	-	-	-
Female	1	2	-	2	-	4	8	5	4	2	2	-
White, both sexes	3	6	2	2	3	7	8	4	3	2	1	-
Male	2	5	2	-	3	3	1	-	-	-	-	-
Female	1	1	-	2	-	4	7	4	3	2	1	-
Black, both sexes	1	1	1	1	1	-	1	1	1	-	1	-
Male	1	-	1	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
North Carolina												
All races, both sexes	2	4	-	3	8	5	2	3	3	1	-	-
Male	2	2	-	1	2	3	1	1	2	-	-	-
Female	-	2	-	2	6	2	1	2	1	1	-	-
White, both sexes	1	1	-	3	7	2	1	2	3	1	-	-
Male	1	1	-	1	2	1	1	1	2	-	-	-
Female	-	-	-	2	5	1	-	1	1	1	-	-
Black, both sexes	1	3	-	-	1	3	1	1	-	-	-	-
Male	1	1	-	-	-	2	-	-	-	-	-	-
Female	-	2	-	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Ohio												
All races, both sexes	2	1	6	3	4	6	4	4	5	4	-	-
Male	2	1	4	2	2	2	2	1	2	3	-	-
Female	-	-	2	1	2	4	2	3	3	1	-	-
White, both sexes	2	1	5	3	3	6	4	4	5	3	-	-
Male	2	1	4	2	2	2	2	1	2	3	-	-
Female	-	-	1	1	1	4	2	3	3	-	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	1	-	2	1	1	2	1	1	1	-	-
Male	1	1	-	2	-	-	1	-	1	-	-	-
Female	1	-	-	-	1	1	1	1	-	1	-	-
White, both sexes	2	1	-	2	1	1	2	1	1	1	-	-
Male	1	1	-	2	-	-	1	-	1	-	-	-
Female	1	-	-	-	1	1	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Oregon												
All races, both sexes	-	-	-	-	1	5	1	1	2	1	-	-
Male	-	-	-	-	-	4	-	1	2	1	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	5	1	1	2	1	-	-
Male	-	-	-	-	-	4	-	1	2	1	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	6	3	1	5	5	7	6	4	5	1	-
Male	2	5	1	-	1	2	4	4	1	-	1	-
Female	1	1	2	1	4	3	3	2	3	5	-	-
White, both sexes	3	4	3	1	2	4	7	6	3	5	1	-
Male	2	4	1	-	-	1	4	4	1	-	1	-
Female	1	-	2	1	2	3	3	2	2	5	-	-
Black, both sexes	-	1	-	-	3	1	-	-	1	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	3	2	1	1	1	2	2	-	2	-	-
Male	1	1	2	1	1	-	-	-	-	2	-	-
Female	-	2	-	-	-	1	2	2	-	-	-	-
White, both sexes	-	3	1	1	1	1	1	2	-	-	-	-
Male	-	1	1	1	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	1	1	2	-	-	-	-
Black, both sexes	1	-	1	-	-	-	1	-	-	2	-	-
Male	1	-	1	-	-	-	-	-	-	2	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
South Dakota												
All races, both sexes	-	-	-	-	-	1	-	1	2	1	1	-
Male	-	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	-	-	-	-	-	1	-	1	2	1	1	-
Male	-	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	4	1	1	-	6	3	3	2	1	-	-
Male	1	2	1	1	-	3	3	1	-	-	-	-
Female	1	2	-	-	-	3	-	2	2	1	-	-
White, both sexes	2	4	1	1	-	6	3	3	1	1	-	-
Male	1	2	1	1	-	3	3	1	-	-	-	-
Female	1	2	-	-	-	3	-	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Texas												
All races, both sexes	5	2	5	5	6	11	8	9	9	2	1	-
Male	3	1	3	2	4	6	3	2	5	-	-	-
Female	2	1	2	3	2	5	5	7	4	2	1	-
White, both sexes	5	1	2	4	6	8	7	6	9	1	-	-
Male	3	-	2	2	4	5	2	1	5	-	-	-
Female	2	1	-	2	2	3	5	5	4	1	-	-
Black, both sexes	-	1	3	-	-	3	1	2	-	1	1	-
Male	-	1	1	-	-	1	1	1	-	-	-	-
Female	-	-	2	-	-	2	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	2	2	-	2	4	5	4	2	-	1	-
Male	1	1	2	-	2	3	2	2	-	-	1	-
Female	-	1	-	-	-	1	3	2	2	-	-	-
White, both sexes	1	1	1	-	1	2	3	3	1	-	-	-
Male	1	1	1	-	1	2	2	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
Black, both sexes	-	1	-	-	1	2	2	1	1	-	1	-
Male	-	-	-	-	1	1	-	1	-	-	1	-
Female	-	1	-	-	-	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Washington												
All races, both sexes	2	1	3	2	3	-	3	1	2	-	-	-
Male	1	1	2	1	1	-	1	-	1	-	-	-
Female	1	-	1	1	2	-	2	1	1	-	-	-
White, both sexes	2	1	3	1	3	-	2	1	2	-	-	-
Male	1	1	2	1	1	-	-	-	1	-	-	-
Female	1	-	1	-	2	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	2	1	-	2	1	3	-	2	2	-	-
Male	-	1	1	-	2	1	1	-	-	-	-	-
Female	1	1	-	-	-	-	2	-	2	2	-	-
White, both sexes	1	2	1	-	2	1	3	-	2	2	-	-
Male	-	1	1	-	2	1	1	-	-	-	-	-
Female	1	1	-	-	-	-	2	-	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Wisconsin												
All races, both sexes	-	1	-	2	4	5	2	4	3	3	-	-
Male	-	1	-	2	2	4	-	4	1	1	-	-
Female	-	-	-	-	2	1	2	-	2	2	-	-
White, both sexes	-	1	-	2	4	4	2	4	3	3	-	-
Male	-	1	-	2	2	3	-	4	1	1	-	-
Female	-	-	-	-	2	1	2	-	2	2	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-